

# Interoceptive awareness and yoga in patients with eating disorder: a preliminary descriptive analysis.

Marc Toutain, Anne-Laure Moscone, Pascale Leconte, Antoine Gauthier

#### ▶ To cite this version:

Marc Toutain, Anne-Laure Moscone, Pascale Leconte, Antoine Gauthier. Interoceptive awareness and yoga in patients with eating disorder: a preliminary descriptive analysis.. The Motion Thickness Workshop, Jun 2022, Caen, France. Unpublished, 2022, 10.13140/RG.2.2.19473.51042. hal-04355639

#### HAL Id: hal-04355639 https://hal.science/hal-04355639v1

Submitted on 10 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Interoceptive awareness and yoga in patients with eating disorder: a preliminary descriptive analysis.

Marc Toutain<sup>1</sup>, Anne-Laure Moscone<sup>23</sup>, Pascale Leconte<sup>1</sup>, Antoine Gauthier<sup>1</sup> 1. UMR-S 1075 Inserm/Unicaen COMETE – Mobilités : Vieillissement, Pathologies, Santé 2. EA 4532 Université Paris-Saclay CIAMS – Complexité, Innovation, Activités Motrices et Sportives 3. CHU Paul-Brousse AP-HP à Villejuif (94), Université Paris-Saclay


## **Introduction**

Eating Disorders (ED), such as anorexia, bulimia and binge eating, affect about 10% of young people (Treasure, 2020). They are characterized by a deviant relationship with food and body, resulting in pathological eating and weight control behaviors. Physiological and psycho-cognitive comorbidities are commonly associated with ED, including an interoceptive deficit (Demartini et al., 2021).

The interoception can be considered as the interpretation of internal and external stimuli producing an overall representation of the state of the body, linked to a cognitive and emotional context and including conscious and unconscious aspects (Figure 1, Berntson & Khalsa, 2021). The conscious aspect is the interoceptive awareness, it is the ability to identify, access, understand, and respond appropriately to the patterns of internal signals, such as digestive system, muscle tension, heart rate, etc. (Price & Hooven, 2018).


Interoceptive awareness (IA) can be improved with mind and body practices like meditation or yoga (Demartini et al., 2021). To our knowledge, the effect of yoga on IA has not been studied in ED patients.

The aim of this study is to assess the effect of yoga on interoceptive awareness and ED symptoms in a clinical sample. We hypothesize that yoga will improve interoceptive awareness and ED symptoms.



### Figure 1. The Continuum of Interoception.

"Interoception is a neural process that traverses sensors, pathways, networks, circuits, and awareness. Each component contributes to the next, although most processing of interoceptive signals occurs beyond the conscious awareness of the organism. Further, interoceptive awareness relies on circuit-based representations of the internal bodily self (e.g., 'my body': manikin at top right) and differs substantially from other circuit-based representations of other living objects (e.g., 'dog', 'plant' etc.)" (Berntson & Khalsa, 2021)



# Results


**MAIA + EAT-26** 


Currently, four participants have completed the protocol. The sample mean age was 21,5 ( $\pm$ 3,69) years, the mean body mass index was 19,47 (±1,51) kg/m² and three were suffering of anorexia and one of bulimia.


Analysis


A descriptive analysis showed that the total MAIA score improved after the 8 weeks of yoga, with a mean pre-yoga score of  $61/160~(\pm 29)$  and a mean post-yoga score of  $90/160~(\pm 18)$ . The subscale of emotional awareness seemed to be an important part of this improvement with a mean pre-yoga score of 1,75/5 ( $\pm$ 1,31) and a mean post-yoga score of 4/5 ( $\pm$ 0,36).


In addition, the EAT-26 total score showed a great diminution with a mean pre-yoga score of  $41,5/78 \ (\pm 16,3)$  and a mean post-yoga score of  $27/78 \ (\pm 18)$ . Two subscales seemed to explain this improvement: the dieting subscale with a mean pre-yoga score of  $27/78 \ (\pm 10)$  and a mean postyoga score of 18,75/39 (±10) and the bulimia and food preoccupation subscale with a mean preyoga score of  $9,75/18 \ (\pm 2,21)$  and a mean post-yoga score of  $4,75/78 \ (\pm 3,86)$ .











# References

Berntson, G. G., & Khalsa, S. S. (2021). Neural Circuits of Interoception. Trends in Neurosciences, 44(1), 17-28. https://doi.org/10.1016/j.tins.2020.09.011

Borden, A., & Cook-Cottone, C. (2020). Yoga and eating disorder prevention and treatment: A comprehensive review and meta-analysis. Eating Disorders, 28(4), 400-437. https://doi.org/10.1080/10640266.2020.1798172

Brown, T. A., Berner, L. A., Jones, M. D., Reilly, E. E., Cusack, A., Anderson, L. K., Kaye, W. H., & Wierenga, C. E. (2017). Psychometric Evaluation and Norms for the Multidimensional Assessment of Interoceptive Awareness (MAIA) in a Clinical Eating Disorders Sample. European Eating Disorders Review: The Journal of the Eating Disorders Association, 25(5), 411-416. https://doi.org/10.1002/erv.2532

Demartini, B., Goeta, D., Marchetti, M., Bertelli, S., Anselmetti, S., Cocchi, A., Ischia, M., & Gambini, O. (2021). The effect of a single yoga class on interoceptive accuracy in patients affected by anorexia nervosa and in healthy controls: A pilot study. Eating and Weight Disorders, 26(5), 1427-1435. https://doi.org/10.1007/s40519-020-00950-3

Price, C. J., & Hooven, C. (2018). Interoceptive Awareness Skills for Emotion Regulation: Theory and Approach of Mindful Awareness in Body-Oriented Therapy (MABT). Frontiers in Psychology, 9, 798.

https://doi.org/10.3389/fpsyg.2018.00798

Treasure, J. (2020). Eating disorders. *Medicine*, 48(11), 727-731. <a href="https://doi.org/10.1016/j.mpmed.2020.08.001">https://doi.org/10.1016/j.mpmed.2020.08.001</a>

# Discussion and conclusion

The MAIA score increase should demonstrate a positive effect of yoga on IA. The strong improvement in the emotional awareness seems to indicate that yoga can specifically target this IA dimension. However our sample showed a lower mean (1,75/5) and a higher standard deviation ( $\pm 1,31$ ) compared to other adult ED sample in the literature with a mean of 2,96/5 ( $\pm 1,23$ ) for this subscale which could be a bias linked to the small size of the sample (Brown & al., 2017).

The EAT-26 decrease is in accordance with the literature and should reveal an improvement of ED symptoms after yoga therapeutic intervention (Borden & Cook-Cottone, 2020).

The sample size was small and additional data are needed for inferential analysis as well as a control group to assess the actual contribution of yoga to these results. Nevertheless this first descriptive analysis suggest a positive effect of yoga on IA for ED outpatients





PUBLIQUE '



