

# Widening exposome exploration by means of complementary HRMS analytical platforms: use case on pesticide exposure

Elodie Mirmont, Tarek Moufawad, Fatima-Zahra Alem, Cécile Chevrier, Nathalie Bonvallot, Emmanuelle Kesse-Guyot, Julia Baudry, Emmanuelle Bichon, Ingrid Guiffard, Jean-Philippe Antignac, et al.

## ▶ To cite this version:

Elodie Mirmont, Tarek Moufawad, Fatima-Zahra Alem, Cécile Chevrier, Nathalie Bonvallot, et al.. Widening exposome exploration by means of complementary HRMS analytical platforms: use case on pesticide exposure. 39ème édition des Journées Françaises de Spectrométrie de Masse, Sep 2023, Marseille, France. hal-04357806

HAL Id: hal-04357806

https://hal.science/hal-04357806

Submitted on 21 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Widening exposome exploration by means of complementary HRMS analytical platforms: use case on pesticide exposure

Elodie Mirmont<sup>1</sup>, Tarek Moufawad<sup>2</sup>, Fatima-Zahra Alem<sup>3</sup>, Cécile Chevrier<sup>3</sup>, Nathalie Bonvallot<sup>3</sup>, Emmanuelle Kesse-Guyot<sup>4</sup>, Julia Baudry<sup>4</sup>, Emmanuelle Bichon<sup>2</sup>, Ingrid Guiffard<sup>2</sup>, Jean-Philippe Antignac<sup>2</sup>, Arthur David<sup>3</sup>, Emilien Jamin<sup>1</sup>, Laurent Debrauwer<sup>1</sup>

#### Introduction

Human exposure to food and environmental contaminants (such as pesticides) is generally estimated by indirect methods.

Targeted bio-monitoring is a way to assess the internal exposure on a limited number of substances, needing large sample amounts for analyses as rigorous and sensitive as possible. Thus, the human health risks associated with multi-exposure to complex mixtures currently remain under-explored.

In this context, the SCREENPEST project aims at providing a unified methodology for large-scale characterisation of human internal exposure to pesticides, via a "suspect profiling" approach, using urine.



# **Objectives**

Develop and set-up an harmonised analytical workflow including sample preparation, instrumental measurement and data processing. Increase the coverage on human exposure to pesticides by a nontargeted multiplexed approach.

Test the method developed through urine samples from cohorts to study possible links between exposure and adverse health effects.

Provide extended information capabilities for early warning on emerging substances and help for decision making.

**PFLAGIE** 

# **Developed workflow**



Suspects list elaboration

Analytical methods optimisation (separation/ detection)

Method development

HRMS spectral library generation

Development of non discriminating extraction techniques Application on cohort urine samples

- → Mother-child cohort (France, 2002-2006)
- Association between the exposure to pesticides of pregnant women and biometric data at birth
- Access to 168 samples

**NutriNet-Santé** 



Large list of standards (ca. 200 pesticides and phase I/II metabolites), covering a wide range of physicochemical properties (logKow:[-4.71;8.38], MM/g.mol<sup>-1</sup>: [72;720]) for analytical developments.

Standard selection: commercial availability, bibliography, pesticide use in France (2019), epidemiologists expertise and available data from the HMB4EU project (10.1021/acs.analchem.2c00061)



Optimisation of 3 HRMS techniques to get the best separation and detection of the suspects by both LC and GC.

Characterisation of the suspects by HRMS




Optimisation of sample preparation: various concentration levels (20, 50 and 100 ng mL-1) spiked in pooled urine (5 replicates).

General population (France, since 2009) and the onset of type 2 diabetes


Association between the exposure to pesticides

→ Access to 200 samples

Performance comparison of two non discriminating extraction techniques (recovery and matrix effect)



## **Extraction and HRMS analysis**





GC - Orbitrap

Reversed Phase (C18) LC - QToF

HILIC - LTQ-Orbitrap XL

Analytical challenges to cover a wide range of compounds with:



- Different physicochemical properties :
  - Mol. weight [72;720]
  - Polarity / Log Kow [-4.71;8.38]
- Poorly characterised biotransformation products
- Low levels concentration

### **Data treatment - Bioinformatic tools**



Extended HRMS spectral library of reference pesticides and metabolites.



Complemented in house targeted and suspect screening chemo-informatics (XenoScreener / PeakForest / MetaboHUB W4M)



Improved annotation identification capacities on large range of pesticides



Statistical analysis and data interpretation (in progress)



















