

Interactome expertise: an integrative approach to evaluate solar radiation impact on skin ecosystem and a new SPF50+ sunscreen efficacy

Carine Jacques, D. Bacqueville, Emilien Jamin, Martine Maitre, Anais Noustens, Isabelle Jouanin, Aimée Simcic-Mori, Daniel Redoulès, Laurent Debrauwer, Sandrine Bessou-Touya, et al.

▶ To cite this version:

Carine Jacques, D. Bacqueville, Emilien Jamin, Martine Maitre, Anais Noustens, et al.. Interactome expertise: an integrative approach to evaluate solar radiation impact on skin ecosystem and a new SPF50+ sunscreen efficacy. 25th world congress of dermatology, Jul 2023, Singapore, Singapore. hal-04357867

HAL Id: hal-04357867

https://hal.science/hal-04357867

Submitted on 21 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Interactome expertise: an integrative approach to evaluate solar radiation impact on skin ecosystem and a new SPF50+ sunscreen efficacy



C. Jacques¹, D. Bacqueville¹, E. L. Jamin².³, M. Maitre¹, A. Noustens¹, I. Jouanin².³, A. Simcic-Mori¹, G. Doat⁴, L. Debrauwer².³, S. Bessou-Touya¹ & H. Duplan¹

¹ Pierre Fabre Dermo-cosmétique, Centre R&D Pierre Fabre, Toulouse, France

²³ Metatoul-AXIOM platform, MetaboHUB, Toxalim, UMR1331 Toxalim, INRAE, Toulouse, France

⁴ Laboratoires dermatologiques Avène, Lavaur, France

ID: 2584

INTRODUCTION

For a deeper understanding of sun exposure impact, we propose a new global and integrative vision, called Interactome expertise. We consider the skin as a complex ecosystem compsoed of the skin tissue, its skin surface lipids, its microbiota; and all these elements are linked and interdependent. Submitted to endogenous and exogenous factors, this ecosystem reacts to maintain skin equilibrium. To evaluate the impact of external factors like solar radiation, the cutaneous ecosystem has to be studied as a whole to understand how they interact with each other and contribute to reply to external stress. The consequences of solar radiation have been previously documented both at genetic and proteomic levels. However, only few studies describe solar radiation contribution to biochemical changes that result in skin metabolome alterations, and interactions between skin and its microbiota have not been described.

MATERIALS & METHODS

We developed an innovative *in vitro* model with RHE colonized with microbiota and sebum (N=18) to reproduce skin ecosystem complexity. This skin ecosystem model was exposed to simulated solar radiation (SSR) with and without application of a new SPF50+ sunscreen.

A multi-omics approach was used to evaluate SSR effect and SPF50+ sunscreen protection. Metabolomics profiles were performed with NMR and UHPLC-HRMS. Data were processed with XCMS using W4M and filtered using quality control samples. Identified discriminant metabolites according to in-house databases were mapped on metabolic pathways (Metexplore). Metagenomics analyses were performed from genomic DNA extract samples. Intergenic spacers 1 (ITSI) of the ribosomal RNA gene, or VI-V3 region of 16S gene was sequenced by high-speed sequencing. Sequences were compared to international data bases and differential analysis was used to identify differences of genus abundance.




Fig.1. In vitro protocol and analysis

RESULTATS




Fig.2. Discriminant and significant metabolites heatmap




Fig.3. Pathways significantly impacted after SSR exposure

Up to 50 metabolites were significantly modified (Fig.2) link to several pathways after solar radiation (Fig.3). The skin is under high oxidative stress with modification of glutathione, purine pathway and urea cycle. Other pathways have been highlighted, linked to skin microbiota, like BCAAs cycle, lipid metabolism and tryptophan pathway. With the new SPF50+ sunscreen is applied, skin ecosystem metabolome was preserved, and interactions (Host/Microbiota) were protected from negative effect of solar radiation.

From 16S analysis, the two main anaerobic bacteria of the genus *Burkholderia* and *Cutibacterium* became more abundant after SSR exposure but not when SPF50+ sunscreen is used (Fig.4).

From ITS analysis, *Malassezia* was the most abundant genus and became less abundant after SSR. These modification were significantly reduced when SPF50+ sunscreen is used (Fig.4).




Fig.4. Bacteria and fungi population diversity

CONCLUSION

This multi-omics approach identified a highly accurate metabolomic signature of the skin ecosystem after solar radiation and the interactions between the skin and cutaneous microbiota leading to oxidative stress/inflammation, immunosuppression and modification of microbial diversity.

The SPF50+ sunscreen, in addition to protecting against the common negative effects of sun exposure, also preserved host-microbial interactions and microbial diversity from the deleterious effects of solar radiation.