



**HAL**  
open science

# Three remarks on the convergence of some discretized second order gradient-like systems

Mohamed Ali Jendoubi, Morgan Pierre

► **To cite this version:**

Mohamed Ali Jendoubi, Morgan Pierre. Three remarks on the convergence of some discretized second order gradient-like systems. 2023. hal-04360196

**HAL Id: hal-04360196**

**<https://hal.science/hal-04360196v1>**

Preprint submitted on 21 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Three remarks on the convergence of some discretized second order gradient-like systems

Mohamed Ali JENDOUBI

*Université de Carthage,*

*Institut Préparatoire aux Etudes Scientifiques et Techniques,*

*B.P. 51 2070 La Marsa, Tunisia.*

and

*Laboratoire Équations aux Dérivées Partielles, LR03ES04,*

*Université Tunis El Manar,*

*Faculté des sciences de Tunis,*

*2092 El Manar, Tunisia.*

mohamedali.jendoubi@ipest.ucar.tn

Morgan PIERRE

*Laboratoire de Mathématiques et Applications,*

*Université de Poitiers, CNRS,*

*F-86073 Poitiers, France*

*The first author dedicates this paper to the memory of his colleague Naceur Manai.*

## **Abstract**

We study several discretizations of a second order gradient-like system with damping. We first consider an explicit scheme with a linear damping in finite dimension. We prove that every solution converges if the nonlinearity satisfies a global Lojasiewicz inequality. Convergence rates are also established. In the case of a strong nonlinear damping, we prove convergence of every solution for a fully implicit scheme in the one-dimensional case, even if the nonlinearity does not satisfy a Lojasiewicz inequality. The optimality of the damping is also established. Numerical simulations illustrate the theoretical results.

**Mathematics Subject Classification 2010 (MSC2010):** 65K05, 65L20, 90C26, 37N40, 26E05, 40A05.

**Key words:** descent methods, real analytic functions, Lojasiewicz gradient inequality, single limit-point convergence, stability, variable time-step discretization, explicit scheme.

## 1 Introduction

In this paper, we study the asymptotic behaviour of solutions to several discretizations of the second order ordinary differential system

$$u'' + \|u'\|^\alpha u' + \nabla F(u) = 0, \quad t \geq 0, \quad (1)$$

where  $F : \mathbb{R}^d \rightarrow \mathbb{R}$  is a  $C^1$  function and  $\alpha \in [0, 1]$ .

The asymptotic behaviour of solutions to (1) has been extensively studied. A crucial tool is the energy estimate

$$\frac{d}{dt} E(t) = -\|u'(t)\|^{2+\alpha} \leq 0,$$

where  $E$  is the energy defined for every solution  $u$  by

$$E(t) = \frac{1}{2} \|u'(t)\|^2 + F(u(t)).$$

The system (1) is known as a gradient-like system. In particular, the  $\omega$ -limit set of a bounded solution consists only of equilibrium points. A natural question is the study of convergence of such a solution.

Haraux [10] proved under quite general assumptions on  $F$  that if  $d = 1$  and  $0 \leq \alpha < 1$ , every bounded solution converges to a critical point of  $F$  as time goes to infinity. This is no longer true if the damping is too weak ( $\alpha \geq 1$ ), as pointed out in [10], or if  $d \geq 2$ , as proved in [14].

However, if  $F$  satisfies additional properties, more can be said. If  $F$  is real analytic,  $\alpha = 0$  and  $d$  is arbitrary, Haraux and Jendoubi [12] proved that every bounded solution to (1) converges to a single point. Their result is based on the celebrated Lojasiewicz inequality [15, 16]. A similar convergence result was obtained by Chergui [7] in the case  $0 \leq \alpha < 1$ , where  $\alpha$  depends also on the global Lojasiewicz exponent of the function  $F$ . We refer the reader to the book [11] for details on this matter. Alvarez [3] proved convergence to an equilibrium for convex functions in the case  $\alpha = 0$  (see also [6]). Generalizations of (1) have been considered in [5].

Discretizations of the system (1) have also been studied. In [3] mentioned above, Alvarez also studied the asymptotic behaviour for convex functions of a fully implicit discretization in the case  $\alpha = 0$ . Convergence to equilibrium was also proved in [9] for a fully implicit scheme in the case of real analytic functions and linear damping ( $\alpha = 0$ ). A semi-implicit scheme was considered in [2] for functions satisfying a Lojasiewicz inequality. Horsin and Jendoubi [13] studied the nonlinear damping  $0 < \alpha < 1$  for a fully implicit scheme. Related schemes were considered in [4].

Our main purpose in this paper is to study an explicit discretization of (1) for a linear damping and a function  $F$  which satisfies a global Lojasiewicz inequality. In contrast to the

fully implicit discretization, we do not prove that the energy is nonincreasing but instead, we prove that a modified energy is nonincreasing. This is achieved in Section 2 and the proof of convergence is based on an angle condition introduced in [1]. Convergence rates are also established.

In Section 3, we consider an implicit discretization of (1) in the one-dimensional case ( $d = 1$ ) and we prove that the sequence generated by the scheme converges for  $0 \leq \alpha < 1$ , without assuming that  $F$  satisfies a Lojasiewicz inequality. The optimality of  $\alpha$  is established in Section 4, where a non convergence result is shown to hold for  $\alpha = 1$  and  $d = 1$ , even if  $F$  satisfies a Lojasiewicz inequality with optimal exponent  $\theta = 1/2$ . We note that altogether, in Sections 2-4, the discrete case mimicks the situation for the continuous system (1). Numerical simulations illustrate the theoretical results in the last section.

## 2 Convergence to equilibrium for an explicit scheme

In this section, we prove convergence to equilibrium for an explicit discretization of the system (1). We first recall a general convergence result.

### 2.1 A discrete angle condition

Let  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}$  be a  $C^1$  function,  $\sigma > 0$  and let us consider a sequence  $(x_n)$  satisfying for all  $n \in \mathbb{N}$

$$\Phi(x_n) - \Phi(x_{n+1}) \geq \sigma \|\nabla \Phi(x_n)\| \|x_{n+1} - x_n\|, \quad (2)$$

$$[\Phi(x_{n+1}) = \Phi(x_n)] \implies [x_{n+1} = x_n]. \quad (3)$$

**Theorem 1 (Absil et al. [1])** *We assume that there exists  $\theta \in (0, \frac{1}{2}]$  such that*

$$\begin{aligned} \forall a \in \mathbb{R}^N \exists c_a > 0 \exists r_a > 0 / \forall x \in \mathbb{R}^N : \\ \|x - a\| < r_a \implies \|\nabla \Phi(x)\| \geq c_a |\Phi(x) - \Phi(a)|^{1-\theta}. \end{aligned} \quad (4)$$

*Let  $(x_n)$  be a sequence satisfying (2) and (3). Then either  $\lim_{n \rightarrow +\infty} \|x_n\| = +\infty$ , or there exists  $x^\infty \in \mathbb{R}^N$  such that  $\nabla \Phi(x^\infty) = 0$  and*

$$\lim_{n \rightarrow +\infty} x_n = x^\infty.$$

*Precisely, in this case we have*

$$\|x_n - x^\infty\| = \begin{cases} O(e^{-cn}) & \text{for some } c > 0 \text{ if } \theta = \frac{1}{2} \\ O(n^{-\frac{\theta}{1-2\theta}}) & \text{if } \theta \in (0, \frac{1}{2}). \end{cases} \quad (5)$$

**Remark 2** If there exists  $M > 0$  such that  $\forall n \in \mathbb{N}, \|x_n\| \leq M$ , then the assumption (4) may merely apply to those  $a \in \mathbb{R}^N$  such that  $\|a\| \leq M$ .

## 2.2 The explicit scheme

We consider a sequence  $(u_n, v_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}^d \times \mathbb{R}^d$  satisfying

$$\begin{cases} \frac{u_{n+1} - u_n}{\Delta t} = v_n \\ \frac{v_{n+1} - v_n}{\Delta t} = -v_n - \nabla F(u_n) \\ u_0, v_0 \in \mathbb{R}^d \end{cases} \quad (6)$$

where  $\alpha > 0$  and  $F : \mathbb{R}^d \rightarrow \mathbb{R}$  is a  $C^1$  function such that

$$\exists L_F > 0 / \forall u, v \in \mathbb{R}^d \quad \|\nabla F(u) - \nabla F(v)\| \leq L_F \|u - v\|, \quad (7)$$

$$\exists c_1, c_2 > 0 / \forall u \in \mathbb{R}^d \quad F(u) \geq c_1 \|u\|^2 - c_2. \quad (8)$$

It is easy to check (see for example [2, 13]) that hypothesis (7) on  $F$  implies the following inequalities ( $\langle \cdot, \cdot \rangle$  denoting the standard scalar product on  $\mathbb{R}^d$ )

$$\forall u, v \in \mathbb{R}^d \quad \langle \nabla F(u) - \nabla F(v), u - v \rangle \geq -L_F \|u - v\|^2, \quad (9)$$

$$\forall u, v \in \mathbb{R}^d \quad F(v) \geq F(u) + \langle \nabla F(u), v - u \rangle - \frac{L_F}{2} \|u - v\|^2. \quad (10)$$

The existence and uniqueness of a sequence satisfying (6) are obvious.

**Proposition 3** *For any  $(u_0, v_0) \in \mathbb{R}^{2d}$  the sequence  $(u_n, v_n)$  given by (6) is well defined.*

We define

$$E(u, v) = \frac{1}{2} \|v\|^2 + F(u).$$

**Proposition 4** *Assume  $F$  satisfies (7). Let  $(u_n, v_n)$  be a sequence satisfying (6), then we have*

$$E(u_{n+1}, v_{n+1}) - E(u_n, v_n) \leq -\Delta t \left[ 1 - \left( 1 + \frac{3L_F}{2} \right) \Delta t \right] \|v_n\|^2 + (\Delta t)^2 \|\nabla F(u_n)\|^2.$$

*Proof.* By taking the scalar product of the second relation of (6) with  $\Delta t v_n$ , it comes

$$\begin{aligned} \langle \frac{v_{n+1} - v_n}{\Delta t}, \Delta t v_n \rangle &= \langle -v_n - \nabla F(u_n), \Delta t v_n \rangle, \\ \langle v_n, v_{n+1} \rangle - \|v_n\|^2 &= -\Delta t \|v_n\|^2 - \langle \nabla F(u_n), \Delta t v_n \rangle, \\ \frac{1}{2} \|v_{n+1}\|^2 - \frac{1}{2} \|v_n\|^2 - \frac{1}{2} \|v_{n+1} - v_n\|^2 &= -\Delta t \|v_n\|^2 - \langle \nabla F(u_n), u_{n+1} - u_n \rangle. \end{aligned}$$

Then we obtain

$$\begin{aligned} & E(u_{n+1}, v_{n+1}) - E(u_n, v_n) - \frac{1}{2} \|v_{n+1} - v_n\|^2 \\ & \leq -\Delta t \|v_n\|^2 + F(u_{n+1}) - F(u_n) - \langle \nabla F(u_n), u_{n+1} - u_n \rangle \\ & \leq -\Delta t \|v_n\|^2 + F(u_{n+1}) - F(u_n) - \langle \nabla F(u_{n+1}), u_{n+1} - u_n \rangle \\ & \quad + \langle \nabla F(u_{n+1}) - \nabla F(u_n), u_{n+1} - u_n \rangle. \end{aligned}$$

By using (7) and (10), we get

$$\begin{aligned}
& E(u_{n+1}, v_{n+1}) - E(u_n, v_n) \\
& \leq -\Delta t \|v_n\|^2 + \frac{1}{2} \|v_{n+1} - v_n\|^2 + \frac{L_F}{2} \|u_{n+1} - u_n\|^2 + L_F \|u_{n+1} - u_n\|^2 \\
& \leq -\Delta t \left[ 1 - \frac{3L_F}{2} \Delta t \right] \|v_n\|^2 + \frac{1}{2} \|v_{n+1} - v_n\|^2.
\end{aligned}$$

Since

$$\frac{1}{2} \|v_{n+1} - v_n\|^2 = \frac{(\Delta t)^2}{2} \|v_n + \nabla F(u_n)\|^2 \leq (\Delta t)^2 \|v_n\|^2 + (\Delta t)^2 \|\nabla F(u_n)\|^2,$$

we deduce

$$E(u_{n+1}, v_{n+1}) - E(u_n, v_n) \leq -\Delta t \left[ 1 - \left( 1 + \frac{3L_F}{2} \right) \Delta t \right] \|v_n\|^2 + (\Delta t)^2 \|\nabla F(u_n)\|^2.$$

Let  $\mathcal{S} = \{a \in \mathbb{R}^d / \nabla F(a) = 0\}$ .

Now we assume also that there exists  $\theta \in (0, \frac{1}{2}]$  such that

$$\forall a \in \mathcal{S} \exists \delta_a > 0 \exists \nu_a > 0 / \forall u \in \mathbb{R}^d : \|u - a\| < \delta_a \implies \|\nabla F(u)\| \geq \nu_a |F(u) - F(a)|^{1-\theta}. \quad (11)$$

□

**Proposition 5** ([15, 16, 8, 7]) *Assumption (11) is satisfied if one of the following two cases holds:*

- $F$  is a polynomial, or
- $F$  is analytic and  $\mathcal{S}$  is compact.

The first part of this proposition is a result of D'Acunto and Kurdyka [8]. The proof of the second part can be found in [7].

**Theorem 6** *Let  $F : \mathbb{R}^d \rightarrow \mathbb{R}$  be a  $C^2$  function satisfying (7), (8) and (11). Assume also that  $\Delta t$  is small enough. Let  $(u_n, v_n)$  be a sequence satisfying (6). Then  $(u_n, v_n)$  is bounded and there exists  $a \in \mathcal{S}$  such that*

$$\lim_{n \rightarrow +\infty} \|v_n\| + \|u_n - a\| = 0.$$

*In addition as  $n \rightarrow +\infty$  we have*

$$\|v_n\| + \|u_n - a\| = \begin{cases} O(e^{-cn}) & \text{for some } c > 0 \text{ if } \theta = \frac{1}{2} \\ O(n^{-\frac{\theta}{1-2\theta}}) & \text{if } \theta \in (0, \frac{1}{2}). \end{cases} \quad (12)$$

**Remark 7** We are able to give an upper bound of  $\Delta t$  namely

$$0 < \Delta t < \min \left( \frac{1}{4 + 6L_F}, \frac{1}{4 + 3L_F(1 + \eta)}, \frac{\eta}{2 + L_F\eta} \right)$$

with  $\eta = \min \left( \frac{1}{2}, \frac{c_1}{2L_F^2}, \frac{1}{2L_F+1}, \frac{1}{2C} \right)$ , where  $C = \max \left( 1, \max_{\|u\| \leq R} \|\nabla^2 F(u)\| \right)$  and  $R$  is given by (17).

*Proof.* Let  $\varepsilon$  be a positive real. We define for all  $u, v \in \mathbb{R}^d$

$$\Phi_\varepsilon(u, v) = E(u, v) + \varepsilon \langle \nabla F(u), v \rangle.$$

First of all we will prove that for  $\varepsilon$  small enough,  $\Phi_\varepsilon$  will be coercive. In fact, using (7), we see that

$$\forall u \in \mathbb{R}^d \quad \|\nabla F(u)\| \leq L_F \|u\| + \|\nabla F(0)\|,$$

and then

$$\forall u \in \mathbb{R}^d \quad \|\nabla F(u)\|^2 \leq 2L_F^2 \|u\|^2 + 2\|\nabla F(0)\|^2. \quad (13)$$

Now for all  $(u, v) \in \mathbb{R}^d$ , we have by Cauchy-Schwarz inequality

$$\begin{aligned} \Phi_\varepsilon(u, v) &\geq \frac{1}{2} \|v\|^2 + F(u) - \frac{\varepsilon}{2} \|v\|^2 - \frac{\varepsilon}{2} \|\nabla F(u)\|^2 \\ &\stackrel{\text{by (8) and (13)}}{\geq} \left( \frac{1}{2} - \frac{\varepsilon}{2} \right) \|v\|^2 + (c_1 - \varepsilon L_F^2) \|u\|^2 - c_2 - \varepsilon \|\nabla F(0)\|^2. \end{aligned}$$

We impose to  $\varepsilon$  to be such that  $\varepsilon \leq \min \left( \frac{1}{2}, \frac{c_1}{2L_F^2} \right)$ . We see that with this choice, we have

$$\Phi_\varepsilon(u, v) \geq \frac{1}{4} \|v\|^2 + \frac{c_1}{2} \|u\|^2 - c_2 - \frac{1}{4} \|\nabla F(0)\|^2, \quad (14)$$

so that  $\Phi_\varepsilon$  is coercive.

Next, we prove that for  $\varepsilon$  small enough,  $(\Phi_\varepsilon(u_n, v_n))$  is nonincreasing. Let us define  $x_n = (u_n, v_n)$ . According to Proposition 4, for all  $n \in \mathbb{N}$  :

$$\begin{aligned} &\Phi_\varepsilon(x_{n+1}) - \Phi_\varepsilon(x_n) \\ &\leq -\Delta t \left[ 1 - \left( 1 + \frac{3L_F}{2} \right) \Delta t \right] \|v_n\|^2 + (\Delta t)^2 \|\nabla F(u_n)\|^2 \\ &\quad + \varepsilon \underbrace{\left[ \langle \nabla F(u_{n+1}), v_{n+1} \rangle - \langle \nabla F(u_n), v_n \rangle \right]}_{T_n}. \end{aligned}$$

We have

$$\begin{aligned}
T_n &= \langle \nabla F(u_{n+1}) - \nabla F(u_n) + \nabla F(u_n), v_n - \Delta t v_n - \Delta t \nabla F(u_n) \rangle - \langle \nabla F(u_n), v_n \rangle \\
&= \langle \nabla F(u_{n+1}) - \nabla F(u_n), v_n - \Delta t v_n - \Delta t \nabla F(u_n) \rangle - \Delta t \|\nabla F(u_n)\|^2 \\
&\quad - \Delta t \langle \nabla F(u_n), v_n \rangle \\
&\leq \|\nabla F(u_{n+1}) - \nabla F(u_n)\| \|v_n - \Delta t v_n - \Delta t \nabla F(u_n)\| - \Delta t \|\nabla F(u_n)\|^2 \\
&\quad - \Delta t \langle \nabla F(u_n), v_n \rangle \\
&\stackrel{\text{by (7)}}{\leq} L_F \|u_{n+1} - u_n\| \|v_n - \Delta t v_n - \Delta t \nabla F(u_n)\| - \Delta t \|\nabla F(u_n)\|^2 - \Delta t \langle \nabla F(u_n), v_n \rangle \\
&\stackrel{\text{by (6)}}{\leq} L_F \Delta t \|v_n\| \|v_n - \Delta t v_n - \Delta t \nabla F(u_n)\| - \Delta t \|\nabla F(u_n)\|^2 - \Delta t \langle \nabla F(u_n), v_n \rangle \\
&\leq L_F \Delta t \|v_n\|^2 + L_F (\Delta t)^2 \|v_n\|^2 + L_F (\Delta t)^2 \|v_n\| \|\nabla F(u_n)\| - \Delta t \|\nabla F(u_n)\|^2 + \\
&\quad \Delta t \|\nabla F(u_n)\| \|v_n\| \\
&\leq \Delta t \left[ \left( L_F + \frac{1}{2} \right) + \frac{3}{2} L_F \Delta t \right] \|v_n\|^2 - \frac{\Delta t}{2} (1 - L_F \Delta t) \|\nabla F(u_n)\|^2
\end{aligned} \tag{15}$$

where we used the triangle and the Cauchy-Schwarz inequalities. Then we get

$$\begin{aligned}
&\Phi_\varepsilon(x_{n+1}) - \Phi_\varepsilon(x_n) \\
&\leq -\Delta t \left[ 1 - \left( 1 + \frac{3L_F}{2} \right) \Delta t \right] \|v_n\|^2 + (\Delta t)^2 \|\nabla F(u_n)\|^2 \\
&\quad + \varepsilon \Delta t \left[ \left( L_F + \frac{1}{2} \right) + \frac{3}{2} L_F \Delta t \right] \|v_n\|^2 - \varepsilon \frac{\Delta t}{2} (1 - L_F \Delta t) \|\nabla F(u_n)\|^2 \\
&\leq -\Delta t \left[ 1 - \varepsilon \left( L_F + \frac{1}{2} \right) - \left( 1 + \frac{3L_F}{2} (1 + \varepsilon) \right) \Delta t \right] \|v_n\|^2 \\
&\quad - \Delta t \left[ \frac{\varepsilon}{2} - (1 + \frac{\varepsilon}{2} L_F) \Delta t \right] \|\nabla F(u_n)\|^2
\end{aligned}$$

We choose  $\varepsilon \leq \bar{\varepsilon} = \min \left( \frac{1}{2}, \frac{c_1}{2L_F^2}, \frac{1}{2L_F+1} \right)$ . Then we obtain

$$\begin{aligned}
&\Phi_\varepsilon(x_{n+1}) - \Phi_\varepsilon(x_n) \\
&\leq -\Delta t \left[ \frac{1}{2} - \left( 1 + \frac{3L_F}{2} (1 + \varepsilon) \right) \Delta t \right] \|v_n\|^2 - \Delta t \left[ \frac{\varepsilon}{2} - (1 + \frac{\varepsilon}{2} L_F) \Delta t \right] \|\nabla F(u_n)\|^2.
\end{aligned}$$

We assume that  $\Delta t$  is small enough such that  $\frac{1}{2} - (1 + \frac{9L_F}{4}) \Delta t > 0$  (then  $\frac{1}{2} - (1 + \frac{3L_F}{2} (1 + \varepsilon)) \Delta t > 0$  since  $\varepsilon \leq \frac{1}{2}$ ) and  $\frac{\varepsilon}{2} - (1 + \frac{\varepsilon}{2} L_F) \Delta t > 0$ .

Then there is a constant  $\delta = \delta(\Delta t, L_F) > 0$  such that

$$\forall n \in \mathbb{N} \quad \Phi_\varepsilon(x_{n+1}) - \Phi_\varepsilon(x_n) \leq -\delta (\|v_n\|^2 + \|\nabla F(u_n)\|^2). \tag{16}$$

Then the sequence  $(\Phi_\varepsilon(x_n))$  is nonincreasing. Using (14), we see that  $(u_n)$  and  $(v_n)$  are



bounded:

$$\begin{aligned}
& \Phi_\varepsilon(x_n) \leq \Phi_\varepsilon(x_0) \\
\implies & \frac{1}{4}\|v_n\|^2 + \frac{c_1}{2}\|u_n\|^2 - c_2 - \frac{1}{4}\|\nabla F(0)\|^2 \leq \frac{1}{2}\|v_0\|^2 + F(u_0) + \varepsilon < \nabla F(u_0), v_0 > \\
\implies & \frac{1}{4}\|v_n\|^2 + \frac{c_1}{2}\|u_n\|^2 \leq \frac{1}{2}\|v_0\|^2 + F(u_0) + \|\nabla F(u_0)\|\|v_0\| + c_2 + \frac{1}{4}\|\nabla F(0)\|^2 \\
\implies & \min\left(\frac{1}{8}, \frac{c_1}{4}\right)[\|v_n\| + \|u_n\|]^2 \leq \frac{1}{2}\|v_0\|^2 + F(u_0) + \|\nabla F(u_0)\|\|v_0\| + c_2 + \frac{1}{4}\|\nabla F(0)\|^2 \\
\implies & \|v_n\| + \|u_n\| \leq R
\end{aligned}$$

where

$$R = \left[ \frac{1}{\min\left(\frac{1}{8}, \frac{c_1}{4}\right)} \left( \frac{1}{2}\|v_0\|^2 + F(u_0) + \|\nabla F(u_0)\|\|v_0\| + c_2 + \frac{1}{4}\|\nabla F(0)\|^2 \right) \right]^{\frac{1}{2}}. \quad (17)$$

At this stage, it is important to note that if we choose  $\varepsilon$  smaller, (16) remains true. Note also that  $R$  is independent of  $\Delta t$  and  $\varepsilon$ .

Obviously we have  $\sum (\|v_n\|^2 + \|\nabla F(u_n)\|^2)$  converges. Then  $v_n \rightarrow 0$  and the  $\omega$ -limit set  $\omega((u_n)_{n \in \mathbb{N}}) := \{a \in \mathbb{R}^d : \exists n_k \rightarrow \infty / u_{n_k} \rightarrow a\}$  is a nonempty compact connected subset of  $\mathcal{S}$ .

Let us show that  $(x_n) = (u_n, v_n)$  satisfies (2) and (3) with the function  $\Phi_\varepsilon$ . Indeed, if for some  $n$  we have  $\Phi_\varepsilon(x_{n+1}) = \Phi_\varepsilon(x_n)$ , then by (16) we deduce that  $v_n = 0$  and  $\nabla F(u_n) = 0$ . Using system (6), we get that  $u_{n+1} = u_n$  and  $v_{n+1} = v_n$ , that is  $x_{n+1} = x_n$  and (3) is satisfied. On the other hand, a simple computation gives

$$\nabla \Phi_\varepsilon(u, v) = \begin{pmatrix} \nabla F(u) + \varepsilon \nabla^2 F(u) \cdot v \\ v + \varepsilon \nabla F(u) \end{pmatrix}.$$

Since  $(u_n)$  and  $(v_n)$  are bounded, there exists a constant  $\eta_1 > 0$  such that

$$\forall n \in \mathbb{N} \quad \|\nabla \Phi_\varepsilon(x_n)\| \leq \eta_1 [\|v_n\| + \|\nabla F(u_n)\|]. \quad (18)$$

On the other hand, there exists a constant  $\eta_2 > 0$  such that

$$\begin{aligned}
\|x_{n+1} - x_n\| &= \|(u_{n+1} - u_n, v_{n+1} - v_n)\| \\
&= \|(\Delta t v_n, -\Delta t v_n - \Delta t \nabla F(u_n))\| \\
&\leq \eta_2 [\|v_n\| + \|\nabla F(u_n)\|].
\end{aligned} \quad (19)$$

By combining (16) – (18) and (19), we get that  $(\Phi_\varepsilon(x_n))$  satisfies (2) with  $\sigma = \frac{\delta}{2\eta_1\eta_2}$ . Now we will show that under the hypothesis (11) on  $F$ ,  $\Phi_\varepsilon$  satisfies (4).

From the remark 2, let  $B = \{(u, v) \in \mathbb{R}^d \times \mathbb{R}^d, \|u\| + \|v\| \leq R\}$  where  $R$  is given by (17). Clearly  $B$  contains the sequence  $(u_n, v_n)$ . Now, for all  $(u, v) \in B$

$$\begin{aligned}
& \|\nabla\Phi_\varepsilon(u, v)\| \\
&= \|\nabla F(u) + \varepsilon\nabla^2 F(u) \cdot v\| + \|v + \varepsilon\nabla F(u)\| \\
&\geq \|\nabla F(u)\| - \varepsilon\|\nabla^2 F(u) \cdot v\| + \|v\| - \varepsilon\|\nabla F(u)\| \\
&\geq (1 - \varepsilon \max_{\|u\| \leq R} \|\nabla^2 F(u)\|)\|v\| + (1 - \varepsilon)\|\nabla F(u)\| \\
&\geq (1 - \varepsilon C)(\|v\| + \|\nabla F(u)\|)
\end{aligned}$$

where  $C = \max\left(1, \max_{\|u\| \leq R} \|\nabla^2 F(u)\|\right)$ .

By possibly taking  $\varepsilon > 0$  smaller ( $\varepsilon = \min(\bar{\varepsilon}, \frac{1}{2C})$ ), there exists  $\rho > 0$  such that

$$\forall (u, v) \in B \quad \|\nabla\Phi_\varepsilon(u, v)\| \geq \rho(\|v\| + \|\nabla F(u)\|). \quad (20)$$

If  $(a, b)$  is not a critical point of  $\Phi_\varepsilon$ , then  $\Phi_\varepsilon$  satisfies (4) with  $\theta = \frac{1}{2}$  as best exponent, thanks to the continuity of  $\Phi_\varepsilon$ .

Let  $(a, b) \in B$  be a critical point of  $\Phi_\varepsilon$ . Then  $\nabla F(a) = 0$  and  $b = 0$ . From (11)

$$\exists \delta_a > 0 \exists \nu_a > 0 / \forall u \in \mathbb{R}^d : \|u - a\| < \delta_a \implies \|\nabla F(u)\| \geq \nu_a |F(u) - F(a)|^{1-\theta}. \quad (21)$$

On the other hand, by using the Cauchy-Schwarz inequality, we get

$$\begin{aligned}
[\Phi_\varepsilon(u, v) - \Phi_\varepsilon(a, 0)]^{1-\theta} &= \left[\frac{1}{2}\|v\|^2 + F(u) - F(a) + \varepsilon \langle \nabla F(u), v \rangle\right]^{1-\theta} \\
&\leq \|v\|^{2(1-\theta)} + |F(u) - F(a)|^{1-\theta} + \|\nabla F(u)\|^{1-\theta} \|v\|^{1-\theta}.
\end{aligned} \quad (22)$$

Thanks to Young's inequality we obtain

$$\|\nabla F(u)\|^{1-\theta} \|v\|^{1-\theta} \leq \|\nabla F(u)\| + \|v\|^{\frac{1-\theta}{\theta}}.$$

Then (22) becomes

$$[\Phi_\varepsilon(u, v) - \Phi_\varepsilon(a, 0)]^{1-\theta} \leq \|v\|^{2(1-\theta)} + |F(u) - F(a)|^{1-\theta} + \|\nabla F(u)\| + \|v\|^{\frac{1-\theta}{\theta - \alpha(1-\theta)}}.$$

Since  $2(1 - \theta)$  and  $\frac{1-\theta}{\theta}$  are bigger than 1, using also (21), we get for all  $(u, v) \in B$  with  $\|v\| \leq 1$  and  $\|u - a\| < \delta_a$

$$\begin{aligned}
[\Phi_\varepsilon(u, v) - \Phi_\varepsilon(a, 0)]^{1-\theta} &\leq \|v\| + |F(u) - F(a)|^{1-\theta} + \|\nabla F(u)\| + \|v\| \\
&\leq \left(2 + \frac{1}{\nu_a}\right) (\|\nabla F(u)\| + \|v\|) \\
&\leq \frac{1}{\rho} \left(2 + \frac{1}{\nu_a}\right) \|\nabla\Phi_\varepsilon(u, v)\| \quad \text{by (20)}.
\end{aligned}$$

Therefore  $\Phi_\varepsilon$  satisfies (4). Theorem 1 is proved.  $\square$

### 3 A convergence result in the one dimensional case

We consider the case of a strong nonlinear damping ( $0 \leq \alpha < 1$ ) in the one-dimensional situation ( $d = 1$ ). This time, we use an implicit discretization of (1). The proof is adapted from the continuous case. We recall it for the reader's convenience.

#### 3.1 The continuous version

Let  $u \in W^{1,\infty}(\mathbb{R})$  be a bounded solution of the ODE

$$u'' + |u'|^\alpha u' + f(u) = 0,$$

where  $f \in C^1(\mathbb{R}, \mathbb{R})$ . We know that (see, e.g., [11])

$$\omega(u, u') \subset f^{-1}(\{0\}) \times \{0\},$$

where  $\omega(u, u')$  is the  $\omega$ -limit set of the solution defined by

$$\omega(u, u') = \{(u^*, v^*) \in \mathbb{R}^2, \exists t_n \rightarrow +\infty, (u(t_n), u'(t_n)) \rightarrow (u^*, v^*) \text{ as } n \rightarrow +\infty\}.$$

**Theorem 8 ([10, 11])** *There exists  $a \in f^{-1}(\{0\})$  such that*

$$\lim_{t \rightarrow \infty} |u'(t)| + |u(t) - a| = 0.$$

*Proof.* For the sake of completeness, we give the proof here. If  $\omega(u, u')$  is not reduced to a single point, then there are  $a, b \in f^{-1}(\{0\})$  ( $a < b$ ) such that  $\omega(u, u') = [a, b] \times \{0\}$ . Let  $c = \frac{a+b}{2}$ . Since  $\lim_{t \rightarrow \infty} u'(t) = 0$  and  $(c, 0) \in \omega(u, u')$ , then there exists  $T > 0$  such that

$$\begin{aligned} |u(T) - c| &< \frac{b-a}{8}, \\ \frac{|u'(t)|^{1-\alpha}}{1-\alpha} &< \frac{b-a}{8} \quad \forall t \geq T. \end{aligned}$$

We will prove that

$$\forall t \geq T \quad u(t) \in [a, b].$$

Assume that this is not the case and let

$$\theta = \inf\{t \geq T, u(t) \notin [a, b]\}.$$

Clearly we have  $\theta < \infty$  and

$$\forall t \in [T, \theta] \quad u(t) \in [a, b].$$

Then

$$\forall t \in [T, \theta] \quad u''(t) + |u'(t)|^\alpha u'(t) = 0.$$

If there is a  $s \in [T, \theta]$  such that  $u'(s) = 0$ , then  $u'(t) = 0$  for all  $t \in [T, \theta]$ , in this case we conclude easily. Otherwise, solving this ODE, we get

$$\forall t \in [T, \theta] \quad |u'(t)| = \frac{1}{[|u'(T)|^{-\alpha} + \alpha(t - T)]^{\frac{1}{\alpha}}}. \quad (23)$$

$$\begin{aligned} |u(\theta) - u(T)| &\leq \int_T^\theta |u'(t)| dt \\ &\leq \int_T^\theta \frac{dt}{[|u'(T)|^{-\alpha} + \alpha(t - T)]^{\frac{1}{\alpha}}} \\ &\leq \frac{|u'(T)|^{1-\alpha}}{1-\alpha} < \frac{b-a}{8}. \end{aligned}$$

$$\begin{aligned} |u(\theta) - c| &\leq |u(\theta) - u(T)| + |u(T) - c| \\ &< \frac{b-a}{8} + \frac{b-a}{8} = \frac{b-a}{4}, \end{aligned}$$

which contradicts the definition of  $\theta$  and then  $\theta = \infty$ .

Now from (23), we deduce that  $u' \sim \frac{1}{\alpha^{\frac{1}{\alpha}} t^{\frac{1}{\alpha}}}$ , and then  $u' \in L^1(\mathbb{R}_+)$ . This proves that  $u$  converges and the  $\omega$ -limit set  $\omega(u, u')$  is reduced to a single point.  $\square$

### 3.2 The discrete version

The scheme reads:

$$\begin{cases} \frac{u_{n+1} - u_n}{\Delta t} = v_{n+1} \\ \frac{v_{n+1} - v_n}{\Delta t} = -|v_{n+1}|^\alpha v_{n+1} - f(u_{n+1}) \\ u_0, v_0 \in \mathbb{R} \end{cases} \quad (24)$$

and  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\exists c_F > 0 / \forall u, v \in \mathbb{R}^d \quad (f(u) - f(v))(u - v) \geq -c_F |u - v|^{\alpha+2}, \quad (25)$$

$$\exists L_F > 0 / \forall u, v \in \mathbb{R}^d \quad |f(u) - f(v)| \leq L_F \|u - v\|. \quad (26)$$

Let  $\mathcal{S} = \{a \in \mathbb{R}^d / f(a) = 0\}$  and  $F$  be an antiderivative of  $f$ . We define

$$E(u, v) = \frac{1}{2}|v|^2 + F(u).$$

We recall the following results.

**Proposition 9** [13] Assume that  $F$  is of class  $C^1(\mathbb{R}^d)$ , coercive and that (25) and (26) hold, then for any  $(u_0, v_0) \in \mathbb{R}^{2d}$ , provided  $\Delta t$  is small enough, the sequence  $(u_n, v_n)$  given by (24) is well defined, and we have

$$\forall n \in \mathbb{N} \quad E(u_{n+1}, v_{n+1}) \leq E(u_n, v_n). \quad (27)$$

Moreover we have

$$E(u_{n+1}, v_{n+1}) - E(u_n, v_n) \leq -\Delta t \left[ 1 - \frac{c_F}{2} (\Delta t)^{\alpha+1} \right] |v_{n+1}|^{\alpha+2}.$$

**Corollary 10** [13] Let  $F$  satisfy (25) and assume that  $0 < \Delta t < \left(\frac{2}{c_F}\right)^{\frac{1}{\alpha+1}}$ . Let  $(u_n, v_n)$  be a sequence satisfying (24). If  $(u_n)$  is bounded, then  $\lim_{n \rightarrow \infty} E(u_n, v_n)$  exists and  $v_n \rightarrow 0$ .

Moreover the  $\omega$ -limit set  $\omega((u_n)_{n \in \mathbb{N}})$  given by

$$\omega((u_n)_{n \in \mathbb{N}}) = \{a \in \mathbb{R} : \exists n_k \rightarrow \infty / u_{n_k} \rightarrow a\}$$

is a nonempty compact connected subset of  $\mathcal{S}$ .

The main result of this paper is the following

**Theorem 11** Under the assumptions of the corollary 10, there exists  $a \in \mathcal{S}$  such that  $u^n \rightarrow a$ .

*Proof.* As in the continuous case, by connectedness we have either  $\omega((u_n)_{n \in \mathbb{N}}) = \{a\} \times \{0\}$  for some  $a \in f^{-1}(\{0\})$  and the result is established, or  $\omega((u_n)_{n \in \mathbb{N}}) = [a, b] \times \{0\}$  for some  $a, b \in f^{-1}(\{0\})$  ( $a < b$ ). In this case, let  $c = \frac{a+b}{2}$ . Using corollary 10, we can choose  $n_0$  large enough such that

$$|u_{n_0} - c| < \frac{b-a}{8}, \quad (28)$$

$$\Delta t |v^{n_0}| + \frac{2^{\alpha+1}}{1-\alpha} |v^{n_0}| < \frac{b-a}{8} \quad \forall n \geq n_0, \quad (29)$$

$$\Delta t |v_n|^\alpha \leq 1, \quad \forall n \geq n_0. \quad (30)$$

Assume by contradiction that the set  $\{n \geq n_0 : u^{n+1} \notin [a, b]\}$  is nonempty and let

$$n_1 = \inf \{n \geq n_0 : u^{n+1} \notin [a, b]\}.$$

Note that  $n_1 > n_0 + 2$ . In fact by (29), we have  $\Delta t |v_{n_0+1}| < \frac{b-a}{8}$  and  $\Delta t |v_{n_0+2}| < \frac{b-a}{8}$ . Using (24) and triangle inequality, we find

$$\begin{aligned} |u_{n_0+1} - c| &\leq |u_{n_0+1} - u_{n_0}| + |u_{n_0} - c| \\ &= \Delta t |v_{n_0+1}| + |u_{n_0} - c| < \frac{b-a}{8} + \frac{b-a}{8} = \frac{b-a}{4}; \\ |u_{n_0+2} - c| &\leq |u_{n_0+2} - u_{n_0+1}| + |u_{n_0+1} - c| \\ &= \Delta t |v_{n_0+2}| + |u_{n_0} - c| < \frac{b-a}{8} + \frac{b-a}{4} = 3 \frac{b-a}{8}. \end{aligned}$$

It is clear that

$$\forall n \in \{n_0, \dots, n_1 - 1\}, u^{n+1} \in [a, b]$$

and then  $f(u^{n+1}) = 0$  since  $[a, b] \subset \mathcal{S}$ . Using (24), we get

$$\forall n \in \{n_0, \dots, n_1 - 1\}, \frac{v_{n+1} - v_n}{\Delta t} = -|v_{n+1}|^\alpha v_{n+1}.$$

which can also be written

$$\forall n \in \{n_0, \dots, n_1 - 1\}, v_{n+1}(1 + \Delta t |v_{n+1}|^\alpha) = v_n.$$

Obviously we have

$$\forall n \in \{n_0, \dots, n_1 - 1\}, |v_n| - |v_{n+1}| = \Delta t |v_{n+1}|^{\alpha+1} \geq 0. \quad (31)$$

Now assume that there exists  $p \in \{n_0, \dots, n_1 - 1\}$  such that  $v^p = 0$ . Then for all  $n \in \{n_0, \dots, n_1 - 1\}$ ,  $v^n = 0$ . In this case  $n_1 = \infty$  and  $(u^n)$  is a constant sequence.

Otherwise for all  $n \in \{n_0, \dots, n_1 - 1\}$ ,  $v^n \neq 0$ . Now from (30) and (31) we deduce that

$$\forall n \in \{n_0, \dots, n_1 - 1\}, |v_n| \leq 2|v_{n+1}|. \quad (32)$$

Then we get from (31)

$$\begin{aligned} \Delta t &= \frac{|v_n| - |v_{n+1}|}{|v_{n+1}|^{\alpha+1}} \\ &\stackrel{\text{by (31) and (32)}}{\leq} 2^{\alpha+1} \frac{|v_n| - |v_{n+1}|}{|v_n|^{\alpha+1}} \\ &\leq 2^{\alpha+1} \int_{|v_{n+1}|}^{|v_n|} \frac{dt}{t^{\alpha+1}} \\ &= \frac{2^{\alpha+1}}{\alpha} [-|v^n|^{-\alpha} + |v^{n+1}|^{-\alpha}]. \end{aligned}$$

Summing from  $n_0$  to  $n \in \{n_0, \dots, n_1 - 1\}$ , we obtain

$$\frac{\alpha}{2^{\alpha+1}} \Delta t (n - n_0) \leq |v^{n+1}|^{-\alpha} - |v^{n_0}|^{-\alpha}$$

or

$$|v^{n+1}| \leq \frac{1}{[|v^{n_0}|^{-\alpha} + \frac{\alpha}{2^{\alpha+1}} \Delta t (n - n_0)]^{\frac{1}{\alpha}}}. \quad (33)$$

Now let  $n \in \{n_0, \dots, n_1 - 1\}$ , we have

$$\begin{aligned}
|u^{n+1} - u^{n_0}| &\leq \sum_{k=n_0}^n |u^{k+1} - u^k| \\
&\stackrel{\text{by (24)}}{=} \Delta t \sum_{k=n_0}^n |v^{k+1}| \\
&\stackrel{\text{by (33)}}{\leq} \Delta t \sum_{k=n_0}^n \frac{1}{\left[|v^{n_0}|^{-\alpha} + \frac{\alpha}{2^{\alpha+1}} \Delta t (k - n_0)\right]^{\frac{1}{\alpha}}} \\
&\leq \Delta t |v^{n_0}| + \Delta t \int_{n_0}^n \frac{dx}{\left[|v^{n_0}|^{-\alpha} + \frac{\alpha}{2^{\alpha+1}} \Delta t (x - n_0)\right]^{\frac{1}{\alpha}}} \\
&\leq \Delta t |v^{n_0}| + \frac{2^{\alpha+1}}{1 - \alpha} |v^{n_0}| \\
&\stackrel{\text{by (29)}}{<} \frac{b - a}{8}. \tag{34}
\end{aligned}$$

In particular with  $n = n_1 - 1$ , we get  $|u^{n_1} - u^{n_0}| < \frac{b-a}{8}$ . By the triangle inequality we deduce

$$\begin{aligned}
|u^{n_1} - c| &\leq |u^{n_1} - u^{n_0}| + |u^{n_0} - c| \\
&\stackrel{\text{by (28)}}{<} \frac{b - a}{8} + \frac{b - a}{8} = \frac{b - a}{4},
\end{aligned}$$

and this contradicts  $u^{n_1} \notin [a, b]$ . Then the set  $\{n \geq n_0 : u^{n+1} \notin [a, b]\}$  is empty and  $n_1 = \infty$ . Now (34) holds for all  $n \geq n_0$  :

$$|u^{n+1} - u^{n_0}| < \frac{b - a}{4}$$

and by triangle inequality, we get

$$\forall n \geq n_0 \quad |u^{n+1} - c| \leq |u^{n+1} - u^{n_0}| + |u^{n_0} - c| < \frac{b - a}{4} + \frac{b - a}{8} = \frac{3(b - a)}{8}.$$

By choosing a subsequence such that  $u^{n_k} \rightarrow b$ , we obtain

$$|b - c| \leq \frac{3(b - a)}{8},$$

a contradiction. We conclude that  $\omega((u^n))$  is a reduced to a single point  $u^*$  and that  $u^n \rightarrow u^*$ .  $\square$

## 4 A non convergence result

We adapt to the discrete case a result from [10, 11]. Let  $\Delta t > 0$  denote the time step. We consider a sequence defined by

$$\left\{ \begin{array}{l} \frac{u_{n+1} - u_n}{\Delta t} = v_{n+1}, \\ \frac{v_{n+1} - v_n}{\Delta t} = -|v_{n+1}|v_{n+1} - f(u_{n+1}), \\ u_0, v_0 \in \mathbb{R}, \end{array} \right. \quad (35)$$

where  $f : \mathbb{R} \rightarrow \mathbb{R}$  is locally Lipschitz continuous and satisfies

$$f \text{ is nondecreasing,} \quad (36)$$

$$f(s) < 0, \quad \forall s < a, \quad (37)$$

$$f(s) = 0, \quad \forall s \in [a, b], \quad (38)$$

$$f(s) > 0, \quad \forall s > b. \quad (39)$$

Here  $a, b \in \mathbb{R}$  are such that  $a < b$ .

We denote by  $F$  an antiderivative of  $f$ . The assumptions on  $f$  imply that  $F$  is convex and that there exist two positive constants  $c_1, c_2$  such that

$$F(s) \geq c_1|s| - c_2, \quad \forall s \in \mathbb{R}. \quad (40)$$

Since  $F$  is convex, we have

$$F(u) - F(\tilde{u}) \geq f(\tilde{u})(u - \tilde{u}), \quad \forall u, \tilde{u} \in \mathbb{R}. \quad (41)$$

Since  $F$  is coercive, we have:

**Proposition 12** *For every  $(u_0, v_0) \in \mathbb{R}^2$ , there exists at least one sequence  $(u_n, v_n)$  which complies with (35).*

*Proof.* Assume by induction that for some  $n \in \mathbb{N}$ , the values  $(u_n, v_n)$  are defined in  $\mathbb{R}^2$ . We consider the function

$$G_n : u \mapsto \frac{(u - u_n)^2}{2\Delta t^2} - \frac{uv_n}{\Delta t} + \frac{|u - u_n|^3}{3\Delta t^2} + F(u).$$

It is clear that  $G_n$  is continuous on  $\mathbb{R}$  and  $G_n(u) \rightarrow +\infty$  as  $|u| \rightarrow +\infty$  (cf. (40)). Thus,  $G_n$  has at least one minimizer  $u^*$  in  $\mathbb{R}$ , which solves  $G'_n(u^*) = 0$ , that is

$$\frac{u^* - u_n}{\Delta t^2} - \frac{v_n}{\Delta t} + \frac{|u^* - u_n|(u^* - u_n)}{\Delta t^2} + f(u^*) = 0.$$

We set  $u_{n+1} = u^*$  and  $v_{n+1} = (u_{n+1} - u_n)/\Delta t$ . We see that  $u_{n+1}, v_{n+1}$  complies with (35) and we proceed by induction.  $\square$



**Remark 13** Let  $(u_0, v_0) \in \mathbb{R}^2$ . By a fixed point argument, it is possible to prove that if  $\Delta t$  is small enough, there exists a unique sequence  $(u_n, v_n)$  which complies with (35) (see, e.g., [13]).

**Theorem 14** Assume that  $f$  satisfies (36)-(39). For every non stationary sequence  $(u_n, v_n)$  which complies with (35), there exist a subsequence  $(u_{n_k}, v_{n_k})$  such that  $u_{n_k} < a$  for all  $k \in \mathbb{N}$  and a subsequence  $(u_{n_{k'}}, v_{n_{k'}})$  such that  $u_{n_{k'}} > b$  for all  $k' \in \mathbb{N}$ .

*Proof.* We multiply the second equation in (35) by  $\Delta t v_{n+1}$ . We find

$$(v_{n+1} - v_n)v_{n+1} + \Delta t |v_{n+1}|^3 = -f(u_{n+1})(u_{n+1} - u_n).$$

By (41),

$$\frac{1}{2}|v_{n+1}|^2 - \frac{1}{2}|v_n|^2 + \frac{1}{2}|v_{n+1} - v_n|^2 + \Delta t |v_{n+1}|^3 \leq F(u_n) - F(u_{n+1}).$$

Thus,

$$\Delta t |v_{n+1}|^3 \leq E(u_n, v_n) - E(u_{n+1}, v_{n+1}), \quad (42)$$

where

$$E(u, v) := \frac{1}{2}|v|^2 + F(u).$$

The sequence  $(E_n) = (E(u_n, v_n))$  is nonincreasing and since  $E(u, v)$  is coercive by (40), the sequence  $(u_n, v_n)$  is bounded. Moreover,  $(E_n)$  has a limit as  $n$  tends to  $+\infty$ , so  $|v_{n+1}| \rightarrow 0$  by (42).

We assume that  $u_n \geq a$  for all  $n \geq n_0$ , for some  $n_0 \in \mathbb{N}$ . We must prove that  $(u_n)$  is stationary.

- If  $v_n \geq 0$  for all  $n \geq n_0$ , then  $(u_n)$  is nondecreasing and since  $(u_n)$  is bounded, it converges to some value  $c \in [a, b]$ . In particular,  $u_n \in [a, b]$  for all  $n \geq n_0$  and the second equation in (35) becomes

$$\frac{v_{n+1} - v_n}{\Delta t} = -|v_{n+1}|v_{n+1} \iff v_{n+1}(1 + \Delta t |v_{n+1}|) = v_n. \quad (43)$$

If  $v_n = 0$  for all  $n \geq n_0$ , then  $(u_n)$  is stationary. Otherwise, there exists  $n_1 \geq n_0$  such that  $v_{n_1} > 0$ . Equation (43) shows that  $v_{n_1+1} > 0$  and  $v_{n_1} > v_{n_1+1}$  and by induction,  $v_n > v_{n+1} > 0$  for all  $n \geq n_1$ . We have

$$\frac{1}{v_{n+1}} - \frac{1}{v_n} = \frac{v_n - v_{n+1}}{v_n v_{n+1}} \leq \frac{v_n - v_{n+1}}{v_{n+1}^2} = \Delta t, \quad \forall n \geq n_1.$$

On summing this inequality, we find

$$\sum_{k=n_1}^n \left( \frac{1}{v_{k+1}} - \frac{1}{v_k} \right) \leq (n - n_1 + 1)\Delta t,$$

that is

$$\frac{1}{v_{n+1}} - \frac{1}{v_{n_1}} \leq (n - n_1 + 1)\Delta t.$$

Thus,

$$v_{n+1} \geq \frac{1}{(n - n_1 + 1)\Delta t + 1/v_{n_1}},$$

and so

$$u_{n+1} - u_n \geq \frac{\Delta t}{(n - n_1 + 1)\Delta t + 1/v_{n_1}}, \quad \forall n \geq n_1.$$

Thus, the series  $\sum_{n=n_1}^{\infty} (u_{n+1} - u_n)$  diverges and  $(u_n)$  tends to  $+\infty$ . This is absurd since  $(u_n)$  is bounded.

- If there exists  $n_1 \geq n_0$  such that  $v_{n_1} < 0$ , the second equation in (35) yields

$$\frac{v_{n+1} - v_n}{\Delta t} + |v_{n+1}|v_{n+1} = -f(u_{n+1}) \leq 0$$

that is

$$v_{n+1}(1 + \Delta t|v_{n+1}|) = v_n - \Delta t f(u_{n+1}), \quad \forall n \geq n_1.$$

In particular,  $v_{n_1+1} < 0$  and by induction,  $v_n < 0$  for all  $n \geq n_1$ . Moreover,

$$|v_{n+1}|(1 + \Delta t|v_{n+1}|) \geq |v_n|, \quad \forall n \geq n_1.$$

If  $|v_n| \geq |v_{n+1}|$ , then

$$\frac{1}{|v_{n+1}|} - \frac{1}{|v_n|} = \frac{|v_n| - |v_{n+1}|}{|v_n||v_{n+1}|} \leq \Delta t.$$

If  $|v_n| < |v_{n+1}|$ , then

$$\frac{1}{|v_{n+1}|} - \frac{1}{|v_n|} < 0.$$

In both cases, we have

$$\frac{1}{|v_{n+1}|} - \frac{1}{|v_n|} \leq \Delta t, \quad \forall n \geq n_1.$$

By summing on  $n$ , we find as previously that

$$|v_{n+1}| \geq \frac{1}{(n - n_1 + 1)\Delta t + 1/|v_{n_1}|},$$

Thus,

$$-(u_{n+1} - u_n) \geq \frac{\Delta t}{(n - n_1 + 1)\Delta t + 1/|v_{n_1}|}, \quad \forall n \geq n_1.$$

This shows that  $\sum_{n=n_1}^{\infty} (u_{n+1} - u_n)$  diverges and so  $(u_n)$  converges to  $-\infty$ . Again, we obtain a contradiction since  $(u_n)$  is bounded.

This proves the existence of the subsequence  $(u_{n_k})$ . The existence of the subsequence  $(u_{n_{k'}})$  follows similarly by reversing all signs.  $\square$

**Remark 15** *The oscillations described in Theorem 14 occur even if  $F$  satisfies a Lojasiewicz inequality. Indeed, we may consider the function  $f$  defined by*

$$f(s) = \begin{cases} s - a & \forall s < a, \\ 0 & \forall s \in [a, b], \\ s - b & \forall s > b. \end{cases}$$

*A Taylor expansion near  $a$  and  $b$  shows that  $F$  satisfies the Lojasiewicz inequality with optimal exponent  $\theta = 1/2$ .*

## 5 Numerical simulations

We have performed numerical simulations with the `Scilab` software.

### 5.1 Slow oscillations for a weak damping




Figure 1: An oscillatory sequence  $(u_n)$  (in black) versus time with  $f$  defined by (44). The two horizontal blue lines represent the values of  $a$  and  $b$ .

We first consider a sequence defined by (35) (the case  $\alpha = 1$ ) with  $f$  defined by

$$f(s) = \begin{cases} -(a-s)^2 & \forall s < a, \\ 0 & \forall s \in [a, b], \\ (s-b)^2 & \forall s > b. \end{cases} \quad (44)$$

and  $a = 0$ ,  $b = 0.2$ . The initial datum is  $(u_0, v_0) = (-0.5, 0)$ . The time step is  $\Delta t = 0.01$  and we have computed 100 000 iterations so that the final time is 1000.

Figure 1 shows the oscillatory behaviour described in Theorem 14. This is described as *slow oscillations in the case of a weak damping* in [10].

## 5.2 Convergence to equilibrium for a “strong” damping




Figure 2: A convergent sequence  $(u_n)$  (in black) versus time with  $f$  defined by (44) in the case  $\alpha = 0.6$ . The two horizontal blue lines represent the values of  $a$  and  $b$ .

Next, we consider a sequence defined by (24) with  $\alpha = 0.6$ . The function  $f$  and the initial condition are the same as previously. The time step is  $\Delta t = 0.01$  and we have computed 100 000 iterations. Figure 2 illustrates the convergence of the sequence proved in Theorem 11.




Figure 3: Behaviour of the sequence  $(u^n) = (u_1^n, u_2^n)$  for the explicit scheme (6) with  $f$  defined by (45). The starting point is  $u^0 = (3, 2)$ . The blue line is the ellipse (46).




Figure 4: Energy vs. time (left) and modified energy vs. time (right).

### 5.3 Convergence to equilibrium for a system

We consider the sequence  $(u_n, v_n)$  defined by the explicit scheme (6). The function  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  is

$$F(x, y) = \begin{cases} \frac{1}{2}(x^2 + 2y^2 - 1)^2 & \text{if } x^2 + 2y^2 - 1 \geq 0, \\ 0 & \text{if } x^2 + 2y^2 - 1 < 0. \end{cases} \quad (45)$$

We note that  $F$  is a convex function and its set of critical point is the ellipse

$$\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : x^2 + 2y^2 - 1 \leq 0\}. \quad (46)$$

Moreover,  $F$  satisfies the Lojasiewicz inequality since it is a polynomial outside  $\mathcal{S}$ .

Figure 3 illustrates the convergence of the sequence  $(u^n)$  to a critical point which belongs to the interior of  $\mathcal{S}$ . The initial condition is  $u^0 = (3, 2)$ ,  $v^0 = (0, 0)$ , the time step is  $\Delta t = 0.01$  and we have performed 1000 iterations. The value of  $u^n$  is approximately  $(-0.68, -0.17)$ .

Figure 4 (left) shows the energy  $E^n$  versus time  $t_n = n\Delta t$  over the first 300 iterations. We see especially at the beginning that  $E^n$  is not monotonic. In contrast, the modified energy  $\Phi_\epsilon$  shown in Figure 4 (right) is decreasing. The value chosen for  $\epsilon$  is 0.01.

## Acknowledgment

The first author wishes to thank the department of mathematics of the university of Poitiers where this work has been initiated.

## References

- [1] P.-A. Absil., R. Mahony and B. Andrews, Convergence of the iterates of descent methods for analytic cost functions. *SIAM J. Optim.* **16** (2005), no. 2, 531–547.
- [2] N. Alaa, and M. Pierre, Convergence to equilibrium for discretized gradient-like systems with analytic features. *IMA J. Numer. Anal.* **33** (2013), 1291–1321.
- [3] F. Alvarez, On the minimizing property of a second order dissipative system in Hilbert spaces. *SIAM J. Control Optim.*, **38** (2000), no. 4, 1102–1119.
- [4] H. Attouch, J. Bolte and B. F. Svaiter, Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forward–backward splitting, and regularized Gauss–Seidel methods. *Math. Program.* **137** (2013), 91–129.
- [5] H. Attouch, R. I. Boç and E. R. Csetnek, Fast optimization via inertial dynamics with closed-loop damping. *J. Eur. Math. Soc.* **25** (2023), no. 05, 1985–2056.
- [6] H. Attouch, X. Goudou and P. Redont, The heavy ball with friction method, I. The continuous dynamical system: global exploration of the local minima of a real-valued function by asymptotic analysis of a dissipative dynamical system. *Commun. Contemp. Math.*, **02** (2000), no. 01, 1–34.
- [7] L. Chergui, Convergence of global and bounded solutions of a second order gradient like system with nonlinear dissipation and analytic nonlinearity. *J. Dynam. Differential Equations* **20** (2008), 643–652.

- [8] D. D’Acunto and K. Kurdyka, Explicit bounds for the Łojasiewicz exponent in the gradient inequality for polynomials. *Ann. Pol. Math.* **87**, (2005) 51–61.
- [9] M. Grasselli and M. Pierre, Convergence to equilibrium of solutions of the backward Euler scheme for asymptotically autonomous second-order gradient-like systems. *Commun. Pure Appl. Anal.* **11** (2012), 2393–2416.
- [10] A. Haraux, Asymptotics for some nonlinear O.D.E. of the second order. (English) *Zbl 0603.34032 Nonlinear Anal., Theory Methods Appl.* **10**, 1347-1355 (1986).
- [11] A. Haraux and M.A. Jendoubi. The convergence problem for dissipative autonomous systems - classical methods and recent advances. *SpringerBriefs in Mathematics, Springer, Bcam, 2015*.
- [12] ———, Convergence of solutions of second-order gradient-like systems with analytic nonlinearities. *J. Differential Equations* **144** (1998), 313–320.
- [13] T. Horsin and M.A. Jendoubi, Asymptotics for some discretizations of dynamical systems, application to second order systems with non-local nonlinearities. *Commun. Pure Appl. Anal.* **21**, 999–1025 (2022).
- [14] M.A. Jendoubi and P. Polacik, Non-stabilizing solutions of semilinear hyperbolic and elliptic equations with damping. *Proc. R. Soc. Edinb., Sect. A, Math.* **133**, 1137–1153 (2003).
- [15] S. Łojasiewicz, Une propriété topologique des sous ensembles analytiques réels. *Colloques internationaux du C.N.R.S #117. Les équations aux dérivées partielles* (1963).
- [16] ———, *Ensembles semi-analytiques, I.H.E.S. notes* (1965).