



HAL
open science

PROBABILITY-GRAPHONS: LIMITS OF LARGE DENSE WEIGHTED GRAPHS

Romain Abraham, Jean-François Delmas, Julien Weibel

► **To cite this version:**

Romain Abraham, Jean-François Delmas, Julien Weibel. PROBABILITY-GRAPHONS: LIMITS OF LARGE DENSE WEIGHTED GRAPHS. 2023. hal-04361443

HAL Id: hal-04361443

<https://hal.science/hal-04361443>

Preprint submitted on 22 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROBABILITY-GRAPHONS: LIMITS OF LARGE DENSE WEIGHTED GRAPHS

ROMAIN ABRAHAM, JEAN-FRANÇOIS DELMAS, AND JULIEN WEIBEL

ABSTRACT. We introduce probability-graphons which are probability kernels that generalize graphons to the case of weighted graphs. Probability-graphons appear as the limit objects to study sequences of large weighted graphs whose distribution of subgraph sampling converge. The edge-weights are taken from a general Polish space, which also covers the case of decorated graphs. Here, graphs can be either directed or undirected. Starting from a distance d_m inducing the weak topology on measures, we define a cut distance on probability-graphons, making it a Polish space, and study the properties of this cut distance. In particular, we exhibit a tightness criterion for probability-graphons related to relative compactness in the cut distance. We also prove that under some conditions on the distance d_m , which are satisfied for some well-know distances like the Prohorov distance, and the Fortet-Mourier and Kantorovitch-Rubinstein norms, the topology induced by the cut distance on the space of probability-graphons is independent from the choice of d_m . Eventually, we prove that this topology coincides with the topology induced by the convergence in distribution of the sampled subgraphs.

1. INTRODUCTION

1.1. Motivation and literature review. Networks appear naturally in a wide variety of context, including for example: biological networks [BS02, LL09], epidemics processes [DM10, KMS17], electrical power grids [AAN04] and social networks [AB02, New03]. Most of those problems involve large dense graphs, that is graphs that have a large number of vertices and a number of edges that scales as the square of the number of vertices. Those graphs are too large to be represented entirely in the targeted applications. The idea is then to go from a combinatorial representation given by the graph to an infinite continuum representation.

In the case of unweighted graphs (i.e. graphs without edge-weights), a theory was developed to study the asymptotic behaviour of large dense graphs, with the limit objects being the so-called *graphons*. The properties of graphons were studied in a series of articles started by [LS06, FLS06, BJR10, BCL⁺08, BCL⁺12]. We shall refer to the monograph [Lov12] which exposes in details the theory of graphons developed in this series of articles. Graphons can be used to define models of random graphs with latent vertex-type variables (called *W*-random graphs) generalizing the Erdős-Rényi graph and the stochastic block model (SBM). The space of graphons can be equipped with the so-called *cut distance*, making it a compact space, and whose topology is that of the convergence in distribution for all sampled subgraphs, or equivalently of the convergence for subgraph homomorphism densities.

In recent years, graphons have been used in several application context: non-parametric estimation methods and algorithms for massive networks [BC17], SIS epidemic models [DDZ22], the study of transferability properties for Graph Neural Networks [KV23]. Furthermore, there

Date: December 22, 2023.

2020 Mathematics Subject Classification. 05C80, 60B10.

Key words and phrases. random graphs, stochastic networks, graphons, probability-graphons, dense graph limits, weighted graphs, decorated graphs.

has been recent developments in the study of mean-field systems using graphons: stochastic games and their Nash equilibria [LS22], opinion dynamic on a graphon [AN22], cooperative multi-agent reinforcement learning [HWYZ23], to cite a few.

However, most real-world phenomenon on the above networks involve weighted networks, where each edge in the graph carries additional information such as intensity or frequency of interaction, or transfer capacity.

There exists many models of random weighted graphs. For example configuration models with edges having independent exponential weights have been considered in [BVDHH10, ADL13, AL15], see also [Gar09, HG13] where the distribution of the weight of an edge depends on the types of its end-points. Random geometric graphs with vertices and edges having independent Gaussian weights have been considered in [AMGM18].

Weighted SBMs (sometimes also called labeled SBMs), in which each edge independently receives a random weight whose distribution depends on the community labels of its end-points, have been studied to solve community detection in [LMX13] (see also [XML14] for more general models where vertex-labels come from a compact space), and exact community recovery in [JL15], and to get bounds on the number of misclassified vertices in [YP16, XJL20]. Note that weighted SBMs correspond to a special case of the probability-graphons we study in this article where the space of vertex-labels is finite (they correspond to the stepfunction probability-graphons we define in Section 3).

Concomitantly to our work, in [AD23], the authors studied mean-field equations on large real-weighted graphs modeling interactions with a probability kernel from $[0, 1]^2$ to $\mathcal{M}_1(\mathbb{R})$ the set of probability measures on \mathbb{R} , but they did not study the topological properties of the set of those probability kernels. Recently, in [HV23], the authors studied the limit of the total weight of the minimum spanning tree (MST) for a sequence of random weighted graphs. Following what has been done for the uniform spanning tree in [HNT18, ANS22], one expects the local and scaling limits of the MST to be directly constructed from the limit of the random weighted graphs.

Motivated by those examples, we shall consider *probability-graphons* as possible limits of large weighted graphs; they are defined as maps from $[0, 1]^2$ to the space of probability measures $\mathcal{M}_1(\mathbf{Z})$ on a Polish space \mathbf{Z} . When \mathbf{Z} is compact, this question has been considered in [LS10] and in [Lov12, Section 17.1] using convergence of homomorphism densities of subgraphs decorated with real functions defined on \mathbf{Z} , but the metric properties of the set of probability-graphons \mathcal{W}_1 have only been established when \mathbf{Z} is finite, see [FOSU16]. The work [KLS22] is an extension of [LS10] where $\mathcal{M}_1(\mathbf{Z})$ is replaced by the dual space \mathcal{Z} of a separable Banach space \mathcal{B} . As $\mathcal{M}_1(\mathbf{Z})$ is a subset of the dual of $C_b(\mathbf{Z})$, this approach covers our setting when $C_b(\mathbf{Z})$ is separable, that is, \mathbf{Z} compact (see Section 2 below). The norm introduced on the space of \mathcal{Z} -valued graphons therein implies the convergence of homomorphism densities of \mathcal{B} -decorated sub-graphs, however there is no equivalence *a priori*.

In this paper we study the topological property of the space of probability-graphons \mathcal{W}_1 when \mathbf{Z} is a general Polish space: the space \mathcal{W}_1 is a Polish topological space and we give “natural” cut distances on \mathcal{W}_1 which are complete. One of the main difficulty is that the space of probability measures $\mathcal{M}_1(\mathbf{Z})$ can be endowed with many distances which induce the topology of weak convergence, each of them giving rise to a different cut distance on \mathcal{W}_1 . We prove that the topology induced on \mathcal{W}_1 does not depend on the initial choice of the distance on $\mathcal{M}_1(\mathbf{Z})$, provided this distance satisfies some simple general conditions. However, we stress that not all of these cut distances are complete. We also check that this topology characterizes the convergence in distribution of the sampled subgraphs with random weights on the edges or equivalently the convergence of the homomorphism densities of $C_b(\mathbf{Z})$ -decorated subgraphs. Similarly to the graphon setting, we prove the convergence in distribution of large sampled weighted subgraphs

from a probability-graphon W to itself. We also provide a tightness criterion for studying the convergence of weighted graphs towards probability-graphons.

In conclusion, we believe that the unified framework developed here is easy-to-work-with and will allow to use probability-graphons to study large (random) weighted graphs.

1.2. New contribution. Through the article, *measure* will always be used to denote a positive measure.

1.2.1. Definition of probability-graphons. In this article, we define an analogue of graphons for weighted graphs, which we call *probability-graphons*, and study their properties. To avoid any confusion, in the rest of the article we say *real-valued graphons* instead of graphons. We consider the general case where weighted graphs take their edge-weights in a Polish space \mathbf{Z} (e.g. \mathbb{Z} , \mathbb{R} or \mathbb{R}^d), which thus also covers the case of decorated graphs, multi-graphs (graphs with possibly multiple edges between two vertices) and dynamical graphs (where edge-weights evolve over time).

We define a *probability-graphon* as a probability kernel $W : [0, 1]^2 \rightarrow \mathcal{M}_1(\mathbf{Z})$, where $\mathcal{M}_1(\mathbf{Z})$ is the space of probability measures on \mathbf{Z} . A probability-graphon can be interpreted as follows: for two “vertex type” x and y in $[0, 1]$, the weight z of an edge between two vertices of type x and y is distributed as the probability measure $W(x, y; dz)$. In particular, the special case $\mathbf{Z} = \{0, 1\}$ allows to recover real-valued graphons: as any real-valued graphon $w : [0, 1]^2 \rightarrow [0, 1]$ can be represented as a probability-graphon $W(x, y; \cdot) = w(x, y)\delta_1 + (1 - w(x, y))\delta_0$, where δ_z denote the Dirac mass located at z . Let us mention that it is possible to define the probability-graphons on a more general probability space $(\Omega, \mathcal{A}, \mu)$ than $[0, 1]$ for the vertex-types, see Remark 3.4 for details. In this article, we also define and study the properties of *signed measure-valued kernels* which are bounded (in total mass/total variation norm) measurable functions $W : [0, 1]^2 \mapsto \mathcal{M}_\pm(\mathbf{Z})$ whose values are signed measures, but for brevity we mainly focus on probability-graphons in this introduction.

As probability-graphons are measurable functions, we identify probability-graphons that are equal for almost every $(x, y) \in [0, 1]^2$, and we denote by \mathcal{W}_1 the space of probability-graphons. Moreover, as we consider weighted graphs that are unlabeled (that is vertices are unordered), we need to consider probability-graphons up to “relabeling”: for a measure-preserving map $\varphi : [0, 1] \rightarrow [0, 1]$ (relabeling map for probability-graphons), we define $W^\varphi(x, y; \cdot) = W(\varphi(x), \varphi(y); \cdot)$; we say that two probability-graphons are weakly isomorphic if there exists measure-preserving maps $\varphi, \psi : [0, 1] \rightarrow [0, 1]$ such that $U^\varphi = W^\psi$ for a.e. $(x, y) \in [0, 1]^2$. We denote by $\widetilde{\mathcal{W}}_1$ the space of probability-graphons where we identify probability-graphons that are weakly isomorphic.

We can always assume that weighted graphs are complete graphs by adding all missing edges and giving them a weight/decoration ∂ which is a cemetery point added to \mathbf{Z} . Any weighted graph G can be represented as a probability-graphon W_G in the following way: denote by n the number of vertices of G and divide the unit interval $[0, 1]$ into n intervals I_1, \dots, I_n of equal lengths, then W_G is defined for $(x, y) \in I_i \times I_j$ as $W_G(x, y; \cdot) = \delta_{M(i, j)}$, where $M(i, j)$ is the weight on the edge (i, j) in G . Note that weighted graphs can be either directed or undirected, in the case of undirected weighted graphs their limit objects are symmetric probability-graphons, that is probability-graphons W such that $W(x, y; \cdot) = W(y, x; \cdot)$.

1.2.2. The cut distance for probability-graphons and its properties. While there is a usual distance on the field of reals \mathbb{R} , this is not the case for probability measures, measures or signed measures endowed with the weak topology. Some commonly used distances include the Prohorov distance $d_{\mathcal{P}}$ which can be defined on measures, and the Kantorovitch-Rubinstein norm $\|\cdot\|_{\text{KR}}$ (sometimes also called the bounded Lipschitz norm) and the Fortet-Mourier norm $\|\cdot\|_{\text{FM}}$ defined on signed measures but metrizing the weak topology on measures. (Note that in general the weak topology

is not metrizable on signed measures, see Section 2 below.) We also use a norm $\|\cdot\|_{\mathcal{F}}$ based on a convergence determining sequence $\mathcal{F} \subset C_b(\mathbf{Z})$. See Section 3.8 for definition of those distances. To define an analogue of the cut norm for probability-graphons, we first need to choose a distance d_m that metrizes the weak topology on the space of sub-probability measures $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (i.e. measures with total mass at most 1); we then define the *cut distance* $d_{\square, m}$ for probability-graphons as:

$$d_{\square, m}(U, W) = \sup_{S, T \subset [0, 1]} d_m\left(U(S \times T; \cdot), W(S \times T; \cdot)\right),$$

where the supremum is taken over all measurable subsets S and T of $[0, 1]$, and where $W(S \times T; \cdot) = \int_{S \times T} W(x, y; \cdot) dx dy$ is a sub-probability measure and similarly for U . Moreover, if the distance d_m is derived from a norm N_m defined on the space of signed measures $\mathcal{M}_{\pm}(\mathbf{Z})$, then the cut distance $d_{\square, m}$ derives from the *cut norm* $N_{\square, m}$ defined on signed measure-valued kernels:

$$N_{\square, m}(W) = \sup_{S, T \subset [0, 1]} N_m\left(W(S \times T; \cdot)\right).$$

We then define the unlabeled *cut distance* $\delta_{\square, m}$ on the space of unlabeled probability-graphons $\widetilde{\mathcal{W}}_1$ as:

$$\delta_{\square, m}(U, W) = \inf_{\varphi} d_{\square, m}(U, W^{\varphi}) = \min_{\varphi, \psi} d_{\square, m}(U^{\varphi}, W^{\psi}),$$

where the infimum is taken over all measure-preserving maps φ and ψ , see Proposition 3.18 for alternative expressions of $\delta_{\square, m}$ (including proof that the minimum exist for the second expression) and see Theorem 3.17 that states that $\delta_{\square, m}$ is indeed a distance on $\widetilde{\mathcal{W}}_1$. In Proposition 4.13, we prove an equivalent of the weak regularity lemma for probability-graphons.

An interesting fact is that under some conditions on d_m , the topology induced by the associated cut distance $\delta_{\square, m}$ does not depend on the particular choice of d_m . The following proposition is a particular case of Theorem 5.5 together with Corollary 4.14.

Proposition 1.1. *The cut distances $\delta_{\square, \mathcal{P}}$, $\delta_{\square, KR}$, $\delta_{\square, FM}$ and $\delta_{\square, \mathcal{F}}$ induce the same topology on the space of probability-graphons $\widetilde{\mathcal{W}}_1$.*

Recall that \mathbf{Z} is a Polish space. We now state that $\widetilde{\mathcal{W}}_1$ is also Polish for the distance $\delta_{\square, \mathcal{P}}$ (but not for $\delta_{\square, \mathcal{F}}$!), and we refer to Theorem 5.10 for other distances.

Theorem 1.2. *The space of probability-graphons $(\widetilde{\mathcal{W}}_1, \delta_{\square, \mathcal{P}})$ is a Polish metric space.*

We prove an analogue of Prohorov's theorem with a tightness criterion for probability-graphons. We say that a subset of probability-graphons $\mathcal{K} \subset \widetilde{\mathcal{W}}_1$ is *tight* if the set of probability measures $\{M_W : W \in \mathcal{K}\}$ is tight (in the sense of probability measures), where $M_W(\cdot) = W([0, 1]^2; \cdot)$. The next result is consequence of Theorem 5.1 as well as Corollary 4.14.

Theorem 1.3 (Compactness property). *Consider the topology on $\widetilde{\mathcal{W}}_1$ from Proposition 1.1.*

- (i) *If a sequence of elements of $\widetilde{\mathcal{W}}_1$ is tight, then it has a converging subsequence.*
- (ii) *If \mathbf{Z} is compact, then the space $\widetilde{\mathcal{W}}_1$ is compact.*

1.2.3. *Sampling from probability-graphons and its link with the cut distance.* Finally, we link the topology of the cut distance $\delta_{\square, m}$ with subgraph sampling. The probability-graphons allow to define models of random weighted graphs (the W -random graph model) which generalize weighted SBM random graphs, and which plays the role of sampled subgraphs for probability-graphons. The W -random graph (or sampled subgraph of size k) $\mathbb{G}(k, W)$ has two parameters, a number of vertices k and a probability-graphon W for edge-weights, and is defined as follows: first let X_1, \dots, X_k be k independent random “vertex-types” uniformly distributed over $[0, 1]$;

then given X_1, \dots, X_k , each edge receive a weight independently, where the weight of the edge (i, j) is distributed as $W(X_i, X_j; \cdot)$.

We also provide the a.s. convergence of sampled subgraphs for the topology from Proposition 1.1, see Theorem 6.13 together with Corollary 5.6.

Theorem 1.4. *Let W be a probability-graphon. Then, a.s. the sequence of sampled subgraphs $(\mathbb{G}(k, W))_{k \in \mathbb{N}^*}$ converges to W for the topology from Proposition 1.1.*

To prove this theorem, we adapt the proof scheme of [Lov12, Sections 10.5 and 10.6] relying on the first and second sampling lemmas for real-valued graphons. The proof is done using the cut distance $\delta_{\square, \mathcal{F}}$ because of the good approximations properties of $\|\cdot\|_{\mathcal{F}}$.

In the case of unweighted graphs, the homomorphism numbers $\text{hom}(F, G)$ count the number of occurrence of a graph F (often called a *motif* or a *graphlet*) as an induced subgraph of G , and their normalized counterparts, the homomorphism densities $t(F, G)$ allow to characterize a graph (up to relabeling and twin-vertices expansion), and also characterize the topology on real-valued graphons. In the case of weighted graphs and probability-graphons, we need to replace absence/presence of edges (which is 0-1 valued) by test functions from $C_b(\mathbf{Z})$ decorating the edges. Hence, we define the *homomorphism density* of a \mathcal{G} -graph F^g which is a finite graph $F = (V, E)$ whose edges are decorated with a family of functions $g = (g_e)_{e \in E}$ from a subset $\mathcal{G} \subset C_b(\mathbf{Z})$ (in practice, we only consider the cases $\mathcal{G} = C_b(\mathbf{Z})$ or $\mathcal{G} = \mathcal{F} \subset C_b(\mathbf{Z})$ a convergence determining sequence), in a probability-graphon W as:

$$t(F^g, W) = M_W^F(g) := \int_{[0,1]^V} \prod_{(i,j) \in E} W(x_i, x_j; g_{i,j}) \prod_{i \in V} dx_i,$$

where $W(x, y; f) = \int_{\mathbf{Z}} f(z) W(x, y; dz)$. Moreover, M_W^F defines a measure on \mathbf{Z}^E (which we still denote by M_W^F) which is defined by $M_W^F(\otimes_{e \in E} g_e) = M_W^F(g)$ for $g = (g_e)_{e \in E}$. Note that when F is the complete graph with k vertices, M_W^F is the joint measure of all the edge-weights of the random graph $\mathbb{G}(k, W)$, and thus characterizes the random graph $\mathbb{G}(k, W)$.

In the counting Lemma 7.5 and the weak counting Lemma 7.7, we prove that the cut norm $\|\cdot\|_{\square, \mathcal{F}}$ allows to control the homomorphism densities. Conversely, in the inverse counting Lemma 7.8, we prove that the cut norm $\|\cdot\|_{\square, \mathcal{F}}$ can be controlled by the homomorphism densities. In particular, the topology of the cut distance turns out to be exactly the topology of convergence in distribution for sampled subgraphs of any given size; the next result is a direct consequence of Theorem 7.11

Theorem 1.5 (Characterization of the topology). *Let $(W_n)_{n \in \mathbb{N}}$ and W be unlabeled probability-graphons from $\tilde{\mathcal{W}}_1$. The following properties are equivalent:*

- (i) $(W_n)_{n \in \mathbb{N}}$ converges to W for the topology from Proposition 1.1.
- (ii) $\lim_{n \rightarrow \infty} t(F^g, W_n) = t(F^g, W)$ for all $C_b(\mathbf{Z})$ -graph F^g .
- (iii) $\lim_{n \rightarrow \infty} t(F^g, W_n) = t(F^g, W)$ for all \mathcal{F} -graph F^g , for some convergence determining sequence \mathcal{F} .
- (iv) For all $k \geq 2$, the sequence of sampled subgraphs $(\mathbb{G}(k, W_n))_{n \in \mathbb{N}}$ converges in distribution to $\mathbb{G}(k, W)$.

Now, we can turn back to the initial problem of finding a limit object for a convergent sequence of weighted graphs $(G_n)_{n \in \mathbb{N}}$; here convergent means that for all $k \geq 2$, the sequence $(\mathbb{G}(k, G_n) = \mathbb{G}(k, W_{G_n}))_{n \in \mathbb{N}}$ of sampled subgraphs of size k (defined above) converges in distribution (to some limit random graph). Note that the tightness criterion for a sequence of probability-graphons $(W_n)_{n \in \mathbb{N}}$ can be equivalently rephrased as tightness of the sequence $(\mathbb{G}(2, W_n))_{n \in \mathbb{N}}$ of sampled subgraphs of size 2. Hence, the convergence in distribution of the sequence $(\mathbb{G}(2, G_n))_{n \in \mathbb{N}}$ implies its tightness, and thus the tightness of the sequence of probability-graphons $(W_{G_n})_{n \in \mathbb{N}}$. Then,

Theorem 1.3 guarantees the existence of a probability-graphon W which is a sub-sequential limit of the sequence $(W_{G_n})_{n \in \mathbb{N}}$ in the cut distance $\delta_{\square, \mathcal{F}}$, and then Theorem 1.5 guarantees that for all $k \geq 2$, the sequence $(\mathbb{G}(k, G_n))_{n \in \mathbb{N}}$ converges in distribution to $\mathbb{G}(k, W)$.

As a consequence, probability-graphons are precisely the limit objects for sequences of weighted graphs $(G_n)_{n \in \mathbb{N}}$ (and also for random weighted graphs) whose number of vertices goes to infinity (otherwise the limit would simply be a weighted graph) and such that for each size $k \geq 2$, the sequence of sampled subgraphs $(\mathbb{G}(k, G_n))_{n \in \mathbb{N}}$ converges in distribution.

Remark 1.6 (Extension to vertex-weights). The framework we have developed for probability-graphons could easily be extended to add weights on the vertices, or equivalently to allow for self-loops (i.e. edges linking a vertex to itself). In this case, weighted graphs and probability-graphons have a two-variable kernel (probability-graphon) W^e for edge-weights as before, and a one-variable kernel $W^v : [0, 1] \rightarrow \mathcal{M}_1(\mathbf{Z})$ for vertex-weights. Note that this implies, as expected, that the same measure-preserving map $\varphi : [0, 1] \rightarrow [0, 1]$ must be used for both kernels W^v and W^e when relabeling.

1.3. Organization of the paper. The rest of the paper is organized as follows. In Section 2, we define some notations used throughout the paper, and remind some properties of the weak topology on the space of signed measures. In Section 3, we define probability-graphons and signed-measure valued kernels, we then define the cut distance and the cut norm and study their properties, and we also give some example of distances with the Prohorov distance $d_{\mathcal{P}}$, the Kanrorovitch-Rubinstein and Fortet-Mourier norms $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$, and the norm $\|\cdot\|_{\mathcal{F}}$ based on a convergence determining sequence. In Section 4, we define the steppings of a probability-graphon (which are stepfunction approximations corresponding to conditional expectations on $[0, 1]^2$), we define the tightness criterion for probability-graphons, and we prove the weak regularity property of the cut distance. In Section 5, we prove the theorem linking the tightness criterion with relative compactness for the cut distance, we prove that under some conditions the topology of the cut distance does not depend on the choice of the initial distance d_m , and we prove that the space of probability-graphons with the cut distance is a Polish space. In Section 6, we define the subgraph $\mathbb{G}(k, W)$ sampled from a probability-graphon W , we then prove approximation bound in the cut norm $\|\cdot\|_{\square, \mathcal{F}}$ between probability-graphons and their sampled subgraphs. In Section 7, we prove the counting lemmas linking the cut distance with the homomorphism densities, and prove that the topology induced by the cut distance coincides with the topology of convergence in distribution for all the sampled subgraphs.

CONTENTS

1. Introduction	1
1.1. Motivation and literature review	1
1.2. New contribution	3
1.3. Organization of the paper	6
2. Notations and topology on the space of signed measures	7
3. Measured-valued graphons and the cut distance	11
3.1. Definition of measure-valued graphons	11
3.2. The cut distance	13
3.3. Graphon relabeling, invariance and smoothness properties	14
3.4. The unlabeled cut distance	16
3.5. Weak isomorphism	16
3.6. The cut norm for stepfunctions	17
3.7. The supremum in S and T in the cut distance $d_{\square, m}$	18
3.8. Examples of distance d_m	19

4.	Tightness and weak regularity	21
4.1.	Approximation by stepfunctions	22
4.2.	Tightness	23
4.3.	Weak regularity	24
4.4.	A stronger weak regularity lemma for $d_{\square, \mathcal{F}}$	27
5.	Compactness and completeness of \mathcal{W}_1	29
5.1.	Tightness criterion and compactness	29
5.2.	Equivalence of topologies induced by $\delta_{\square, m}$	30
5.3.	Completeness	32
6.	Sampling from probability-graphons	35
6.1.	$\mathcal{M}_1(\mathbf{Z})$ -Graphs and weighted graphs	35
6.2.	W -random graphs	36
6.3.	Estimation of the distance by sampling	36
6.4.	The distance between a probability-graphon and its sample	39
7.	The Counting Lemmas and the topology of probability-graphons	40
7.1.	The homomorphism densities	40
7.2.	The Counting Lemma	41
7.3.	The Inverse Counting Lemma	42
7.4.	Subgraph sampling and the topology of probability-graphons	44
8.	Proofs of Theorem 5.1 and Theorem 5.5	45
	Acknowledgement	55
	References	56

2. NOTATIONS AND TOPOLOGY ON THE SPACE OF SIGNED MEASURES

Through the article, *measure* will always be used to denote a positive measure.

Let $\mathbb{N} = \mathbb{Z}_+$ be the set of non-negative integers, $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$ the set of positive integers, and, for $n \in \mathbb{N}^*$, we define the integer set $[n] = \{1, \dots, n\}$. For $k \in \mathbb{N}^*$, the set $[0, 1]^k$ is endowed with the Borel σ -field and the Lebesgue measure λ_k ; and we write λ for λ_k when the context is clear. The supremum of a real-valued function f defined on $[0, 1]^k$ is denoted by $\|f\|_\infty = \sup_{x \in [0, 1]^k} f(x)$.

Let d be a distance on a topological space (X, \mathcal{O}) .

- (i) The distance d is continuous w.r.t. the topology \mathcal{O} if the identity map from (X, \mathcal{O}) to (X, d) is continuous.
- (ii) The distance d is sequentially continuous w.r.t. the topology \mathcal{O} if for any sequence $(x_n)_{n \in \mathbb{N}}$ in X which converges to some limit x for the topology \mathcal{O} , we also have that $\lim_{n \rightarrow \infty} d(x_n, x) = 0$.

Let d and d' be two distances on a space X . We say that d' is continuous (resp. uniformly continuous) w.r.t. d if the identity map from (X, d) to (X, d') is continuous (resp. uniformly continuous).

Remark 2.1. If the topology \mathcal{O} is metrizable (i.e. can be generated by a distance on the space X), then the topology on X induced by the distance d is equivalent to \mathcal{O} if and only if for every sequence with values in X , convergence for d is equivalent to convergence for \mathcal{O} (see [Eng89, Theorem 4.1.2]). Moreover, when the topology is metrizable, then topological notions and their sequential counterparts coincides (e.g. compact and sequentially compact sets, closed and sequentially closed sets, see [Eng89, Proposition 4.1.1 and Theorem 4.1.17]).

Remark 2.2. For a function, continuity always implies sequential continuity; and the converse is also true when the topology is metrizable.

A map $\varphi : \Omega_1 \rightarrow \Omega_2$ between two probability spaces $(\Omega_i, \mathcal{A}_i, \pi_i)$, $i = 1, 2$, is measure-preserving if it is measurable and if for every $A \in \mathcal{A}_2$, $\pi_2(A) = \pi_1(\varphi^{-1}(A))$. In this case, for every measurable non-negative function $f : \Omega_2 \rightarrow \mathbb{R}$, we have:

$$(1) \quad \int_{\Omega_1} f(\varphi(x)) \pi_1(dx) = \int_{\Omega_2} f(x) \pi_2(dx).$$

We denote by $S_{[0,1]}$ the set of bijective measure-preserving maps from $[0, 1]$ with the Lebesgue measure to itself, and by $\tilde{S}_{[0,1]}$ the set of measure-preserving maps from $[0, 1]$ with the Lebesgue measure to itself.

Let $(\mathbf{Z}, \mathcal{O}_{\mathbf{Z}})$ be some (non-empty) Polish space, and let $\mathcal{B}(\mathbf{Z})$ be the Borel σ -field on \mathbf{Z} generated by the topology $\mathcal{O}_{\mathbf{Z}}$. We denote by $C_b(\mathbf{Z})$ the space of real-valued continuous bounded functions on $(\mathbf{Z}, \mathcal{O}_{\mathbf{Z}})$. We denote by $\mathcal{M}_{\pm}(\mathbf{Z})$ the space of finite signed measures on $(\mathbf{Z}, \mathcal{B}(\mathbf{Z}))$; $\mathcal{M}_+(\mathbf{Z})$ the subspace of measures; $\mathcal{M}_{\leq 1}(\mathbf{Z})$ the subspace of measures with total mass at most 1; and $\mathcal{M}_1(\mathbf{Z})$ the subspace of probability measures. We have:

$$\mathcal{M}_1(\mathbf{Z}) \subset \mathcal{M}_{\leq 1}(\mathbf{Z}) \subset \mathcal{M}_+(\mathbf{Z}) \subset \mathcal{M}_{\pm}(\mathbf{Z}).$$

For a signed measure $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$, we remind the definition of the Hahn-Jordan decomposition $\mu = \mu^+ - \mu^-$ where $\mu^+, \mu^- \in \mathcal{M}_+(\mathbf{Z})$ are mutually singular measures (that is $\mu^+(A) = 0$ and $\mu^-(A^c) = 0$ for some measurable set A), as well as the total variation measure of μ which is defined as $|\mu| = \mu^+ + \mu^- \in \mathcal{M}_+(\mathbf{Z})$. Note that for a measure $\mu \in \mathcal{M}_+(\mathbf{Z})$, we simply have $|\mu| = \mu$. For a signed-measure $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$ and a real-valued measurable function f defined on \mathbf{Z} , we write $\mu(f) = \langle \mu, f \rangle = \int f d\mu = \int_{\mathbf{Z}} f(x) \mu(dx)$ the integral of f w.r.t. μ whenever it is well defined. For a signed measure $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$, we denote by $\|\mu\|_{\infty} = \mu^+(\mathbf{Z}) + \mu^-(\mathbf{Z})$ its total mass, which is also equal to the supremum of $\mu(f)$ over all measurable functions f with values in $[-1, 1]$.

We endow $\mathcal{M}_{\pm}(\mathbf{Z})$ with the topology of weak convergence, that is the smallest topology for which the maps $\mu \mapsto \mu(f)$ are continuous for all $f \in C_b(\mathbf{Z})$. In particular, a sequence of signed measures $(\mu_n)_{n \in \mathbb{N}}$ weakly converges to some $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$ if and only if, for every function $f \in C_b(\mathbf{Z})$, we have $\lim_{n \rightarrow +\infty} \mu_n(f) = \mu(f)$. Let us recall that $\mathcal{M}_+(\mathbf{Z})$ and $\mathcal{M}_1(\mathbf{Z})$ endowed with the topology of weak convergence are Polish spaces.

Remark 2.3 (The weak topology on $\mathcal{M}_{\pm}(\mathbf{Z})$). The topology of weak convergence on the set of signed measures $\mathcal{M}_{\pm}(\mathbf{Z})$ is equivalent to the weak-* topology on $\mathcal{M}_{\pm}(\mathbf{Z})$ seen as a subspace of the topological dual of $C_b(\mathbf{Z})$ (see the paragraph after Definition 3.1.1 in [Bog18]). As usual in probability theory, this topology will be simply called the weak topology (this is also consistent with [Bog18]).

We recall that a sequence of $[0, 1]$ -valued functions $\mathcal{F} = (f_k)_{k \in \mathbb{N}}$ in $C_b(\mathbf{Z})$, with $f_0 = \mathbf{1}$ the constant function equal to one, is:

- (i) **Separating** if for every measures μ, ν from $\mathcal{M}_{\pm}(\mathbf{Z})$ (or equivalently just from $\mathcal{M}_+(\mathbf{Z})$) such that for every $k \in \mathbb{N}$, $\mu(f_k) = \nu(f_k)$, then $\mu = \nu$.
- (ii) **Convergence determining** if for every $(\mu_n)_{n \in \mathbb{N}}$ and μ measures from $\mathcal{M}_+(\mathbf{Z})$ such that we have $\lim_{n \rightarrow +\infty} \mu_n(f_k) = \mu(f_k)$ for all $k \in \mathbb{N}$, then $(\mu_n)_{n \in \mathbb{N}}$ weakly converges to μ .

Notice that a convergence determining sequence is also separating. A sequence of functions is separating if and only if it separates the points of \mathbf{Z} (see [EK09, Theorem 3.4.5]). There always exists a convergence determining sequence on Polish spaces, see [Bog18, Corollary 2.2.6]

or the proof of Proposition 3.4.4 in [EK09] (which are stated for probability measures but can be extended to finite positive measures as we required that $\mathbf{1}$ belongs to \mathcal{F}). Note that there does not exist a convergence determining sequence for $\mathcal{M}_\pm(\mathbf{Z})$ as the weak topology is not metrizable on $\mathcal{M}_\pm(\mathbf{Z})$ (see Remark 2.6 below).

Remark 2.4 (The Borel σ -field on $\mathcal{M}_\pm(\mathbf{Z})$). By [Bog18, Corollary 5.1.9], the Borel σ -field on $\mathcal{M}_\pm(\mathbf{Z})$, associated with the weak topology, is countably generated and can be generated by either:

- the family of maps $\mu \mapsto \mu(f_n)$ where the sequence $(f_n)_{n \in \mathbb{N}}$ of functions from $C_b(\mathbf{Z})$ is separating;
- the family of maps $\mu \mapsto \mu(B)$ where $B \in \mathcal{A}$ and the subset $\mathcal{A} \subset \mathcal{B}(\mathbf{Z})$ is countable and generates the whole σ -field $\mathcal{B}(\mathbf{Z})$ (such subset \mathcal{A} always exists, see [Bog07b, Corollary 6.7.5]).

Note that the Borel σ -field of a Polish space is generated by any family of Borel functions that separates points (see [Bog07b, Theorem 6.8.9]).

Furthermore, the maps $\mu \mapsto \mu^+$ and $\mu \mapsto \mu^-$ (and thus also $\mu \mapsto |\mu|$) are measurable (see [DF64, Theorem 2.8] and Remark 2.4). As a consequence, the map $\mu \mapsto \|\mu\|_\infty$ is also measurable (in fact it is even lower semicontinuous by [Bog18, Theorem 2.7.4]). Note that $\mathcal{M}_1(\mathbf{Z})$ and $\mathcal{M}_+(\mathbf{Z})$ are closed, and thus measurable, subsets of $\mathcal{M}_\pm(\mathbf{Z})$.

We define the following two important properties for subsets of signed measures, which are related to relative compactness (see Lemma 2.8 below).

Definition 2.5. Let $\mathcal{M} \subset \mathcal{M}_\pm(\mathbf{Z})$ be a subset of signed measures.

(i) The set \mathcal{M} is bounded (in total variation) if:

$$\sup_{\mu \in \mathcal{M}} \|\mu\|_\infty < +\infty.$$

(ii) The set \mathcal{M} is tight if for all $\varepsilon > 0$, there exists a compact set $K \subset \mathbf{Z}$ such that:

$$\sup_{\mu \in \mathcal{M}} |\mu|(K^c) \leq \varepsilon.$$

Remark 2.6 (On the compact sets and metrizability of the weak topology). Recall that \mathbf{Z} is a Polish space. We stress that the weak topology on signed measures is not metrizable unless it coincides with the strong topology (see [Var58, Theorem 4.1]), which happens only when the initial space \mathbf{Z} is finite (see [Bog18, Proposition 3.1.8]).

Moreover, the closed norm ball $\{\mu \in \mathcal{M}_\pm(\mathbf{Z}) : \|\mu\|_\infty \leq 1\}$ of $\mathcal{M}_\pm(\mathbf{Z})$ is metrizable if and only if \mathbf{Z} is compact (see [Bog18, Proposition 3.1.8 and Theorem 3.1.9]).

Let $\mathcal{M} \subset \mathcal{M}_\pm(\mathbf{Z})$. The following properties are equivalent (see [Bog18, Theorems 2.3.4 and 3.1.9]):

- (i) \mathcal{M} is weakly compact (i.e. \mathcal{M} is compact for the weak topology);
- (ii) \mathcal{M} is sequentially weakly compact (that is every sequence $(\mu_n)_{n \in \mathbb{N}}$ in \mathcal{M} has a subsequence that converges to some limit $\mu \in \mathcal{M}$);
- (iii) \mathcal{M} is compact for the sequential weak topology (for which sets are closed if and only if they are closed under weak convergence).

Moreover, when any of those is true, \mathcal{M} is tight, bounded, and metrizable in the weak topology. Furthermore, the Kantorovitch-Rubinstein and Fortet-Mouriet norms $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$ (defined in Section 3.8.2) can be used to generate the weak topology on a weakly compact set (see [Bog18, Remark 3.2.5]).

Nevertheless, the weak topology on the unit sphere $\{\mu \in \mathcal{M}_\pm(\mathbf{Z}) : \|\mu\|_\infty = 1\}$ of $\mathcal{M}_\pm(\mathbf{Z})$ is always metrizable with a complete metric, making the unit sphere a Polish space, however, the

Kantorovitch-Rubinstein and Fortet-Mouriet norms $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$ do not provide a complete metrization in this case (see [Bog18, Theorem 3.2.8]).

Remark 2.7 (On the compactness of $\mathcal{M}_1(\mathbf{Z})$). Let \mathcal{M} be either $\mathcal{M}_1(\mathbf{Z})$, $\mathcal{M}_{\leq 1}(\mathbf{Z})$ or the closed norm ball $\{\mu \in \mathcal{M}_{\pm}(\mathbf{Z}) : \|\mu\|_{\infty} \leq 1\}$ of $\mathcal{M}_{\pm}(\mathbf{Z})$. Then, \mathcal{M} is weakly compact if and only if \mathbf{Z} is compact.

We give a short proof of this statement. As $\mathcal{M}_1(\mathbf{Z})$ is closed in $\mathcal{M}_{\pm}(\mathbf{Z})$ for the weak topology, if \mathcal{M} is weakly compact, then $\mathcal{M}_1(\mathbf{Z})$ is also weakly compact, and thus \mathbf{Z} is compact by [Var58, Theorem 3.4]. Conversely, if \mathbf{Z} is compact, then by [Bog18, Theorem 1.3.3], we know that $\mathcal{M}_{\pm}(\mathbf{Z})$ (endowed with the weak topology) is the topological dual space of $C_b(\mathbf{Z})$ (endowed with the uniform convergence topology), thus using Banach-Alaoglu theorem (see [Bog18, Theorem 1.3.6]), we get that the closed unit norm-ball of $\mathcal{M}_{\pm}(\mathbf{Z})$, and thus \mathcal{M} , are compact for the weak topology.

We recall the following result, which is an equivalent of Prohorov's theorem for signed measures.

Lemma 2.8 (Prohorov's theorem for signed measures, [Bog18, Theorems 2.3.4 and 3.1.9]). *Let \mathbf{Z} be a Polish space, and let $\mathcal{M} \subset \mathcal{M}_{\pm}(\mathbf{Z})$ be a subset of signed measures on \mathbf{Z} . Then the following conditions are equivalent:*

- (i) \mathcal{M} is relatively sequentially compact, that is every sequence $(\mu_n)_{n \in \mathbb{N}}$ in \mathcal{M} contains a subsequence which weakly converges in $\mathcal{M}_{\pm}(\mathbf{Z})$.
- (ii) \mathcal{M} is relatively compact for the weak topology, that is the closure of \mathcal{M} is compact for the weak topology.
- (iii) The family \mathcal{M} is tight and bounded.

Remark 2.9 (On the weak sequential topology). When the space \mathbf{Z} is infinite, the weak topology does not coincide with the weak sequential topology on $\mathcal{M}_{\pm}(\mathbf{Z})$ (but recall from Remark 2.6 that their compact sets are the same). Recall that if the space \mathbf{Z} is compact, then the unit norm ball of $\mathcal{M}_{\pm}(\mathbf{Z})$ is metrizable, and thus the weak topology and the weak sequential topology coincide on it. However, if the space \mathbf{Z} is non-compact, then the weak topology and the weak sequential topology do not coincide on the unit norm ball of $\mathcal{M}_{\pm}(\mathbf{Z})$.

We give a short proof of those statements according to \mathbf{Z} being compact or not.

- (i) Remind that when \mathbf{Z} is an infinite compact space (for instance $\mathbf{Z} = [0, 1]$), the Banach space $C_b(\mathbf{Z})$ is infinite-dimensional and separable (using Stone-Weierstrass theorem), and its topological dual is $(C_b(\mathbf{Z}))^* = \mathcal{M}_{\pm}(\mathbf{Z})$ (see [Bog18, Theorem 1.3.3]). Thus, using [HS96, Theorem 2.5], we get the existence of a countable subset which is weak sequentially closed yet weak dense in $\mathcal{M}_{\pm}(\mathbf{Z})$. In particular, the weak sequential topology and the weak topology do not coincide on $\mathcal{M}_{\pm}(\mathbf{Z})$.
- (ii) Assume that the space \mathbf{Z} is non-compact. Thus, \mathbf{Z} contains a countable closed subset F whose points are at mutual distances uniformly bounded away from zero. By [Bog18, Remark 3.1.7], the weak topology on $\mathcal{M}_{\pm}(F)$ for a closed subset F coincides with the trace of the weak topology on the whole space. By [Bog18, Section 3.1, p. 102], $\mathcal{M}_{\pm}(F)$ is homeomorphic to ℓ^1 both endowed with their weak topology, weak convergence on ℓ^1 is equivalent to norm convergence, and the weak topology on ℓ^1 is not sequential, even on the unit norm ball. Hence, the weak topology on $\mathcal{M}_{\pm}(\mathbf{Z})$ is not sequential, even on the unit norm ball.

We define the notion of a quasi-convex distance, which generalizes the convexity of a norm.

Definition 2.10 (Quasi-convex distance). *Let (X, d) be a metric space which is a convex subset of a vector space. The distance d is quasi-convex if for all $x_1, x_2, y_1, y_2 \in X$ and all $\alpha \in [0, 1]$, we have:*

$$d(\alpha x_1 + (1 - \alpha)x_2, \alpha y_1 + (1 - \alpha)y_2) \leq \max(d(x_1, y_1), d(x_2, y_2)).$$

In particular, any distance (on a convex subset of a vector space) which derive from a norm is quasi-convex.

Lemma 2.11. *Let d_m be distance on $\mathcal{M}_\epsilon(\mathbf{Z})$ with $\epsilon \in \{+, \pm\}$ which is quasi-convex and sequentially continuous with respect to the weak topology. Then, d_m is uniformly continuous with respect to $\|\cdot\|_\infty$ on $\mathcal{M}_\epsilon(\mathbf{Z})$.*

Proof. We shall simply consider the case $\mathcal{M} = \mathcal{M}_+(\mathbf{Z})$, the other case being simpler. We first check that for all $\mu \in \mathcal{M}$ and $\varepsilon > 0$, there exists $\eta > 0$ such that for all $\nu \in \mathcal{M}$, we have that $\|\mu - \nu\|_\infty < \eta$ implies $d_m(\mu, \nu) < \varepsilon$. As d_m is sequentially continuous w.r.t. the weak topology, it is also (sequentially) continuous w.r.t. the strong topology. Let $\mu \in \mathcal{M}$ and $\varepsilon > 0$. Then, the set $\{\nu \in \mathcal{M} : d_m(\mu, \nu) < \varepsilon\}$ is an open set of \mathcal{M} containing μ both for d_m and for the strong topology. Thus, it contains a neighborhood of μ for the strong topology $\{\nu \in \mathcal{M} : \|\mu - \nu\|_\infty < \eta\}$ for $\eta > 0$ small enough. This proves the claim.

As d_m is quasi-convex and \mathcal{M} is a cone, for $\mu, \nu \in \mathcal{M}$ we have:

$$d_m(\mu, \mu + \nu) = d_m\left(\frac{1}{2} \cdot (2\mu + 0), \frac{1}{2} \cdot (2\mu + 2\nu)\right) \leq \max(d_m(2\mu, 2\mu), d_m(0, 2\nu)) = d_m(0, 2\nu).$$

Let $\varepsilon > 0$ be fixed. We choose $\eta \in (0, 1)$ such that $\|\nu\|_\infty < \eta$, with $\nu \in \mathcal{M}$, implies $d_m(0, \nu) < \varepsilon$. Let $\mu, \nu \in \mathcal{M}$ be such that $\|\mu - \nu\|_\infty < \eta/2$. Let $\lambda' = \mu + \nu$ and f (resp. g) the density of μ (resp. ν) with respect to λ' . We set $\pi = \min(f, g)\lambda'$, $\mu' = (f - g)_+\lambda'$ and $\nu' = (f - g)_-\lambda'$ so that $\pi, \mu', \nu' \in \mathcal{M}$, $\mu = \pi + \mu'$ and $\nu = \pi + \nu'$. Since $\mu' - \nu' = \mu - \nu$ and μ' and ν' are mutually singular, we deduce that $\|\mu'\|_\infty + \|\nu'\|_\infty < \eta/2$. We get:

$$\begin{aligned} d_m(\mu, \nu) &= d_m(\pi + \mu', \pi + \nu') \leq d_m(\pi, \pi + \mu') + d_m(\pi, \pi + \nu') \\ &\leq d_m(0, 2\mu') + d_m(0, 2\nu') \\ &\leq 2\varepsilon. \end{aligned}$$

Hence, the distance d_m is uniformly continuous with respect to $\|\cdot\|_\infty$ on \mathcal{M} . \square

3. MEASURED-VALUED GRAPHONS AND THE CUT DISTANCE

In Section 3.1, we introduce the measure-valued graphons, which are a generalization of real-valued graphons (i.e. $[0, 1]$ -valued measurable functions defined on $[0, 1]^2$). We refer to the monography [Lov12] on real-valued graphons for more details. In Sections 3.2, 3.3 and 3.4, we introduce the cut distance, and its unlabeled variant, on the space of measure-valued graphons which are analogous to the ones for real-valued graphons (see [Lov12, Chapter 8]). In Section 3.5, we define a weak isomorphism relation for measure-valued graphons based on this distance. Then, in Section 3.6, we give an alternative combinatorial formulation of the cut distance for stepfunctions.

3.1. Definition of measure-valued graphons. We start by defining measure-valued kernels and graphons which are a generalization of real-valued kernels and graphons. Recall that \mathbf{Z} is a Polish space and $\mathcal{M}_\pm(\mathbf{Z})$ is the space of *finite* signed measures.

Definition 3.1 (Signed measure-valued kernels). *A signed measure-valued kernel or $\mathcal{M}_\pm(\mathbf{Z})$ -valued kernel is a map W from $[0, 1]^2$ to $\mathcal{M}_\pm(\mathbf{Z})$, such that:*

- (i) W is a signed-measure in dz : for every $(x, y) \in [0, 1]^2$, $W(x, y; \cdot)$ belongs to $\mathcal{M}_\pm(\mathbf{Z})$.
- (ii) W is measurable in (x, y) : for every measurable set $A \subset \mathbf{Z}$, the function $(x, y) \mapsto W(x, y; A)$ defined on $[0, 1]^2$ is measurable.
- (iii) W is bounded:

$$(2) \quad \|W\|_\infty := \sup_{x, y \in [0, 1]} \|W(x, y; \cdot)\|_\infty < +\infty.$$

We denote by \mathcal{W}_1 (resp. $\mathcal{W}_{\leq 1}$, resp. \mathcal{W}_+ , resp. \mathcal{W}_{\pm}) the space of probability measure-valued kernels or simply probability-graphons (resp. sub-probability measure-valued kernels, resp. measure-valued kernels, resp. signed measure-valued kernels), where we identify kernels that are equal a.e. on $[0, 1]^2$, with respect to the Lebesgue measure. Then, (2) should be read with an essential supremum instead of a supremum. In what follows, we always assume for simplicity that we choose representatives of measure-valued kernels such that $\|W\|_{\infty}$ is also the essential supremum of $(x, y) \mapsto \|W(x, y; \cdot)\|_{\infty}$.

For $\mathcal{M} \subset \mathcal{M}_{\pm}(\mathbf{Z})$, we denote by $\mathcal{W}_{\mathcal{M}}$ the subset of signed measure-valued kernel $W \in \mathcal{W}_{\pm}$ which are \mathcal{M} -valued: $W(x, y; \cdot) \in \mathcal{M}$ for every $(x, y) \in [0, 1]^2$.

Remark 3.2 (On real-valued kernels). Let $\mathbf{Z} = \{0, 1\}$ be equipped with the discrete topology. Every real-valued graphon w can be represented using a probability-graphon W defined for every $x, y \in [0, 1]$ by $W(x, y; dz) = w(x, y)\delta_1(dz) + (1 - w(x, y))\delta_0(dz)$, where δ_z is the Dirac mass located at z . In particular we have that $w(x, y) = W(x, y; \{1\})$ for $x, y \in [0, 1]$.

Let $W \in \mathcal{W}_{\pm}$ be a signed measure-valued kernel. Define the map $W^+ : [0, 1]^2 \rightarrow \mathcal{M}_+(\mathbf{Z})$ to be the positive part of W , i.e. for every $(x, y) \in [0, 1]^2$, $W^+(x, y; \cdot)$ is the positive part of the measure $W(x, y; \cdot)$. Similarly define $W^- : [0, 1]^2 \rightarrow \mathcal{M}_+(\mathbf{Z})$ the negative part of W ; and then define $|W| = W^+ + W^-$ the total variation of W and $\|W\| = |W|(\mathbf{Z})$ the total mass of W .

Lemma 3.3 (The positive part W^+ of a kernel). *The maps W^+ , W^- and $|W|$ are all measure-valued kernels, and the map $\|W\| : (x, y) \mapsto \|W(x, y; \cdot)\|_{\infty}$ is measurable.*

Proof. The statements for $|W|$ and $\|W\|$ are immediate consequences of the statements for W^+ and W^- ; and as the proof for W^+ and W^- are similar, we only need to prove that W^+ is a measure-valued kernel. It is immediate that W^+ is bounded and that for every $(x, y) \in [0, 1]^2$, $W^+(x, y; \cdot)$ is a measure in $\mathcal{M}_+(\mathbf{Z})$. Thus, we are left to prove the measurability of W^+ in (x, y) . By [DF64, Proposition 2.1] and Remark 2.4, a signed measure-valued kernel U is measurable in (x, y) (i.e. for every $A \in \mathcal{B}(\mathbf{Z})$, the map $(x, y) \mapsto U(x, y; A)$ is measurable) if and only if the map $(x, y) \mapsto U(x, y; \cdot)$ is measurable from $[0, 1]^2$ (with its Borel σ -field) to $\mathcal{M}_{\pm}(\mathbf{Z})$ equipped with the Borel σ -field generated by the weak topology. By [DF64, Theorem 2.8], the map $\mu \mapsto \mu^+$, that associate to a signed measure the positive part of its Hahn-Jordan decomposition, is measurable from $\mathcal{M}_{\pm}(\mathbf{Z})$ to $\mathcal{M}_+(\mathbf{Z})$ both endowed with the Borel σ -field generated by the weak topology. Considering the composition of W and $\mu \mapsto \mu^+$, we get that W^+ is measurable in (x, y) and is thus a measure-valued kernel. \square

Remark 3.4 (Probability-graphons $W : \Omega \times \Omega \rightarrow \mathcal{M}_1(\mathbf{Z})$). Similarly to the case of real-valued graphons, it is possible to replace the vertex-type space $[0, 1]$ by any standard probability space $(\Omega, \mathcal{A}, \pi)$ that might be more appropriate to represent vertex-types for some applications, and to consider probability-graphons of the form $W : \Omega \times \Omega \rightarrow \mathcal{M}_1(\mathbf{Z})$. We recall that a standard probability space $(\Omega, \mathcal{A}, \pi)$ is a probability space such that there exists a measure-preserving map $\varphi : [0, 1] \rightarrow \Omega$, where $[0, 1]$ is endowed with the Borel σ -field and the Lebesgue measure. In particular, every Polish space endowed with its Borel σ -field is a standard probability space. As an example, the space $[0, 1]^2$ equipped with the Borel σ -field and the Lebesgue measure λ_2 is a standard probability space; we will reuse this fact later.

Using the measure preserving map φ , it is then possible to consider an unlabeled version W^{φ} of W constructed on $\Omega' = [0, 1]$, and to modify the definition of the cut distance $\delta_{\square, m}$ similarly as in [Jan13, Theorem 6.9] to allow each probability-graphons to be constructed on different standard probability spaces. For simplicity, in this article we only consider the equivalent case where all probability-graphons are constructed on $\Omega = [0, 1]$.

Remark 3.5 (Symmetric kernels). We shall consider non-symmetric measure-valued kernels and probability-graphons in order to handle directed graphs whose adjacency matrices are thus

a priori non-symmetric. We say that a measure-valued kernel or graphon W is symmetric if for a.e. $x, y \in [0, 1]$, $W(x, y; \cdot) = W(y, x; \cdot)$.

We define stepfunctions measure-valued kernel which are often used for approximation.

Definition 3.6 (Signed measure-valued stepfunctions). *A signed measure-valued kernel $W \in \mathcal{W}_\pm$ is a stepfunction if there exists a finite partition of $[0, 1]$ into measurable (possibly empty) sets, say $\mathcal{P} = \{S_1, \dots, S_k\}$, such that W is constant on the sets $S_i \times S_j$, for $1 \leq i, j \leq k$. We say that W and the partition \mathcal{P} are adapted to each other. We write $|\mathcal{P}| = k$ the number of elements of the partition \mathcal{P} .*

3.2. The cut distance. We define a distance and a norm on signed measure-valued graphons and kernels, called the *cut distance* and the *cut norm* respectively which are analogous to the cut norm for real-valued graphons and kernels, see [Lov12, Chapter 8]. For a signed measure-valued kernel $W \in \mathcal{W}_\pm$ and a measurable subsets $A \subset [0, 1]^2$, we denote by $W(A; \cdot)$ the signed measure on \mathbf{Z} defined by:

$$W(A; \cdot) = \int_A W(x, y; \cdot) \, dx dy.$$

Definition 3.7 (The cut distance $d_{\square, m}$). *Let d_m be a quasi-convex distance on \mathcal{M} a convex subset of $\mathcal{M}_\pm(\mathbf{Z})$ containing the zero measure. The associated cut distance $d_{\square, m}$ is the function defined on \mathcal{W}_\pm^2 by:*

$$(3) \quad d_{\square, m}(U, W) = \sup_{S, T \subset [0, 1]} d_m\left(U(S \times T; \cdot), W(S \times T; \cdot)\right),$$

where the supremum is taken over all measurable subsets S and T of $[0, 1]$.

Notice that the right-hand side of (3) is well defined as \mathcal{M} contains the zero measure (and thus if U belongs to $\mathcal{W}_\mathcal{M}$ then $U(A; \cdot)$ belongs to \mathcal{M}).

Definition 3.8 (The cut norm $N_{\square, m}$). *The cut norm $N_{\square, m}$ associated with a norm N_m on $\mathcal{M}_\pm(\mathbf{Z})$ is the function defined on \mathcal{W}_\pm by:*

$$N_{\square, m}(W) = \sup_{S, T \subset [0, 1]} N_m\left(W(S \times T; \cdot)\right),$$

where the supremum is taken over all measurable subsets S and T of $[0, 1]$.

The next proposition states that the cut distance (resp. norm) is indeed a distance (resp. norm); its extension to distances on $\mathcal{M}_+(\mathbf{Z})$ and $\mathcal{M}_\pm(\mathbf{Z})$ is immediate.

Proposition 3.9 ($d_{\square, m}$ is a distance, $N_{\square, m}$ is a norm). *The cut distance $d_{\square, m}$ associated with a distance d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$) is a distance on \mathcal{W}_1 (resp. \mathcal{W}_+). The cut norm $N_{\square, m}$ associated with a norm N_m on $\mathcal{M}_\pm(\mathbf{Z})$ is a norm on \mathcal{W}_\pm .*

Moreover, when the distance d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$) derives from a norm N_m on $\mathcal{M}_\pm(\mathbf{Z})$, then the distance $d_{\square, m}$ derives also from the norm $N_{\square, m}$.

Proof. Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (the proof for the case $\mathcal{M}_+(\mathbf{Z})$ is similar). It is clear that $d_{\square, m}$ is symmetric and satisfies the triangular inequality. Thus, we only need to prove that $d_{\square, m}$ is separating. Let U and W be two probability-graphons such that $d_{\square, m}(U, W) = 0$. Then, for every measurable subsets $S, T \subset [0, 1]$, we have $U(S \times T; \cdot) = W(S \times T; \cdot)$. Let $\mathcal{F} = (f_k)_{k \in \mathbb{N}}$ be a separating sequence. For every $k \in \mathbb{N}$, and for every measurable subsets $S, T \subset [0, 1]$, we have that $U(S \times T; f_k) = W(S \times T; f_k)$. This implies that $\int U(x, y, f_k) \, dx dy = \int W(x, y, f_k) \, dx dy$ for all $k \in \mathbb{N}$. Hence, we deduce that for all $k \in \mathbb{N}$, $U(x, y; f_k) = W(x, y; f_k)$ for almost every $(x, y) \in [0, 1]^2$. Thus, $U(x, y; \cdot) = W(x, y; \cdot)$ for almost every $(x, y) \in [0, 1]^2$. This implies that $d_{\square, m}$ is separating on \mathcal{W}_1 , and thus a distance on \mathcal{W}_1 .

The proof for the cut norm is similar. The proof of the last part of the proposition is clear. \square

3.3. Graphon relabeling, invariance and smoothness properties. The analogue of graph relabelings for graphons are measure-preserving maps. Recall the definition of a measure-preserving map from Section 2, and in particular (1). Recall $\tilde{S}_{[0,1]}$ denotes the set of measure-preserving (measurable) maps from $[0, 1]$ to $[0, 1]$ endowed with the Lebesgue measure, and $S_{[0,1]}$ denotes its subset of bijective maps.

The relabeling of a signed measure-valued kernel W by a measure-preserving map φ , is the signed measure-valued kernel W^φ defined for every $x, y \in [0, 1]$ and every measurable set $A \subset \mathbf{Z}$ by:

$$W^\varphi(x, y; A) = W(\varphi(x), \varphi(y); A) \quad \text{for } x, y \in [0, 1] \text{ and } A \subset \mathbf{Z} \text{ measurable.}$$

We say that a subset $\mathcal{K} \subset \mathcal{W}_\pm$ is *uniformly bounded* if:

$$(4) \quad \sup_{W \in \mathcal{K}} \|W\|_\infty < +\infty.$$

Definition 3.10 (Invariance and smoothness of a distance on kernels). *Let d be a distance on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_\pm). We say that the distance d is:*

- (i) **Invariant:** *if $d(U, W) = d(U^\varphi, W^\varphi)$ for every bijective measure-preserving map $\varphi \in S_{[0,1]}$ and $U, V \in \mathcal{W}_1$ (resp. U, V belongs to \mathcal{W}_+ or \mathcal{W}_\pm).*
- (ii) **Smooth:** *if a.e. weak convergence implies convergence for d , that is, if $(W_n)_{n \in \mathbb{N}}$ and W are kernels from \mathcal{W}_1 (resp. kernels from \mathcal{W}_+ or \mathcal{W}_\pm that are uniformly bounded and) such that for a.e. $(x, y) \in [0, 1]^2$, $W_n(x, y; \cdot)$ weakly converges to $W(x, y; \cdot)$ as $n \rightarrow \infty$, then $\lim_{n \rightarrow \infty} d(W_n, W) = 0$.*

We say that a norm N on \mathcal{W}_\pm is *invariant* (resp. *smooth*) if its associated distance d on \mathcal{W}_\pm is *invariant* (resp. *smooth*).

We shall see in Section 3.8 some examples of distances d_m for which the associated cut distance $d_{\square, m}$ is invariant and smooth. The invariance property from Definition 3.10 is always satisfied by the cut distance, and thus also by the cut norm.

Lemma 3.11 ($d_{\square, m}$ is invariant). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$, resp. $\mathcal{M}_\pm(\mathbf{Z})$). Then the cut distance $d_{\square, m}$ on \mathcal{W}_1 (resp. \mathcal{W}_+ , resp. \mathcal{W}_\pm) is invariant.*

Proof. For a signed measure-valued kernel W , a bijective measure-preserving map $\varphi \in S_{[0,1]}$, and measurable sets $S, T \subset [0, 1]$, we have thanks to (1):

$$\int_{S \times T} W^\varphi(x, y; \cdot) \, dx dy = \int_{S \times T} W(\varphi(x), \varphi(y); \cdot) \, dx dy = \int_{\varphi(S) \times \varphi(T)} W(x, y; \cdot) \, dx dy.$$

Hence, taking the supremum over every measurable sets $S, T \subset [0, 1]$, we get that the cut distance $d_{\square, m}$ is invariant. \square

When a smooth distance on \mathcal{W}_1 or \mathcal{W}_+ derives from a distance on $\mathcal{M}_1(\mathbf{Z})$ or $\mathcal{M}_+(\mathbf{Z})$, we have the following result.

Lemma 3.12 (Smoothness and the weak topology). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_\pm(\mathbf{Z})$) such that the distance $d_{\square, m}$ on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_\pm) is smooth. Then, the distance d_m is continuous w.r.t. the weak topology on $\mathcal{M}_1(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$).*

Proof. Let $(\mu_n)_{n \in \mathbb{N}}$, and μ be measures from $\mathcal{M}_1(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$) such that $(\mu_n)_{n \in \mathbb{N}}$ weakly converges to μ . Consider the constant measure-valued graphons (resp. kernels) $W_n \equiv \mu_n$, $n \in \mathbb{N}$, and $W \equiv \mu$. Then, for every $x, y \in [0, 1]$, $W_n(x, y; \cdot)$ weakly converges to $W(x, y; \cdot)$ as $n \rightarrow \infty$. As the distance $d_{\square, m}$ is smooth, we get that $\lim_{n \rightarrow \infty} d_{\square, m}(W_n, W) = 0$. Considering $S = T = [0, 1]$ in the cut distance, we deduce that $\lim_{n \rightarrow \infty} d_m(\mu_n, \mu) = 0$. \square

The next lemma is a partial converse of Lemma 3.12, it gives sufficient conditions for $d_{\square, m}$ to be smooth. Remind the definition of a quasi-convex distance in Definition 2.10.

Proposition 3.13 ($d_{\square, m}$ is smooth). *Let d_m be distance on $\mathcal{M}_\epsilon(\mathbf{Z})$ with $\epsilon \in \{+, \pm\}$ which is quasi-convex and sequentially continuous w.r.t. the weak topology (on $\mathcal{M}_\epsilon(\mathbf{Z})$). Then, the cut distance $d_{\square, m}$ is smooth.*

Moreover, for all $U, W \in \mathcal{W}_\epsilon$, and for all measurable $A \subset [0, 1]^2$, we have:

$$(5) \quad d_m(U(A; \cdot), W(A; \cdot)) \leq \operatorname{essup}_{(x, y) \in A} d_m(U(x, y; \cdot), W(x, y; \cdot)).$$

To prove Proposition 3.13, we first need to prove the following lemma for approximation by \mathcal{M} -valued kernels taking finitely many values.

Lemma 3.14. *Let $W \in \mathcal{W}_\pm$ and a subset $A \subset [0, 1]^2$. There exists a sequence $(W_n)_{n \in \mathbb{N}}$ in \mathcal{W}_\pm such that $(W_n(A; \cdot))_{n \in \mathbb{N}}$ weakly converges to $W(A; \cdot)$ and for all $n \in \mathbb{N}$, W_n is finitely valued and takes its values in $\{W(x, y; \cdot) : (x, y) \in A\}$.*

Proof. By scaling, we may assume that $\|W\|_\infty \leq 1$. Let $(f_k)_{k \in \mathbb{N}}$ be a convergence determining sequence with $f_0 = \mathbb{1}$ and f_k takes values in $[0, 1]$. Thus, for all $(x, y) \in [0, 1]^2$, $\epsilon \in \{\pm 1\}$ and $k \in \mathbb{N}$, we have $W_\epsilon(x, y; f_k) \in [0, 1]$. For all $n \in \mathbb{N}$, let $(C_{n, i})_{1 \leq i \leq d_n}$ be a partition of $[0, 1]^{2(n+1)}$ into $d_n = n^{2(n+1)}$ hypercubes of edge-length $r_n = 1/n$. Then, for all $n \in \mathbb{N}$ and $i \in [d_n]$, define $B_{n, i} = A \cap (W_+(\cdot; (f_i)_{0 \leq i \leq n}, W_-(\cdot; (f_i)_{0 \leq i \leq n})^{-1}(C_{n, i}))$; thus we get a partition $(B_{n, i})_{1 \leq i \leq d_n}$ of A . If $B_{n, i} \neq \emptyset$, fix some $\mu_{n, i} \in \{W(x, y; \cdot) : (x, y) \in B_{n, i}\}$. If $A \neq [0, 1]^2$, fix some $\mu_\partial \in \{W(x, y; \cdot) : (x, y) \in [0, 1]^2 \setminus A\}$. For $n \in \mathbb{N}$, we define $W_n = \mathbb{1}_{A^c} \mu_\partial + \sum_{i=1}^{d_n} \mathbb{1}_{B_{n, i}} \mu_{n, i}$, which is finitely valued and takes its values in $\{W(x, y; \cdot) : (x, y) \in A\}$.

Let $k \in \mathbb{N}$ and $\epsilon \in \{\pm\}$. For all $n \geq k$, we have:

$$|W_\epsilon(A; f_k) - (W_n)_\epsilon(A; f_k)| \leq \sum_{i=1}^{d_n} \int_{B_{n, i}} |W_\epsilon(x, y; f_k) - (\mu_{n, i})_\epsilon| \, dx dy \leq \frac{1}{n}.$$

As $(f_k)_{k \in \mathbb{N}}$ is convergence determining, this implies that $((W_n)_\epsilon(A; \cdot))_{n \in \mathbb{N}}$ weakly converges to $W_\epsilon(A; \cdot)$ for $\epsilon \in \{\pm\}$. Hence, $(W_n(A; \cdot))_{n \in \mathbb{N}}$ weakly converges to $W(A; \cdot)$. \square

Proof of Proposition 3.13. As d_m is quasi-convex, (5) is immediate when U and W take only finitely many values. Now, assume that U and W are arbitrary $\mathcal{M}_\epsilon(\mathbf{Z})$ -valued kernels. Let $\epsilon > 0$. As d_m is sequentially continuous w.r.t. the weak topology, using Lemma 3.14, there exist two $\mathcal{M}_\epsilon(\mathbf{Z})$ -valued kernel U' and W' such that $d_m(U'(A; \cdot), U(A; \cdot)) < \epsilon$ and U' is finitely valued and takes its values in $\{U(x, y; \cdot) : (x, y) \in A\}$, and similarly for W' and W . Thus, we have:

$$d_m(U(A; \cdot), W(A; \cdot)) \leq 2\epsilon + \operatorname{essup}_{(x, y) \in A} d_m(U(x, y; \cdot), W(x, y; \cdot)),$$

and this being true for all $\epsilon > 0$, we get (5).

Let $(W_n)_{n \in \mathbb{N}}$ and W be $\mathcal{M}_\epsilon(\mathbf{Z})$ -valued kernels which are uniformly bounded by some constant $C < \infty$ and such that for a.e. $(x, y) \in [0, 1]^2$, the sequence $((W_n(x, y; \cdot))_{n \in \mathbb{N}}$ converges to $W(x, y; \cdot)$ for the weak topology, and thus also for d_m . Let $\epsilon > 0$ and $S, T \subset [0, 1]$. As d_m is quasi-convex and sequentially continuous w.r.t. the weak topology, using Lemma 2.11, there exists $\eta > 0$ such that for all $\mu, \nu \in \mathcal{M}_\epsilon(\mathbf{Z})$, we have that $\|\mu - \nu\|_\infty < \eta$ implies $d_m(\mu, \nu) < \epsilon$. For all $n \in \mathbb{N}$, define the measurable set:

$$A_n = \{(x, y) \in S \times T : d_m(W_n(x, y; \cdot), W(x, y; \cdot)) < \epsilon\}.$$

By assumption, we have that $\lim_{n \rightarrow \infty} \lambda(A_n) = \lambda(S \times T)$. Let $N \in \mathbb{N}$ be such that for $n \geq N$, we have $\lambda((S \times T) \setminus A_n) < \eta/C$. Let $n \geq N$. Remark that $W_n((S \times T) \setminus A_n; \cdot)$ and $W((S \times T) \setminus A_n; \cdot)$

have total mass at most $C\lambda(A_n^c) < \eta$. Thus, we have that $d_m(W_n(A_n; \cdot), W_n(S \times T; \cdot)) < \varepsilon$ and $d_m(W(A_n; \cdot), W(S \times T; \cdot)) < \varepsilon$. Hence, using (5) we get that:

$$\begin{aligned} d_m(W_n(S \times T; \cdot), W(S \times T; \cdot)) &\leq 2\varepsilon + d_m(W_n(A_n; \cdot), W(A_n; \cdot)) \\ &\leq 2\varepsilon + \operatorname{esssup}_{(x,y) \in A_n} d_m(W_n(x, y; \cdot), W(x, y; \cdot)) \\ &\leq 3\varepsilon. \end{aligned}$$

Taking the supremum over $S, T \subset [0, 1]$, we get $d_{\square, m}(W_n, W) \leq 3\varepsilon$. This being true for all $\varepsilon > 0$, we conclude that $(W_n)_{n \in \mathbb{N}}$ converges to W for $d_{\square, m}$, and thus $d_{\square, m}$ is smooth. \square

3.4. The unlabeled cut distance. We can now define the cut distance for unlabeled graphons.

Definition 3.15 (The unlabeled cut distance $\delta_{\square, m}$). *Set $\mathcal{K} \in \{\mathcal{W}_1, \mathcal{W}_+, \mathcal{W}_\pm\}$. Let d be an invariant distance on the kernel space \mathcal{K} . The premetric δ_{\square} on \mathcal{K} , also called the cut distance, is defined by:*

$$(6) \quad \delta_{\square}(U, W) = \inf_{\varphi \in S_{[0,1]}} d(U, W^\varphi) = \inf_{\varphi \in S_{[0,1]}} d(U^\varphi, W).$$

Notice that δ_{\square} satisfies the symmetry property (as d is invariant) and the triangular inequality. Hence, δ_{\square} induces a distance (that we still denote by δ_{\square}) on the quotient space $\tilde{\mathcal{K}}_d = \mathcal{K} / \sim_d$ of kernels in \mathcal{K} associated with the equivalence relation \sim_d defined by $U \sim_d W$ if and only if $\delta_{\square}(U, W) = 0$.

When the metric $d = d_{\square, m}$ on $\mathcal{K} = \mathcal{W}_1$ (resp. \mathcal{W}_+ , resp. \mathcal{W}_\pm) derives from a metric d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$, resp. $\mathcal{M}_\pm(\mathbf{Z})$), and is thus invariant thanks to Lemma 3.11, we write $\delta_{\square, m}$ for δ_{\square} and $\tilde{\mathcal{K}}_m$ for $\tilde{\mathcal{K}}_{d_{\square, m}}$. We shall see in Theorem 5.5 and Corollary 5.6 that under some conditions, different choices of distance d_m , which induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, lead to the same quotient space, then simply denoted by $\tilde{\mathcal{W}}_1$, with the same topology.

3.5. Weak isomorphism. Similarly to Theorem 8.13 in [Lov12], when the distance d_m is such that $d_{\square, m}$ is invariant and smooth, we can rewrite the cut distance $\delta_{\square, m}$ as a minimum instead of an infimum using measure-preserving maps, see the last equality in (7).

We introduce a weak isomorphism relation that allows to “un-label” probability-graphons.

Definition 3.16 (Weak isomorphism). *We say that two signed measure-valued kernels U and W are weakly isomorphic (and we note $U \sim W$) if there exists two measure-preserving maps $\varphi, \psi \in \tilde{S}_{[0,1]}$ such that $U^\varphi(x, y; \cdot) = W^\psi(x, y; \cdot)$ for a.e. $x, y \in [0, 1]$.*

We denote by $\tilde{\mathcal{W}}_\pm = \mathcal{W}_\pm / \sim$ (resp. $\tilde{\mathcal{W}}_1 = \mathcal{W}_1 / \sim$) the space of unlabeled signed measure-valued kernels (resp. probability-graphons) i.e. the space of signed measure-valued kernels (resp. probability-graphons) where we identify signed measure-valued kernels (resp. probability-graphons) that are weakly isomorphic.

Notice that $U \sim W$ implies that $\|U\|_\infty = \|W\|_\infty$ (we recall that signed measure-valued kernels are only defined for a.e. $x, y \in [0, 1]$ and that $\|W\|_\infty$ in (2) is an essup in general). In particular, the notion of uniformly bounded subset defined in (4) naturally extends to $\tilde{\mathcal{W}}_\pm$. The last part of this section is devoted to the proof of the following key result.

Theorem 3.17 (Weak isomorphism and δ_{\square}). *Let d be a distance defined on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_\pm) which is invariant and smooth. Then, two kernels are weakly isomorphic, i.e. $U \sim W$, if and only if $U \sim_d W$, i.e. $\delta_{\square}(U, W) = 0$.*

Furthermore, the map δ_{\square} is a distance on $\tilde{\mathcal{W}}_1 = \tilde{\mathcal{W}}_{1,d}$ (resp. $\tilde{\mathcal{W}}_+ = \tilde{\mathcal{W}}_{+,d}$ or $\tilde{\mathcal{W}}_\pm = \tilde{\mathcal{W}}_{\pm,d}$).

As a first step in the proof of Theorem 3.17, following [Lov12], we give a nice description of δ_{\square} using couplings. We say that a measure μ on $[0, 1]^2$ is a coupling measure on $[0, 1]^2$ (between two copies of $[0, 1]$ each equipped with the Lebesgue measure) if the projection maps on each components $\tau, \rho : [0, 1]^2 \rightarrow [0, 1]$ (where $[0, 1]^2$ is equipped with the measure μ and $[0, 1]$ with the Lebesgue measure λ) are measure-preserving. Thus for every kernel W on $([0, 1], \mathcal{B}([0, 1]), \lambda)$, the function W^{τ} is a kernel on the probability space $([0, 1]^2, \mathcal{B}([0, 1]^2), \mu)$, and similarly for the projection ρ .

Let φ be a given measure-preserving map from $[0, 1]$ with the Lebesgue measure to $[0, 1]^2$ with a coupling measure μ . For an invariant distance d on \mathcal{W}_1 (resp. \mathcal{W}_{\pm}), we define a distance, say d^{μ} , on kernels on $([0, 1]^2, \mathcal{B}([0, 1]^2), \mu)$ by:

$$d^{\mu}(U', W') = d(U'^{\varphi}, W'^{\varphi}).$$

It is easy to see that, for U and W kernels on $[0, 1]$, we have $d^{\mu}(U^{\tau}, W^{\tau}) = d(U, W)$ as d is invariant and $\tau \circ \varphi$ is a measure-preserving map from $[0, 1]$ to itself; and similarly $d^{\mu}(U^{\rho}, W^{\rho}) = d(U, W)$.

A straightforward adaptation of the proof of [Lov12, Theorem 8.13] gives the next result.

Proposition 3.18 (Minima in the cut distance δ_{\square}). *Let d be a distance defined on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}) which is invariant and smooth. Then, we have the following alternative formulations for the cut distance δ_{\square} on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}):*

$$\begin{aligned} \delta_{\square}(U, W) &= \inf_{\varphi \in \mathcal{S}_{[0,1]}} d(U, W^{\varphi}) = \inf_{\varphi \in \tilde{\mathcal{S}}_{[0,1]}} d(U, W^{\varphi}) \\ (7) \quad &= \inf_{\psi \in \mathcal{S}_{[0,1]}} d(U^{\psi}, W) = \inf_{\psi \in \tilde{\mathcal{S}}_{[0,1]}} d(U^{\psi}, W) \\ &= \inf_{\varphi, \psi \in \mathcal{S}_{[0,1]}} d(U^{\psi}, W^{\varphi}) = \min_{\varphi, \psi \in \tilde{\mathcal{S}}_{[0,1]}} d(U^{\psi}, W^{\varphi}), \end{aligned}$$

and

$$\delta_{\square}(U, W) = \min_{\mu} d^{\mu}(U^{\tau}, W^{\rho})$$

where μ range over all coupling measures on $[0, 1]^2$.

Proof of Theorem 3.17. We deduce from the last equality in (7) that $\delta_{\square}(U, W) = 0$ if and only if there exist measure-preserving maps $\varphi, \psi \in \tilde{\mathcal{S}}_{[0,1]}$ such that $U^{\psi}(x, y; \cdot) = W^{\varphi}(x, y; \cdot)$ for a.e. $x, y \in [0, 1]$. This gives that the equivalence relations \sim_d and \sim are the same. \square

3.6. The cut norm for stepfunctions. For a quasi-convex distance d_m , the cut distance d_m for stepfunctions can be reformulated using a finite combinatorial optimization. For a collection of subsets \mathcal{P} , denote by $\sigma(\mathcal{P})$ the σ -field generated by \mathcal{P} .

Lemma 3.19 (Combinatorial optimization of quasi-convex d_m for stepfunctions). *Let d_m be a quasi-convex distance on \mathcal{M} a convex subset of $\mathcal{M}_{\pm}(\mathbf{Z})$ containing the zero measure. Let $U, W \in \mathcal{W}_{\mathcal{M}}$ be \mathcal{M} -valued stepfunctions adapted to the same finite partition \mathcal{P} . Then, there exists $S, T \in \sigma(\mathcal{P})$ such that:*

$$d_{\square, m}(U, W) = d_m(U(S \times T; \cdot), W(S \times T; \cdot)).$$

Proof. Let $\mathcal{P} = \{S_1, \dots, S_k\}$ with $k = |\mathcal{P}|$ the size of the partition \mathcal{P} . First, remark that the quantity $d_{\square, m}(U, W) = d_m(U(S' \times T'; \cdot), W(S' \times T'; \cdot))$ depends on S' and T' only through the values of $\lambda(S' \cap S_i)$ and $\lambda(T' \cap S_i)$ for $1 \leq i \leq k$. Thus, the cut distance between U and W can be reformulated as:

$$d_{\square, m}(U, W) = \sup_{0 \leq \alpha_i, \beta_i \leq \lambda(S_i); 1 \leq i \leq k} d_m \left(\sum_{1 \leq i, j \leq k} \alpha_i \beta_j \mu_{i, j}(\cdot), \sum_{1 \leq i, j \leq k} \alpha_i \beta_j \nu_{i, j}(\cdot) \right),$$

where $\mu_{i,j}$ (resp. $\nu_{i,j}$) is the constant value of $U(x, y; \cdot)$ (resp. $W(x, y; \cdot)$) when $x \in S_i$ and $y \in S_j$. Moreover, when we fix the value of $\beta = (\beta_i)_{1 \leq i \leq k}$, the quantity

$$d_m \left(\sum_{1 \leq i, j \leq k} \alpha_i \beta_j \mu_{i,j}(\cdot), \sum_{1 \leq i, j \leq k} \alpha_i \beta_j \nu_{i,j}(\cdot) \right)$$

is a quasi-convex function of $\alpha = (\alpha_i)_{1 \leq i \leq k}$, and thus realizes its maximum on the extremal points of the hypercube $\prod_{i=1}^k [0, \lambda(S_i)]$, i.e. when α_i equals 0 or $\lambda(S_i)$ for every $1 \leq i \leq k$. By symmetry, a similar argument holds for β . The cut distance can thus be reformulated as the combinatorial optimization:

$$d_{\square, m}(U, W) = \max_{I, J \subset [k]} d_m \left(\sum_{i \in I, j \in J} \mu_{i,j}(\cdot), \sum_{i \in I, j \in J} \nu_{i,j}(\cdot) \right).$$

Let $I, J \subset [k]$ that maximizes this combinatorial optimization, and take $S = \cup_{i \in I} S_i$ and $T = \cup_{j \in J} S_j$ to conclude. \square

3.7. The supremum in S and T in the cut distance $d_{\square, m}$. In this section, we prove that the supremum in the cut distance $d_{\square, m}$ is achieved by some subsets $S, T \subset [0, 1]$.

For $W \in \mathcal{M}_{\pm}(\mathbf{Z})$ and $f, g : [0, 1] \rightarrow [0, 1]$ measurable, we define the signed measure:

$$W(f \otimes g; \cdot) = \int_{[0, 1]^2} W(x, y; \cdot) f(x) g(y) \, dx dy.$$

Remark that if we have $W \in \mathcal{W}_{\epsilon}$ with $\epsilon \in \{1, \leq 1, +, \pm\}$, then we have $W(f \otimes g; \cdot) \in \mathcal{M}_{\epsilon}(\mathbf{Z})$.

Lemma 3.20 (The supremum in the cut distance $d_{\square, m}$ for quasi-convex distance d_m). *Let d_m be a quasi-convex distance on $\mathcal{M}_{\epsilon}(\mathbf{Z})$ with $\epsilon \in \{+, \pm\}$ that is sequentially continuous w.r.t. the weak topology. Let $U, W \in \mathcal{W}_{\epsilon}$. Then, there exist measurable subsets $S, T \subset [0, 1]$ such that $f = \mathbf{1}_S$ and $g = \mathbf{1}_T$ achieve the supremum in:*

$$\sup_{f, g} d_m \left(U(f \otimes g; \cdot), W(f \otimes g; \cdot) \right)$$

where the supremum is taken over measurable functions f, g from $[0, 1]$ to itself.

Proof. Define the map $\Psi : (f, g) \mapsto d_m(U(f \otimes g; \cdot), W(f \otimes g; \cdot))$, and denote $C = \sup_{f, g} \Psi(f, g)$, where the supremum is taken over measurable functions f, g from $[0, 1]$ to itself. Let $(f_n)_{n \in \mathbb{N}}$ and $(g_n)_{n \in \mathbb{N}}$ be sequences of measurable functions from $[0, 1]$ to itself such that $\lim_{n \rightarrow \infty} \Psi(f_n, g_n) = C$. As the unit ball of $L^{\infty}([0, 1], \lambda)$ is compact for the weak-* topology (with primal space $L^1([0, 1], \lambda)$), upon taking subsequences, we may assume that $(f_n)_{n \in \mathbb{N}}$ (resp. $(g_n)_{n \in \mathbb{N}}$) weak-* converges to some f (resp. g) which take values in $[0, 1]$. Thus, $(f_n \otimes g_n)_{n \in \mathbb{N}}$ weak-* converges to $f \otimes g$ in $L^{\infty}([0, 1]^2, \lambda_2)$. In particular, for every $h \in C_b(\mathbf{Z})$, as $W[h]$ is a real-valued kernel, this implies that $\lim_{n \rightarrow \infty} W(f_n \otimes g_n; h) = W(f \otimes g; h)$. This being true for every $h \in C_b(\mathbf{Z})$, we get that the sequence $(W(f_n \otimes g_n; \cdot))_{n \in \mathbb{N}}$ in $\mathcal{M}_{\epsilon}(\mathbf{Z})$ weakly converges to $W(f \otimes g; \cdot) \in \mathcal{M}_{\epsilon}(\mathbf{Z})$; and similarly for U . As d_m is sequentially continuous w.r.t. the weak topology on $\mathcal{M}_{\epsilon}(\mathbf{Z})$, we get that $C = \lim_{n \rightarrow \infty} \Psi(f_n, g_n) = \Psi(f, g)$.

Now, we show that we can replace the functions f and g by functions that only take the values 0 and 1 (i.e. indicator functions). We first fix g and do this for f . Let X be a random variable uniformly distributed over $[0, 1]$, and consider the random function $\mathbf{1}_{X \leq f}$. Remark that we have $\mathbb{E}[W(\mathbf{1}_{X \leq f} \otimes g; \cdot)] = W(f \otimes g; \cdot)$, and similarly for U . As d_m is quasi-convex and sequentially

continuous w.r.t. the weak topology, we have:

$$\begin{aligned} C &\geq \sup_{x \in [0,1]} d_m(U(\mathbb{1}_{x \leq f} \otimes g; \cdot), W(\mathbb{1}_{x \leq f} \otimes g; \cdot)) \\ &\geq d_m(\mathbb{E}[U(\mathbb{1}_{X \leq f} \otimes g; \cdot)], \mathbb{E}[W(\mathbb{1}_{X \leq f} \otimes g; \cdot)]) \\ &= d_m(U(f \otimes g; \cdot), W(f \otimes g; \cdot)) \\ &= C, \end{aligned}$$

where in the second equality we used the quasi-convex supremum inequality from (5) with the $\mathcal{M}_\epsilon(\mathbf{Z})$ -valued kernels $U'(x, y; \cdot) = U(\mathbb{1}_{x \leq f} \otimes g; \cdot)$ and $W'(x, y; \cdot) = W(\mathbb{1}_{x \leq f} \otimes g; \cdot)$, and $A = [0, 1]^2$. All inequalities being equalities, this imposes:

$$C = \sup_{x \in [0,1]} d_m(U(\mathbb{1}_{x \leq f} \otimes g; \cdot), W(\mathbb{1}_{x \leq f} \otimes g; \cdot)) = \lim_{n \rightarrow \infty} d_m(U(\mathbb{1}_{r_n \leq f} \otimes g; \cdot), W(\mathbb{1}_{r_n \leq f} \otimes g; \cdot)),$$

for some sequence $(x_n)_{n \in \mathbb{N}}$ in $[0, 1]$. Upon taking a subsequence, we may assume that the sequence $(x_n)_{n \in \mathbb{N}}$ monotonically converges to some $x \in [0, 1]$. In particular, the sequence of functions $(\mathbb{1}_{x_n \leq f})_{n \in \mathbb{N}}$ (monotonically) converges to the function $f' = \mathbb{1}_{x \leq f}$ (resp. $f' = \mathbb{1}_{x < f}$) if $(x_n)_{n \in \mathbb{N}}$ is non-decreasing (resp. decreasing), and thus also weak-* converges in $L^\infty([0, 1], \lambda)$. Using, as in the first part of the proof, the sequential continuity of the function Ψ w.r.t. the weak-* topology on $L^\infty([0, 1], \lambda)$, we get that $\Psi(f', g) = d_m(U(f' \otimes g; \cdot), W(f' \otimes g; \cdot)) = C$, that is we can replace f by the indicator function f' . The same argument allows to replace g by an indicator function. \square

3.8. Examples of distance d_m . We consider usual distances and norms on $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_\pm(\mathbf{Z})$ that induce the weak topology on $\mathcal{M}_+(\mathbf{Z})$. All the distances we consider are quasi-convex, and all the norms we consider are sequentially continuous w.r.t. the weak topology on $\mathcal{M}_\pm(\mathbf{Z})$. Thus their associated cut distances are invariant and smooth by Lemma 3.11 and Proposition 3.13. Properties for the cut distances associated with those distances and norms are summarized in Corollaries 4.14 and 5.6.

In this section, we assume that (\mathbf{Z}, d_0) is a Polish metric space, and remind that $\mathcal{B}(\mathbf{Z})$ denotes its Borel σ -field.

3.8.1. The Prohorov distance $d_{\mathcal{P}}$. The Prohorov distance $d_{\mathcal{P}}$ is a complete distance defined on the set of finite measures $\mathcal{M}_+(\mathbf{Z})$ that induces the weak topology (see [Bil99, Theorem 6.8]). It is defined for $\mu, \nu \in \mathcal{M}_+(\mathbf{Z})$ as:

$$(8) \quad d_{\mathcal{P}}(\mu, \nu) = \inf\{\varepsilon > 0 : \forall A \in \mathcal{B}(\mathbf{Z}), \mu(A) \leq \nu(A^\varepsilon) + \varepsilon \text{ and } \nu(A) \leq \mu(A^\varepsilon) + \varepsilon\},$$

where $A^\varepsilon = \{x \in \mathbf{Z} : \exists y \in A, d_0(x, y) < \varepsilon\}$. For probability measures, we only need one inequality in (8) to define the Prohorov distance; however for positive measures we need both inequalities as two arbitrary positive measures might not have the same total mass. For $d_m = d_{\mathcal{P}}$, we use the subscript $m = \mathcal{P}$. We now prove that the Prohorov distance is quasi-convex.

Lemma 3.21. *The Prohorov distance $d_{\mathcal{P}}$ is quasi-convex on $\mathcal{M}_+(\mathbf{Z})$.*

Proof. Let $\mu_1, \mu_2, \nu_1, \nu_2 \in \mathcal{M}_+(\mathbf{Z})$ and let $\alpha \in [0, 1]$. Let $\varepsilon > \max(d_{\mathcal{P}}(\mu_1, \nu_1), d_{\mathcal{P}}(\mu_2, \nu_2))$, then for all $i \in \{1, 2\}$ and $B \in \mathcal{B}(\mathbf{Z})$, we have that $\mu_i(B) \leq \nu_i(B^\varepsilon) + \varepsilon$ and $\nu_i(B) \leq \mu_i(B^\varepsilon) + \varepsilon$. Taking a linear combination of those inequalities, we get that for all $B \in \mathcal{B}(\mathbf{Z})$, we have that $\alpha\mu_1(B) + (1 - \alpha)\mu_2(B) \leq \alpha\nu_1(B^\varepsilon) + (1 - \alpha)\nu_2(B^\varepsilon) + \varepsilon$, and similarly when swapping the role (μ_1, μ_2) and (ν_1, ν_2) . Hence, we get that $d_{\mathcal{P}}(\alpha\mu_1 + (1 - \alpha)\mu_2, \alpha\nu_1 + (1 - \alpha)\nu_2) \leq \varepsilon$, and taking the infimum over ε , we get that $d_m(\alpha\mu_1 + (1 - \alpha)\mu_2, \alpha\nu_1 + (1 - \alpha)\nu_2) \leq \max(d_m(\mu_1, \nu_1), d_m(\mu_2, \nu_2))$. \square

3.8.2. *The Kantorovitch-Rubinshtein and Fortet-Mourier norms.* The Kantorovitch-Rubinshtein norm $\|\cdot\|_{\text{KR}}$ (sometimes also called the bounded Lipschitz distance) and the Fortet-Mourier norm $\|\cdot\|_{\text{FM}}$ are two norms defined on $\mathcal{M}_{\pm}(\mathbf{Z})$ that induce the weak topology on $\mathcal{M}_{+}(\mathbf{Z})$ (see Section 3.2 in [Bog18] for definition and properties of those norms). They are defined for $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$ by:

$$\begin{aligned}\|\mu\|_{\text{KR}} &= \sup \left\{ \int_{\mathbf{Z}} f \, d\mu : f \text{ is 1-Lipschitz and } \|f\|_{\infty} \leq 1 \right\}, \\ \|\mu\|_{\text{FM}} &= \sup \left\{ \int_{\mathbf{Z}} f \, d\mu : f \text{ is Lipschitz and } \|f\|_{\infty} + \text{Lip}(f) \leq 1 \right\},\end{aligned}$$

where $\|f\|_{\infty} = \sup_{x \in \mathbf{Z}} |f(x)|$ is the infinite norm and $\text{Lip}(f)$ is the smallest constant $L > 0$ such that f is L -Lipschitz. Those two norms are metrically equivalent, see beginning of Section 3.2 in [Bog18]:

$$(9) \quad \|\mu\|_{\text{FM}} \leq \|\mu\|_{\text{KR}} \leq 2\|\mu\|_{\text{FM}}.$$

Note that we have $\|\mu\|_{\text{KR}} \leq \|\mu\|_{\infty}$, and thus those two norms are sequentially continuous w.r.t. the weak topology on $\mathcal{M}_{\pm}(\mathbf{Z})$.

An easy adaptation of the proof for Theorem 3.2.2 in [Bog18] gives the following comparison between $d_{\mathcal{P}}$, $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$.

Lemma 3.22 (Comparison of $d_{\mathcal{P}}$, $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$). *Let $\mu, \nu \in \mathcal{M}_{+}(\mathbf{Z})$. Then, we have:*

$$\frac{d_{\mathcal{P}}(\mu, \nu)^2}{1 + d_{\mathcal{P}}(\mu, \nu)} \leq \|\mu - \nu\|_{\text{FM}} \leq \|\mu - \nu\|_{\text{KR}} \leq (2 + \min(\mu(\mathbf{Z}), \nu(\mathbf{Z}))) d_{\mathcal{P}}(\mu, \nu).$$

In particular, the Prohorov distance $d_{\mathcal{P}}$ is uniformly continuous w.r.t. $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$ on $\mathcal{M}_{+}(\mathbf{Z})$; and $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$ are uniformly continuous w.r.t. $d_{\mathcal{P}}$ on $\mathcal{M}_{\leq 1}(\mathbf{Z})$.

For the special choice $N_{\text{m}} = \|\cdot\|_{\text{KR}}$ (resp. $N_{\text{m}} = \|\cdot\|_{\text{FM}}$), we use the subscript $\text{m} = \text{KR}$ (resp. $\text{m} = \text{FM}$).

3.8.3. *A norm based on a convergence determining sequence.* From a convergence determining sequence $\mathcal{F} = (f_k)_{k \in \mathbb{N}}$, where $f_0 = \mathbf{1}$ and $f_k \in C_b(\mathbf{Z})$ takes values in $[0, 1]$, we define a norm on $\mathcal{M}_{\pm}(\mathbf{Z})$ metrizing the weak topology on $\mathcal{M}_{+}(\mathbf{Z})$, for $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$, by:

$$(10) \quad \|\mu\|_{\mathcal{F}} = \sum_{k \in \mathbb{N}} 2^{-k} |\mu(f_k)|.$$

Note that we have $\|\mu\|_{\mathcal{F}} \leq 2\|\mu\|_{\infty}$, and thus $\|\cdot\|_{\mathcal{F}}$ is sequentially continuous w.r.t. the weak topology on $\mathcal{M}_{\pm}(\mathbf{Z})$. For the special choice $N_{\text{m}} = \|\cdot\|_{\mathcal{F}}$, we use the subscript $\text{m} = \mathcal{F}$.

Even though the norm $\|\cdot\|_{\mathcal{F}}$ is not complete when \mathbf{Z} is not compact (see Lemma 3.23 below), the cut norm $\|\cdot\|_{\square, \mathcal{F}}$ and the cut distance $\delta_{\square, \mathcal{F}}$ will turn out to be very useful in Sections 6 and 7 to link the topology of the cut distance to the homomorphism densities. Recall $d_{\mathcal{F}}$ is the distance derived from the norm $\|\cdot\|_{\mathcal{F}}$.

Lemma 3.23 ($d_{\mathcal{F}}$ is not complete in general). *Let \mathcal{F} be a convergence determining sequence. Then, the distance $d_{\mathcal{F}}$ is complete over $\mathcal{M}_1(\mathbf{Z})$ if and only if $\mathcal{M}_1(\mathbf{Z})$ is a compact space, i.e. , if and only if \mathbf{Z} is compact.*

Proof. Theorem 3.4 in [Var58] states that \mathbf{Z} is compact if and only if $\mathcal{M}_1(\mathbf{Z})$ is compact. When this is the case, any distance metrizing the weak topology on $\mathcal{M}_1(\mathbf{Z})$ is complete.

Reciprocally, assume that $d_{\mathcal{F}}$ is a complete metric over $\mathcal{M}_1(\mathbf{Z})$ and write $\mathcal{F} = (f_m)_{m \in \mathbb{N}}$. Let $(\mu_n)_{n \in \mathbb{N}}$ be an arbitrary sequence of probability measures from $\mathcal{M}_1(\mathbf{Z})$. For every $m \in \mathbb{N}$, as f_m takes values in $[0, 1]$, we have for every $n \in \mathbb{N}$ that $\mu_n(f_m) \in [0, 1]$. Hence, using a diagonal

extraction, there exists a subsequence $(\mu_{n_k})_{k \in \mathbb{N}}$ of the sequence $(\mu_n)_{n \in \mathbb{N}}$ such that for every $m \in \mathbb{N}$, the sequence $(\mu_{n_k}(f_m))_{k \in \mathbb{N}}$ converges, that is, $(\mu_{n_k})_{k \in \mathbb{N}}$ is a Cauchy sequence for the distance $d_{\mathcal{F}}$. As we assumed the distance $d_{\mathcal{F}}$ to be complete, this implies that the sequence $(\mu_n)_{n \in \mathbb{N}}$ has a convergent subsequence. The sequence $(\mu_n)_{n \in \mathbb{N}}$ being arbitrary, we conclude that the space $\mathcal{M}_1(\mathbf{Z})$ is sequentially compact, and thus compact by Remark 2.6. \square

For $W \in \mathcal{W}_{\pm}$ and $f \in C_b(\mathbf{Z})$, we denote by $W[f]$ the real-valued kernel defined by:

$$(11) \quad W[f](x, y) = W(x, y; f) = \int_{\mathbf{Z}} f(z) W(x, y; dz).$$

We denote by $\|\cdot\|_{\square, \mathbb{R}}$ (resp. $\|\cdot\|_{\square, \mathbb{R}}^+$) the cut norm (resp. one-sided version of the cut norm) for real-valued kernels defined as:

$$(12) \quad \|w\|_{\square, \mathbb{R}} = \sup_{S, T \subset [0, 1]} \left| \int_{S \times T} w(x, y) dx dy \right| \quad \text{and} \quad \|w\|_{\square, \mathbb{R}}^+ = \sup_{S, T \subset [0, 1]} \int_{S \times T} w(x, y) dx dy,$$

where w is a real-valued kernel w (see [Lov12, Section 8.2], resp. [Lov12, Section 10.3], for definition and properties of those objects).

The following two remarks link the cut norm $\|\cdot\|_{\square, \mathcal{F}}$ of a signed measure-valued kernel W with the cut norm $\|\cdot\|_{\square, \mathbb{R}}$ of the real-valued kernels $W[f]$ for some particular choices of functions $f \in C_b(\mathbf{Z})$. We will reuse those facts in Section 6.

Remark 3.24 (Link between $\|\cdot\|_{\square, \mathcal{F}}$ and $\|\cdot\|_{\square, \mathbb{R}}^+$). For $\mu \in \mathcal{M}_{\pm}(\mathbf{Z})$ we have:

$$(13) \quad \|\mu\|_{\mathcal{F}} = \sup_{\varepsilon \in \{\pm 1\}^{\mathbb{N}}} \sum_{n \in \mathbb{N}} 2^{-n} \varepsilon_n \mu(f_n) = \sup_{\varepsilon \in \{\pm 1\}^{\mathbb{N}}} \mu \left(\sum_{n \in \mathbb{N}} 2^{-n} \varepsilon_n f_n \right),$$

with $\varepsilon = (\varepsilon_n)_{n \in \mathbb{N}}$. Hence, for a signed measure-valued kernel $W \in \mathcal{W}_{\pm}$, we have:

$$(14) \quad \|W\|_{\square, \mathcal{F}} = \sup_{\varepsilon \in \{\pm 1\}^{\mathbb{N}}} \sup_{S, T \subset [0, 1]} W \left(S \times T; \sum_{n \in \mathbb{N}} 2^{-n} \varepsilon_n f_n \right) = \sup_{\varepsilon \in \{\pm 1\}^{\mathbb{N}}} \left\| W \left[\sum_{n \in \mathbb{N}} 2^{-n} \varepsilon_n f_n \right] \right\|_{\square, \mathbb{R}}^+.$$

Remark 3.25 (Inequality with $\|\cdot\|_{\square, \mathcal{F}}$ and $\|\cdot\|_{\square, \mathbb{R}}$). For a signed measure-valued kernel W , we have:

$$(15) \quad \begin{aligned} \|W\|_{\square, \mathcal{F}} &= \sup_{S, T \subset [0, 1]} \sum_{n=0}^{\infty} 2^{-n} \left| \int_{S \times T} W(x, y, f_n) dx dy \right| \\ &\leq \sum_{n=0}^{\infty} 2^{-n} \sup_{S, T \subset [0, 1]} \left| \int_{S \times T} W(x, y, f_n) dx dy \right| \\ &= \sum_{n=0}^{\infty} 2^{-n} \|W[f_n]\|_{\square, \mathbb{R}}. \end{aligned}$$

4. TIGHTNESS AND WEAK REGULARITY

In this section, using a conditional expectation approach as in [Lov12, Chapter 9], we provide approximations of signed measure-valued kernels and probability-graphons by stepfunctions with an explicit bound on the quality of the approximation. This procedure takes into account that signed measure-valued kernels are infinite-dimensional valued.

4.1. Approximation by stepfunctions. We start by introducing the partitioning of a signed measure-valued kernel.

Definition 4.1 (The stepping operator). *Let $W \in \mathcal{W}_\pm$ be a signed measure-valued kernel and $\mathcal{P} = \{S_1, \dots, S_k\}$ be a finite partition of $[0, 1]$. We define the kernel stepfunction $W_{\mathcal{P}}$ adapted to the partition \mathcal{P} by averaging W over the partition subsets:*

$$W_{\mathcal{P}}(x, y; \cdot) = \frac{1}{\lambda(S_i)\lambda(S_j)} W(S_i \times S_j; \cdot) \quad \text{for } x \in S_i, y \in S_j,$$

when $\lambda(S_i) \neq 0$ and $\lambda(S_j) \neq 0$, and $W_{\mathcal{P}}(x, y; \cdot) = 0$ the null measure otherwise. We call the map $W \mapsto W_{\mathcal{P}}$ defined on \mathcal{W}_\pm the stepping operator (associated with the finite partition \mathcal{P}).

Since the signed measure-valued kernel are defined up to an a.e. equivalence, the value of $W_{\mathcal{P}}(x, y; \cdot)$ for $x \in S_i, y \in S_j$ when $\lambda(S_i)\lambda(S_j)$ is unimportant.

Remark 4.2 (Link with conditional expectation). The stepfunction $W_{\mathcal{P}}$ can be viewed as the conditional expectation of W w.r.t. the (finite) sigma-field $\sigma(\mathcal{P} \times \mathcal{P})$ on $[0, 1]^2$, where $W : [0, 1]^2 \rightarrow \mathcal{M}_\pm(\mathbf{Z})$ is seen as a random signed measure in $\mathcal{M}_\pm(\mathbf{Z})$ and the probability measure on $[0, 1]^2$ is the Lebesgue measure.

Remark 4.3 (Steppings are convex stable). Let $\mathcal{M} \subset \mathcal{M}_\pm(\mathbf{Z})$ be a convex subset of measures, for instance \mathcal{M} is $\mathcal{M}_1(\mathbf{Z})$, $\mathcal{M}_{\leq 1}(\mathbf{Z})$, $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_\pm(\mathbf{Z})$. Whenever $W \in \mathcal{W}_\pm$ is a \mathcal{M} -valued kernel, then by simple computation its stepping $W_{\mathcal{P}}$ is also a \mathcal{M} -valued kernel.

In the following remark, we give a characterization of refining partitions that generate the Borel σ -field of $[0, 1]$.

Remark 4.4 (On refining partitions that generates the Borel σ -field). Let $(\mathcal{P}_k)_{k \in \mathbb{N}}$ be a sequence of refining partitions of $[0, 1]$. It generates the Borel σ -field of $[0, 1]$ (that is, $\{S : S \in \mathcal{P}_k, k \in \mathbb{N}\}$ generates the Borel σ -field of $[0, 1]$) if and only if $(\mathcal{P}_k)_{k \in \mathbb{N}}$ separates points (that is, for every distinct $x, y \in [0, 1]$, there exists $k \in \mathbb{N}$ such that x and y belong to different classes of \mathcal{P}_k).

Indeed, assume that $(\mathcal{P}_k)_{k \in \mathbb{N}}$ separates points, and consider the countable family of Borel-measurable functions $\mathcal{F} = \{\mathbf{1}_S : S \in \mathcal{P}_k, k \in \mathbb{N}\}$ which separates points. Thus, by [Bog07b, Theorem 6.8.9] (remark that a Polish space is a Souslin space, see [Bog07b, Definition 6.6.1]), the family \mathcal{F} generates the Borel σ -field of $[0, 1]$. This implies that the family of Borel sets $\{S : S \in \mathcal{P}_k, k \in \mathbb{N}\}$ generates the Borel σ -field of $[0, 1]$.

Conversely, assume there exist $x, y \in [0, 1]$ which are not separated by $(\mathcal{P}_k)_{k \in \mathbb{N}}$, i.e. for all $k \in \mathbb{N}$, x and y belong to the same class of \mathcal{P}_k . This implies that the set $\{x\}$ does not belong to the σ -field generated by $(\mathcal{P}_k)_{k \in \mathbb{N}}$, and thus $(\mathcal{P}_k)_{k \in \mathbb{N}}$ does not generate the Borel σ -field of $[0, 1]$.

Recall the definition of the norm $\|\cdot\|_\infty$ on \mathcal{W}_\pm defined in (2). The following lemma allows to approximate any signed measure-valued kernel by its steppings.

Lemma 4.5 (Approximation using the stepping operator). *Let $W \in \mathcal{W}_\pm$ be a signed measure-valued kernel (which is bounded by definition). Let $(\mathcal{P}_n)_{n \in \mathbb{N}}$ be a refining sequence of finite partitions of $[0, 1]$ that generates the Borel σ -field on $[0, 1]$. Then, the sequence $(W_{\mathcal{P}_n})_{n \in \mathbb{N}}$ is uniformly bounded by $\|W\|_\infty$, and weakly converges to W almost everywhere (on $[0, 1]^2$).*

Proof. Set $W_n = W_{\mathcal{P}_n}$ for $n \in \mathbb{N}$. By definition of the stepping operator, we have for every $n \in \mathbb{N}$ and every $(x, y) \in [0, 1]^2$ that the total mass of $W_n(x, y; \cdot)$ is upper bounded by $\|W\|_\infty$.

Recall that for $W \in \mathcal{W}_\pm$ and $f \in C_b(\mathbf{Z})$, the real-valued kernel $W[f]$ is defined by (11). First assume that $W \in \mathcal{W}_+$. Let $\mathcal{F} = (f_k)_{k \in \mathbb{N}}$ be a convergence determining sequence, with by convention $f_0 = \mathbf{1}$. For every $k \in \mathbb{N}$ and $n \in \mathbb{N}$, an immediate computation gives $W_n[f_k] = (W[f_k])_{\mathcal{P}_n}$. For every $k \in \mathbb{N}$, as $W[f_k]$ is a real-valued kernel, we can apply the closed martingale

theorem (as $(W[f_k])_{\mathcal{P}_n}$ can be viewed as a conditional expectation, see Remark 4.2), and we get that $\lim_{n \rightarrow \infty} W_n[f_k] = W[f_k]$ almost everywhere, since $(\mathcal{P}_n)_{n \in \mathbb{N}}$ generates the Borel σ -field. Hence, as the sequence $(f_k)_{k \in \mathbb{N}}$ is convergence determining, the sequence $(W_n)_{n \in \mathbb{N}}$ weakly converges to W almost everywhere.

Now, for $W \in \mathcal{W}_{\pm}$, write $W = W^+ - W^-$ where $W^+, W^- \in \mathcal{W}_+$ (see Lemma 3.3). By linearity of the stepping operator, remark that we have $W_n = (W^+)_{\mathcal{P}_n} - (W^-)_{\mathcal{P}_n}$ for all $n \in \mathbb{N}$. By the first case, we have that the sequence $((W^+)_{\mathcal{P}_n})_{n \in \mathbb{N}}$ weakly converges a.e. to W^+ , and similarly for $((W^-)_{\mathcal{P}_n})_{n \in \mathbb{N}}$ and W^- . Hence, the sequence $(W_n)_{n \in \mathbb{N}}$ weakly converges to W almost everywhere. \square

We first provide a separability result on the space of probability-graphons.

Proposition 4.6 (Separability of \mathcal{W}_1 and $\widetilde{\mathcal{W}}_1$). *Let d be a smooth distance on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}). Then, the space (\mathcal{W}_1, d) (resp. (\mathcal{W}_+, d) or (\mathcal{W}_{\pm}, d)) is separable.*

If furthermore d is invariant (which implies that δ_{\square} is a distance), then the space $(\widetilde{\mathcal{W}}_1, \delta_{\square})$ (resp. $(\widetilde{\mathcal{W}}_+, \delta_{\square})$ or $(\widetilde{\mathcal{W}}_{\pm}, \delta_{\square})$) is separable.

Proof. We shall consider the space of probability-graphons \mathcal{W}_1 , as the proofs for \mathcal{W}_+ and \mathcal{W}_{\pm} are similar. Applying Lemma 4.5 with the sequence of dyadic partitions, for every probability-graphon W , we can find a sequence of probability-graphon stepfunctions adapted to finite dyadic partitions and converging to W almost everywhere on $[0, 1]^2$.

As the space \mathbf{Z} is separable, the space of probability measures $\mathcal{M}_1(\mathbf{Z})$ is also separable for the weak topology (see [Bil99, Theorem 6.8]). Let $\mathcal{A} \subset \mathcal{M}_1(\mathbf{Z})$ be an at most countable dense (for the weak topology) subset. Then, for any stepfunction $W \in \mathcal{W}_1$ adapted to a finite dyadic partition, we can approach it everywhere on $[0, 1]^2$ by a sequence of \mathcal{A} -valued stepfunctions adapted to the same finite dyadic partition.

Hence, for every $W \in \mathcal{W}_1$, there exists a sequence $(W_n)_{n \in \mathbb{N}}$ in the countable set of \mathcal{A} -valued stepfunctions adapted to a finite dyadic partition that converges to W almost everywhere on $[0, 1]^2$. As d is smooth, we get that this convergence also holds for d . Thus, the space (\mathcal{W}_1, d) is complete.

Remind that by Theorem 3.17, when the distance d is invariant and smooth, then the premetric δ_{\square} is a distance on $\widetilde{\mathcal{W}}_1$. In that case, convergence for d implies convergence for δ_{\square} , and thus the space $(\widetilde{\mathcal{W}}_1, \delta_{\square})$ is also separable. \square

4.2. Tightness. Similarly to the case of signed measures (remind Lemma 2.8), we introduce a tightness criterion for signed measure-valued kernels that characterizes relative compactness, see Proposition 4.8 below. For a signed measure-valued kernel $W \in \mathcal{W}_{\pm}$, we define the measure $M_W \in \mathcal{M}_+(\mathbf{Z})$ by:

$$(16) \quad M_W(dz) = |W|([0, 1]^2; dz) = \int_{[0, 1]^2} |W|(x, y; dz) \, dx dy,$$

where for every $x, y \in [0, 1]$, $|W|(x, y; \cdot)$ is the total variation of $W(x, y; \cdot)$ (see Lemma 3.3). In particular, if W is a probability-graphon then M_W is a probability measure from $\mathcal{M}_1(\mathbf{Z})$. Notice also that if W and U are weakly isomorphic, then $M_W = M_U$, so that the application $W \mapsto M_W$ can be seen as a map from $\widetilde{\mathcal{W}}_1$ (resp. $\widetilde{\mathcal{W}}_{\pm}$) to $\mathcal{M}_1(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$).

Definition 4.7 (Tightness criterion). *A subset $\mathcal{K} \subset \mathcal{W}_{\pm}$ (resp. $\mathcal{K} \subset \widetilde{\mathcal{W}}_{\pm}$) is said to be tight if the subset of measures $\{M_W : W \in \mathcal{K}\} \subset \mathcal{M}_+(\mathbf{Z})$ is tight.*

The following proposition shows the equivalence between a global tightness criterion and a local tightness criterion. Recall that uniformly bounded subsets of $\widetilde{\mathcal{W}}_{\pm}$ are discussed after Definition 3.16. Recall also λ_2 is the Lebesgue measure on $[0, 1]^2$.

Proposition 4.8 (Alternative tightness criterion). *Let $\mathcal{K} \subset \mathcal{W}_\pm$ (or $\mathcal{K} \subset \widetilde{\mathcal{W}}_\pm$) be a uniformly bounded subset of signed measure-valued kernels. The set \mathcal{K} is tight if and only if for every $\varepsilon > 0$, there exists a compact set $K \subset \mathbf{Z}$, such that for every $W \in \mathcal{K}$ we have:*

$$(17) \quad \lambda_2\left(\{(x, y) \in [0, 1]^2 : |W|(x, y; K^c) \leq \varepsilon\}\right) > 1 - \varepsilon.$$

Proof. As the left hand side of (17) is invariant by relabeling, it is enough to do the proof for \mathcal{W}_\pm . Let $\mathcal{K} \subset \mathcal{W}_\pm$ be uniformly bounded and set $C = \sup_{W \in \mathcal{K}} \|W\|_\infty < \infty$. Assume that for every $\varepsilon > 0$, there exists a compact set $K \subset \mathbf{Z}$, such that (17) holds for every $W \in \mathcal{K}$. Let $1 > \varepsilon > 0$. Thus, there exists a compact subset $K \subset \mathbf{Z}$ such that for every $W \in \mathcal{K}$ there exists a subset $A_W \subset [0, 1]^2$ with (Lebesgue) measure at least $1 - \varepsilon$, such that for every $(x, y) \in A_W$, we have $|W|(x, y; K^c) \leq \varepsilon$. We have that for all $W \in \mathcal{K}$:

$$M_W(K^c) = \int_{[0, 1]^2} |W|(x, y; K^c) \, dx dy \leq \|W\|_\infty \lambda_2(A_W^c) + \varepsilon \lambda_2(A_W) \leq (C + 1)\varepsilon.$$

Hence, the subset of measures $\{M_W : W \in \mathcal{K}\} \subset \mathcal{M}_+(\mathbf{Z})$ is tight, that is \mathcal{K} is tight.

Conversely, suppose that \mathcal{K} is tight. Let $\varepsilon > 0$. There exists a compact set $K \subset \mathbf{Z}$ such that for every $W \in \mathcal{K}$, we have $M_W(K^c) < \varepsilon^2$. For $W \in \mathcal{K}$, define $A_W = \{(x, y) \in [0, 1]^2 : |W|(x, y; K^c) \leq \varepsilon\}$. We have:

$$\varepsilon^2 > M_W(K^c) = \int_{[0, 1]^2} |W|(x, y; K^c) \, dx dy \geq \varepsilon \lambda_2(A_W^c).$$

Hence, $\lambda_2(A_W) > 1 - \varepsilon$, and consequently Equation (17) holds. \square

We end this section on a continuity result of the map $W \mapsto M_W$.

Lemma 4.9 (Regularity of the map $W \mapsto M_W$). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$). Then the map $W \mapsto M_W$ is 1-Lipschitz, and thus continuous, from $(\widetilde{\mathcal{W}}_{1,m}, \delta_{\square,m})$ (resp. $(\widetilde{\mathcal{W}}_{+,m}, \delta_{\square,m})$) to $(\mathcal{M}_1(\mathbf{Z}), d_m)$ (resp. $(\mathcal{M}_+(\mathbf{Z}), d_m)$).*

Proof. Taking $S = T = [0, 1]$ in Definition (3) of $d_{\square,m}$, we get that $d_m(M_U, M_W) \leq d_{\square,m}(W, U)$. As $M_{U\varphi} = M_U$ for any measure-preserving map φ thanks to (1), we deduce from Definition (6) of $\delta_{\square,m}$ that $d_m(M_U, M_W) \leq \delta_{\square,m}(U, W)$. \square

4.3. Weak regularity. We shall consider the following extra regularities of distances on the set of signed measure-valued kernels w.r.t. the stepping operator. For a finite partition \mathcal{P} , denote by $|\mathcal{P}|$ the size of the partition \mathcal{P} , i.e. the number of sets composing \mathcal{P} .

Definition 4.10 (Regularities of distances). *Let d be a distance on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_\pm).*

- (i) **Weak regularity.** *The distance d is weakly regular if whenever the subset \mathcal{K} of \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_\pm) is tight (resp. tight and uniformly bounded), then for every $\varepsilon > 0$, there exists $m \in \mathbb{N}^*$, such that for every kernel $W \in \mathcal{K}$, and for every finite partition \mathcal{Q} of $[0, 1]$, there exists a finite partition \mathcal{P} of $[0, 1]$ that refines \mathcal{Q} such that:*

$$|\mathcal{P}| \leq m|\mathcal{Q}| \quad \text{and} \quad d(W, W_{\mathcal{P}}) < \varepsilon.$$

- (ii) **Regularity w.r.t. the stepping operator.** *The distance d is regular w.r.t. the stepping operator if (resp. for any finite constant $C \geq 0$) there exists a finite constant $C_0 > 0$ such that for every W, U in \mathcal{W}_1 (resp. in \mathcal{W}_+ or \mathcal{W}_\pm , with $\|W\|_\infty \leq C$ and $\|U\|_\infty \leq C$) and every finite partition \mathcal{P} of $[0, 1]$, then we have:*

$$(18) \quad d(W, W_{\mathcal{P}}) \leq C_0 d(W, U_{\mathcal{P}}).$$

We say that a norm N on \mathcal{W}_\pm is weakly regular (resp. regular w.r.t. the stepping operator) if its associated distance d on \mathcal{W}_\pm is weakly regular (resp. regular w.r.t. the stepping operator).

The weak regularity property is an analogue to the weak regularity lemma for real-valued graphons (see [Lov12, Lemma 9.15]). If a distance d is weakly regular, then for a subset $\mathcal{K} \subset \mathcal{M}_{\pm}(\mathbf{Z})$ which is tight and uniformly bounded, every \mathcal{K} -valued kernel can be approximated by a stepfunction with a uniform bound. The regularity w.r.t. the stepping operator states that the stepping operator gives an almost optimal way to approximate a signed measure-valued kernel using stepfunctions adapted to a given partition.

4.3.1. *An example of cut distance regular w.r.t. the stepping operator.* Remind the definition of a quasi-convex distance in Definition 2.10. We first show that the stepping operator is 1-Lipschitz for the cut distance $d_{\square, m}$ when the distance d_m is quasi-convex.

Lemma 4.11 (The stepping operator is 1-Lipschitz). *Let d_m be a quasi-convex distance on \mathcal{M} a convex subset of $\mathcal{M}_{\pm}(\mathbf{Z})$ containing the zero measure. Then, the stepping operator associated with a given finite partition of $[0, 1]$ is 1-Lipschitz on $\mathcal{W}_{\mathcal{M}}$ for the cut distance $d_{\square, m}$.*

Proof. Let $U, W \in \mathcal{W}_{\mathcal{M}}$ be \mathcal{M} -valued kernels, and let \mathcal{P} be a finite measurable partition of $[0, 1]$. As $U_{\mathcal{P}}$ and $W_{\mathcal{P}}$ are stepfunctions adapted to the same partition, and as d_m is quasi-convex, we can use Lemma 3.19 to get for some $S, T \in \sigma(\mathcal{P})$ that:

$$d_{\square, m}(U_{\mathcal{P}}, W_{\mathcal{P}}) = d_m(U_{\mathcal{P}}(S \times T; \cdot), W_{\mathcal{P}}(S \times T; \cdot)) = d_m(U(S \times T; \cdot), W(S \times T; \cdot)) \leq d_{\square, m}(U, W),$$

where the second equality comes from the fact that the integrals are equal as $S, T \in \sigma(\mathcal{P})$ and thus the integration is over full steps of the partition. Hence, the stepping operator is 1-Lipschitz on $\mathcal{W}_{\mathcal{M}}$ for the cut distance d_m . \square

For a quasi-convex distance d_m , the cut distance $d_{\square, m}$ is regular w.r.t. the stepping operator with $C_0 = 2$ in (18) (and one can take $C = +\infty$ in Definition 4.10 (ii)).

Lemma 4.12 ($d_{\square, m}$ is regular w.r.t. the stepping operator). *Let d_m be a quasi-convex distance on \mathcal{M} a convex subset of $\mathcal{M}_{\pm}(\mathbf{Z})$ containing the zero measure. Let $W, U \in \mathcal{W}_{\epsilon}$ be $\mathcal{W}_{\mathcal{M}}$ -valued kernels, and let \mathcal{P} be a finite partition of $[0, 1]$. Then, we have:*

$$d_{\square, m}(W, W_{\mathcal{P}}) \leq 2d_{\square, m}(W, U_{\mathcal{P}}).$$

Proof. The proof is similar to the proof of [Lov12, Lemma 9.12]. As d_m is quasi-convex, using Lemma 4.11, we get:

$$d_{\square, m}(W, W_{\mathcal{P}}) \leq d_{\square, m}(W, U_{\mathcal{P}}) + d_{\square, m}(U_{\mathcal{P}}, W_{\mathcal{P}}) \leq 2d_{\square, m}(W, U_{\mathcal{P}}).$$

\square

4.3.2. *An example of weakly regular cut distance.* We have the following general result. Recall Definitions 3.10 and 4.10 on distances and norms on \mathcal{W}_{ϵ} , with $\epsilon \in \{+, \pm\}$, being invariant, smooth, weakly regular and regular w.r.t. the stepping operator.

Proposition 4.13 (Weak regularity of $d_{\square, m}$). *Let d_m be a quasi-convex distance on $\mathcal{M}_{\epsilon}(\mathbf{Z})$, with $\epsilon \in \{+, \pm\}$, which is sequentially continuous w.r.t. the weak topology. Then, the cut distance $d_{\square, m}$ on \mathcal{W}_{ϵ} is invariant, smooth, weakly regular and regular w.r.t. the stepping operator.*

Using results from Section 3.8, we directly get the following weak regularity of the cut distance $d_{\square, \mathcal{P}}$ and the cut norms $\|\cdot\|_{\square, \mathcal{F}}$, $\|\cdot\|_{\square, KR}$ and $\|\cdot\|_{\square, FM}$.

Corollary 4.14 (Weak regularity of usual distances and norms). *The cut norms $\|\cdot\|_{\square, \mathcal{F}}$, $\|\cdot\|_{\square, KR}$ and $\|\cdot\|_{\square, FM}$ (resp. the cut distance $d_{\square, \mathcal{P}}$) on \mathcal{W}_{\pm} (resp. \mathcal{W}_{+}) are invariant, smooth, weakly regular and regular w.r.t. the stepping operator.*

Proof of Proposition 4.13. We deduce from Lemmas 3.11 and 4.12, Proposition 3.13 and that the cut distance $d_{\square, m}$ on \mathcal{W}_ϵ is invariant, smooth and regular w.r.t. the stepping operator. We are left to prove that $d_{\square, m}$ is weakly regular on \mathcal{W}_ϵ . We prove it by considering in the first step the case \mathbf{Z} compact and in a second step the general case \mathbf{Z} Polish.

Step 1. We assume \mathbf{Z} compact. As in the definition of weak regularity, let $\mathcal{K} \subset \mathcal{W}_\epsilon$ be a subset of $\mathcal{M}_\epsilon(\mathbf{Z})$ -valued kernels that is tight and uniformly bounded by some finite constant C . Let $\mathcal{M} \subset \mathcal{M}_\epsilon(\mathbf{Z})$ be the subset of elements of $\mathcal{M}_\epsilon(\mathbf{Z})$ with total mass at most C ; in particular \mathcal{M} is a convex set containing 0 and $\mathcal{K} \subset \mathcal{W}_\mathcal{M}$. As \mathbf{Z} is compact, from Remarks 2.6 and 2.7, we know that the weak topology is metrizable on \mathcal{M} and that \mathcal{M} is compact, and thus sequentially weakly compact. Hence, as d_m is sequentially continuous w.r.t. the weak topology on $\mathcal{M}_\epsilon(\mathbf{Z})$, we have that (\mathcal{M}, d_m) is sequentially compact, and thus compact.

Denote by $B(\mu, r) = \{\nu \in \mathcal{M} : d_m(\mu, \nu) < r\}$ the open ball centered at $\mu \in \mathcal{M}$ with radius $r > 0$. Let $\varepsilon > 0$. As \mathcal{M} is compact, there exist $\mu_1, \dots, \mu_n \in \mathcal{M}$, $n \in \mathbb{N}^*$, such that $\mathcal{M} = \cup_{i=1}^n B(\mu_i, \varepsilon)$. For $1 \leq i \leq n$, define $A_i = B(\mu_i, \varepsilon) \setminus \cup_{j < i} B(\mu_j, \varepsilon)$, so that $\{A_1, \dots, A_n\}$ is a finite partition (with possibly some empty sets) of \mathcal{M} .

Every \mathcal{M} -valued kernel W can be approximated by a $\{\mu_1, \dots, \mu_n\}$ -valued kernel U defined for every $(x, y) \in [0, 1]^2$ by $U(x, y; \cdot) = \mu_i$ for i such that $W(x, y; \cdot) \in A_i$. Thus, by construction, we have that for every $(x, y) \in [0, 1]^2$, $d_m(W(x, y; \cdot), U(x, y; \cdot)) < \varepsilon$. Applying the quasi-convex supremum inequality from (5) to W and U , we get that:

$$d_{\square, m}(W, U) \leq \operatorname{esssup}_{(x, y) \in [0, 1]^2} d_m(W(x, y; \cdot), U(x, y; \cdot)) \leq \varepsilon.$$

Then, as the stepping operator is 1-Lipschitz for the cut norm, see Lemma 4.11, we have for any finite partition \mathcal{P} of $[0, 1]$ that:

$$(19) \quad \begin{aligned} d_{\square, m}(W, W_{\mathcal{P}}) &\leq d_{\square, m}(W, U) + d_{\square, m}(U, U_{\mathcal{P}}) + d_{\square, m}(U_{\mathcal{P}}, W_{\mathcal{P}}) \\ &\leq 2\varepsilon + d_{\square, m}(U, U_{\mathcal{P}}). \end{aligned}$$

Hence, to get the weak regularity property for \mathcal{M} -valued kernels, we are left to prove it for the much smaller set of \mathcal{V} -valued kernels, where \mathcal{V} is the convex hull of $\{\mu_1, \dots, \mu_n\}$.

As d_m is quasi-convex and sequentially continuous w.r.t. the weak topology, using Lemma 2.11, there exists $\eta > 0$ such that for all $\mu, \nu \in \mathcal{M}_\epsilon(\mathbf{Z})$, we have that $\|\mu - \nu\|_\infty < \eta$ implies that $d_m(\mu, \nu) \leq \varepsilon$.

As \mathcal{V} is a subset of a vector space with finite dimension n , the norm $\|\cdot\|_\infty$ seen over \mathcal{V} is equivalent to the L_1 -norm $\mu = \sum_{i=1}^n \alpha_i \mu_i \mapsto \|\alpha\|_1 = \sum_{i=1}^n |\alpha_i|$. We can now see \mathcal{V} -valued kernel as \mathbb{R}^n -valued graphon with a cut norm derived from the L_1 -norm $\|\cdot\|_1$, and in this case the proof for the weak regularity Lemma 9.9 in [Lov12] can easily be adapted. Hence, we have the weak regularity property for \mathcal{V} -valued kernels: there exists $m \in \mathbb{N}^*$, such that for every \mathcal{V} -valued kernel U' , and for every finite partition \mathcal{Q} of $[0, 1]$ there exists a finite partition \mathcal{P} of $[0, 1]$ that refines \mathcal{Q} , and such that $|\mathcal{P}| \leq m|\mathcal{Q}|$ and $\sup_{S, T \subset [0, 1]} \|(U' - U'_{\mathcal{P}})(S \times T; \cdot)\|_\infty < \eta$, and thus $d_{\square, m}(U', U'_{\mathcal{P}}) \leq \varepsilon$.

Taking $U' = U$ in (19), we get that $d_{\square, m}(W, W_{\mathcal{P}}) \leq 3\varepsilon$ and $|\mathcal{P}| \leq m|\mathcal{Q}|$. This concludes the proof of the lemma when \mathbf{Z} is compact.

Step 2. We consider the general case \mathbf{Z} Polish. We now prove that $d_{\square, m}$ is weakly regular on \mathcal{W}_ϵ . Let $\mathcal{K} \subset \mathcal{W}_\epsilon$ be a subset of $\mathcal{M}_\epsilon(\mathbf{Z})$ -valued kernels that is tight and uniformly bounded, and denote by $C = \sup_{W \in \mathcal{K}} \|W\|_\infty < \infty$.

Let $\varepsilon > 0$. As d_m is quasi-convex and sequentially continuous w.r.t. the weak topology, using Lemma 2.11, there exists $\eta > 0$ such that for all $\mu, \nu \in \mathcal{M}_\epsilon(\mathbf{Z})$, we have that $\|\mu - \nu\|_\infty < \eta$ implies that $d_m(\mu, \nu) < \varepsilon$. Without loss of generality, we assume that $\eta \leq \varepsilon$. Let $\eta_C = \min(\eta, \eta/C)$.

As \mathcal{K} is tight, using Proposition 4.8, there exists a compact set $K \subset \mathbf{Z}$, such that for every $W \in \mathcal{K}$ the subset $A_W = \{(x, y) \in [0, 1]^2 : |W|(x, y; K^c) \leq \eta_C/2\}$ has Lebesgue measure at least $1 - \eta_C/2$. Let $W \in \mathcal{K}$, and define the signed measure-valued kernel U by: $U(x, y; \cdot) = W(x, y; \cdot \cap K)$ for every $(x, y) \in A_W$, and $U(x, y; \cdot) = 0$ otherwise. Let $S, T \subset [0, 1]$. We have:

$$\begin{aligned} \|(W - U)(S \times T; \cdot)\|_\infty &\leq \int_{S \times T} \|W(x, y; \cdot) - U(x, y; \cdot)\|_\infty \, dx dy \\ &\leq \int_{A_W \cap (S \times T)} |W|(x, y; K^c) \, dx dy + \int_{A_W^c \cap (S \times T)} \|W(x, y; \cdot)\|_\infty \, dx dy \\ &\leq \eta_C/2 + C \cdot \eta_C/2 \\ &\leq \eta. \end{aligned}$$

Thus, we have that $d_m(W(S \times T; \cdot), U(S \times T; \cdot)) < \varepsilon$. Since this holds for all $S, T \subset [0, 1]$, we get that $d_{\square, m}(W, U) \leq \varepsilon$.

Notice that the $\mathcal{M}_\pm(\mathbf{Z})$ -valued kernel U is also a $\mathcal{M}_\pm(K)$ -valued kernel, where $K \subset \mathbf{Z}$ is a compact set, and that $\|U\|_\infty \leq \|W\|_\infty \leq C$. Further remark that, using Lemma 4.11, for every $W \in \mathcal{K}$ and every finite partition \mathcal{P} of $[0, 1]$, we have that:

$$\begin{aligned} d_{\square, m}(W, W_{\mathcal{P}}) &\leq d_{\square, m}(W, U) + d_{\square, m}(U, U_{\mathcal{P}}) + d_{\square, m}(U_{\mathcal{P}}, W_{\mathcal{P}}) \\ &\leq 2\varepsilon + d_{\square, m}(U, U_{\mathcal{P}}). \end{aligned}$$

Hence, to get the weak regularity property for $d_{\square, m}$ on \mathcal{K} (see Definition 4.10 (i)), it is enough to prove that $d_{\square, m}$ restricted to $\mathcal{M}_\varepsilon(K)$ -valued kernels is weakly regular, which is true by Step 1. As a consequence, we get that $d_{\square, m}$ on \mathcal{W}_ε is weakly regular. \square

4.4. A stronger weak regularity lemma for $d_{\square, \mathcal{F}}$. In this subsection, we prove a stronger version of the weak regularity lemma for the special case of the cut distance $d_{\square, \mathcal{F}}$. We shall use this result for the proof of the second sampling Lemma 6.12.

Let $\mathcal{F} = (f_n)_{n \in \mathbb{N}}$, with $f_0 = \mathbf{1}$ and f_n takes values in $[0, 1]$, be a convergence determining sequence, which is assumed fixed in this section.

4.4.1. Comparison between $\|\cdot\|_{\square, \mathcal{F}}$ and an euclidian norm. To better understand the stepping operator, we introduce a scalar product over signed measure-valued kernels. The link between this scalar product and the norm $\|\cdot\|_{\square, \mathcal{F}}$ is given by Lemma 4.15. We define the scalar product $\langle \cdot, \cdot \rangle_{\mathcal{F}}$ on signed measure-valued kernels for $U, W \in \mathcal{W}_\pm$ by:

$$\langle U, W \rangle_{\mathcal{F}} = \sum_{n \geq 0} 2^{-n} \langle U[f_n], W[f_n] \rangle,$$

where for all n the scalar product taken for $U[f_n]$ and $W[f_n]$ is the usual scalar product in $L^2([0, 1]^2, \lambda_2)$ for real-valued kernels:

$$\langle U[f_n], W[f_n] \rangle = \int_{[0, 1]^2} U[f_n](x, y) W[f_n](x, y) \, dx dy.$$

The scalar product $\langle \cdot, \cdot \rangle_{\mathcal{F}}$ induces a norm on \mathcal{W}_\pm which we denote by $\|\cdot\|_{2, \mathcal{F}}$.

Let \mathcal{P} be a finite partition of $[0, 1]$. As the stepping operator for measurable real-valued L^2 functions on $[0, 1]^2$ is a linear projection, and is idempotent and symmetric, and by definition of the scalar product $\langle \cdot, \cdot \rangle_{\mathcal{F}}$ for signed measure-valued kernels, we have that the stepping operator for signed measure-valued kernels is linear, idempotent and symmetric for $\langle \cdot, \cdot \rangle_{\mathcal{F}}$. Moreover, the stepping operator is the orthogonal projection for $\langle \cdot, \cdot \rangle_{\mathcal{F}}$ onto the space of stepfunctions with steps in \mathcal{P} .

Note that for a probability-graphon $W \in \mathcal{W}_1$, we have $\|W\|_{2, \mathcal{F}} \leq \sqrt{2}$ as each f_n takes values in $[0, 1]$. The following technical lemma gives a comparison between $\|\cdot\|_{\square, \mathcal{F}}$ and $\|\cdot\|_{2, \mathcal{F}}$.

Lemma 4.15 (Comparison between $\|\cdot\|_{\square, \mathcal{F}}$ and $\|\cdot\|_{2, \mathcal{F}}$). *For a signed measure-valued kernel $W \in \mathcal{W}_{\pm}$, we have $\|W\|_{\square, \mathcal{F}} \leq \sqrt{2}\|W\|_{2, \mathcal{F}}$.*

Proof. Let $S, T \subset [0, 1]$ be measurable subsets. By the Cauchy-Schwarz inequality, we have $|\langle W[f_n], \mathbf{1}_{S \times T} \rangle|^2 \leq \|W[f_n]\|_2^2 = \langle W[f_n], W[f_n] \rangle$ for every $n \geq 0$. Using this inequality along with Jensen's inequality, we get for every $S, T \subset [0, 1]$ that:

$$\begin{aligned} \left(\sum_{n \geq 0} 2^{-n} |W(S \times T, f_n)| \right)^2 &= \left(\sum_{n \geq 0} 2^{-n} \langle W[f_n], \mathbf{1}_{S \times T} \rangle \right)^2 \\ &\leq \sum_{n \geq 0} 2^{-n+1} |\langle W[f_n], \mathbf{1}_{S \times T} \rangle|^2 \\ &\leq \sum_{n \geq 0} 2^{-n+1} \langle W[f_n], W[f_n] \rangle \\ &= 2(\|W\|_{2, \mathcal{F}})^2. \end{aligned}$$

Taking the supremum over every measurable subsets $S, T \subset [0, 1]$ gives the desired inequality. \square

4.4.2. *The weak regularity lemma for $\|\cdot\|_{\square, \mathcal{F}}$.* The following lemma gives an explicit bound on the approximation of a signed measure-valued kernel, say W , by its steppings $W_{\mathcal{P}}$, with \mathcal{P} a finite partition on $[0, 1]$. Its proof is a straightforward adaptation of the proof of the weak regularity lemma for real-valued graphons in [Lov12, Lemma 9.9].

Lemma 4.16 (Weak regularity lemma for $\|\cdot\|_{\square, \mathcal{F}}$, simple formulation). *For every signed measure-valued kernel $W \in \mathcal{W}_{\pm}$ and $k \geq 1$, there exists a finite partition \mathcal{P} of $[0, 1]$ such that $|\mathcal{P}| = k$ and:*

$$\|W - W_{\mathcal{P}}\|_{\square, \mathcal{F}} \leq \frac{\sqrt{8}}{\sqrt{\log(k)}} \|W\|_{2, \mathcal{F}}.$$

In particular, if $W \in \mathcal{W}_1$ is a probability-graphon, (as $\|W\|_{2, \mathcal{F}} \leq \sqrt{2}$) we have:

$$\|W - W_{\mathcal{P}}\|_{\square, \mathcal{F}} \leq \frac{4}{\sqrt{\log(k)}}.$$

It is possible in the weak regularity lemma to ask for extra requirements, for instance to start from an already existing partition, or to ask the partition to be balanced, as stated in the following lemma. The proof is a straightforward adaptation of the proof of [Lov12, Lemma 9.15].

Lemma 4.17 (Weak regularity lemma for $\|\cdot\|_{\square, \mathcal{F}}$, with extra requirements). *Let $W \in \mathcal{W}_1$ be a probability-graphon, and let $1 \leq m < k$.*

(i) *For every partition \mathcal{Q} of $[0, 1]$ into m classes, there is a partition \mathcal{P} with k classes refining \mathcal{Q} and such that:*

$$\|W - W_{\mathcal{P}}\|_{\square, \mathcal{F}} \leq \frac{4}{\sqrt{\log(k/m)}}.$$

(ii) *For every partition \mathcal{Q} of $[0, 1]$ into m classes, there is an equipartition (i.e. a finite partition into classes with the same measure) \mathcal{P} of $[0, 1]$ into k classes and such that:*

$$\|W - W_{\mathcal{P}}\|_{\square, \mathcal{F}} \leq 2\|W - W_{\mathcal{Q}}\|_{\square, \mathcal{F}} + \frac{2m}{k}.$$

5. COMPACTNESS AND COMPLETENESS OF \mathcal{W}_1

In Section 5.1, we link the tightness criterion for measure-valued kernels with the relative compactness w.r.t. the cut distance $\delta_{\square, m}$. In Section 5.2, we compare the topologies induced by the cut distance $\delta_{\square, m}$ for different choice of the distance d_m , and state that under some conditions on d_m , those topologies coincide. In Section 5.3, we investigate the completeness of \mathcal{W}_1 endowed with the cut distance $\delta_{\square, m}$ and prove that the space of probability-graphons $\widetilde{\mathcal{W}}_1$ is a Polish space (Theorem 5.10), and that it is compact if and only if \mathbf{Z} is compact (Corollary 5.13). The technical proofs are postponed to Section 8.

5.1. Tightness criterion and compactness. Let $\mathcal{M} \subset \mathcal{M}_{\pm}(\mathbf{Z})$ be a subset of signed measures on \mathbf{Z} . Recall that $\mathcal{W}_{\mathcal{M}} \subset \mathcal{W}_{\pm}$ denote the subset of signed measure-valued kernels which are \mathcal{M} -valued. In this section, we shall denote by $\widetilde{\mathcal{W}}_{\mathcal{M}}$ the quotient of $\mathcal{W}_{\mathcal{M}}$ identifying signed measure-valued kernels that are weakly isomorphic.

Remind from Definition 3.15 and Theorem 3.17 that for an invariant, smooth and weakly regular distance d on \mathcal{W}_1 (resp. \mathcal{W}_+ , \mathcal{W}_{\pm}), δ_{\square} is defined as $\delta_{\square}(U, W) = \inf_{\varphi \in S_{[0,1]}} d(U, W^{\varphi})$, and is a distance on $\widetilde{\mathcal{W}}_1$ (resp. $\widetilde{\mathcal{W}}_+$, $\widetilde{\mathcal{W}}_{\pm}$).

We are now ready to formulate the important following theorem, which relates tightness with compactness and convergence for signed measure-valued kernels. We prove this theorem in Section 8.

Theorem 5.1 (Compactness theorem for $\widetilde{\mathcal{W}}_1$). *Let d be an invariant, smooth and weakly regular distance on \mathcal{W}_1 (resp. \mathcal{W}_{\pm}).*

- (i) *If a sequence of elements of \mathcal{W}_1 or $\widetilde{\mathcal{W}}_1$ (resp. \mathcal{W}_{\pm} or $\widetilde{\mathcal{W}}_{\pm}$) is tight (resp. tight and uniformly bounded), then it has a subsequence converging for δ_{\square} .*
- (ii) *If $\mathcal{M} \subset \mathcal{M}_1(\mathbf{Z})$ (resp. $\mathcal{M} \subset \mathcal{M}_{\pm}(\mathbf{Z})$) is convex and compact (resp. sequentially compact) for the weak topology, then the space $(\widetilde{\mathcal{W}}_{\mathcal{M}}, \delta_{\square})$ is convex and compact.*
- (iii) *If \mathbf{Z} is compact, then the space $(\widetilde{\mathcal{W}}_1, \delta_{\square})$ is compact.*

We deduce from this theorem a characterization of relative compactness for subsets of probability-graphons.

Proposition 5.2 (Characterization of relative compactness). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_{\pm}(\mathbf{Z})$) that induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$). Assume that the distance $d_{\square, m}$ on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}) is (invariant) smooth and weakly regular.*

- (i) *If a sequence of elements of \mathcal{W}_1 or $\widetilde{\mathcal{W}}_1$ (resp. \mathcal{W}_+ or $\widetilde{\mathcal{W}}_+$) is converging for $\delta_{\square, m}$, then it is tight.*
- (ii) *Let \mathcal{K} be a subset of $\widetilde{\mathcal{W}}_1$ (resp. a uniformly bounded subset of $\widetilde{\mathcal{W}}_+$). Then, the set \mathcal{K} is relatively compact for $\delta_{\square, m}$ if and only if it is tight.*
- (iii) *Let \mathcal{M} be a subset of $\mathcal{M}_+(\mathbf{Z})$ which is bounded, convex and closed for the weak topology. Then the set $\widetilde{\mathcal{W}}_{\mathcal{M}}$ is convex and closed in $\widetilde{\mathcal{W}}_+$.*

Remark that convergence for $\delta_{\square, m}$ does not necessarily imply tightness on \mathcal{W}_{\pm} or on $\widetilde{\mathcal{W}}_{\pm}$.

Proof. We consider the case where d_m is a distance on $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_{\pm}(\mathbf{Z})$, the case with $\mathcal{M}_{\leq 1}(\mathbf{Z})$ is similar.

We prove Point (i). Let $(W_n)_{n \in \mathbb{N}}$ be a convergent sequence of \mathcal{W}_+ (and thus of $\widetilde{\mathcal{W}}_+$) for $\delta_{\square, m}$. We deduce from the continuity of the map $W \mapsto M_W$, see Lemma 4.9, that the sequence $(M_{W_n})_{n \in \mathbb{N}}$ is converging for d_m , and thus is tight as d_m induces the weak topology on $\mathcal{M}_+(\mathbf{Z})$. Then, by definition the sequence $(W_n)_{n \in \mathbb{N}}$ is tight.

We prove Point (ii). If $\mathcal{K} \subset \widetilde{\mathcal{W}}_+$ is tight and uniformly bounded, then by Theorem 5.1 (i) every sequence in \mathcal{K} has a subsequence converging for $\delta_{\square, m}$, which implies that \mathcal{K} is relatively compact in the metric space $(\widetilde{\mathcal{W}}_+, \delta_{\square, m})$ (see Remark 2.1).

Conversely, assume that $\mathcal{K} \subset \widetilde{\mathcal{W}}_+$ is uniformly bounded and relatively compact for $\delta_{\square, m}$. Define $\mathcal{M} = \{M_W : W \in \mathcal{K}\} \subset \mathcal{M}_+(\mathbf{Z})$. By Lemma 4.9, the mapping $W \mapsto M_W$ is continuous from $(\widetilde{\mathcal{W}}_+, \delta_{\square, m})$ to $(\mathcal{M}_+(\mathbf{Z}), d_m)$. Hence, as d_m induces the weak topology on $\mathcal{M}_+(\mathbf{Z})$, the set \mathcal{M} is also relatively compact in $\mathcal{M}_+(\mathbf{Z})$ for the weak topology. As the space \mathbf{Z} is Polish, applying Lemma 2.8, we get that $\mathcal{M} \subset \mathcal{M}_+(\mathbf{Z})$ is tight, and by Definition 4.7, the set $\mathcal{K} \subset \widetilde{\mathcal{W}}_+$ is tight.

We postpone the proof of Point (iii) to Section 8 on page 54. \square

5.2. Equivalence of topologies induced by $\delta_{\square, m}$. The following lemma allows to show a first result on equivalence of the topologies induced by the cut distance $\delta_{\square, m}$ for different distances d_m , where the sub-script m is used to distinguish different distances. Its proof is given below. Remind from Theorem 3.17 that $d_{\square, m}$ must be smooth for $\delta_{\square, m}$ to be a distance.

Lemma 5.3 (Comparison of topologies induced by $d_{\square, m}$ and $\delta_{\square, m}$). *Let d_m and $d_{m'}$ be two distances on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that $d_{m'}$ is uniformly continuous w.r.t. d_m (in particular, d_m induces a finer topology than $d_{m'}$ on $\mathcal{M}_{\leq 1}(\mathbf{Z})$). Then, we have the following properties.*

- (i) *The distance $d_{\square, m'}$ is uniformly continuous w.r.t. $d_{\square, m}$ on \mathcal{W}_1 . In particular $d_{\square, m}$ induces a finer topology than $d_{\square, m'}$ on \mathcal{W}_1 .*
- (ii) *If the distance $d_{\square, m}$ on \mathcal{W}_1 is smooth, then the distance $d_{\square, m'}$ is also smooth and $\delta_{\square, m'}$ is uniformly continuous w.r.t. $\delta_{\square, m}$. In particular, $\delta_{\square, m}$ induces a finer topology than $\delta_{\square, m'}$ on $\widetilde{\mathcal{W}}_1$.*
- (iii) *If the distance $d_{\square, m}$ on \mathcal{W}_1 is weakly regular, then the distance $d_{\square, m'}$ is also weakly regular.*
- (iv) *Assume that the distance $d_{m'}$ induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, and that the distance $d_{\square, m}$ is smooth and weakly regular. In particular, the distance d_m also induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$. Then, the distances $\delta_{\square, m}$ and $\delta_{\square, m'}$ induce the same topology on $\widetilde{\mathcal{W}}_1$.*

We will see some application of Lemma 5.3 in Corollary 5.6 below.

Remark 5.4 (Extension to \mathcal{W}_{\pm} for topology comparisons). In Lemma 5.3 (i)-(iii), one can replace \mathcal{W}_1 and $\widetilde{\mathcal{W}}_1$ by \mathcal{W}_+ and $\widetilde{\mathcal{W}}_+$ or by \mathcal{W}_{\pm} and $\widetilde{\mathcal{W}}_{\pm}$ as soon as the distances d_m and $d_{m'}$ are defined on $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_{\pm}(\mathbf{Z})$; in this case comparisons of topologies only apply on uniformly bounded subsets. In Lemma 5.3 (iv), one can replace $\widetilde{\mathcal{W}}_1$ by $\widetilde{\mathcal{W}}_{\mathcal{M}}$ with a bounded subset $\mathcal{M} \subset \mathcal{M}_+(\mathbf{Z})$ as soon as the distances d_m and $d_{m'}$ are defined on $\mathcal{M}_+(\mathbf{Z})$.

Proof of Lemma 5.3. We prove Point (i). Let $\varepsilon > 0$. As $d_{m'}$ is uniformly continuous w.r.t. d_m , there exists $\eta > 0$ such that for every $\mu, \nu \in \mathcal{M}_{\leq 1}(\mathbf{Z})$, if $d_m(\mu, \nu) < \eta$, then $d_{m'}(\mu, \nu) < \varepsilon$. Let $U, W \in \mathcal{W}_1$ such that $d_{\square, m}(U, W) < \eta$. Then, for every subsets $S, T \subset [0, 1]$, we have:

$$d_{m'}(U(S \times T; \cdot), W(S \times T; \cdot)) < \varepsilon.$$

Thus, $d_{\square, m'}(U, W) \leq \varepsilon$. Hence, $d_{\square, m'}$ is uniformly continuous w.r.t. $d_{\square, m}$.

We prove Point (ii). Assume that $d_{\square, m}$ is smooth. Let $(W_n)_{n \in \mathbb{N}}$ and W be probability-graphons such that $W_n(x, y; \cdot)$ weakly converges to $W(x, y; \cdot)$ for almost every $x, y \in [0, 1]$. Since the cut distance $d_{\square, m}$ is smooth, we get that $d_{\square, m}(W_n, W) \rightarrow 0$. As $d_{\square, m'}$ is uniformly continuous (and thus also continuous) w.r.t. $d_{\square, m}$, we have that $d_{\square, m'}(W_n, W) \rightarrow 0$. Hence, $d_{\square, m'}$ is smooth.

Furthermore, let $\varepsilon > 0$. Let $\eta > 0$ be such that for every $\mu, \nu \in \mathcal{M}_{\leq 1}(\mathbf{Z})$, $d_m(\mu, \nu) < \eta$ implies $d_{m'}(\mu, \nu) < \varepsilon$. For every $U, W \in \mathcal{W}_1$ such that $\delta_{\square, m}(U, W) < \eta$, there exists $\varphi \in \mathcal{S}_{[0, 1]}$

such that $d_{\square,m}(U, W^\varphi) < \eta$, which implies that $d_{\square,m'}(U, W^\varphi) < \varepsilon$, which then implies that $\delta_{\square,m'}(U, W) < \varepsilon$. That is, $\delta_{\square,m'}$ is uniformly continuous w.r.t. $\delta_{\square,m}$.

We prove Point (iii). Assume that $d_{\square,m}$ is weakly regular. Let $\mathcal{K} \subset \mathcal{W}_1$ be tight. Let $\varepsilon > 0$. As $d_{\square,m'}$ is uniformly continuous w.r.t. $d_{\square,m}$, there exists $\eta > 0$ such that for every $U, W \in \mathcal{W}_1$, if $d_{\square,m}(U, W) < \eta$, then $d_{\square,m'}(U, W) < \varepsilon$. Since $d_{\square,m}$ is weakly regular, there exists $m \in \mathbb{N}^*$, such that for every probability-graphon $W \in \mathcal{K}$, and for every finite partition \mathcal{Q} of $[0, 1]$, there exists a finite partition \mathcal{P} of $[0, 1]$ that refines \mathcal{Q} and such that $|\mathcal{P}| \leq m|\mathcal{Q}|$ and $d_{\square,m}(W, W_{\mathcal{P}}) < \eta$; and thus we also have $d_{\square,m'}(W, W_{\mathcal{P}}) < \varepsilon$. Hence, $d_{\square,m'}$ is weakly regular.

We prove Point (iv). Assume that $d_{m'}$ induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ and that $d_{\square,m}$ is smooth and weakly regular. In particular, the topology induced by d_m is finer than the topology induced by $d_{m'}$, i.e. finer than the weak topology. As $d_{\square,m}$ is smooth, by Lemma 3.12, d_m is continuous w.r.t. the weak topology (i.e. the weak topology is finer than the topology induced by d_m), and thus d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$. By Points (ii) and (iii), we get that $d_{\square,m'}$ is also smooth and weakly regular. By Point (ii), the distance $\delta_{\square,m}$ induces a finer topology than $\delta_{\square,m'}$ on $\widetilde{\mathcal{W}}_1$.

We now prove that the topology of $\delta_{\square,m'}$ is finer than the topology of $\delta_{\square,m}$. Let $(W_n)_{n \in \mathbb{N}}$ and W be probability-graphons in $\widetilde{\mathcal{W}}_1$, such that W_n converges to W for $\delta_{\square,m'}$. By Proposition 5.2 (i), we deduce that the sequence $(W_n)_{n \in \mathbb{N}}$ is tight. As $d_{\square,m}$ is smooth and weakly regular, Theorem 5.1 gives that every subsequence $(W_{n_k})_{k \in \mathbb{N}}$ of the sequence $(W_n)_{n \in \mathbb{N}}$ has a further subsequence $(W_{n'_k})_{k \in \mathbb{N}}$ that converges for $\delta_{\square,m}$ to a limit, say $U \in \widetilde{\mathcal{W}}_1$. Since $\delta_{\square,m}$ is finer than $\delta_{\square,m'}$, we deduce that $(W_{n'_k})_{k \in \mathbb{N}}$ converges also to U for $\delta_{\square,m'}$; but, as a subsequence, it also converges to W for $\delta_{\square,m'}$. As $\delta_{\square,m'}$ is a distance on \mathcal{W}_1 thanks to Theorem 3.17, we get $U = W$. Hence, every subsequence of $(W_n)_{n \in \mathbb{N}}$ has a further subsequence that converges to W for $\delta_{\square,m}$, therefore the whole sequence itself converges to W for $\delta_{\square,m}$. Consequently, $\delta_{\square,m'}$ is finer than $\delta_{\square,m}$, and thus those two distances induce the same topology on $\widetilde{\mathcal{W}}_1$. \square

The following theorem states that the topology induced by $\delta_{\square,m}$ does not depend on d_m under some hypothesis. We prove this theorem in Section 8. Recall that under suitable conditions satisfied in the next theorem, the quotient space $\widetilde{\mathcal{W}}_1$ does not depend on the choice of the distance d_m , see Theorem 3.17.

Theorem 5.5 (Equivalence of topologies induced by $\delta_{\square,m}$ on $\widetilde{\mathcal{W}}_1$). *The topology on the space probability-graphon $\widetilde{\mathcal{W}}_1$ induced by the distance $\delta_{\square,m}$ does not depend on the choice of the distance d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, as long as d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ and the cut distance $d_{\square,m}$ on \mathcal{W}_1 is (invariant) smooth, weakly regular and regular w.r.t. the stepping operator.*

Remind from Proposition 4.13 that when the distance d_m is quasi-convex and continuous w.r.t. the weak topology on $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_\pm(\mathbf{Z})$, then the cut distance $d_{\square,m}$ is invariant, smooth, weakly regular and regular w.r.t. the stepping operator. This is in particular the case of $d_{\mathcal{P}}$, $\|\cdot\|_{\mathcal{F}}$, $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\text{FM}}$.

The next corollary is an immediate consequence of Lemma 3.22, Corollary 4.14, Lemma 5.3 and Theorem 5.5. This corollary gathers results comparing the topology induced by the cut distances associated with the distances introduced in Section 3.8. It is yet unclear if the distances $d_{\square,\mathcal{F}}$ induces the same topology on the space of labeled probability-graphons \mathcal{W}_1 as the one induced by $d_{\square,\mathcal{P}}$, $d_{\square,\text{FM}}$ or $d_{\square,\text{KR}}$.

Corollary 5.6 (Topological equivalence of the cut distances associated to $d_{\mathcal{P}}$, $\|\cdot\|_{\text{FM}}$, $\|\cdot\|_{\text{KR}}$ and $\|\cdot\|_{\mathcal{F}}$). *The cut distances $d_{\square,\mathcal{P}}$ on \mathcal{W}_+ and $d_{\square,\text{KR}}$, $d_{\square,\text{FM}}$ and $d_{\square,\mathcal{F}}$ on \mathcal{W}_\pm are invariant, smooth, weakly regular and regular w.r.t. the stepping operator. Moreover, we have the following comparison between the distances introduced in Section 3.8.*

- (i) The cut norms $\|\cdot\|_{\square, FM}$ and $\|\cdot\|_{\square, KR}$ (resp. the cut distances $\delta_{\square, FM}$ and $\delta_{\square, KR}$) are metrically equivalent on \mathcal{W}_{\pm} (resp. $\widetilde{\mathcal{W}}_{\pm}$).
- (ii) The cut distances $\delta_{\square, FM}$, $\delta_{\square, KR}$ and $\delta_{\square, \mathcal{P}}$ (resp. $d_{\square, FM}$, $d_{\square, KR}$ and $d_{\square, \mathcal{P}}$) are uniformly continuous w.r.t. one another, and thus induce the same topology on $\widetilde{\mathcal{W}}_1$ (resp. \mathcal{W}_1) and on every uniformly bounded subset of $\widetilde{\mathcal{W}}_+$ (resp. \mathcal{W}_+).
- (iii) The cut distances $\delta_{\square, FM}$, $\delta_{\square, KR}$, $\delta_{\square, \mathcal{P}}$ and $\delta_{\square, \mathcal{F}}$, for every choice of the convergence determining sequence \mathcal{F} , induce the same topology on $\widetilde{\mathcal{W}}_1$.

Proof. The first part of the corollary is a re-statement of Corollary 4.14. Point (i) is an immediate consequence of (9).

We now prove Point (ii). Thanks to (9) and Point (i), it is enough to consider only the Prohorov and the Kantorovitch-Rubinshtein distances. As $d_{\mathcal{P}}$ is uniformly continuous w.r.t. d_{KR} (see Lemma 3.22), applying Lemma 5.3 (remind Corollary 4.14) with Remark 5.4 in mind, we get that $\delta_{\square, \mathcal{P}}$ (resp. $d_{\square, \mathcal{P}}$) is uniformly continuous w.r.t. $\delta_{\square, KR}$ (resp. $d_{\square, KR}$) on every uniformly bounded subset of $\widetilde{\mathcal{W}}_+$ (resp. \mathcal{W}_+). As d_{KR} is also uniformly continuous w.r.t. $d_{\mathcal{P}}$ (see Lemma 3.22), applying again Lemma 5.3, we have that $\delta_{\square, KR}$ (resp. $d_{\square, KR}$) is uniformly continuous w.r.t. $\delta_{\square, \mathcal{P}}$ (resp. $d_{\square, \mathcal{P}}$) on every uniformly bounded subset of $\widetilde{\mathcal{W}}_+$ (resp. \mathcal{W}_+).

Point (iii) is an immediate consequence of Corollary 4.14 and Theorem 5.5, together with Point (ii). \square

Remark 5.7 (Extension to uniformly bounded subsets of $\widetilde{\mathcal{W}}_+$). In Theorem 5.5 and also in Corollary 5.6 (iii), one can replace $\widetilde{\mathcal{W}}_1$ by $\widetilde{\mathcal{W}}_{\mathcal{M}}$ with a bounded subset $\mathcal{M} \subset \mathcal{M}_+(\mathbf{Z})$ as soon as the distance d_m is defined on $\mathcal{M}_+(\mathbf{Z})$. (One has in mind the case $\mathcal{M} = \mathcal{M}_{\leq 1}(\mathbf{Z})$.) This can be seen by an easy modification in the proof of Theorem 5.5. Alternatively, this can be seen using scaling to reduce the case of general \mathcal{M} to the case of $\mathcal{M}_{\leq 1}(\mathbf{Z})$, and then adding a cemetery point (for missing mass of measures) to \mathbf{Z} to further reduce to the case of $\mathcal{M}_1(\mathbf{Z})$.

5.3. Completeness. Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ or $\mathcal{M}_+(\mathbf{Z})$. We shall consider a slight modification of the cut distances $d_{\square, m}$ and $\delta_{\square, m}$ to achieve completeness. Recall the measure $M_W \in \mathcal{M}_+(\mathbf{Z})$ defined by (16) associated to $W \in \mathcal{W}_+$.

Definition 5.8 (The cut distances $d_{\square, m}^c$ and $\delta_{\square, m}^c$). Let d_m and d^c be two distances on $\mathcal{M}_{\epsilon}(\mathbf{Z})$ with $\epsilon \in \{\leq 1, +\}$. We define the cut distance $d_{\square, m}^c$ on the space of $\mathcal{M}_{\epsilon}(\mathbf{Z})$ -valued kernels \mathcal{W}_{ϵ} as:

$$d_{\square, m}^c(U, W) = d_{\square, m}(U, W) + d^c(M_U, M_W),$$

and the cut (pseudo-)distance $\delta_{\square, m}^c$ on the space of unlabeled $\mathcal{M}_{\epsilon}(\mathbf{Z})$ -valued kernels $\widetilde{\mathcal{W}}_{\epsilon}$ as:

$$\delta_{\square, m}^c(U, W) = \inf_{\varphi \in \mathcal{S}_{[0,1]}} d_{\square, m}^c(U, W^{\varphi}) = \delta_{\square, m}(U, W) + d^c(M_U, M_W).$$

Notice that by Lemma 3.11 and the definition of M_W , the distance $d_{\square, m}^c$ is invariant.

Lemma 5.9 (Topological equivalence of $\delta_{\square, m}$ and $\delta_{\square, m}^c$). Let d_m and d^c be two distances on $\mathcal{M}_{\epsilon}(\mathbf{Z})$, with $\epsilon \in \{\leq 1, +\}$, such that d^c is continuous w.r.t. d_m and that $d_{\square, m}$ is (invariant and) smooth on \mathcal{W}_{ϵ} . Then, the cut distance $d_{\square, m}^c$ is invariant and smooth and $\delta_{\square, m}^c$ is a distance on $\widetilde{\mathcal{W}}_{\epsilon}$. Moreover, the distances $d_{\square, m}$ and $d_{\square, m}^c$ (resp. $\delta_{\square, m}$ and $\delta_{\square, m}^c$) induce the same topology on the space \mathcal{W}_{ϵ} (resp. $\widetilde{\mathcal{W}}_{\epsilon}$).

Proof. Let $(W_n)_{n \in \mathbb{N}}$ and W be elements of $\mathcal{W}_{\leq 1}$ such that $(W_n(x, y; \cdot))_{n \in \mathbb{N}}$ weakly converges to $W(x, y; \cdot)$ for almost every $x, y \in [0, 1]$. Since the distance $d_{\square, m}$ is smooth, we have that $\lim_{n \rightarrow \infty} d_{\square, m}(W_n, W) = 0$. Using Lemma 4.9 on the continuity of the map $W \mapsto M_W$ and that d^c is continuous w.r.t. d_m , we obtain that $\lim_{n \rightarrow \infty} d_{\square, m}^c(W_n, W) = 0$. This gives that the

distance $d_{\square,m}^c$ is smooth. Since we have already seen that $d_{\square,m}^c$ is invariant, we deduce from Theorem 3.17 that $\delta_{\square,m}^c$ is a distance on $\widetilde{\mathcal{W}}_1$.

We now prove that the two distances $d_{\square,m}$ and $\delta_{\square,m}^c$ induce the same topology (which implies that this is also true for $\delta_{\square,m}$ and $\delta_{\square,m}^c$). As $d_{\square,m} \leq d_{\square,m}^c$, convergence for $d_{\square,m}^c$ implies convergence for $d_{\square,m}$. Conversely, let $(W_n)_{n \in \mathbb{N}}$ be a sequence in \mathcal{W}_ϵ that converges for $d_{\square,m}$ to a limit, say $W \in \mathcal{W}_\epsilon$. Using again Lemma 4.9 and the continuity of d^c w.r.t. d_m , we obtain that $\lim_{n \rightarrow \infty} d^c(M_{W_n}, M_W) = 0$. This clearly implies that the sequence $(W_n)_{n \in \mathbb{N}}$ converges to W for $d_{\square,m}^c$. Then, the two distances have the same convergent sequences and thus induce the same topology (see Remark 2.1). \square

Recall \mathbf{Z} is a Polish space. We already proved in Proposition 4.6 that the space $(\widetilde{\mathcal{W}}_1, \delta_{\square,m})$ is separable; and we now investigate completeness of this space.

Theorem 5.10 ($\widetilde{\mathcal{W}}_1$ is a Polish space). *Let d_m and d^c be two distances on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that d^c induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, d^c is complete and continuous w.r.t. d_m , and $d_{\square,m}$ is (invariant) smooth and weakly regular on \mathcal{W}_1 . Then, the space $(\widetilde{\mathcal{W}}_1, \delta_{\square,m}^c)$ is a Polish metric space.*

Note that the assumptions in Theorem 5.10 imply that d_m also induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$. Indeed, as d^c is continuous w.r.t. d_m , the topology induced by d_m is finer than the topology induced by d^c , i.e. finer than the weak topology. As $d_{\square,m}$ is smooth, by Lemma 3.12, d_m is continuous w.r.t. the weak topology (i.e. the weak topology is finer than the topology induced by d_m), and thus d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$.

Also note that Theorem 5.10 can easily be extended to $\mathcal{W}_{\leq 1}$ or the space of unlabeled \mathcal{M} -valued kernels $\widetilde{\mathcal{W}}_{\mathcal{M}}$ when \mathcal{M} is a bounded convex closed subset of $\mathcal{M}_+(\mathbf{Z})$.

Proof. From Lemma 5.9, we have that $\delta_{\square,m}^c$ is a distance on $\widetilde{\mathcal{W}}_1$ which induces the same topology as $\delta_{\square,m}$, and from Proposition 4.6, we have that $(\widetilde{\mathcal{W}}_1, \delta_{\square,m})$, and thus $(\widetilde{\mathcal{W}}_1, \delta_{\square,m}^c)$, is separable. To get that this latter space is Polish, we are left to prove that the distance $\delta_{\square,m}^c$ is complete.

Let $(W_n)_{n \in \mathbb{N}}$ be a sequence of probability-graphons that is Cauchy for $\delta_{\square,m}^c$. By definition of the cut distance $\delta_{\square,m}^c$, the sequence of probability measures $(M_{W_n})_{n \in \mathbb{N}}$ is Cauchy in $\mathcal{M}_1(\mathbf{Z})$ for the complete distance d^c . Thus, the sequence $(M_{W_n})_{n \in \mathbb{N}}$ is weakly convergent as d^c induces the weak topology, which implies that it is tight (see Lemma 2.8). Hence, by definition, the sequence of probability-graphons $(W_n)_{n \in \mathbb{N}}$ is tight. By Theorem 5.1 (i), there exists a subsequence $(W_{n_k})_{k \in \mathbb{N}}$ that converges for $\delta_{\square,m}$ to a limit, say $W \in \widetilde{\mathcal{W}}_1$. This subsequence also converges for $\delta_{\square,m}^c$ to W as $\delta_{\square,m}$ and $\delta_{\square,m}^c$ induce the same topology. Finally, because the sequence $(W_n)_{n \in \mathbb{N}}$ is Cauchy for $\delta_{\square,m}^c$ and has a subsequence converging to W for $\delta_{\square,m}^c$, the whole sequence must also converge to W for $\delta_{\square,m}^c$. Consequently, the distance $\delta_{\square,m}^c$ is complete. \square

The following lemma shows that every probability measure can be represented as a constant probability-graphon.

Lemma 5.11 ($\mathcal{M}_1(\mathbf{Z})$ seen as a closed subset of \mathcal{W}_1). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that $d_{\square,m}$ is (invariant and) smooth on \mathcal{W}_1 . Then, the map $\mu \mapsto W_\mu \equiv \mu$ is an injection from $(\mathcal{M}_1(\mathbf{Z}), d_m)$ to $(\widetilde{\mathcal{W}}_1, \delta_{\square,m})$ with a closed range and continuous inverse.*

Proof. For any $\mu \in \mathcal{M}_1(\mathbf{Z})$ consider the constant probability-graphon $W_\mu \equiv \mu$, and notice that $M_{W_\mu} = \mu$, that $W_\mu(S \times T; \cdot) = \lambda(S)\lambda(T)\mu$ for all measurable $S, T \subset [0, 1]$, and that $W_\mu^\varphi = W_\mu$

for any measure-preserving map φ . This readily implies that for $\mu \in \mathcal{M}_1(\mathbf{Z})$ and $W \in \mathcal{W}_1$:

$$(20) \quad \delta_{\square, m}(W_\mu, W) = d_{\square, m}(W_\mu, W) = \sup_{S, T \subset [0, 1]} d_m(\lambda(S)\lambda(T)\mu, W(S \times T; \cdot)) \geq d_m(\mu, M_W).$$

In particular, taking $W = W_\nu$ for $\nu \in \mathcal{M}_1(\mathbf{Z})$ we get that $\delta_{\square, m}(W_\mu, W_\nu) \geq d_m(\mu, \nu)$. This implies that the map $\mathcal{I} : \mu \mapsto W_\mu \equiv \mu$ is an injection, and its inverse, given by the map $W_\mu \mapsto \mu$, is 1-Lipschitz.

Let $(\mu_n)_{n \in \mathbb{N}}$ be a sequence in $\mathcal{M}_1(\mathbf{Z})$ such that the sequence $(W_{\mu_n})_{n \in \mathbb{N}}$ converges for $\delta_{\square, m}$ to a limit, say W . We deduce from (20) that $(\mu_n)_{n \in \mathbb{N}}$ converges for d_m to $\mu = M_W$ and that for all measurable $S, T \subset [0, 1]$, $(\lambda(S)\lambda(T)\mu_n)_{n \in \mathbb{N}}$ converges for d_m to $W(S \times T; \cdot)$. This implies that $W(S \times T; \cdot) = \lambda(S)\lambda(T)\mu(\cdot)$ for all measurable $S, T \subset [0, 1]$, that is, $W = W_\mu$. This implies that the image by \mathcal{I} of any closed subset of $\mathcal{M}_1(\mathbf{Z})$ is a closed subset of \mathcal{W}_1 , and thus the range of \mathcal{I} is closed. \square

Remark 5.12 (Extension to isometric representation of $\mathcal{M}_1(\mathbf{Z})$). If the distance d_m , in addition to the hypothesis of Lemma 5.11, is sub-homogeneous, that is, for all $\mu, \nu \in \mathcal{M}_1(\mathbf{Z})$ we have $d_m(\mu, \nu) = \sup_{r \in [0, 1]} d_m(r\mu, r\nu)$ (which is the case if d_m is quasi-convex), then we deduce from (20) that the map $\mu \mapsto W_\mu \equiv \mu$ is isometric from $(\mathcal{M}_1(\mathbf{Z}), d_m)$ to $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$.

We now state characterization of compactness and completeness for the space of probability-graphons. Recall \mathbf{Z} is a Polish space.

Corollary 5.13 (Characterization of compactness and completeness for $\widetilde{\mathcal{W}}_1$). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, which induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, and such that $d_{\square, m}$ is (invariant) smooth and weakly regular on \mathcal{W}_1 . We have the following properties.*

- (i) \mathbf{Z} is compact $\iff (\mathcal{M}_{\leq 1}(\mathbf{Z}), d_m)$ is compact $\iff (\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is compact.
- (ii) If $(\mathcal{M}_{\leq 1}(\mathbf{Z}), d_m)$ is complete then $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is complete.
- (iii) Assume furthermore that d_m is sub-homogeneous (see Remark 5.12). If $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is complete, then $(\mathcal{M}_1(\mathbf{Z}), d_m)$ is complete.

Proof. We prove Point (i). From Remark 2.7, we already know that \mathbf{Z} is compact if and only if $\mathcal{M}_{\leq 1}(\mathbf{Z})$ is weakly compact, i.e. compact for d_m as d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$.

Now, assume that $(\mathcal{M}_{\leq 1}(\mathbf{Z}), d_m)$ is compact. Applying Theorem 5.1 (iii), we get that the space $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is also compact.

Conversely, assume that $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is compact. By Lemma 4.9, the mapping $W \mapsto M_W$ is continuous from $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ to $(\mathcal{M}_1(\mathbf{Z}), d_m)$, and as $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is compact its image through this mapping is also compact. To conclude, it is enough to check that this mapping is surjective. But this is clear as the image of the constant probability-graphon $W_\mu \equiv \mu$ is $M_{W_\mu} = \mu$. Hence, $(\mathcal{M}_1(\mathbf{Z}), d_m)$ (and thus $(\mathcal{M}_{\leq 1}(\mathbf{Z}), d_m)$) is compact.

We prove Point (ii). Assume that $(\mathcal{M}_{\leq 1}(\mathbf{Z}), d_m)$ is complete. Thus, we can choose $d^c = d_m$ in Definition 5.8, and apply Theorem 5.10 to get that $(\widetilde{\mathcal{W}}_1, \delta_{\square, m}^c)$ is complete. As $d^c = d_m$, we have $\delta_{\square, m} \leq \delta_{\square, m}^c \leq 2\delta_{\square, m}$. Hence, $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is also complete.

We prove Point (iii). Assume that $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is complete. Let $(\mu_n)_{n \in \mathbb{N}}$ be a Cauchy sequence of probability measures in $(\mathcal{M}_1(\mathbf{Z}), d_m)$. By Remark 5.12, the sequence of constant probability-graphons $(W_{\mu_n})_{n \in \mathbb{N}}$ is also Cauchy for $\delta_{\square, m}$. As $(\widetilde{\mathcal{W}}_1, \delta_{\square, m})$ is complete, there exists a probability-graphon $W \in \widetilde{\mathcal{W}}_1$ such that $(W_{\mu_n})_{n \in \mathbb{N}}$ converges to W for the cut distance $\delta_{\square, m}$. Thanks to Lemma 5.11, W is constant equal to some $\mu \in \mathcal{M}_1(\mathbf{Z})$, and $(\mu_n)_{n \in \mathbb{N}}$ converges to μ for d_m . Hence, $(\mathcal{M}_1(\mathbf{Z}), d_m)$ is complete. \square

6. SAMPLING FROM PROBABILITY-GRAPHONS

Measured-valued graphons allow to define models for generating random weighted graphs that are more general than the models based on real-valued graphons. We prove that the weighted graphs sampled from probability-graphons are close to their original model for the cut distance $\delta_{\square, \mathcal{F}}$, where $\mathcal{F} = (f_k)_{k \in \mathbb{N}}$ (with $f_0 = \mathbf{1}$) is a convergence determining sequence.

It would have been more natural to work in Sections 6 and 7 with the Kantorovitch-Rubinshtein norm or the Fortet-Mourier norm that both treats all test functions in a uniform manner. Unfortunately, the supremum in the definition of both of this norms does not behave well regarding the probabilities and expectations of graphs sampled from probability-graphons. We need in our proofs (and in particular that of the First Sampling Lemma 6.7 below) to consider simultaneously only a finite number of test functions in order to control the probability of failure for our stochastic bounds.

6.1. $\mathcal{M}_1(\mathbf{Z})$ -Graphs and weighted graphs. A graph $G = (V, E)$ is composed of a finite set of vertices $V(G) = V$, and a set of edges $E(G) = E$ which is a subset of $V \times V$ avoiding the diagonal. When its set of edges $E(G)$ is symmetric, we say that G is *symmetric* or *non-oriented*. We denote by $v(G) = |V(G)|$ the number of vertices of this graph, and by $e(G) = |E(G)|$ its number of edges.

Definition 6.1 (\mathcal{X} -graphs). *Let \mathcal{X} be a non-empty set. A \mathcal{X} -graph is a triplet $G = (V, E, \Phi)$ where (V, E) is a graph and $\Phi : E \rightarrow \mathcal{X}$ is a map that associates a decoration $x = \Phi(e) \in \mathcal{X}$ to each edge $e \in E$. When $\mathcal{X} = \mathbf{Z}$, we say that G is a weighted graph.*

Furthermore, the graph G is said to be symmetric if (V, E) is a symmetric graph and if Φ is a symmetric function, i.e. for every edge $(x, y) \in E$, we have $(y, x) \in E$ and $\Phi(x, y) = \Phi(y, x)$.

Remark 6.2 ($\mathcal{M}_1(\mathbf{Z})$ -Graphs as probability-graphons). Any labeled $\mathcal{M}_1(\mathbf{Z})$ -graph G can be naturally represented as an $\mathcal{M}_1(\mathbf{Z})$ -valued graphon, which we denote by W_G , in the following way. Let $G = (V, E, M)$ be a $\mathcal{M}_1(\mathbf{Z})$ -graph, with $v(G) = n \in \mathbb{N}^*$. Denote by $V = [n] = \{1, \dots, n\}$ the vertices of G . Consider intervals of length $1/n$: for $1 \leq i \leq n$, let $J_i = ((i-1)/n, i/n]$. We then define the $\mathcal{M}_1(\mathbf{Z})$ -valued graphon stepfunction W_G associated with the $\mathcal{M}_1(\mathbf{Z})$ -graph G by:

$$\forall (i, j) \in E, \quad \forall (x, y) \in J_i \times J_j, \quad W_G(x, y; dz) = \Phi(i, j)(dz);$$

and $W_G(x, y; dz)$ equals the Dirac mass at ∂ otherwise, where ∂ is an element of \mathbf{Z} used as a cemetery point for missing edges in graphs.

In this section, we investigate weighted graphs sampled from probability-graphons. Hence, using the cemetery point argument in the remark above, we only consider complete graphs for the rest of this section.

Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$. If G and H have the same vertex-set, the cut distance between them is defined as the cut distance between their associated graphons:

$$d_{\square, m}(G, H) = d_{\square, m}(W_G, W_H).$$

When G and H does not have the same vertex-sets, as the numbering of the vertices in Remark 6.2 is arbitrary, we must consider the unlabeled cut distance between them defined as the cut distance between their associated graphons:

$$\delta_{\square, m}(G, H) = \delta_{\square, m}(W_G, W_H).$$

Remind that when the distance d_m derives from a norm N_m on $\mathcal{M}_{\pm}(\mathbf{Z})$, Lemma 3.19 applies, and the cut distance $d_{\square, m}(G, H)$ can be rewritten as a combinatorial optimization over whole steps.

Remark 6.3 (Weighted graphs as $\mathcal{M}_1(\mathbf{Z})$ -graphs). We will sometimes need to interpret a weighted graph G as a $\mathcal{M}_1(\mathbf{Z})$ -graph where a weight x on an edge is replaced by δ_x the Dirac mass located at x .

Notation 6.4 (The real-weighted graph $G[f]$). For a $\mathcal{M}_1(\mathbf{Z})$ -graph (resp. weighted graph) G and a function $f \in C_b(\mathbf{Z})$, we denote by $G[f]$ the real-weighted graph with the same vertex set and edge set as G , and where the edge (i, j) has weight $\Phi_{G[f]}(i, j) = \Phi_G(i, j; f) = \int_{\mathbf{Z}} f(z) \Phi_G(i, j; dz)$ (resp. $\Phi_{G[f]}(i, j) = f(\Phi_G(i, j))$), where Φ_G is the decoration of the $\mathcal{M}_1(\mathbf{Z})$ -graph G .

6.2. W -random graphs. Let W be a probability-graphon, and $x = (x_1, \dots, x_n)$, $n \in \mathbb{N}^*$, be a sequence of points from $[0, 1]$. We define the $\mathcal{M}_1(\mathbf{Z})$ -graph $\mathbb{H}(x, W)$ as the complete graph whose vertex set is $[n] = \{1, \dots, n\}$, and with each edge (i, j) decorated by the probability measure $W(x_i, x_j; dz)$.

Let H be any $\mathcal{M}_1(\mathbf{Z})$ -graph. We can define from H a random weighted (directed) graph $\mathbb{G}(H)$ whose vertex set $V(H)$ and edge set $E(H)$ are the same as H , and with each edge (i, j) having a random weight $\beta_{i,j}$ distributed according to the probability distribution decorating the edge (i, j) in H , all the weights being independent from each other. For the special case where $H = \mathbb{H}(x, W)$, we simply note $\mathbb{G}(x, W) = \mathbb{G}(\mathbb{H}(x, W))$.

An important special case is when the sequence X is chosen at random: $X = (X_i)_{1 \leq i \leq n}$ where the X_i are independent and uniformly distributed on $[0, 1]$. For this special case, we simply note $\mathbb{H}(n, W) = \mathbb{H}(X, W)$ and $\mathbb{G}(n, W) = \mathbb{G}(X, W)$, that are conditionally on $X = x$, distributed respectively as $\mathbb{H}(x, W)$ and $\mathbb{G}(x, W)$. The random graphs $\mathbb{H}(n, W)$ and $\mathbb{G}(n, W)$ are called W -random graphs.

Remark 6.5 (The case of symmetric graphons). In the special case where W is a symmetric probability-graphon, the $\mathcal{M}_1(\mathbf{Z})$ -graph $\mathbb{H}(x, W)$ is also symmetric. From a symmetric $\mathcal{M}_1(\mathbf{Z})$ -graph H , the random weighted graph $\mathbb{G}(H)$ is not necessarily symmetric, but we can define a random symmetric weighted graph $\mathbb{G}^{\text{sym}}(H)$ whose vertex set $V(H)$ and $E(H)$ are the same as H , and with independent weights $\beta_{i,j} = \beta_{j,i}$ on each edge $(i, j) = (j, i)$ distributed according to $\Phi_H(i, j; \cdot)$. For $H = \mathbb{H}(x, W)$ we simply note $\mathbb{G}^{\text{sym}}(x, W)$ and $\mathbb{G}^{\text{sym}}(n, W)$.

For a weighted graph G , and for $1 \leq k \leq v(G)$, we can define the random weighted graph $\mathbb{G}(k, G)$ as being the sub-graph of G induced by a uniform random subset of k distinct vertices from G . Then, upper bounding by the probability that a uniformly-chosen map $[k] \rightarrow V(G)$ is non-injective, we get the following bound on the total variation distance between the graphs obtained from G and its associated graphon W_G :

$$d_{\text{var}}(\mathbb{G}(k, G), \mathbb{G}(k, W_G)) \leq \binom{k}{2} \frac{1}{v(G)},$$

where d_{var} is the total variation distance between probability measures.

6.3. Estimation of the distance by sampling.

6.3.1. The first sampling lemma. In this subsection, we link sampling from graphons with the cut distance. This result is the equivalent of Lemma 10.6 in [Lov12]. The main consequence of the following lemma is that the cut distance $d_{\square, \mathcal{F}}$ between two probability-graphons can be estimated by sampling.

Notation 6.6 (The random stepfunction W_X). For a measure-valued kernel W (resp. a real-valued kernel w) and a vector $X = (X_i)_{1 \leq i \leq k}$ composed of k independent random variables uniformly distributed over $[0, 1]$, we denote by $W_X = W_{\mathbb{H}(k, W)}$ (resp. w_X) the random measure-valued (resp. real-valued) stepfunction with k steps of size $1/k$, and where the step (i, j) has value $W(X_i, X_j; \cdot)$ (resp. $w(X_i, X_j)$).

Lemma 6.7 (First Sampling Lemma). *Let \mathcal{F} be a convergence determining sequence. Let $k \in \mathbb{N}^*$, and $U, W \in \mathcal{W}_1$ be two probability-graphons, and let X be a random vector uniformly distributed over $[0, 1]^k$. Then with probability at least $1 - 4k^{1/4} e^{-\sqrt{k}/10}$, we have:*

$$-\frac{2}{k^{1/4}} \leq \|U_X - W_X\|_{\square, \mathcal{F}} - \|U - W\|_{\square, \mathcal{F}} \leq \frac{9}{k^{1/4}}.$$

An immediate consequence of Lemma 6.7 is that the decorated graphs with probability measures on their edges $\mathbb{H}(k, U)$ and $\mathbb{H}(k, W)$ can be coupled in order that $d_{\square, \mathcal{F}}(\mathbb{H}(k, U), \mathbb{H}(k, W))$ is close to $d_{\square, \mathcal{F}}(U, W)$ with high probability.

To prove the first sampling lemma, we first need to prove the following lemma which states that the cut norm $\|\cdot\|_{\square, \mathcal{F}}$ can be approximated by the maximum of the one-sided cut norm using a finite number of function. Remind from Remark 3.24 the definition of the one-sided version of the cut norm $\|\cdot\|_{\square, \mathbb{R}}^+$.

Lemma 6.8 (Approximation bound with $\|\cdot\|_{\square, \mathcal{F}}$ and $\|\cdot\|_{\square, \mathbb{R}}^+$). *Let $U, W \in \mathcal{W}_1$ and let $N \in \mathbb{N}$. For every $\varepsilon = (\varepsilon_n)_{1 \leq n \leq N} \in \{\pm 1\}^N$, define $g_{N, \varepsilon} = \sum_{n=1}^N 2^{-n} \varepsilon_n f_n$. Then, we have:*

$$\|U - W\|_{\square, \mathcal{F}} - 2^{-N} \leq \max_{\varepsilon \in \{\pm 1\}^N} \|(U - W)[g_{N, \varepsilon}]\|_{\square, \mathbb{R}}^+ \leq \|U - W\|_{\square, \mathcal{F}}.$$

Proof. First remark that for $n \in \mathbb{N}$, f_n takes values in $[0, 1]$, and thus $U[f_n] - W[f_n]$ takes values in $[-1, 1]$. Remind that $f_0 = \mathbf{1}$, and thus $U[f_0] - W[f_0] \equiv 0$. Upper bounding integrals by 1 for indices $n > N$, we get:

$$\|U - W\|_{\square, \mathcal{F}} \leq \sup_{S, T \subset [0, 1]} \sum_{n=1}^N 2^{-n} \left| \int_{S \times T} (U - W)[f_n](x, y) \, dx dy \right| + 2^{-N}.$$

And adding the non-negative terms for $n > N$, we get:

$$\sup_{S, T \subset [0, 1]} \sum_{n=1}^N 2^{-n} \left| \int_{S \times T} (U - W)[f_n](x, y) \, dx dy \right| \leq \|U - W\|_{\square, \mathcal{F}}.$$

Using the same idea as in (13) and (14), we get:

$$\sup_{S, T \subset [0, 1]} \sum_{n=1}^N 2^{-n} \left| \int_{S \times T} (U - W)[f_n](x, y) \, dx dy \right| = \max_{\varepsilon \in \{\pm 1\}^N} \|(U - W)[g_{N, \varepsilon}]\|_{\square, \mathbb{R}}^+,$$

which concludes the proof. \square

Proof of Lemma 6.7. Remark that for $f \in C_b(\mathbf{Z})$ and $W \in \mathcal{W}_{\pm}$, we have $(W_X)[f] = (W[f])_X$, and we thus write $W[f]_X$ without any ambiguity.

Assume that $k \geq 2^4$ (otherwise the lower bound in the lemma is trivial). Set $N = \lceil \log_2(k^{1/4}) \rceil$, so that $2^{-1}k^{-1/4} < 2^{-N} \leq k^{-1/4}$. Let $\varepsilon \in \{\pm 1\}^N$. Remark that as the f_n take values in $[0, 1]$, the real-valued kernels $(U - W)[f_n]$ take values in $[-1, 1]$, and thus the real-valued kernel $(U - W)[g_{N, \varepsilon}]$ also take values in $[-1, 1]$. Applying Lemma 10.7 in [Lov12] to the real-valued kernel $(U - W)[g_{N, \varepsilon}]$, we get with probability at least $1 - 2e^{-\sqrt{k}/10}$ that:

$$(21) \quad -\frac{3}{k} \leq \|(U - W)[g_{N, \varepsilon}]_X\|_{\square, \mathbb{R}}^+ - \|(U - W)[g_{N, \varepsilon}]\|_{\square, \mathbb{R}}^+ \leq \frac{8}{k^{1/4}},$$

where remind that $\|\cdot\|_{\square, \mathbb{R}}^+$ is the one-sided version of the cut norm for real-valued kernels defined in (12). Hence, with probability at least $1 - 2^{N+1} e^{-\sqrt{k}/10} \geq 1 - 4k^{1/4} e^{-\sqrt{k}/10}$, we have that the

bounds in (21) holds for every $\varepsilon \in \{\pm 1\}^N$ simultaneously; and when all of this holds, applying Lemma 6.8 to U, W and to U_X, W_X , we get:

$$\begin{aligned} \|U_X - W_X\|_{\square, \mathcal{F}} &\leq \max_{\varepsilon \in \{\pm 1\}^N} \|(U - W)[g_{N, \varepsilon}]_X\|_{\square, \mathbb{R}}^+ + 2^{-N} \\ &\leq \max_{\varepsilon \in \{\pm 1\}^N} \|(U - W)[g_{N, \varepsilon}]\|_{\square, \mathbb{R}}^+ + \frac{9}{k^{1/4}} \\ &\leq \|U - W\|_{\square, \mathcal{F}} + \frac{9}{k^{1/4}}, \end{aligned}$$

and similarly:

$$\begin{aligned} \|U - W\|_{\square, \mathcal{F}} &\leq \max_{\varepsilon \in \{\pm 1\}^N} \|(U - W)[g_{N, \varepsilon}]\|_{\square, \mathbb{R}}^+ + 2^{-N} \\ &\leq \max_{\varepsilon \in \{\pm 1\}^N} \|(U - W)[g_{N, \varepsilon}]_X\|_{\square, \mathbb{R}}^+ + \frac{1}{k^{1/4}} + \frac{3}{k} \\ &\leq \|U_X - W_X\|_{\square, \mathcal{F}} + \frac{2}{k^{1/4}}. \end{aligned}$$

This concludes the proof. \square

6.3.2. Approximation with random weighted graphs. As a consequence of the First Sampling Lemma 6.7, we get that the cut distance between the sampled graphs $\mathbb{H}(k, U)$ and $\mathbb{H}(k, W)$ (with the proper coupling) is close to the cut distance between the probability-graphons U and W . The following lemma states that if k is large enough, then $\mathbb{G}(k, W)$ is close to $\mathbb{H}(k, W)$ in the cut distance $d_{\square, \mathcal{F}}$, and thus the cut distance between the random weighted graphs $\mathbb{G}(k, U)$ and $\mathbb{G}(k, W)$ is also close to $d_{\square, \mathcal{F}}(U, W)$.

Recall from Section 6.2 the definition of the random weighted graph $\mathbb{G}(H)$ when H is a $\mathcal{M}_1(\mathbf{Z})$ -graph. Following Remarks 6.3 and 6.2, we shall see the weighted graph $\mathbb{G}(H)$ as a $\mathcal{M}_1(\mathbf{Z})$ -graph or even as a probability-graphon.

Lemma 6.9 (Bound in probability for $d_{\square, \mathcal{F}}(\mathbb{G}(H), H)$). *For every $\mathcal{M}_1(\mathbf{Z})$ -graph H with k vertices, and for every $\varepsilon \geq 10/\sqrt{k}$, we have:*

$$\mathbb{P}\left(d_{\square, \mathcal{F}}(\mathbb{G}(H), H) > 2\varepsilon\right) \leq e^{-\varepsilon^2 k^2}.$$

Remark 6.10 (Bound in expectation for $d_{\square, \mathcal{F}}(\mathbb{G}(H), H)$). Remind that $d_{\square, \mathcal{F}}(\mathbb{G}(H), H) \leq 1$. Applying Lemma 6.9 with $\varepsilon = 10/\sqrt{k}$, we get the following bound on the expectation of $d_{\square, \mathcal{F}}(\mathbb{G}(H), H)$:

$$\mathbb{E}[d_{\square, \mathcal{F}}(\mathbb{G}(H), H)] \leq \frac{20}{\sqrt{k}} + e^{-100k} < \frac{21}{\sqrt{k}}.$$

Proof of Lemma 6.9. Let H and ε be as in the lemma. Assume that $\varepsilon \leq 1/2$ (otherwise the probability to bound in the lemma is null). To simplify the notations, denote by $G = \mathbb{G}(H)$ through this proof. Define $N = \lceil \log_2(\varepsilon^{-1}) \rceil$, so that $\sum_{n=N+1}^{\infty} 2^{-n} \leq \varepsilon$. Upper bounding by 1 the terms for $n > N$ in (15), we get for $U, W \in \mathcal{W}_1$:

$$d_{\square, \mathcal{F}}(U, W) \leq \sum_{n=1}^N 2^{-n} \|U[f_n] - W[f_n]\|_{\square, \mathbb{R}} + \varepsilon,$$

where remind that $\|\cdot\|_{\square, \mathbb{R}}$ is the cut norm for real-valued kernels defined in (12). Using this equation with the graphs G and H , we get:

$$(22) \quad \begin{aligned} \mathbb{P}(d_{\square, \mathcal{F}}(G, H) > 2\varepsilon) &\leq \mathbb{P}\left(\sum_{n=1}^N 2^{-n} d_{\square, \mathbb{R}}(G[f_n], H[f_n]) > \varepsilon\right) \\ &\leq \sum_{n=1}^N \mathbb{P}(d_{\square, \mathbb{R}}(G[f_n], H[f_n]) > \varepsilon), \end{aligned}$$

where $d_{\square, \mathbb{R}}$ denotes the cut distance associated to the cut norm $\|\cdot\|_{\square, \mathbb{R}}$ for real-valued graphons and kernels. Remark that for every $n \in \mathbb{N}$, $H[f_n]$ and $G[f_n]$ are real-weighted graphs with weights in $[0, 1]$. Thus, by a straightforward adaptation of the proof of [Lov12, Lemma 10.11], we get:

$$(23) \quad \forall n \in [N], \quad \mathbb{P}(d_{\square}(G[f_n], H[f_n]) > \varepsilon) \leq 2 \cdot 4^k e^{-2\varepsilon^2 k^2}.$$

Combining (22) and (23), we get for $\varepsilon > 10/\sqrt{k}$:

$$\mathbb{P}(d_{\square, \mathcal{F}}(G, H) > 2\varepsilon) \leq 2N4^k e^{-2\varepsilon^2 k^2} \leq e^{-\varepsilon^2 k^2},$$

where the last bound derives from simple calculus. This concludes the proof. \square

We can apply the First Sampling Lemma 6.7 along with Lemma 6.9 to get the following lemma, equivalent of the first sampling lemma for the random weighted graph $\mathbb{G}(k, W)$:

Corollary 6.11 (First Sampling Lemma for $\mathbb{G}(k, W)$). *Let $U, W \in \mathcal{W}_1$ be two probability-graphons, and $k \in \mathbb{N}^*$. Then, we can couple the random weighted graphs $\mathbb{G}(k, U)$ and $\mathbb{G}(k, W)$ such that with probability at least $1 - (4k^{1/4} + 1)e^{-\sqrt{k}/10}$, we have:*

$$|d_{\square, \mathcal{F}}(\mathbb{G}(k, U), \mathbb{G}(k, W)) - d_{\square, \mathcal{F}}(U, W)| \leq \frac{13}{k^{1/4}}.$$

Proof. Assume that $k \geq 13^4$ (otherwise the bound in the corollary is trivial). Then, we have with probability at least $1 - 4k^{1/4}e^{-\sqrt{k}/10} - 2e^{-100k} > 1 - (4k^{1/4} + 1)e^{-\sqrt{k}/10}$:

$$\begin{aligned} |d_{\square, \mathcal{F}}(\mathbb{G}(k, U), \mathbb{G}(k, W)) - d_{\square, \mathcal{F}}(U, W)| &\leq |d_{\square, \mathcal{F}}(\mathbb{G}(k, U), \mathbb{G}(k, W)) - d_{\square, \mathcal{F}}(\mathbb{H}(k, U), \mathbb{H}(k, W))| \\ &\quad + |d_{\square, \mathcal{F}}(\mathbb{H}(k, U), \mathbb{H}(k, W)) - d_{\square, \mathcal{F}}(U, W)| \\ &\leq d_{\square, \mathcal{F}}(\mathbb{G}(k, U), \mathbb{H}(k, U)) \\ &\quad + d_{\square, \mathcal{F}}(\mathbb{G}(k, W), \mathbb{H}(k, W)) + \frac{9}{k^{1/4}} \\ &\leq \frac{40}{\sqrt{k}} + \frac{9}{k^{1/4}} \\ &\leq \frac{13}{k^{1/4}}, \end{aligned}$$

where we used the upper bound from the First Sampling Lemma 6.7 (which gives the coupling with the same random vector X to define both graphs $U_X = \mathbb{H}(k, U)$ and $W_X = \mathbb{H}(k, W)$) for the second inequality, the upper bound from Lemma 6.9 with $\varepsilon = 10/\sqrt{k}$ with both U and W for the third inequality, and that $\frac{1}{\sqrt{k}} \leq \frac{1}{13k^{1/4}}$ for the last inequality. \square

6.4. The distance between a probability-graphon and its sample. In this section, we present the Second Sampling Lemma, that shows that a sampled $\mathcal{M}_1(\mathbf{Z})$ -graph is close to its original probability-graphon with high probability. Note that we use the unlabeled cut distance $\delta_{\square, \mathcal{F}}$ rather than $d_{\square, \mathcal{F}}$ as the sample points are unordered. The bound on the distance is much weaker than the one in the First Sampling Lemma 6.7, but nevertheless goes to 0 as the sample size increases.

The proof is a straightforward adaptation of the proof of [Lov12, Lemma 10.16] (replacing the weak regularity lemma and the first sampling lemma by their counterparts for probability-graphons, that is Lemmas 4.17 and 6.7; the sample concentration theorem for real-valued graphons can easily be adapted to probability-graphons).

Lemma 6.12 (Second Sampling Lemma). *Let \mathcal{F} be a convergence determining sequence. Let $W \in \widetilde{\mathcal{W}}_1$ be a probability-graphon and $k \in \mathbb{N}^*$. Then, with probability at least $1 - \exp(-k/(2 \ln(k)))$ we have:*

$$\delta_{\square, \mathcal{F}}(\mathbb{H}(k, W), W) \leq \frac{21}{\sqrt{\ln(k)}} \quad \text{and} \quad \delta_{\square, \mathcal{F}}(\mathbb{G}(k, W), W) \leq \frac{22}{\sqrt{\ln(k)}}.$$

In the above lemma, the asymmetric random graph $\mathbb{G}(k, W)$ can be replaced by the symmetric random graph $\mathbb{G}^{\text{sym}}(k, W)$ without changing the proof. Similarly, the results in Section 6.3.2 can be reformulated with symmetric random graphs $\mathbb{G}^{\text{sym}}(k, W)$ and $\mathbb{G}^{\text{sym}}(H)$ (but with a slight modification of the proof for Lemma 6.9 to symmetrize the random variable $X_{i,j}$ and with the upper bound $e^{-\varepsilon^2 k^2/2}$, see also [Lov12, Lemma 10.11]).

As an immediate consequence of Lemma 6.12 and of the Borel-Cantelli lemma, we get the convergence of the sampled subgraphs for the cut distance $\delta_{\square, \mathcal{F}}$.

Theorem 6.13 (Convergence of sampled subgraphs). *Let \mathcal{F} be a convergence determining sequence. Let $W \in \widetilde{\mathcal{W}}_1$ be a probability-graphon. Then, a.s. the sequence of sampled subgraphs $(\mathbb{G}(k, W))_{k \in \mathbb{N}^*}$ converges to W for the cut distance $\delta_{\square, \mathcal{F}}$, and thus for any cut distance $\delta_{\square, m}$ from Theorem 5.5.*

7. THE COUNTING LEMMAS AND THE TOPOLOGY OF PROBABILITY-GRAPHONS

In this section, we introduce the homomorphism densities for probability-graphons, and then we link those to the cut distance $\delta_{\square, \mathcal{F}}$ through the Counting Lemma and the Inverse Counting Lemma. Those results are analogous to the case of real-valued graphons, see [Lov12, Chapter 7] for the definition of homomorphism densities and [Lov12, Chapter 10] for the Counting Lemma and Inverse Counting Lemma. The main differences with [Lov12] are: the decoration of the edges of the graphs with functions from $C_b(\mathbf{Z})$; the Counting Lemma for the decorations belonging only in the convergence determining sequence \mathcal{F} ; the more technical proof of the Inverse Counting Lemma. Note that we need to work with $\delta_{\square, \mathcal{F}}$ here as the proof of the Inverse Counting Lemma relies on the second sampling Lemma 6.12.

7.1. The homomorphism densities. In the case of non-weighted graphs, the homomorphism densities $t(F, G)$ allow to characterize a graph (up to twin-vertices expansion), and also allow to define a topology for real-valued graphons. In the case of weighted graphs and probability-graphons, we need to replace the absence/presence of edges (which is 0-1 valued) by test functions from $C_b(\mathbf{Z})$ decorating each edge.

In this section, we often need to fix the underlying (directed) graph structure $F = (V, E)$ (which may be incomplete) of a $C_b(\mathbf{Z})$ -graph and to vary only the $C_b(\mathbf{Z})$ -decorating functions $g = (g_e)_{e \in E}$, thus we will write $F^g = (V, E, g)$ for a $C_b(\mathbf{Z})$ -graph. Moreover, when there exists a convergence determining sequence \mathcal{F} such that $g_e \in \mathcal{F}$ for every edge $e \in E$, we say that F^g is a \mathcal{F} -graph and use the same notation conventions.

Definition 7.1 (Homomorphism density). *We define the homomorphism density of a $C_b(\mathbf{Z})$ -graph F^g in a signed measure-valued kernel $W \in \mathcal{W}_{\pm}$ as:*

$$(24) \quad t(F^g, W) = M_W^F(g) = \int_{[0,1]^{V(F)}} \prod_{(i,j) \in E(F)} W(x_i, x_j; g_{i,j}) \prod_{i \in V(F)} dx_i.$$

Moreover, M_W^F defines a measure on \mathbf{Z}^E (which we still denote by M_W^F) which is characterized by $M_W^F(\otimes_{e \in E} g_e) = M_W^F(g)$ for $g = (g_e)_{e \in E}$.

Remark 7.2 (Invariance under relabeling of homomorphism densities). Let $\varphi : [0, 1] \rightarrow [0, 1]$ be a measure-preserving map. As $\varphi^{\otimes k} : (x_1, \dots, x_k) \mapsto (\varphi(x_1), \dots, \varphi(x_k))$ is a measure-preserving map on $[0, 1]^k$, applying the transfer formula (see (1)), we get that for every $C_b(\mathbf{Z})$ -graph F^g and every signed measure-valued kernel $W \in \mathcal{W}_\pm$, we have $t(F^g, W^\varphi) = t(F^g, W)$. Thus $t(F^g, \cdot)$ can be extended to $\widetilde{\mathcal{W}}_\pm$.

Remark 7.3. When $W \in \mathcal{W}_\pm$ is a measure-valued kernel, and F is the graph with two vertices and one edge, we get that $M_W^F = M_W$ the measure defined in (16).

Remark 7.4 (Adding missing edges to F). When we work with probability-graphons, we can always assume the graph F to be complete, by adding the missing edges (i, j) and decorating them with the constant function $g_{(i, j)} = \mathbb{1}$.

For a finite weighted graph G , we define the *homomorphism density* of the $C_b(\mathbf{Z})$ -graph F^g in G as $t(F^g, G) = t(F^g, W_G)$ (remind from Remark 6.2 the definition of W_G), that is:

$$t(F^g, G) = \frac{1}{v(G)^k} \sum_{(x_1, \dots, x_k) \in V(G)^k} \prod_{(i, j) \in E(F)} g_{(i, j)}(\Phi_G(x_i, x_j)),$$

where $k = v(F)$ and $\Phi_G(x_i, x_j)$ is the weight of the directed edge from x_i to x_j .

7.2. The Counting Lemma. The following lemma links the homomorphism densities with the cut distance $\delta_{\square, \mathcal{F}}$ for some convergence determining sequence $\mathcal{F} = (f_n)_{n \in \mathbb{N}}$ (with $f_0 = \mathbb{1}$ and f_n takes values in $[0, 1]$). This lemma is a generalization to probability-graphons of the Counting Lemma for real-valued graphons (see Lemmas 10.22 and 10.23 from [Lov12]). Recall that by Remark 7.2, $t(F^g, \cdot)$ is defined on $\widetilde{\mathcal{W}}_\pm$.

Lemma 7.5 (Counting Lemma). *Let $\mathcal{F} = (f_n)_{n \in \mathbb{N}}$ be a convergence determining sequence (with $f_0 = \mathbb{1}$ and f_n takes values in $[0, 1]$). Let F^g be a \mathcal{F} -graph, and for every edge $e \in E(F)$, let $n_e \in \mathbb{N}$ be such that $g_e = f_{n_e}$. Then, for every probability-graphons $W, W' \in \widetilde{\mathcal{W}}_1$, we have:*

$$|t(F^g, W) - t(F^g, W')| \leq \left(\sum_{e \in E(F)} 2^{n_e} \right) \delta_{\square, \mathcal{F}}(W, W').$$

Remark 7.6 ($W \mapsto t(F^g, W)$ is Lipschitz). The Lipschitz constant given by the lemma is too large to be useful in practical cases. Nevertheless, the homomorphism density function $W \mapsto t(F^g, W)$ is Lipschitz on the space of unlabeled probability-graphons $\widetilde{\mathcal{W}}_1$ equipped with the cut distance $\delta_{\square, \mathcal{F}}$.

Proof of Lemma 7.5. To do this proof, we will apply Lemma 10.24 from [Lov12], which applies to graphs F whose edges are decorated with (possibly different) real-valued graphons $w = (w_e : e \in E(F))$, and the associated homomorphism density is defined as

$$(25) \quad t(F, w) = \int_{[0, 1]^{V(F)}} \prod_{(i, j) \in E(F)} w_e(x_i, x_j) \prod_{i \in V(F)} dx_i.$$

Remind from (11) that for a probability-graphon $W \in \mathcal{W}_1$ and a function $f \in \mathcal{F}$ (which is $[0, 1]$ -valued by our definition of convergence determining sequences), we have that $W[f]$ is a real-valued graphon. Define the collections of real-valued graphons $w = (W[g_e] : e \in E(F))$ and $w' = (W'[g_e] : e \in E(F))$. Notice from (24) and (25) that we have $t(F, w) = t(F^g, W)$ and

$t(F, w') = t(F^g, W')$. Applying [Lov12, Lemma 10.24] to the graph F and edge-decorations w and w' , we get:

$$|t(F^g, W) - t(F^g, W')| = |t(F, w) - t(F, w')| \leq \sum_{e \in E(F)} \|W[g_e] - W'[g_e]\|_{\square, \mathbb{R}},$$

where the norm $\|\cdot\|_{\square, \mathbb{R}}$ in the upper bound is the cut norm for real-valued graphons (see (12) for definition of this object). For $e \in E(F)$, by definition of the cut distance $d_{\square, \mathcal{F}}$ and using (10), we have:

$$\|W[g_e] - W'[g_e]\|_{\square, \mathbb{R}} \leq 2^{n_e} d_{\square, \mathcal{F}}(W, W').$$

Hence, combining all those upper bounds, we get the bound in the lemma but with $d_{\square, \mathcal{F}}$ instead of $\delta_{\square, \mathcal{F}}$. Since $t(F^g, \cdot)$ is invariant under relabeling by Remark 7.2, taking the infimum over all relabelings allows to replace $d_{\square, \mathcal{F}}$ by $\delta_{\square, \mathcal{F}}$ and to get the bound in the lemma. \square

We have just seen that homomorphism densities defined using only functions from \mathcal{F} are Lipschitz. We are going to see that the other homomorphism densities are nevertheless continuous.

Lemma 7.7 (Weak Counting Lemma). *Let \mathcal{F} be a convergence determining sequence (with $f_0 = \mathbf{1}$). Let $(W_n)_{n \in \mathbb{N}}$ and W be probability-graphons such that $\lim_{n \rightarrow \infty} t(F^g, W_n) = t(F^g, W)$ for all \mathcal{F} -graph F^g (which in particular the case if $\lim_{n \rightarrow \infty} \delta_{\square, \mathcal{F}}(W_n, W) = 0$ by the Counting Lemma 7.5). Then, for every $C_b(\mathbf{Z})$ -graph F^g we have:*

$$t(F^g, W_n) \xrightarrow[n \rightarrow \infty]{} t(F^g, W).$$

Proof. Let $F = (V, E)$ be some fixed (directed) graph. By assumption, we have for all edge-decorations $g = (g_e)_{e \in E}$ in \mathcal{F} that $\lim_{n \rightarrow \infty} M_{W_n}^F(\otimes_{e \in E} g_e) = M_W^F(\otimes_{e \in E} g_e)$ (see Definition 7.1). By [EK09, Chapter 3, Proposition 4.6], $\mathcal{F}^{\otimes E}$ is a (countable) convergence determining family on $\mathcal{M}_+(\mathbf{Z}^E)$. Thus, the sequence of measures $(M_{W_n}^F)_{n \in \mathbb{N}}$ converges to M_W^F for the weak topology on $\mathcal{M}_+(\mathbf{Z}^E)$. And in particular, for every edge-decoration function $g = (g_e)_{e \in E}$ (here for every $e \in E$, $g_e \in C_b(\mathbf{Z})$ is arbitrary) we have $M_{W_n}^F(\otimes_{e \in E} g_e) = t(F^g, W_n) \rightarrow t(F^g, W) = M_W^F(\otimes_{e \in E} g_e)$ as $n \rightarrow \infty$. This being true for all choices of the graph F , it concludes the proof. \square

7.3. The Inverse Counting Lemma. The goal of this subsection is to establish a converse to the Counting Lemma: if two probability-graphons are close in terms of homomorphism densities, then they are close w.r.t. the cut distance $\delta_{\square, \mathcal{F}}$.

Lemma 7.8 (Inverse Counting Lemma). *Let $\mathcal{F} = (f_n)_{n \in \mathbb{N}}$ be a convergence determining sequence (with $f_0 = \mathbf{1}$ and f_n takes values in $[0, 1]$). Let $U, W \in \mathcal{W}_1$ be two probability-graphons, and let $k, n_0 \in \mathbb{N}^*$. Assume that we have $|t(F^g, U) - t(F^g, W)| \leq 2^{-k - n_0 k^2}$ for every (complete) $C_b(\mathbf{Z})$ -graph F^g with k vertices and such that the edge-decoration functions $g = (g_e)_{e \in E(F)}$ are products (without repetition) of the functions $(f_n)_{1 \leq n \leq n_0}$ and $(1 - f_n)_{1 \leq n \leq n_0}$. Then, we have:*

$$\delta_{\square, \mathcal{F}}(U, W) \leq \frac{44}{\sqrt{\log(k)}} + 2^{-n_0}.$$

To prove Lemma 7.8, we first need to prove the special case where the space \mathbf{Z} is finite.

Lemma 7.9 (Inverse Counting Lemma, case with finite space \mathbf{Z}). *Assume that the space \mathbf{Z} is finite with cardinality n_1 , for simplicity say $\mathbf{Z} = [n_1]$. Define the indicator functions $f_n : z \mapsto \mathbf{1}_{\{z=n\}}$ for $n \in [n_1]$, in particular $\mathcal{H} = (f_n)_{1 \leq n \leq n_1}$ is a finite convergence determining sequence. Let $U, W \in \mathcal{W}_1$ be two probability-graphons, and let $k \in \mathbb{N}^*$. Assume that we have $|t(F^g, U) - t(F^g, W)| < 2^{-k - \log_2(n_1)k^2}$ for every (complete) \mathcal{H} -graph F^g with k vertices.*

Then, for any (possibly finite) convergence determining sequence \mathcal{F} , we have:

$$\delta_{\square, \mathcal{F}}(U, W) \leq \frac{44}{\sqrt{\log(k)}}.$$

Abusing notations, we can identify a weight-value $n \in \mathbf{Z}$ with its indicator functions f_n , and doing this identification for edge-decoration functions, we can identify a \mathcal{F} -graph F^g with its corresponding weighted graph. In particular, doing so we get $t(F^g, W) = \mathbb{P}(\mathbb{G}(k, W) = F^g)$ for every \mathcal{F} -graph F^g with k vertices. The proof of Lemma 7.9 is then a straightforward adaptation of the proof of [Lov12, Lemma 10.31 and Lemma 10.32].

Proof of Lemma 7.8. As the functions $(f_n)_{n \in \mathbb{N}}$ take value in $[0, 1]$, for all φ measure-preserving map, for all $S, T \subset [0, 1]$ measurable sets and for all $n \in \mathbb{N}$, we have:

$$\left| U(S \times T; f_n) - W^\varphi(S \times T; f_n) \right| \leq 1.$$

Using this bound, we get the following bound (remind that $f_0 = \mathbb{1}$):

$$(26) \quad \delta_{\square, \mathcal{F}}(U, W) \leq \inf_{\varphi \in \mathcal{S}[0,1]} \sup_{S, T \subset [0,1]} \sum_{n=1}^{n_0} 2^{-n} \left| U(S \times T; f_n) - W^\varphi(S \times T; f_n) \right| + 2^{-n_0}.$$

Hence, for a point $z \in \mathbf{Z}$, the upper bound in (26) uses only the information given by $(f_n(z))_{n \in [n_0]}$. In order to discretize the space $[0, 1]^{n_0}$, we replace a point $p = (p_1, \dots, p_{n_0}) \in [0, 1]^{n_0}$ by a random point $(Y_1, \dots, Y_{n_0}) \in \{0, 1\}^{n_0}$ where the Y_i are independent random variables with Bernoulli distribution of parameter p_i . This leads us to replace a $\mathcal{M}_1(\mathbf{Z})$ -valued kernel W by the $\mathcal{M}_1(\{0, 1\}^{n_0})$ -valued kernel \tilde{W} defined for all $(x, y) \in [0, 1]^2$, and for all $s = (s_1, \dots, s_{n_0}) \in \{0, 1\}^{n_0}$ as:

$$\tilde{W}(x, y; \{s\}) = W(x, y; f^s) \quad \text{where} \quad f^s = \prod_{n=1}^{n_0} f_n^{s_n} (1 - f_n)^{1-s_n}.$$

Fix some enumeration $(s^m)_{m \in [2^{n_0}]}$ of the points in $\{0, 1\}^{n_0}$, and define the indicator functions $\tilde{h}_m : s \mapsto \mathbb{1}_{\{s=s^m\}}$ for $m \in [2^{n_0}]$, in particular $\tilde{\mathcal{H}} = (\tilde{h}_m)_{1 \leq m \leq 2^{n_0}}$ is a finite convergence determining sequence on $\mathcal{M}_+(\{0, 1\}^{n_0})$. Let $F^{\tilde{g}}$ be a $\tilde{\mathcal{H}}$ -graph with vertex set $V(F) = [k]$, and for every edge $e \in E(F)$, let $m_e \in [2^{n_0}]$ be such that $\tilde{g}_e = \tilde{h}_{m_e}$. Define the edge-decoration functions $g = (g_e)_{e \in E(F)}$ for every edge $e \in E(F)$ as $g_e = f^{s^{m_e}}$, then we get:

$$t(F^{\tilde{g}}, \tilde{W}) = \int_{[0,1]^k} \prod_{(i,j) \in E(F)} \tilde{W}(x_i, x_j; \{s^{m_e}\}) \prod_{i=1}^k dx_i = t(F^g, W).$$

Thus, the $\mathcal{M}_1(\{0, 1\}^{n_0})$ -valued graphons \tilde{U} and \tilde{W} inherit the bounds on the homomorphism densities: for every $\tilde{\mathcal{H}}$ -graph $F^{\tilde{g}}$, we have $|t(F^{\tilde{g}}, \tilde{U}) - t(F^{\tilde{g}}, \tilde{W})| \leq 2^{-k-n_0k^2}$.

Define for all $n \in [n_0]$ the function $\tilde{f}_n : s \mapsto \mathbb{1}_{\{s_n=1\}}$, and let $\tilde{\mathcal{F}}$ be the concatenation of $(\tilde{f}_n)_{n \in [n_0]}$ and $\tilde{\mathcal{H}}$, in particular $\tilde{\mathcal{F}}$ is a finite convergence determining sequence on $\mathcal{M}_+(\{0, 1\}^{n_0})$. Finally, as $\delta_{\square, \tilde{\mathcal{F}}}(\tilde{U}, \tilde{W})$ upper bounds the first term in the upper bound of (26), applying Lemma 7.9 with the finite space $\mathbf{Z} = \{0, 1\}^{n_0}$ and $n_1 = 2^{n_0}$, the finite convergence determining sequences $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{H}}$, and the $\mathcal{M}_1(\{0, 1\}^{n_0})$ -valued graphons \tilde{U} and \tilde{W} , we get:

$$\delta_{\square, \mathcal{F}}(U, W) \leq \frac{44}{\sqrt{\ln(k)}} + 2^{-n_0},$$

which concludes the proof. \square

7.4. Subgraph sampling and the topology of probability-graphons. Thanks to the Weak Counting Lemma 7.7 and the Inverse Counting Lemma 7.8, we can formulate a new informative characterization of weak isomorphism, i.e. equality in the space of unlabeled probability-graphons $\widetilde{\mathcal{W}}_1$.

Proposition 7.10 (Characterization of equality for $\delta_{\square, m}$). *Let $U, W \in \mathcal{W}_1$ be two probability-graphons. The following properties are equivalent:*

- (i) $\delta_{\square, m}(U, W) = 0$ for some (and hence for every) choice of the distance d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that the cut distance $d_{\square, m}$ on \mathcal{W}_1 is (invariant) smooth.
- (ii) There exist $\varphi, \psi \in \widetilde{S}_{[0,1]}$ such that $U^\varphi = W^\psi$ almost everywhere on $[0, 1]^2$.
- (iii) $t(F^g, U) = t(F^g, W)$ for all $C_b(\mathbf{Z})$ -graph F^g .
- (iv) $t(F^g, U) = t(F^g, W)$ for all \mathcal{F} -graph F^g .

Proof. The equivalence between Properties (i) and (ii) is a consequence of Proposition 3.18 on the cut distance. Remark 7.2 gives that Property (ii) implies Property (iii). It is clear that Property (iii) implies Property (iv). The Inverse Counting Lemma 7.8 with the Weak Counting Lemma 7.7 give that Property (iv) implies Property (i) (with $d_m = d_{\mathcal{F}}$). Hence, we have the desired equivalence. \square

Thanks to the Weak Counting Lemma 7.7 and the Inverse Counting Lemma 7.8, we get the following characterization of the topology induced by the cut distance $\delta_{\square, m}$ on the space of unlabeled probability-graphons $\widetilde{\mathcal{W}}_1$ in terms of homomorphism densities

Theorem 7.11 (Characterization of the topology induced by $\delta_{\square, m}$). *Let $(W_n)_{n \in \mathbb{N}}$ and W be unlabeled probability-graphons from $\widetilde{\mathcal{W}}_1$. The following properties are equivalent:*

- (i) $\lim_{n \rightarrow \infty} \delta_{\square, m}(W_n, W) = 0$ for some (and hence for every) choice of the distance d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ and the cut distance $d_{\square, m}$ on \mathcal{W}_1 is (invariant) smooth, weakly regular and regular w.r.t. the stepping operator.
- (ii) $\lim_{n \rightarrow \infty} t(F^g, W_n) = t(F^g, W)$ for all $C_b(\mathbf{Z})$ -graph F^g .
- (iii) $\lim_{n \rightarrow \infty} t(F^g, W_n) = t(F^g, W)$ for all \mathcal{F} -graph F^g .
- (iv) For all $k \geq 2$, the sequence of sampled subgraphs $(\mathbb{G}(k, W_n))_{n \in \mathbb{N}}$ converges in distribution to $\mathbb{G}(k, W)$.

In particular, the topology induced by the cut distance $\delta_{\square, \mathcal{F}}$ on the space of unlabeled probability-graphons $\widetilde{\mathcal{W}}_1$ coincides with the topology generated by the homomorphism densities functions $W \mapsto t(F^g, W)$ for all $C_b(\mathbf{Z})$ -graph F^g .

Proof. By Theorem 5.5, convergence for $\delta_{\square, \mathcal{F}}$ is equivalent to convergence for $\delta_{\square, m}$ for every choice of the distance d_m on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ and the cut distance $d_{\square, m}$ on \mathcal{W}_1 is (invariant) smooth, weakly regular and regular w.r.t. the stepping operator. Taking $d_m = d_{\mathcal{P}}$, the Weak Counting Lemma 7.7 gives that Property (i) implies Property (ii). It is clear that Property (ii) implies Property (iii). The Inverse Counting Lemma 7.8 with the Weak Counting Lemma 7.7 give that Property (iii) implies Property (i) (with $d_m = d_{\mathcal{F}}$). Notice that when F is the complete graph with k vertices, M_W^F is the joint measure of all the edge-weights of the random graph $\mathbb{G}(k, W)$, and thus characterizes the distribution random graph $\mathbb{G}(k, W)$. Thus (remind Definition 7.1), Property (ii) and Property (iv) are equivalent. Hence, we have the desired equivalence. \square

Question 7.12 (Do the distances $d_{\square, \mathcal{F}}$ all induce the same topology?). *Even though every distance $\delta_{\square, \mathcal{F}}$ generates the same topology on the space of unlabeled probability-graphons $\widetilde{\mathcal{W}}_1$, it is an open question whether or not this is also the case that every distance $d_{\square, \mathcal{F}}$ induces the same topology on the space of labeled probability-graphons \mathcal{W}_1 .*

The following proposition states that to prove existence of a limit unlabeled probability-graphon it is enough to prove that there exists a convergence determining sequence \mathcal{F} such that for every \mathcal{F} -graph F^g the homomorphism densities $t(F^g, \cdot)$ converge.

Proposition 7.13 (Existence of a limit unlabeled probability-graphon). *Let d_m be a distance on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ such that d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$ and the cut distance $d_{\square, m}$ on \mathcal{W}_1 is (invariant) smooth, weakly regular and regular w.r.t. the stepping operator.*

Let $(W_n)_{n \in \mathbb{N}}$ be sequence of unlabeled probability-graphons in \mathcal{W}_1 that is tight. Let \mathcal{F} be a convergence determining sequence such that for every \mathcal{F} -graph F^g the sequence $(t(F^g, W_n))_{n \in \mathbb{N}}$ converges. Then, there exists an unlabeled probability-graphon $W \in \widetilde{\mathcal{W}}_1$ such that the sequence $(W_n)_{n \in \mathbb{N}}$ converges to W for $\delta_{\square, m}$.

Proof. Since the sequence $(W_n)_{n \in \mathbb{N}}$ is tight, by Theorem 5.1, there exists a subsequence $(W_{n_k})_{k \in \mathbb{N}}$ of the sequence $(W_n)_{n \in \mathbb{N}}$ that converges to some W for $\delta_{\square, m}$. By Theorem 7.11, we have for every \mathcal{F} -graph F^g that $\lim_{k \rightarrow \infty} t(F^g, W_{n_k}) = t(F^g, W)$; and as we already know that the sequence $(t(F^g, W_n))_{n \in \mathbb{N}}$ converges, we have that $\lim_{n \rightarrow \infty} t(F^g, W_n) = t(F^g, W)$. Hence, by Theorem 7.11, we get that the sequence $(W_n)_{n \in \mathbb{N}}$ converges to W for $\delta_{\square, m}$. \square

Remark 7.14. For the special case $\mathbf{Z} = \{0, 1\}$, which is compact, we find back that convergence for real-valued graphons is characterized by the convergence of the homomorphism densities. Notice the tightness condition of Proposition 7.13 is automatically satisfied as \mathbf{Z} is compact.

8. PROOFS OF THEOREM 5.1 AND THEOREM 5.5

We start by proving a lemma that allows to construct a convergent subsequence and its limit kernel for a tight sequence of measure-valued kernels. This lemma is useful for the proofs of both Theorem 5.1 and Theorem 5.5. For the proof of Theorem 5.5, we will also need the convergence to hold simultaneously for two distances δ_{\square} and δ'_{\square} . Remind from Definition 4.1 the definition of the stepfunction $W_{\mathcal{P}}$ for a signed measure-valued kernel W and a finite partition \mathcal{P} of $[0, 1]$. For a finite partition \mathcal{P} of $[0, 1]$, define its diameter as the smallest diameter of its sets, i.e. $\text{diam}(\mathcal{P}) = \min_{S \in \mathcal{P}} \text{diam}(S) = \min_{S \in \mathcal{P}} \sup_{x, y \in S} |x - y|$.

Lemma 8.1 (Convergence using given approximation partitions). *Let d be an invariant smooth distance on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}). Let $(W_n)_{n \in \mathbb{N}}$ be a sequence in \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}) which is tight (resp. uniformly bounded and tight). Further assume that we are given, for every $n, k \in \mathbb{N}$, partitions $\mathcal{P}_{n, k}$ of $[0, 1]$, such that these partitions and the corresponding stepfunctions $W_{n, k} = (W_n)_{\mathcal{P}_{n, k}}$ satisfy the following conditions:*

- (i) *the partition $\mathcal{P}_{n, k+1}$ is a refinement of $\mathcal{P}_{n, k}$,*
- (ii) *$\text{diam}(\mathcal{P}_{n, k}) \leq 2^{-k}$ and $|\mathcal{P}_{n, k}| = m_k$ depends only on k (and not on n),*
- (iii) *$d(W_n, W_{n, k}) \leq 1/(k+1)$.*

Then, there exists a subsequence $(W_{n_{\ell}})_{\ell \in \mathbb{N}}$ of the sequence $(W_n)_{n \in \mathbb{N}}$ and a measure-valued kernel $W \in \mathcal{W}_1$ (resp. $W \in \mathcal{W}_+$ or $W \in \mathcal{W}_{\pm}$) such that $(W_{n_{\ell}})_{\ell \in \mathbb{N}}$ converges to W for δ_{\square} .

Moreover, assume that d' is another invariant smooth distance on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}) such that for every $n \in \mathbb{N}$ and $k \in \mathbb{N}$, $W_{n, k}$ also satisfies:

- (iv) *$d'(W_n, W_{n, k}) \leq 1/(k+1)$.*

Then, there exists a subsequence $(W_{n_{\ell}})_{\ell \in \mathbb{N}}$ of the sequence $(W_n)_{n \in \mathbb{N}}$ and a measure-valued kernel $W \in \mathcal{W}_1$ (resp. $W \in \mathcal{W}_+$ or $W \in \mathcal{W}_{\pm}$) such that $(W_{n_{\ell}})_{\ell \in \mathbb{N}}$ converges to the same measure-valued kernel W simultaneously both for δ_{\square} and for δ'_{\square} , the cut distance associated with d' .

Proof. We adapt here the general scheme from the proof of Theorem 9.23 in [Lov12], but the argument for the convergence of the U_k , defined below, takes into account that measure-valued kernels are infinite-dimensional valued. We set (remind from (2) the definition of $\|\cdot\|_\infty$):

$$C = \sup_{n \in \mathbb{N}} \|W_n\|_\infty < +\infty.$$

The proof is divided into four steps.

Step 1: Without loss of generality, the partitions $\mathcal{P}_{n,k}$ are made of intervals. For every $n \in \mathbb{N}$, we can rearrange the points of $[0, 1]$ by a measure-preserving map so that the partitions $\mathcal{P}_{n,k}$ are made of intervals, and we replace W_n by its rearranged version.

An argument similar to the next lemma is used in the proof in [Lov12, Proof of Theorem 9.23] without any reference. So, we provide a proof and stress that diameters of the partitions shrinking to zero is an important assumption (see Remark 8.3 below).

Lemma 8.2 (Kernel rearrangement with interval partitions). *Let $(\mathcal{P}_k)_{k \in \mathbb{N}}$ be a refining sequence of finite partitions of $[0, 1]$ whose diameter converges to zero. Then, there exist a measure-preserving map $\varphi \in \tilde{S}_{[0,1]}$ and a refining sequence of partitions made of intervals $(\mathcal{Q}_k)_{k \in \mathbb{N}}$ such that for all $k \in \mathbb{N}$, and all set $S \in \mathcal{P}_k$ there exists a set $R \in \mathcal{Q}_k$ such that a.e. $\mathbb{1}_R = \mathbb{1}_{\varphi^{-1}(S)}$.*

In particular, if W is a signed measure-valued kernel, then for $U = W^\varphi$, we have that a.e. $U_{\mathcal{Q}_k} = (W^\varphi)_{\mathcal{Q}_k} = (W_{\mathcal{P}_k})^\varphi$ for all $k \in \mathbb{N}$.

Notice that, according to Remark 4.4, the sequence of refining partition $(\mathcal{P}_k)_{k \in \mathbb{N}}$, with a partition diameter converging to 0, separates points and thus generates the Borel σ -field of $[0, 1]$.

Proof. Consider the infinite Ulam-Harris tree $\mathcal{T}^\infty = \{u_1 \cdots u_k : k \in \mathbb{N}, u_1, \dots, u_k \in \mathbb{N}^*\}$, where for $k = 0$ the empty word $u = \partial$ is called the root node of the tree; for a node $u = u_1 \cdots u_k \in \mathcal{T}^\infty$, we define its height as $h(u) = k$, and if $k > 0$ we define its parent node as $p(u) = u_1 \cdots u_{k-1}$ and we say that u is a child node of $p(u)$. We order vertices on the tree \mathcal{T}^∞ with the lexicographical (total) order $<_{\text{lex}}$. As a first step, we construct a subtree $\mathcal{T} \subset \mathcal{T}^\infty$ that indexes the sets in the partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$, such that for every $k \in \mathbb{N}$, $\mathcal{P}_k = \{S_u : u \in \mathcal{T}, h(u) = k\}$, and such that if $S_v \subset S_u$ with $S_v \in \mathcal{P}_k$ and $S_u \in \mathcal{P}_{k-1}$, then $p(v) = u$.

Without loss of generality, we may assume that $\mathcal{P}_0 = \{[0, 1]\}$, and we label its only set by the empty word ∂ , and we set $S_\partial = [0, 1]$. Then, suppose we have already labeled the sets from $\mathcal{P}_0, \dots, \mathcal{P}_k$, and we proceed to label the sets from \mathcal{P}_{k+1} . Because the partition \mathcal{P}_{k+1} is a refinement of \mathcal{P}_k , we can group the sets of \mathcal{P}_{k+1} by their unique parent set from \mathcal{P}_k , i.e. for every $S_u \in \mathcal{P}_k$, let $\mathcal{O}_u = \{S \in \mathcal{P}_{k+1} : S \subset S_u\}$, then $S_u = \cup_{S \in \mathcal{O}_u} S$. For $S_u \in \mathcal{P}_k$, we fix an arbitrary enumeration of $\mathcal{O}_u = \{S^1, \dots, S^\ell\}$ with $\ell = |\mathcal{O}_u|$, then label the set S^j by u_j , and set $S^j = S_{u_j}$; remark that the parent node of $w = u_j$ is $p(w) = u$, and the height of node w is $h(w) = h(u) + 1 = k + 1$. Hence, we have labeled every set from \mathcal{P}_{k+1} . To finish the construction, we set $\mathcal{T} = \{u : \exists k \in \mathbb{N}, \exists S \in \mathcal{P}_k, S \text{ has label } u\}$.

We now proceed to construct a measure-preserving map ψ such that the image of every set S_u is a.e. equal to an interval, and such that those intervals are ordered w.r.t. to the order of their labels in \mathcal{T} .

Define the map $\sigma : [0, 1] \rightarrow \mathcal{T}^\mathbb{N}$ by $\sigma(x) = (u^k(x))_{k \in \mathbb{N}} \in \mathcal{T}^\mathbb{N}$ where $u^k(x)$ is the only node of \mathcal{T} with height k such that $x \in S_{u^k(x)}$ (and thus $u^{k+1}(x)$ is a child node of $u^k(x)$). Remark that if $u^{k_0}(x) <_{\text{lex}} u^{k_0}(y)$ for some $k_0 \in \mathbb{N}$, then $u^k(x) <_{\text{lex}} u^k(y)$ for every $k \geq k_0$. We extend naturally the total order $<_{\text{lex}}$ from \mathcal{T} to a the total order on $\mathcal{T}^\mathbb{N}$: for $(u^k)_{k \in \mathbb{N}}, (v^k)_{k \in \mathbb{N}} \in \mathcal{T}^\mathbb{N}$, $(u^k)_{k \in \mathbb{N}} <_{\text{lex}} (v^k)_{k \in \mathbb{N}}$ if $u^{k_0} <_{\text{lex}} v^{k_0}$ where k_0 is the smallest k such that $u^k \neq v^k$.

For every $u \in \mathcal{T}$, define:

$$A^-(u) = \bigcup_{v <_{\text{lex}} u : h(v)=h(u)} S_v \quad \text{and} \quad A^+(u) = A^-(u) \cup S_u,$$

and then define $C^-(u) = \lambda(A^-(u))$ and $C^+(u) = \lambda(A^+(u))$. Now, define ψ as, for $x \in [0, 1]$:

$$\psi(x) = \lambda(A^-(x)) \quad \text{where} \quad A^-(x) = \{y \in [0, 1] : \sigma(y) <_{\text{lex}} \sigma(x)\} = \cup_{k \in \mathbb{N}} A^-(u^k(x)).$$

Moreover, as the sequence of partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$ has a diameter that converges to zero, and thus separates points, the map σ is injective. Thus, we also have:

$$\psi(x) = \lambda(A^+(x)) \quad \text{where} \quad A^+(x) = \{y \in [0, 1] : \sigma(y) \leq_{\text{lex}} \sigma(x)\} = A^-(x) \cup \{x\}.$$

Remark that both $A^-(x)$ and $A^+(x)$ are Borel measurable.

Remark that for every $k \in \mathbb{N}$, we have $A^-(u^k(x)) \subset A^-(x) \subset A^+(x) \subset A^+(u^k(x))$. In particular, for every $u \in \mathcal{T}$, we have $\psi(S_u) \subset [C^-(u), C^+(u)]$; however $\psi(S_u)$ is not necessarily an interval, but we shall see that $\lambda(\psi(S_u)) = C^+(u) - C^-(u)$, i.e. $\psi(S_u)$ is a.e. equal to $[C^-(u), C^+(u)]$. Remark that, as the sequence of partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$ is refining, we get that $[C^-(u), C^+(u)] = \cup_{v: p(v)=u} [C^-(v), C^+(v)]$ for every $u \in \mathcal{T} \setminus \{\partial\}$.

As the diameter of the partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$ converges to zero, we have the following alternative formula for ψ :

$$\psi(x) = \lim_{k \rightarrow \infty} C^-(u^k(x)) = \lim_{k \rightarrow \infty} C^+(u^k(x)).$$

For every $k \in \mathbb{N}$, the map $x \mapsto C^-(u^k(x))$ is a simple function (constant on each $S \in \mathcal{P}_k$ and takes finitely-many values), and thus ψ is measurable as a limit of measurable maps.

We outline the rest of the proof. We first prove that ψ is measure-preserving. Secondly, we prove that ψ is a.e. bijective and construct its a.e. inverse map φ . Thirdly, we prove that $(\varphi^{-1}(\mathcal{P}_k))_{k \in \mathbb{N}}$ is a refining sequence of partitions. And lastly, we approximate almost everywhere the sequence of partitions $(\varphi^{-1}(\mathcal{P}_k))_{k \in \mathbb{N}}$ by a sequence of refining partitions composed of intervals.

We now prove that ψ is measure preserving. Remark that $\psi(x)$ is a non-decreasing function of $\sigma(x)$ for the total relation order $<_{\text{lex}}$, i.e. $\psi(y) \leq \psi(x)$ if and only if $\sigma(y) \leq_{\text{lex}} \sigma(x)$. Hence, $\psi^{-1}([0, \psi(x)]) = \{y \in [0, 1] : \sigma(y) \leq_{\text{lex}} \sigma(x)\}$, and we have:

$$\lambda(\psi^{-1}([0, \psi(x)])) = \lambda(\{y \in [0, 1] : \sigma(y) \leq_{\text{lex}} \sigma(x)\}) = \psi(x).$$

Thus, to show that ψ is measure preserving we just need to show that $\psi([0, 1])$ is dense in $[0, 1]$. For every $u \in \mathcal{T}$, as $\psi(S_u) \subset [C^-(u), C^+(u)]$, we know that the interval $[C^-(u), C^+(u)]$ contains at least one point of the form $\psi(x)$. Remark that for all $k \in \mathbb{N}$, we have $[0, 1] = \cup_{u \in \mathcal{T}: h(u)=k} [C^-(u), C^+(u)]$. Hence, as $\lambda([C^-(u), C^+(u)]) = \lambda(S_u) \leq \text{diam}(\mathcal{P}_{h(u)})$ for every $u \in \mathcal{T}$, and as the diameter of the partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$ converges to zero, we know that each interval of positive length contains a point of the form $\psi(x)$ for some $x \in [0, 1]$, which implies that $\psi([0, 1])$ is indeed dense in $[0, 1]$.

We now prove that ψ is a.e. bijective and construct its a.e. inverse map φ . Without loss of generality, assume that there is no set S_u with null measure. Consider two distinct elements $x, y \in [0, 1]$ such that $\sigma(x) <_{\text{lex}} \sigma(y)$. Assume that $\psi(x) = \psi(y)$, and let $N \in \mathbb{N}$ be the last index k such that $u^k(x) = u^k(y)$. Then, for every $k > N$, we have $u^k(x) <_{\text{lex}} u^k(y)$, which implies that $\psi(x) \leq C^+(u^k(x)) \leq C^-(u^k(y)) \leq \psi(y)$; and thus $\psi(x) = \psi(y) = C^+(u^k(x)) = C^-(u^k(y))$, which in turn implies that there is no node of \mathcal{T} between $u^k(x)$ and $u^k(y)$. Remark that this situation is analogous to the terminating decimal versus repeating decimal situation. Hence, we proved that there is no node between $u^{N+1}(x)$ and $u^{N+1}(y)$ and that for every $k > N$, $u^{k+1}(x)$ is the right-most child of $u^k(x)$, and $u^{k+1}(y)$ is the left-most child of $u^k(y)$ (i.e. $u^{k+1}(x) = u^k(x)|\mathcal{O}_{u^k(x)}$ and $u^{k+1}(y) = u^k(y)1$). Remind that the map σ is injective. Putting all of this together, we get that the set $\{(x, y) \in [0, 1] : \psi(x) = \psi(y), x < y\}$ can be indexed by the nodes of \mathcal{T} , and is thus at most countable. Hence, the map ψ is injective on a subset $D \subset [0, 1]$ with measure one (indeed $[0, 1] \setminus D$ is at most countable), and as ψ is measure preserving, we get that

$\psi(D)$ has measure one, and thus ψ is bijective from D to $\psi(D)$, that is, ψ is a.e. bijective. We construct the map φ as the inverse map of ψ for $x \in \psi(D)$ and $\varphi(x) = 0$ for $x \in [0, 1] \setminus \psi(D)$. Without loss of generality, we assume that $0 \notin D$. Thus, φ is the a.e. inverse map of ψ , that is, $\varphi \circ \psi(x) = \psi \circ \varphi(x) = x$ for almost every $x \in [0, 1]$.

We are left to prove that φ is measurable and measure preserving. As we saw that each point $z \in [0, 1]$ as a pre-image $\psi^{-1}(z) = \{x \in [0, 1] : \psi(x) = z\}$ at most countable (indeed of cardinal at most 2), thus [Pur66] insures that ψ is bimeasurable (i.e. ψ is (Borel) measurable and for all Borel set $B \subset [0, 1]$, $\psi(B)$ is also a Borel set). Let $B \subset [0, 1]$ be a Borel set. We have that $\varphi^{-1}(B) = \varphi^{-1}(B \cap D) = \psi(B \cap D)$ is a Borel set, where the first equality uses that $\varphi([0, 1]) = D \cup \{0\}$, the second equality uses that ψ is the inverse of φ on D , and lastly we used that ψ is bimeasurable. We also have that $\varphi^{-1}(B \cup \{0\}) = \varphi^{-1}(B) \cup ([0, 1] \setminus \psi(D))$ is a Borel set. Moreover, we have:

$$\lambda(\varphi^{-1}(B)) = \lambda(\psi(B \cap D)) = \lambda(\psi^{-1}(\psi(B \cap D))) = \lambda(B \cap D) = \lambda(B),$$

where we used that $\varphi^{-1}(B) = \psi(B \cap D)$ for the first equality, that ψ is measure preserving for the second equality, that ψ is bijective from D to $\psi(D)$ for the third equality, and that D has measure one for the last equality. We also have:

$$\lambda(\varphi^{-1}(B \cup \{0\})) = \lambda(\varphi^{-1}(B)) + \lambda([0, 1] \setminus \psi(D)) = \lambda(B) = \lambda(B \cup \{0\}),$$

where we used that $\varphi^{-1}(B) \subset \psi(D)$ and $[0, 1] \setminus \psi(D)$ are disjoint sets for the first equality, that $\lambda(\varphi^{-1}(B)) = \lambda(B)$ and that $\psi(D)$ has measure one for the second equality. Hence, the map φ is measurable and measure preserving.

We now prove that $(\varphi^{-1}(\mathcal{P}_k))_{k \in \mathbb{N}}$ is a refining sequence of partitions. For $k \in \mathbb{N}$, as \mathcal{P}_k is a finite partition of $[0, 1]$, we have that $\varphi^{-1}(\mathcal{P}_k) = \{\varphi^{-1}(S_u) : u \in \mathcal{T}, h(u) = k\}$ is also a finite partition of $[0, 1]$. Moreover, as $(\mathcal{P}_k)_{k \in \mathbb{N}}$ is a refining sequence of partitions, we get that the sequence of partitions $(\varphi^{-1}(\mathcal{P}_k))_{k \in \mathbb{N}}$ is also refining. Remark that the sets $\varphi^{-1}(S_u)$ are not necessarily intervals, they are intervals minus some at most countable sets (this is similar to the unit line minus the Cantor set).

To finish the proof, we are left to construct a refining sequence of partitions made of intervals $(\mathcal{Q}_k)_{k \in \mathbb{N}}$ that agrees almost everywhere with the refining sequence of partitions $(\varphi^{-1}(\mathcal{P}_k))_{k \in \mathbb{N}}$. For $u \in \mathcal{T}$, define $R_u = [C^-(u), C^+(u)[$ (and $R_u = [C^-(u), C^+(u)]$ if u is the unique node such that $v \leq_{\text{lex}} u$ for every $v \in \mathcal{T}$ with $h(v) = h(u)$). As ψ is measure preserving, and as $\psi(S_u) \subset [C^-(u), C^+(u)]$ with $\lambda(S_u) = C^+(u) - C^-(u)$, we get that $\lambda([C^-(u), C^+(u)] \setminus \psi(S_u)) = 0$. As φ is the a.e. inverse map of ψ , we have that a.e. $\mathbb{1}_{\varphi^{-1}(S_u)} = \mathbb{1}_{\psi(S_u)} = \mathbb{1}_{[C^-(u), C^+(u)]} = \mathbb{1}_{R_u}$, i.e. R_u agrees almost everywhere with $\varphi^{-1}(S_u)$. For $k \in \mathbb{N}$, define the finite partition $\mathcal{Q}_k = \{R_u : h(u) = k\}$. Then, by definition of the sets R_u , the sequence of partitions $(\mathcal{Q}_k)_{k \in \mathbb{N}}$ is refining. This concludes the proof. \square

Remark 8.3 (The shrinking diameter assumption is important). Even if it is not stressed in [Lov12, Proof of Theorem 9.23], the measure preserving map φ (a fortiori an a.e. invertible one) in Lemma 8.2 cannot be obtained without any assumption on the refining sequence of partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$ (in our case, we assumed that their diameter converges to zero). Indeed consider the sequence of partitions where for every $k \in \mathbb{N}$, \mathcal{P}_k is composed of the sets:

$$S_{k,j} = [j2^{-k-1}, (j+1)2^{-k-1}[\cup [1/2 + j2^{-k-1}, 1/2 + (j+1)2^{-k-1}[, \quad 0 \leq j < 2^k,$$

i.e. $S_{k,j}$ is the union of two dyadic interval translated by $1/2$, (also add 1 to the set $S_{k,0}$ to get a complete partition). Then, for every $x \in [0, 1/2[$, x and $x + 1/2$ belong to the same set of \mathcal{P}_k for every $k \in \mathbb{N}$; in particular the diameter of the partitions $(\mathcal{P}_k)_{k \in \mathbb{N}}$ does not converge to zero. By contradiction, assume there exist a measure preserving map $\varphi \in \tilde{S}_{[0,1]}$ and a sequence

of interval partitions $(\mathcal{Q}_k)_{k \in \mathbb{N}}$ such that for all $k \in \mathbb{N}$ and all set $S_{k,j} \in \mathcal{P}_k$ with $0 \leq j < 2^k$, there exists a interval set $I_{k,j} \in \mathcal{Q}_k$ such that a.e. $\mathbb{1}_{I_{k,j}} = \mathbb{1}_{\varphi^{-1}(S_{k,j})}$. In particular, the set $I_{k,j}$ must be an interval of length 2^{-k} . Hence, \mathcal{Q}_k is a dyadic partition with stepsize 2^{-k} , and thus the diameter of the partitions $(\mathcal{Q}_k)_{k \in \mathbb{N}}$ converges to zero. For every $x \in [0, 1/2[$, we get that $\text{diam}(\varphi^{-1}(\{x, x + 1/2\})) \leq \text{diam}(\mathcal{Q}_k) = 2^{-k}$ for all $k \in \mathbb{N}$; this implies that $\varphi^{-1}(\{x, x + 1/2\})$ is a singleton, i.e. either $x \notin \varphi([0, 1])$ or $x + 1/2 \notin \varphi([0, 1])$. Hence, we have $\lambda([0, 1/2[\cap\varphi([0, 1])) = \lambda([1/2, 1[\setminus\varphi([0, 1]))$ and $\lambda([0, 1/2[\setminus\varphi([0, 1])) = \lambda([1/2, 1[\cap\varphi([0, 1]))$. As $\lambda([0, 1/2]) = \lambda([0, 1/2[\cap\varphi([0, 1])) + \lambda([0, 1/2[\setminus\varphi([0, 1])) = 1/2$ because φ is measure preserving, we get that $\lambda(\varphi([0, 1])) = \lambda([0, 1/2[\cap\varphi([0, 1])) + \lambda([1/2, 1[\cap\varphi([0, 1])) = 1/2$, which contradicts the fact that φ is measure preserving.

Now, for every $n \in \mathbb{N}$, applying Lemma 8.2 to $(\mathcal{P}_{n,k})_{k \in \mathbb{N}}$ and W_n , we get a measure-preserving map φ_n and a refining sequence of partitions $(\mathcal{P}'_{n,k})_{k \in \mathbb{N}}$ made of intervals such that for all $k \in \mathbb{N}$, and all set $R \in \mathcal{P}'_{n,k}$ there exists a set $S \in \mathcal{P}_{n,k}$ such that a.e. $\mathbb{1}_R = \mathbb{1}_{\varphi_n^{-1}(S)}$. In particular, for all $k \in \mathbb{N}$, the sequence of partitions $(\mathcal{P}_{n,k})_{k \in \mathbb{N}}$ still satisfy (i)–(ii). Set $W'_n = W_n^{\varphi_n}$ and $W'_{n,k} = W_{n,k}^{\varphi_n}$ so that almost everywhere:

$$W'_{n,k} = ((W_n)_{\mathcal{P}_{n,k}})^{\varphi_n} = (W_n^{\varphi_n})_{\mathcal{P}'_{n,k}} = (W')_{\mathcal{P}'_{n,k}}.$$

As d and d' are invariant, we have for every $n, k \in \mathbb{N}$ that $d(W_n, W_{n,k}) = d(W'_n, W'_{n,k})$, and similarly for d' . This insures that the signed measure-valued kernels $(W'_n)_{n \in \mathbb{N}}$ and $(W'_{n,k})_{n \in \mathbb{N}, k \in \mathbb{N}}$, still satisfy (iii)–(iv). Remind that for a measure-valued kernel W and a measure-preserving map φ , $\delta_{\square, m}(W, W^\varphi) = 0$. Hence, we can replace the signed measure-valued kernels $(W_n)_{n \in \mathbb{N}}$ and $(W_{n,k})_{n \in \mathbb{N}, k \in \mathbb{N}}$, by $(W'_n)_{n \in \mathbb{N}}$ and $(W'_{n,k})_{n \in \mathbb{N}, k \in \mathbb{N}}$, and assume that the partitions $\mathcal{P}_{n,k}$ are made of intervals.

Step 2: There exists a subsequence $(W_{n_\ell})_{\ell \in \mathbb{N}}$ such that for every $k \in \mathbb{N}$ and $\epsilon \in \{+, -\}$, the subsequence $(W_{n_\ell, k}^\epsilon)_{\ell \in \mathbb{N}}$ weakly converges, as $\ell \rightarrow \infty$, almost everywhere to a limit, say U_k^ϵ which is a stepfunction adapted to a partition with m_k elements (some elements might be empty sets).

Fix some $k \in \mathbb{N}$. The stepfunctions $(W_{n,k} = (W_n)_{\mathcal{P}_{n,k}})_{n \in \mathbb{N}}$ all have the same number of steps m_k . For $n \in \mathbb{N}$, denote by $\mathcal{P}_{n,k} = \{S_{n,k,i} : 1 \leq i \leq m_k\}$ the interval partition adapted to $W_{n,k}$ where the intervals are order according to the natural order on $[0, 1]$ (note that some intervals might be empty, simply put them at the end). For $n \in \mathbb{N}$ and $1 \leq i \leq m_k$, let $\lambda(S_{n,k,i})$ denote the length of the interval $S_{n,k,i} \in \mathcal{P}_{n,k}$. As the lengths of steps take values in the compact set $[0, 1]$, there exists a subsequence of indices $(n_\ell)_{\ell \in \mathbb{N}}$ such that for every $1 \leq i \leq m_k$, there exists $s_{k,i} \in [0, 1]$ such that $\lim_{\ell \rightarrow \infty} \lambda(S_{n_\ell, k, i}) = s_{k,i}$. Denote by $\mathcal{P}_k = \{S_{k,i} : 1 \leq i \leq m_k\}$ the interval partition composed of m_k intervals where the i -th interval $S_{k,i}$ has length $s_{k,i}$ (note that some intervals might be empty). Up to a diagonal extraction, we can assume that the convergence holds for every $k \in \mathbb{N}$ simultaneously. Remark that for all $n, k \in \mathbb{N}$, the fact that $\mathcal{P}_{n, k+1}$ is a refinement of $\mathcal{P}_{n, k}$ can be simply restated as linear relations on the interval lengths $(\lambda(S_{n, k, i}))_{1 \leq i \leq m_k}$ and $(\lambda(S_{n, k+1, i}))_{1 \leq i \leq m_{k+1}}$. As linear relations are preserved when taking the limit, we get that the partition \mathcal{P}_{k+1} is a refinement of \mathcal{P}_k for all $k \in \mathbb{N}$. We assume from now on that $(W_n)_{n \in \mathbb{N}}$ and $(W_{n,k})_{n \in \mathbb{N}, k \in \mathbb{N}}$, are the corresponding subsequences.

For every $n \in \mathbb{N}$, we decompose $W_n = W_n^+ - W_n^-$ into its positive and negative kernel parts, see Lemma 3.3. For $n, k \in \mathbb{N}$ and $\epsilon \in \{+, -\}$, we define $W_{n,k}^\epsilon = (W_n^\epsilon)_{\mathcal{P}_{n,k}}$. In particular, remark that $W_{n,k} = W_{n,k}^+ - W_{n,k}^-$ and for all $\ell \geq k$, that $W_{n,k}^\epsilon = (W_{n,\ell}^\epsilon)_{\mathcal{P}_{n,k}}$. Let $\epsilon \in \{+, -\}$ and $1 \leq i, j \leq m_k$ such that $s_{k,i} s_{k,j} > 0$ be fixed. For every $n \in \mathbb{N}$, we have on $S_{n,k,i} \times S_{n,k,j}$ that

$W_{n,k}^\varepsilon = \mu_{n,k}^{i,j,\varepsilon} \in \mathcal{M}_+(\mathbf{Z})$ with:

$$\mu_{n,k}^{i,j,\varepsilon}(\cdot) = \frac{1}{\lambda(S_{n,k,i})\lambda(S_{n,k,j})} W_n^\varepsilon(S_{n,k,i} \times S_{n,k,j}; \cdot).$$

We have that:

$$\|\mu_{n,k}^{i,j,\varepsilon}\|_\infty \leq \|W_n\|_\infty \leq C.$$

This gives that the sequence $(\mu_{n,k}^{i,j,\varepsilon})_{n \in \mathbb{N}}$ in $\mathcal{M}_\pm(\mathbf{Z})$ is bounded. We now prove it is tight. Let $\varepsilon > 0$. As $\lim_{n \rightarrow \infty} \lambda(S_{n,k,\ell}) = s_{k,\ell} > 0$ for $\ell = i, j$, we deduce that there exists $c > 0$ such that for every $n \in \mathbb{N}$ large enough and $\ell = i, j$, we have $\lambda(S_{n,k,\ell}) > c$. Set $\varepsilon' = c^2\varepsilon$. As the sequence $(W_n)_{n \in \mathbb{N}}$ in $\widehat{\mathcal{W}}_\pm$ is tight, there exists a compact set $K \subset \mathbf{Z}$ such that for every $n \in \mathbb{N}$, $M_{W_n}(K^c) \leq \varepsilon'$. Hence, for every $n \in \mathbb{N}$ large enough, we have:

$$\mu_{n,k}^{i,j,\varepsilon}(K^c) \leq \frac{1}{\lambda(S_{n,k,i})\lambda(S_{n,k,j})} M_{W_n}(K^c) \leq \varepsilon.$$

This gives that the sequence $(\mu_{n,k}^{i,j,\varepsilon})_{n \in \mathbb{N}}$ in $\mathcal{M}_+(\mathbf{Z})$ is bounded and tight, and thus by Lemma 2.8, it has a convergent subsequence. By diagonal extraction, we can assume there is a subsequence $(W_{n_\ell})_{\ell \in \mathbb{N}}$ such that for all $k \in \mathbb{N}$, all $1 \leq i, j \leq m_k$ such that $s_{k,i}s_{k,j} > 0$, and all $\varepsilon \in \{+, -\}$, the subsequence $(\mu_{n_\ell,k}^{i,j,\varepsilon})_{\ell \in \mathbb{N}}$ weakly converges to a limit say $\mu_k^{i,j,\varepsilon}$. Define the stepfunction $U_k^\varepsilon \in \mathcal{W}_+$ adapted to the partition \mathcal{P}_k which is equal to $\mu_k^{i,j,\varepsilon}$ on $S_{k,i} \times S_{k,j}$ (if $s_{k,i}s_{k,j} = 0$, set $\mu_k^{i,j,\varepsilon} = 0$). We have in particular obtained that, for every $k \in \mathbb{N}$, the subsequence $(W_{n_\ell,k}^\varepsilon)_{\ell \in \mathbb{N}}$ weakly converges a.e. to U_k^ε which is a stepfunction adapted to a partition with m_k elements; this implies that the subsequence $(W_{n_\ell,k})_{\ell \in \mathbb{N}}$ also weakly converges a.e. to $U_k = U_k^+ - U_k^-$. We now assume that $(W_n)_{n \in \mathbb{N}}$ is such a subsequence. With this convention, notice that for all $k, n \in \mathbb{N}$ and $\varepsilon \in \{+, -\}$:

$$(27) \quad \|U_k^\varepsilon\|_\infty \leq \sup_{n \in \mathbb{N}} \|W_{n,k}^\varepsilon\|_\infty \leq \sup_{n \in \mathbb{N}} \|W_n\|_\infty = C < +\infty.$$

Step 3: There exists a subsequence of $(U_k)_{k \in \mathbb{N}}$ which weakly converges to a limit $U \in \mathcal{W}_\pm$ almost everywhere on $[0, 1]^2$. The proof of this step is postponed to the end. Without loss of generality we still write $(U_k)_{k \in \mathbb{N}}$ for this subsequence.

Step 4: We have $\lim_{n \rightarrow \infty} \delta_\square(U, W_n) = \lim_{n \rightarrow \infty} \delta'_\square(U, W_n) = 0$. Let $\varepsilon > 0$. As the cut distances d is smooth, we deduce from Step 3, that for k large enough $d(U, U_k) \leq \varepsilon$. By hypothesis (iii) on the sequence $(W_{n,k})_{n \in \mathbb{N}}$, we also have that for k large enough $d(W_n, W_{n,k}) \leq \varepsilon$. For such large k , as by step 2 the sequence $(W_{n,k})_{n \in \mathbb{N}}$ weakly converges almost everywhere to U_k , and again as the cut distances d is smooth, there is a n_0 such that for every $n \geq n_0$, $d(U_k, W_{n,k}) \leq \varepsilon$. Then for all $n \geq n_0$, we have:

$$\begin{aligned} \delta_\square(U, W_n) &\leq \delta_\square(U, U_k) + \delta_\square(U_k, W_{n,k}) + \delta_\square(W_{n,k}, W_n) \\ &\leq d(U, U_k) + d(U_k, W_{n,k}) + d(W_{n,k}, W_n) \\ &\leq 3\varepsilon. \end{aligned}$$

This gives that $\lim_{n \rightarrow \infty} \delta_\square(W_n, U) = 0$.

If we consider a second distance d' as in the lemma, then similarly $\lim_{n \rightarrow \infty} \delta'_\square(W_n, U) = 0$.

Proof of Step 3. Assume that the claim is true for measure-valued kernels. Then, if $(U_k)_{k \in \mathbb{N}}$ is a sequence of signed-measure valued kernels, applying the claim to $(U_k^\varepsilon)_{k \in \mathbb{N}}$, for $\varepsilon \in \{+, -\}$, we get a measure-valued $U^\varepsilon \in \mathcal{W}_+$ such that the sequence $(U_k^\varepsilon)_{k \in \mathbb{N}}$ weakly converges a.e. to U^ε . Thus, the sequence $(U_k)_{k \in \mathbb{N}}$ weakly converges a.e. to $U = U^+ - U^-$.

Hence, we are left to prove the claim for measure-valued kernels. The proof is divided in four steps. The first three steps also work for signed-measure valued kernels, but the last argument of step 3.d. only works for measures.

Step 3.a: The sequence $(U_k)_{k \in \mathbb{N}}$ inherit the tightness property from the sequence $(W_n)_{n \in \mathbb{N}}$. Let $\varepsilon > 0$. Since the sequence $(W_n)_{n \in \mathbb{N}}$ is tight, there exists a compact set $K \subset \mathbf{Z}$ such that for every $n \in \mathbb{N}$, we have $M_{W_n}(K^c) \leq \varepsilon$. Remark that:

$$M_{W_{n,k}} = \sum_{1 \leq i, j \leq m_k} \lambda(S_{n,k,i}) \lambda(S_{n,k,j}) \mu_{n,k}^{i,j} = M_{W_n} \quad \text{and} \quad M_{U_k} = \sum_{1 \leq i, j \leq m_k} s_{k,i} s_{k,j} \mu_k^{i,j}.$$

For all $k \in \mathbb{N}$ and $1 \leq i, j \leq m_k$, as the sequence $(\mu_{n,k}^{i,j})_{n \in \mathbb{N}}$ weakly converges to $\mu_k^{i,j}$, using [Bog18, Theorem 2.7.4] with the open subset $K^c \subset \mathbf{Z}$, we get that $\mu_k^{i,j}(K^c) \leq \liminf_{n \rightarrow \infty} \mu_{n,k}^{i,j}(K^c)$. As $\lim_{n \rightarrow \infty} \lambda(S_{n,k,i}) = s_{k,i}$ for all $1 \leq i \leq m_k$, and summing those bounds, we get:

$$M_{U_k}(K^c) \leq \liminf_{n \rightarrow \infty} M_{W_{n,k}}(K^c) = \liminf_{n \rightarrow \infty} M_{W_n}(K^c) \leq \varepsilon.$$

Consequently, the sequence $(U_k)_{k \in \mathbb{N}}$ is tight.

Step 3.b: Convergence of the measures \hat{U}_k in $\mathcal{M}_+([0, 1]^2 \times \mathbf{Z})$ defined for $k \in \mathbb{N}$ as:

$$(28) \quad \hat{U}_k(dx, dy, dz) = U_k(x, y; dz) \lambda_2(dx, dy).$$

Since the sequence $(M_{U_k})_{k \in \mathbb{N}}$ in $\mathcal{M}_+(\mathbf{Z})$ is tight, for all $\varepsilon > 0$, there exists a compact set $K \subset \mathbf{Z}$ such that for every $k \in \mathbb{N}$, $M_{U_k}(K^c) \leq \varepsilon$; and thus $\hat{U}_k(\hat{K}^c) = M_{U_k}(K^c) \leq \varepsilon$ where $\hat{K} = [0, 1]^2 \times K$ is a compact subset of $[0, 1]^2 \times \mathbf{Z}$, that is, the sequence $(\hat{U}_k)_{k \in \mathbb{N}}$ in $\mathcal{M}_+([0, 1]^2 \times \mathbf{Z})$ is tight. The sequence $(\hat{U}_k)_{k \in \mathbb{N}}$ is also bounded as $\|\hat{U}_k\|_\infty \leq \|U_k\|_\infty \leq C$ thanks to (27). Hence, using Lemma 2.8, there exists a subsequence $(\hat{U}_{k_\ell})_{\ell \in \mathbb{N}}$ of the sequence $(\hat{U}_k)_{k \in \mathbb{N}}$ that converges to some measure, say \hat{U} , in $\mathcal{M}_+([0, 1]^2 \times \mathbf{Z})$. Remark that, when considering the subsequence of indices $(k_\ell)_{\ell \in \mathbb{N}}$, the subsequences $(W_{n,k_\ell})_{\ell \in \mathbb{N}}$, $n \in \mathbb{N}$, still satisfy properties (i)-(iv) of Lemma 8.1, and for all $\ell \in \mathbb{N}$, the sequence $(W_{n,k_\ell})_{n \in \mathbb{N}}$ still weakly converges to U_{k_ℓ} . Without loss of generality, we now work with this subsequence and thus write k instead of k_ℓ .

Step 3.c: The measure $\hat{U}(dx, dy, dz)$ can be disintegrated w.r.t. $\lambda_2(dx, dy)$ giving us an element of \mathcal{W}_+ . To prove this, we need the following disintegration theorem for measures, see [Kal17, Theorem 1.23] (stated in more the general framework of Borel spaces) which generalizes the disintegration theorem for probability measures [Kal02, Theorem 6.3]. The notation $\mu \sim \nu$ for two measures μ and ν means that $\mu \ll \nu$ and $\nu \ll \mu$, where $\mu \ll \nu$ means that μ is absolutely continuous w.r.t. ν .

Lemma 8.4 (Disintegration theorem for measures, [Kal17, Theorem 1.23]). *Let ρ be a measure on $S \times T$, where S is a measurable space and T a Polish space. Then there exist a measure $\nu \equiv \rho(\cdot \times T)$ on S and a probability kernel $\mu : S \rightarrow \mathcal{M}_1(T)$ such that $\rho = \nu \otimes \mu$ (i.e. $\rho(ds, dt) = \nu(ds) \mu(s; dt)$). Moreover, the measures $\mu_s = \mu(s; \cdot)$ are unique for ν -a.e. $s \in S$.*

Using Lemma 8.4 with $S = [0, 1]^2$ and $T = \mathbf{Z}$, we get that there exists a probability kernel U' in \mathcal{W}_1 such that:

$$\hat{U}(dx, dy, dz) = U'(x, y; dz) \pi(dx, dy),$$

where $\pi = \hat{U}(\cdot \times \mathbf{Z})$ is a measure on $[0, 1]^2$.

We now need to prove that $\pi \ll \lambda_2$. By contradiction, assume this is false, then there exists a measurable set $A \in \mathcal{B}([0, 1]^2)$ such that $\lambda_2(A) = 0$ and $\pi(A) > 0$. As the measure $\int_A U'(x, y; \cdot) \pi(dx, dy)$ is not null, there exists $f \in C_b(\mathbf{Z})$ such that $\int_A U'(x, y; f) \pi(dx, dy) \neq 0$. As the sequence $(\hat{U}_k)_{k \in \mathbb{N}}$ weakly converges to \hat{U} in $\mathcal{M}_+([0, 1]^2 \times \mathbf{Z})$ by step 3.b, we have that the sequence of measures $\hat{U}_k(dx, dy; f) = U_k(x, y; f) \lambda_2(dx, dy)$ weakly converges as $k \rightarrow \infty$ to $\hat{U}(dx, dy; f) = U'(x, y; f) \pi(dx, dy)$ in $\mathcal{M}_+([0, 1]^2)$. Moreover, as the maps $x, y \mapsto U_k(x, y; f)$ are uniformly bounded (by $\|f\|_\infty \|U_k\|_\infty \leq C \|f\|_\infty$, see (27)), they are also uniformly integrable (w.r.t. λ_2), and applying [Bog07a, Corollary 4.7.19] there exist a subsequence $(U_{k_\ell})_{\ell \in \mathbb{N}}$ and a

bounded function g_f on $[0, 1]^2$ such that for every bounded measurable function $h \in L^\infty([0, 1]^2)$, we have:

$$\lim_{\ell \rightarrow \infty} \int U_{k_\ell}(x, y; f) h(x, y) \lambda_2(dx, dy) = \int g_f(x, y) h(x, y) \lambda_2(dx, dy).$$

In particular, the sequence of measures $(U_{k_\ell}(x, y; f) \lambda_2(dx, dy))_{\ell \in \mathbb{N}}$ weakly converges to the measure $g_f(x, y) \lambda_2(dx, dy)$, which imposes the equality between measures:

$$\hat{U}(dx, dy, f) = U'(x, y; f) \pi(dx, dy) = g_f(x, y) \lambda_2(dx, dy).$$

Hence, taking $h = \mathbb{1}_A$, we get:

$$\hat{U}(A, f) = \int_A U'(x, y; f) \pi(dx, dy) = \int_A g_f(x, y) \lambda_2(dx, dy) = 0,$$

which yields a contradiction. Consequently, the measure π is absolutely continuous w.r.t. λ_2 , with density still denoted by π , and we set λ_2 -a.e. on $[0, 1]^2$:

$$(29) \quad U(x, y; dz) = \pi(x, y) U'(x, y; dz) \quad \text{and thus} \quad \hat{U}(dx, dy, dz) = U(x, y; dz) \lambda_2(dx, dy).$$

Step 3.d: The sequence $(U_k)_{k \in \mathbb{N}}$ weakly converges to U almost everywhere on $[0, 1]^2$. Recall that by construction, the stepfunction U_k is adapted to the partition \mathcal{P}_k defined in Step 2, and that \mathcal{P}_{k+1} is a refinement of \mathcal{P}_k . A closer look at Step 2 yields that for all $\ell \geq k$, since $W_{n,k} = (W_{n,\ell})_{\mathcal{P}_{n,k}}$, we also get:

$$(30) \quad U_k = (U_\ell)_{\mathcal{P}_k}.$$

We prove (30) for $\ell = k + 1$, the other cases follow by induction. As $\mathcal{P}_{n,k+1}$ is a refinement of $\mathcal{P}_{n,k}$, we already know that U_k and $(U_{k+1})_{\mathcal{P}_k}$ are both stepfunctions adapted to the finite partition \mathcal{P}_k . Thus, we only need to verify that U_k and $(U_{k+1})_{\mathcal{P}_k}$ take the same value on each step. For every $n \in \mathbb{N}$, the fact that $W_{n,k} = (W_{n,k+1})_{\mathcal{P}_{n,k}}$ implies that for all $1 \leq i, j \leq m_k$ such that $\lambda(S_{n,k,i})\lambda(S_{n,k,j}) > 0$, we have:

$$\mu_{i,j}^{n,k} = \sum_{i' \in I_i, j' \in I_j} \frac{\lambda(S_{n,k+1,i'})\lambda(S_{n,k+1,j'})}{\lambda(S_{n,k,i})\lambda(S_{n,k,j})} \mu_{i',j'}^{n,k+1},$$

and this equation is preserved when taking the limit $n \rightarrow \infty$, which gives us:

$$\mu_{i,j}^k = \sum_{i' \in I_i, j' \in I_j} \frac{s_{k+1,i'} s_{k+1,j'}}{s_{k,i} s_{k,j}} \mu_{i',j'}^{k+1}, \quad \text{for all } 1 \leq i, j \leq m_k \text{ such that } s_{k,i} s_{k,j} > 0.$$

This proves that the stepfunctions U_k and $(U_{k+1})_{\mathcal{P}_k}$ take the same value on each step $S_{k,i} \times S_{k,j}$ with positive size $s_{k,i} s_{k,j} > 0$ (on a step with null size $s_{k,i} s_{k,j} = 0$, U_k and $(U_{k+1})_{\mathcal{P}_k}$ are both equal to the null measure). This gives that $U_k = (U_{k+1})_{\mathcal{P}_k}$.

Let $f \in C_b(\mathbf{Z})$ be a bounded continuous function, and X, Y be independent uniform random variables on $[0, 1]$. Then (30) and (27) imply that the sequence $N^f = (N_k^f = U_k(X, Y; f))_{k \in \mathbb{N}}$ is a martingale bounded by $C\|f\|_\infty$ for the filtration $(\mathcal{F}_k)_{k \in \mathbb{N}}$, where the σ -field \mathcal{F}_k is generated by the events $\{X \in S_{k,i}\} \cap \{Y \in S_{k,j}\}$ for $1 \leq i, j \leq m_k$ and $S_{k,\ell} \in \mathcal{P}_k$. By the martingale convergence theorem, the martingale N^f is almost surely convergent, that is, the sequence $(U_k[f])_{k \in \mathbb{N}}$ converges λ_2 -a.e. to a bounded measurable function u_f . Let $g : [0, 1]^2 \rightarrow \mathbb{R}$ be a

bounded measurable function. We get:

$$\begin{aligned}
\int_{[0,1]^2} g(x,y) U(x,y;f) \lambda_2(\mathrm{d}x\mathrm{d}y) &= \int_{[0,1]^2 \times \mathbf{Z}} g(x,y) f(z) \hat{U}(\mathrm{d}x, \mathrm{d}y, \mathrm{d}z) \\
&= \lim_{k \rightarrow \infty} \int_{[0,1]^2 \times \mathbf{Z}} g(x,y) f(z) \hat{U}_k(\mathrm{d}x, \mathrm{d}y, \mathrm{d}z) \\
&= \lim_{k \rightarrow \infty} \mathbb{E}[g(X,Y) U_k(X,Y;f)] \\
&= \int_{[0,1]^2} g(x,y) u_f(x,y) \lambda_2(\mathrm{d}x\mathrm{d}y),
\end{aligned}$$

where we used the definition (29) of U for the first equality, that $(\hat{U}_k)_{k \in \mathbb{N}}$ weakly converges to \hat{U} for the second, the definition (28) of \hat{U}_k for the third, and the convergence of the martingale N^f for the last. Since g is arbitrary, we deduce that λ_2 -a.e. $U(\cdot, \cdot; f) = u_f$ and thus that the sequence $(U_k[f])_{k \in \mathbb{N}}$ converges λ_2 -a.e. to $U[f]$. Applying this result for all $f \in \mathcal{F} = (f_m)_{m \in \mathbb{N}}$ a convergence determining sequence (with the convention $f_0 = \mathbf{1}$), we deduce that the sequence $(U_k)_{k \in \mathbb{N}}$ weakly converges to U almost everywhere on $[0, 1]^2$. Remind from Section 2 that convergence determining sequences exist only for measures and not for signed measures in general, this is why we worked with measures in Step 3. This ends the proof of Step 3, and thus ends the proof of the lemma. \square

We are now ready to prove Theorem 5.1.

Proof of Theorem 5.1. We first prove Point (i) on \mathcal{W}_\pm (the proof on \mathcal{W}_1 is similar). Since the distance d is weakly regular and the sequence $(W_n)_{n \in \mathbb{N}}$ is uniformly bounded and tight in \mathcal{W}_\pm , we can construct inductively for every $n \in \mathbb{N}$ a sequence $(\mathcal{P}_{n,k})_{k \in \mathbb{N}}$ of partitions of $[0, 1]$ such that hypothesis (i)-(iii) of Lemma 8.1 are satisfied: $\mathcal{P}_{n,k+1}$ being obtained by applying the weak regularity property (see Definition 4.10-(i)) with starting partition $\mathcal{Q}_{n,k} = \mathcal{P}_{n,k} \wedge \mathcal{D}_k$, where \mathcal{D}_k is the dyadic partition with stepsize $2^{-(k+1)}$. (We may assume that the partitions $\mathcal{P}_{n,k}$ for all $n \in \mathbb{N}$ have the same size m_k by adding empty sets.) Then as d is also invariant and smooth on \mathcal{W}_\pm , the first part of Lemma 8.1 directly gives Point (i).

Before proving Point (ii), we first need to prove the following lemma.

Lemma 8.5 (Compactness theorem for $\mathcal{W}_\mathcal{M}$). *Let d be an invariant, smooth and weakly regular distance on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_\pm). Let \mathcal{M} be a convex and weakly closed subset of $\mathcal{M}_1(\mathbf{Z})$ (resp. $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_\pm(\mathbf{Z})$). Let $(W_n)_{n \in \mathbb{N}}$ be a sequence of \mathcal{M} -valued kernels which is tight and uniformly bounded. Then, $(W_n)_{n \in \mathbb{N}}$ has a subsequence that converges for δ_\square to some \mathcal{M} -valued kernel.*

Proof. First remark that, as \mathcal{M} is convex, the image of $\mathcal{W}_\mathcal{M}$ by the stepping operator $W \mapsto W_\mathcal{P}$, where \mathcal{P} is a finite partition of $[0, 1]$, is a subset of $\mathcal{W}_\mathcal{M}$. Hence, a close look at the proof of Lemma 8.1 (the partitions are constructed as in the proof of Point (i) from Theorem 5.1), and using the notation therein, shows that, up to taking subsequences, one can take the stepping kernels $W_{n,k}$ and U_k in $\mathcal{W}_\mathcal{M}$, such that $(U_k)_{k \in \mathbb{N}}$ weakly converges to U a.e. and the subsequence $(W_{n_\ell})_{\ell \in \mathbb{N}}$ converges to U w.r.t. δ_\square . Since $U_k(x, y; \cdot) \in \mathcal{M}$ weakly converges to $U(x, y; \cdot)$ for almost every $x, y \in [0, 1]$ and since \mathcal{M} is weakly closed (and thus sequentially weakly closed), we deduce that $U(x, y; \cdot)$ belongs to \mathcal{M} for almost every $x, y \in [0, 1]$. This means that $U \in \mathcal{W}_\mathcal{M}$. \square

We prove Point (ii) for $\mathcal{M} \subset \mathcal{M}_\pm(\mathbf{Z})$ (the proof for $\mathcal{M} \subset \mathcal{M}_1(\mathbf{Z})$ is identical). The fact that $\mathcal{W}_\mathcal{M}$ and $\widetilde{\mathcal{W}}_\mathcal{M}$ are convex is clear as \mathcal{M} is convex. Let $(W_n)_{n \in \mathbb{N}}$ be a sequence of elements of $\widetilde{\mathcal{W}}_\mathcal{M}$. Since \mathcal{M} is convex, we deduce that $(M_{W_n})_{n \in \mathbb{N}}$ is a sequence in \mathcal{M} . As \mathcal{M} is sequentially compact for the weak topology, \mathcal{M} is tight and bounded by Lemma 2.8, and thus the sequence $(W_n)_{n \in \mathbb{N}}$ is tight and uniformly bounded (remind Definition 4.7). Hence, using Lemma 8.5, we

get that from any sequence in $\widetilde{\mathcal{W}}_{\mathcal{M}}$, we can extract a subsequence which converges for δ_{\square} to an element in $\widetilde{\mathcal{W}}_{\mathcal{M}}$. This implies that $(\widetilde{\mathcal{W}}_{\mathcal{M}}, \delta_{\square})$ is compact.

Point (iii) is a direct consequence of Point (ii) as if \mathbf{Z} is compact, so is $\mathcal{M}_1(\mathbf{Z})$. \square

Proof of Point (iii) from Proposition 5.2. We prove Point (iii). The fact that $\mathcal{W}_{\mathcal{M}}$ and $\widetilde{\mathcal{W}}_{\mathcal{M}}$ are convex is clear as \mathcal{M} is convex. To prove that $\widetilde{\mathcal{W}}_{\mathcal{M}}$ is closed, we consider a sequence $(W_n)_{n \in \mathbb{N}}$ in $\widetilde{\mathcal{W}}_{\mathcal{M}}$ that converges for $\delta_{\square, m}$ to some $W \in \widetilde{\mathcal{W}}_{\pm}$. As $(W_n)_{n \in \mathbb{N}}$ is a Cauchy sequence for $\delta_{\square, m}$, by Lemma 4.9, $(M_{W_n})_{n \in \mathbb{N}}$ is a Cauchy sequence for d_m and thus is tight. Hence, $(W_n)_{n \in \mathbb{N}}$ is uniformly bounded and tight. Applying Lemma 8.5, there exists a subsequence $(W_{n_k})_{k \in \mathbb{N}}$ of the sequence $(W_n)_{n \in \mathbb{N}}$ which converges for $\delta_{\square, m}$ to some \mathcal{M} -valued kernel $U \in \widetilde{\mathcal{W}}_{\mathcal{M}}$. But as a subsequence, $(W_{n_k})_{k \in \mathbb{N}}$ must also converge for $\delta_{\square, m}$ to W . This implies that $W = U$ is a \mathcal{M} -valued kernel. \square

In order to prove Theorem 5.5, we first prove a lemma that allows to construct the partitions needed to use Lemma 8.1.

Lemma 8.6 (Construction of partitions for two distances). *Let d and d' be two distances on \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}) which are invariant, smooth, weakly regular and regular w.r.t. the stepping operator (see Definitions 3.10 and 4.10). Let $(W_n)_{n \in \mathbb{N}}$ be a sequence in \mathcal{W}_1 (resp. \mathcal{W}_+ or \mathcal{W}_{\pm}) which is tight (resp. uniformly bounded and tight). Then, there exists sequences $(\mathcal{P}_{n,k})_{k \in \mathbb{N}}$, $n \in \mathbb{N}$, of partitions of $[0, 1]$ such that hypothesis (i)–(iv) of Lemma 8.1 are satisfied.*

Proof. We prove the result on \mathcal{W}_{\pm} (the proof on \mathcal{W}_1 and \mathcal{W}_+ is similar). To simplify notations, write $d^1 = d$ and $d^2 = d'$. We proceed by induction on $k \in \mathbb{N} \cup \{-1\}$. For every $n \in \mathbb{N}$, set $\mathcal{P}_{n,-1} = \{\{0, 1\}\}$ the trivial partition with size 1. Let $k \in \mathbb{N}$ and assume that we have already constructed partitions $(\mathcal{P}_{n,k-1})_{n \in \mathbb{N}}$ that have the same size m_{k-1} . Now we proceed to construct partitions $(\mathcal{P}_{n,k})_{n \in \mathbb{N}}$ that satisfy hypothesis (i)–(iv).

Set $C = \sup_{n \in \mathbb{N}} \|W_n\|_{\infty}$, which is finite as the sequence $(W_n)_{n \in \mathbb{N}}$ is uniformly bounded. As d^i , with $i = 1, 2$, are regular w.r.t. the stepping operator, there exists a finite constant $C_0 > 0$ such that for every $W, U \in \mathcal{W}_{\pm}$, with $\|W\|_{\infty} \leq C$ and $\|U\|_{\infty} \leq C$, and U a stepfunction adapted to a finite partition \mathcal{Q} :

$$(31) \quad d^i(W, W_{\mathcal{Q}}) \leq C_0 d^i(W, U).$$

Set $\varepsilon = 1/C_0(k+1)$. Since d^i , with $i = 1, 2$, are weakly regular and the sequence $(W_n)_{n \in \mathbb{N}}$ is tight and uniformly bounded, there exists $r_k \in \mathbb{N}^*$, such that for every $n \in \mathbb{N}$, there exists a partition $\mathcal{R}_{n,k}^i$ of $[0, 1]$ that refines $\mathcal{Q}_{n,k} = \mathcal{P}_{n,k-1} \wedge \mathcal{D}_k$, where \mathcal{D}_k is the dyadic partition with stepsize 2^{-k} , such that:

$$(32) \quad |\mathcal{R}_{n,k}^i| \leq r_k |\mathcal{Q}_{n,k}| \leq 2^k r_k |\mathcal{P}_{n,k-1}| \quad \text{and} \quad d^i(W_n, (W_n)_{\mathcal{R}_{n,k}^i}) \leq \varepsilon.$$

(Indeed, a close look at the proof shows that $\mathcal{P}_{n,k-1}$ refines \mathcal{D}_{k-1} by construction, thus $\mathcal{Q}_{n,k}$ cuts each set of $\mathcal{P}_{n,k-1}$ in at most 2 sets, and we get $|\mathcal{Q}_{n,k}| \leq 2|\mathcal{P}_{n,k-1}|$.) Now, let $\mathcal{P}_{n,k}$ be the common refinement of $\mathcal{R}_{n,k}^1$ and $\mathcal{R}_{n,k}^2$; it is a refinement of $\mathcal{P}_{n,k-1}$, has diameter at most 2^{-k} and size:

$$|\mathcal{P}_{n,k}| \leq 2^{2k} r_k^2 |\mathcal{P}_{n,k-1}|^2 = 2^{2k} r_k^2 m_{k-1}^2.$$

If necessary, by completing $\mathcal{P}_{n,k}$ with null sets, we may assume that $|\mathcal{P}_{n,k}| = m_k$, where $m_k = 2^{2k} r_k^2 m_{k-1}^2$. As $(W_n)_{\mathcal{R}_{n,k}^i}$ is a stepfunction adapted to the partition $\mathcal{P}_{n,k}$, we deduce from (31) and (32) that for $i = 1, 2$ and $n \in \mathbb{N}$:

$$d^i(W_n, (W_n)_{\mathcal{P}_{n,k}}) \leq C_0 d^i(W_n, (W_n)_{\mathcal{R}_{n,k}^i}) \leq C_0 \varepsilon = \frac{1}{k+1}.$$

Hence, for every $n \in \mathbb{N}$, the partition $\mathcal{P}_{n,k}$ satisfies the hypothesis (i)-(iv) of Lemma 8.1. Thus, the induction is complete. \square

Proof of Theorem 5.5. Let d_m and $d_{m'}$ be as in Theorem 5.5.

Let $(W_n)_{n \in \mathbb{N}}$ be a sequence of probability-graphons that converges to some $W \in \widetilde{\mathcal{W}}_1$ for $\delta_{\square, m}$. By Lemma 4.9, the sequence of probability measure $(M_{W_n})_{n \in \mathbb{N}}$ converges to M_W for the distance d_m . As d_m induces the weak topology on $\mathcal{M}_{\leq 1}(\mathbf{Z})$, we have that the sequence $(M_{W_n})_{n \in \mathbb{N}}$ is tight, and thus the sequence $(W_n)_{n \in \mathbb{N}}$ is also tight (remind Definition 4.7). The sequence $(W_n)_{n \in \mathbb{N}}$ is also uniformly bounded as a sequence in $\widetilde{\mathcal{W}}_1$. Applying Lemma 8.6 with the distances $d = d_{\square, m}$ and $d' = d_{\square, m'}$, which are invariant, smooth, weakly regular and regular w.r.t. the stepping operator, we get sequences of partitions $(\mathcal{P}_{n,k})_{k \in \mathbb{N}}$, $n \in \mathbb{N}$, that satisfy hypothesis (i)-(iv) of Lemma 8.1. We then deduce from the last part of Lemma 8.1 that any subsequence of $(W_n)_{n \in \mathbb{N}}$ has a further subsequence which converges to the same limit for both $\delta_{\square, m}$ and $\delta_{\square, m'}$, this limit must then be W . This implies that the sequence $(W_n)_{n \in \mathbb{N}}$ converges to W for $\delta_{\square, m'}$.

The role of d_m and $d_{m'}$ being symmetric, we conclude that the distances $\delta_{\square, m}$ and $\delta_{\square, m'}$ induce the same topology on $\widetilde{\mathcal{W}}_1$. \square

ACKNOWLEDGEMENT

We thank Pierre-André Zitt for some helpful discussion in particular on Lemma 3.20.

✂

Index of notation

<p style="text-align: center;">Measures</p> <ul style="list-style-type: none"> - $(\mathbf{Z}, \mathcal{O}_{\mathbf{Z}})$ a topological Polish space - $\mathcal{B}(\mathbf{Z})$ the Borel σ-field induced by $\mathcal{O}_{\mathbf{Z}}$ - $C_b(\mathbf{Z})$ the set of continuous bounded real-valued functions on \mathbf{Z} - <i>measure</i> = positive measure - $\mathcal{M}_{\pm}(\mathbf{Z})$ the set of signed measures on \mathbf{Z} - $\mathcal{M}_+(\mathbf{Z})$ the set of measures on \mathbf{Z} - $\mathcal{M}_1(\mathbf{Z})$ the set of probability measures on \mathbf{Z} - $\mathcal{M}_{\leq 1}(\mathbf{Z})$ the set of sub-probability measures on \mathbf{Z}, i.e. measures with total mass at most 1 - μ^+, μ^- the positive and negative parts of μ from its Hahn-Jordan decomposition - $\mu = \mu_+ + \mu_-$ the total variation measure of μ - $\ \mu\ _{\infty} = \mu (\mathbf{Z})$ the total mass of μ - d_m a distance on either $\mathcal{M}_{\leq 1}(\mathbf{Z})$, $\mathcal{M}_+(\mathbf{Z})$ or $\mathcal{M}_{\pm}(\mathbf{Z})$ 	<ul style="list-style-type: none"> - N_m a norm on $\mathcal{M}_{\pm}(\mathbf{Z})$ - $d_{\mathcal{P}}$ the Prohorov distance - $\ \cdot\ _{\text{KR}}$ the Kantorovitch-Rubinstein norm - $\ \cdot\ _{\text{FM}}$ the Fortet-Mourier norm - $\ \cdot\ _{\mathcal{F}}$ the norm based on a convergence determining sequence \mathcal{F} <hr/> <p style="text-align: center;">Relabelings and partitions</p> <ul style="list-style-type: none"> - $S_{[0,1]}$ the set of bijective measure-preserving maps from $([0, 1], \lambda)$ to itself - $\bar{S}_{[0,1]}$ the set of measure-preserving maps from $([0, 1], \lambda)$ to itself - \mathcal{P} the number of sets in the finite partition \mathcal{P} <hr/> <p style="text-align: center;">Kernels and graphons spaces</p> <ul style="list-style-type: none"> - \mathcal{W}_1 the set of probability-graphons - \mathcal{W}_+ the set of measure-valued kernels - \mathcal{W}_{\pm} the set of signed measure-valued kernels
--	--

<ul style="list-style-type: none"> - $\mathcal{W}_{\mathcal{M}}$ the set of \mathcal{M}-valued kernels with $\mathcal{M} \subset \mathcal{M}_{\pm}(\mathbf{Z})$ - $\widetilde{\mathcal{W}}_1$ the set of unlabeled probability-graphons - $\widetilde{\mathcal{W}}_+$ the set of unlabeled measure-valued kernels - $\widetilde{\mathcal{W}}_{\pm}$ the set of unlabeled signed measure-valued kernels - $\widetilde{\mathcal{W}}_{\mathcal{M}}$ the set of unlabeled \mathcal{M}-valued kernels 	<ul style="list-style-type: none"> - F^g a finite graph whose edges are decorated with functions in $C_b(\mathbf{Z})$ - $t(F^g, W) = M_W^F(g)$ the homomorphism density of F^g in W
<hr/>	
Kernels and graphons	
<ul style="list-style-type: none"> - W^+ and W^- the positive and negative part of $W \in \mathcal{W}_{\pm}$ - $W = W^+ + W^-$ - $W(A; \cdot) = \int_A W(x, y; \cdot) dx dy$ for $A \subset [0, 1]^2$ - $W[f](x, y) = W(x, y; f)$ for $f \in C_b(\mathbf{Z})$ - $W_{\mathcal{P}}$ the stepping of W w.r.t. a partition \mathcal{P} - $\ W\ _{\infty} := \sup_{x, y \in [0, 1]} \ W(x, y; \cdot)\ _{\infty}$ - $M_W(dz) = W ([0, 1]^2; dz)$ - W_G the probability-graphon associated to a $\mathcal{M}_1(\mathbf{Z})$-graph or a weighted graph G - $\mathbb{H}(k, W)$ the $\mathcal{M}_1(\mathbf{Z})$-graph with k vertices sampled from $W \in \mathcal{W}_1$ - $\mathbb{G}(k, W)$ the $\mathcal{M}_1(\mathbf{Z})$-graph with k vertices sampled from $W \in \mathcal{W}_1$ 	<ul style="list-style-type: none"> - $d_{\square, m}$ the cut distance associated to d_m - $N_{\square, m}$ the cut norm associated to N_m - δ_{\square} the unlabeled distance associated to an arbitrary distance d - $\delta_{\square, m}$ the unlabeled cut distance associated to $d_{\square, m}$ or $N_{\square, m}$ - $\ \cdot\ _{\square, \mathbb{R}}$ the cut norm for real-valued kernels - $\ \cdot\ _{\square, \mathbb{R}}^+$ the positive part of the cut norm for real-valued kernels
<hr/>	
Definitions	
<ul style="list-style-type: none"> - W <i>weak isomorphism</i> of kernels and graphons in Definition 3.16 on page 16 - S <i>tightness</i> for sets of kernels or graphons in Definition 4.7 on page 23 - d <i>invariant</i> and <i>smooth</i> for a distance d on graphon spaces in Definition 3.10 on page 14 - d <i>weakly regular</i> and <i>regular w.r.t. the stepping operator</i> for a distance d on graphon spaces in Definition 4.10 on page 24 	

⌘.....

REFERENCES

- [AAN04] Réka Albert, István Albert, and Gary L. Nakarado. Structural vulnerability of the North American power grid. *Physical Review E*, 69(2):025103, February 2004.
- [AB02] Réka Albert and Albert-László Barabási. Statistical mechanics of complex networks. *Reviews of Modern Physics*, 74(1):47–97, January 2002.
- [AD23] Nathalie Ayi and Nastassia Pouradieu. Graph Limit for Interacting Particle Systems on Weighted Random Graphs, July 2023.
- [ADL13] Hamed Amini, Moez Draief, and Marc Lelarge. Flooding in Weighted Sparse Random Graphs. *SIAM Journal on Discrete Mathematics*, 27(1):1–26, January 2013.
- [AL15] Hamed Amini and Marc Lelarge. The diameter of weighted random graphs. *The Annals of Applied Probability*, 25(3), June 2015.
- [AMGM18] L Alonso, J A Méndez-Bermúdez, A González-Meléndrez, and Yamir Moreno. Weighted random-geometric and random-rectangular graphs: Spectral and eigenfunction properties of the adjacency matrix. *Journal of Complex Networks*, 6(5):753–766, October 2018.
- [AN22] Giacomo Aletti and Giovanni Naldi. Opinion dynamics on graphon: The piecewise constant case. *Applied Mathematics Letters*, 133:108227, November 2022.

- [ANS22] Eleanor Archer, Asaf Nachmias, and Matan Shalev. The GHP scaling limit of uniform spanning trees in high dimensions, April 2022.
- [BC17] Christian Borgs and Jennifer Chayes. Graphons: A Nonparametric Method to Model, Estimate, and Design Algorithms for Massive Networks. In *Proceedings of the 2017 ACM Conference on Economics and Computation*, pages 665–672, Cambridge Massachusetts USA, June 2017. ACM.
- [BCL⁺08] C. Borgs, J.T. Chayes, L. Lovász, V.T. Sós, and K. Vesztegombi. Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing. *Advances in Mathematics*, 219(6):1801–1851, December 2008.
- [BCL⁺12] Christian Borgs, Jennifer Chayes, László Lovász, Vera Sós, and Katalin Vesztegombi. Convergent sequences of dense graphs II. Multiway cuts and statistical physics. *Annals of Mathematics*, 176(1):151–219, July 2012.
- [Bil99] Patrick Billingsley. *Convergence of Probability Measures*. Wiley, New York, second edition edition, 1999.
- [BJR10] Béla Bollobás, Svante Janson, and Oliver Riordan. The Cut Metric, Random Graphs, and Branching Processes. *Journal of Statistical Physics*, 140(2):289–335, July 2010.
- [Bog07a] Vladimir I. Bogachev. *Measure Theory Vol. 1*, volume 1. Springer, Berlin ; New York, 2007.
- [Bog07b] Vladimir I. Bogachev. *Measure Theory Vol. 2*, volume 2. Springer, Berlin ; New York, 2007.
- [Bog18] Vladimir I. Bogachev. *Weak Convergence of Measures*. Number volume 234 in Mathematical Surveys and Monographs. American Mathematical Society, Providence, Rhode Island, 2018.
- [BS02] Stefan Bornholdt and Hans Georg Schuster, editors. *Handbook of Graphs and Networks: From the Genome to the Internet*. Wiley, 1 edition, November 2002.
- [BVDHH10] Shankar Bhamidi, Remco Van Der Hofstad, and Gerard Hooghiemstra. First passage percolation on random graphs with finite mean degrees. *The Annals of Applied Probability*, 20(5), October 2010.
- [DDZ22] Jean-François Delmas, Dylan Dronnier, and Pierre-André Zitt. An infinite-dimensional metapopulation SIS model. *Journal of Differential Equations*, 313:1–53, March 2022.
- [DF64] Lester Dubins and David Freedman. Measurable sets of measures. *Pacific Journal of Mathematics*, 14(4):1211–1222, December 1964.
- [DM10] Moez Draief and Laurent Massoulié. *Epidemics and Rumours in Complex Networks*. Number 369 in London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge ; New York, 2010.
- [EK09] Stewart N Ethier and Thomas G Kurtz. *Markov processes: characterization and convergence*, volume 282. John Wiley & Sons, 2009.
- [Eng89] Ryszard Engelking. *General topology*. Number 6 in Sigma series in pure mathematics. Heldermann, Berlin, rev. and completed ed edition, 1989.
- [FLS06] Michael Freedman, László Lovász, and Alexander Schrijver. Reflection positivity, rank connectivity, and homomorphism of graphs. *Journal of the American Mathematical Society*, 20(1):37–51, April 2006.
- [FOSU16] Victor Falgas-Ravry, Kelly O’Connell, Johanna Strömberg, and Andrew Uzzell. Multicolour containers and the entropy of decorated graph limits, July 2016.
- [Gar09] Diego Garlaschelli. The weighted random graph model. *New Journal of Physics*, 11(7):073005, July 2009.
- [HG13] T. R. Hurd and J. P. Gleeson. On Watts’ cascade model with random link weights. *Journal of Complex Networks*, 1(1):25–43, June 2013.
- [HNT18] Jan Hladký, Asaf Nachmias, and Tuan Tran. The local limit of the uniform spanning tree on dense graphs. *Journal of Statistical Physics*, 173(3-4):502–545, November 2018.
- [HS96] A. Humphreys and Stephen Simpson. Separable Banach space theory needs strong set existence axioms. *Transactions of the American Mathematical Society*, 348(10):4231–4255, 1996.
- [HV23] Jan Hladký and Gopal Viswanathan. Random minimum spanning tree and dense graph limits, October 2023.
- [HWYZ23] Yuanquan Hu, Xiaoli Wei, Junji Yan, and Hengxi Zhang. Graphon mean-field control for cooperative multi-agent reinforcement learning. *Journal of the Franklin Institute*, September 2023.
- [Jan13] Svante Janson. Graphons, cut norm and distance, couplings and rearrangements. *New York Journal of Mathematics*, 4, March 2013.
- [JL15] Varun Jog and Po-Ling Loh. Recovering communities in weighted stochastic block models. In *2015 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 1308–1315, Monticello, IL, September 2015. IEEE.
- [Kal02] Olav Kallenberg. *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2002.

- [Kal17] Olav Kallenberg. *Random Measures, Theory and Applications*, volume 77 of *Probability Theory and Stochastic Modelling*. Springer International Publishing, Cham, 2017.
- [KLS22] Dávid Kunszenti-Kovács, László Lovász, and Balázs Szegedy. Multigraph limits, unbounded kernels, and Banach space decorated graphs. *Journal of Functional Analysis*, 282(2):109284, January 2022.
- [KMS17] István Z. Kiss, Joel C. Miller, and Péter L. Simon. *Mathematics of Epidemics on Networks: From Exact to Approximate Models*, volume 46 of *Interdisciplinary Applied Mathematics*. Springer International Publishing, Cham, 2017.
- [KV23] Nicolas Keriven and Samuel Vaiter. What functions can Graph Neural Networks compute on random graphs? The role of Positional Encoding, May 2023.
- [LL09] Theodore Gyle Lewis and T. G. Lewis. *Network Science: Theory and Practice*. Wiley, Hoboken, NJ, 2009.
- [LMX13] Marc Lelarge, Laurent Massoulié, and Jiaming Xu. Reconstruction in the labeled stochastic block model. In *2013 IEEE Information Theory Workshop (ITW)*, pages 1–5, Sevilla, Spain, September 2013. IEEE.
- [Lov12] László Lovász. *Large networks and graph limits*, volume 60. American Mathematical Society, Colloquium Publications, 2012.
- [LS06] László Lovász and Balázs Szegedy. Limits of dense graph sequences. *Journal of Combinatorial Theory, Series B*, 96(6):933–957, November 2006.
- [LS10] László Lovász and Balázs Szegedy. Limits of compact decorated graphs, October 2010.
- [LS22] Daniel Lackner and Agathe Soret. A Label-State Formulation of Stochastic Graphon Games and Approximate Equilibria on Large Networks. *Mathematics of Operations Research*, page moor.2022.1329, November 2022.
- [New03] M. E. J. Newman. The Structure and Function of Complex Networks. *SIAM Review*, 45(2):167–256, January 2003.
- [Pur66] Robert Purves. Bimeasurable functions. *Fund. Math.*, 58:149–157, 1966.
- [Var58] Veeravalli S. Varadarajan. Weak convergence of measures on separable metric spaces. *Sankhyā: The Indian Journal of Statistics (1933-1960)*, 19(1/2):15–22, 1958.
- [XJL20] Min Xu, Varun Jog, and Po-Ling Loh. Optimal rates for community estimation in the weighted stochastic block model. *The Annals of Statistics*, 48(1), February 2020.
- [XML14] Jiaming Xu, Laurent Massoulié, and Marc Lelarge. Edge label inference in generalized stochastic block models: from spectral theory to impossibility results. In Maria Florina Balcan, Vitaly Feldman, and Csaba Szepesvári, editors, *Proceedings of The 27th Conference on Learning Theory*, volume 35 of *Proceedings of Machine Learning Research*, pages 903–920, Barcelona, Spain, 13–15 Jun 2014. PMLR.
- [YP16] Se-Young Yun and Alexandre Proutiere. Optimal cluster recovery in the labeled stochastic block model. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.

ROMAIN ABRAHAM, INSTITUT DENIS POISSON, UNIVERSITÉ D’ORLÉANS, UNIVERSITÉ DE TOURS, CNRS, FRANCE
Email address: `romain.abraham@univ-orleans.fr`

JEAN-FRANÇOIS DELMAS, CERMICS, ECOLE DES PONTS, FRANCE
Email address: `jean-francois.delmas@enpc.fr`

JULIEN WEIBEL, INSTITUT DENIS POISSON, UNIVERSITÉ D’ORLÉANS, UNIVERSITÉ DE TOURS, CNRS, FRANCE
 AND CERMICS, ECOLE DES PONTS, FRANCE
Email address: `julien.weibel@normalesup.org`