



HAL
open science

THREE-TERM ASYMPTOTIC FORMULA FOR LARGE EIGENVALUES OF THE TWO-PHOTON QUANTUM RABI MODEL

Anne Boutet de Monvel, Lech Zielinski

► **To cite this version:**

Anne Boutet de Monvel, Lech Zielinski. THREE-TERM ASYMPTOTIC FORMULA FOR LARGE EIGENVALUES OF THE TWO-PHOTON QUANTUM RABI MODEL. 2023. hal-04363130

HAL Id: hal-04363130

<https://hal.science/hal-04363130>

Preprint submitted on 23 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

THREE-TERM ASYMPTOTIC FORMULA FOR LARGE EIGENVALUES OF THE TWO-PHOTON QUANTUM RABI MODEL

ANNE BOUTET DE MONVEL¹ AND LECH ZIELINSKI²

ABSTRACT. We prove that the spectrum of the two-photon quantum Rabi Hamiltonian consists of two eigenvalue sequences $(E_m^+)_{m=0}^\infty, (E_m^-)_{m=0}^\infty$ satisfying a three-term asymptotic formula with the remainder estimate $O(m^{-1} \ln m)$ when m tends to infinity. By analogy to the one-photon quantum Rabi model, the leading three terms of this asymptotic formula, describe a generalized rotating-wave approximation for large eigenvalues of the two-photon quantum Rabi model.

Keywords: unbounded self-adjoint operators, discrete spectrum, asymptotic distribution of eigenvalues, quantum Rabi model

Mathematics Subject Classification: 47A75, 81Q10, 47B25, 47B36.

1. GENERAL PRESENTATION OF THE PAPER

1.1. **Introduction.** The simplest interaction between a two-level atom and a classical light field is described by the semi-classical Rabi model [22], [23]. The quantum Rabi model (QRM) couples a two-level system (TLS) with a quantized single-mode radiation and is considered as a particularly important model in quantum electrodynamics: we refer to [8] concerning the historical aspects of the QRM and to [21] for a list of recent research works and experimental realizations of the QRM.

The simplest QRM is defined by the one-photon Hamiltonian $\mathbf{H}_{\text{Rabi}}^{(1)}$ given in Definition 1.2(c). The operator $\mathbf{H}_{\text{Rabi}}^{(1)}$ is a self-adjoint operator depending on two real parameters: g (the coupling constant) and Δ (the energy separation in the TLS). Its spectrum is discrete and the fundamental question is how to find a good approximation of the corresponding eigenvalues.

The first step in this direction, is the *rotating-wave approximation* (RWA) introduced in the famous paper of Jaynes and Cummings [20]. However, the RWA is a correct approximation only when g is close to 0 and Δ close to 1 for $\mathbf{H}_{\text{Rabi}}^{(1)}$ given in Definition 1.2(c). The most popular idea of going beyond the limitations of RWA, bears the name of the *generalized rotating-wave approximation* (GRWA) after Irish [18]. The same idea was considered before by Feranchuk, Komarov, Ulyanenko [13], under the name of the *zeroth order approximation of the operator method* (see [14]). According to the GRWA, the spectrum of $\mathbf{H}_{\text{Rabi}}^{(1)}$ is composed of two eigenvalue sequences: $(E_m^+)_{m=0}^\infty$ and $(E_m^-)_{m=0}^\infty$, satisfying $E_m^\pm \approx E_{m,\text{GRWA}}^\pm$ and the formula (25) in [13] claims that

$$E_{m,\text{GRWA}}^\pm \approx m - g^2 \pm \mathfrak{r}_m \text{ as } m \rightarrow \infty$$

Date: December 23, 2023.

holds with

$$\mathfrak{r}_m := \frac{\Delta \cos(4g\sqrt{m} - \frac{\pi}{4})}{2\sqrt{2\pi g\sqrt{m}}}. \quad (1.1)$$

In [3]-[4] we proved that if \mathfrak{r}_m is given by (1.1) and $\varepsilon > 0$, then one has

$$E_m^\pm = m - g^2 \pm \mathfrak{r}_m + O(m^{-\frac{1}{2}+\varepsilon}) \text{ as } m \rightarrow \infty \quad (1.2)$$

and

$$E_{m,\text{GRWA}}^\pm = m - g^2 \pm \mathfrak{r}_m + O(m^{-\frac{1}{2}}) \text{ as } m \rightarrow \infty \quad (1.3)$$

Thus (1.2)-(1.3) allow one to estimate the quality of GRWA for large eigenvalues of the one-photon QRM. In this paper we will prove a similar result for the two-photon QRM defined by the Hamiltonian $\mathbf{H}_{\text{Rabi}}^{(2)}$ given in Definition 1.2(d).

We remark that the two-photon QRM was used in [15] to describe a two-level atom interacting with squeezed light and we refer to [9] for the overview of works concerning the two-photon quantum Rabi model (see also [21]). In what follows, we always assume that the coupling constant satisfies the condition $0 < g < 1/2$. This hypothesis implies that $\mathbf{H}_{\text{Rabi}}^{(2)}$ is self adjoint and has a discrete spectrum. In [6] we proved that the spectrum of $\mathbf{H}_{\text{Rabi}}^{(2)}$ is composed of two eigenvalue sequences: $(E_m^+)_{m=0}^\infty$ and $(E_m^-)_{m=0}^\infty$, satisfying

$$E_m^\pm = \left(m + \frac{1}{2}\right)\sqrt{1-4g^2} - \frac{1}{2} + O(m^{-1/3}) \text{ as } m \rightarrow \infty \quad (1.4)$$

In this paper we will prove the three-term asymptotic formula

$$E_m^\pm = \left(m + \frac{1}{2}\right)\sqrt{1-4g^2} - \frac{1}{2} \pm \mathfrak{r}_m + O(m^{-1} \ln m) \text{ as } m \rightarrow \infty \quad (1.5)$$

where \mathfrak{r}_m is given by (1.20). By analogy to (1.2)-(1.3), the expression

$$\left(m + \frac{1}{2}\right)\sqrt{1-4g^2} - \frac{1}{2} \pm \mathfrak{r}_m$$

can be viewed as a GRWA for large eigenvalues of the two-photon QRM.

1.2. Quantum Rabi Model.

Notation 1.1. (a) In what follows, \mathbb{Z} is the set of integers and $\mathbb{N} := \{n \in \mathbb{Z} : n \geq 0\}$.

(b) We denote by $\ell^2(\mathbb{N})$ the complex Hilbert space of square-summable sequences $x : \mathbb{N} \rightarrow \mathbb{C}$ equipped with the scalar product

$$\langle x, y \rangle_{\ell^2(\mathbb{N})} = \sum_{m=0}^{\infty} \overline{x(m)}y(m) \quad (1.6)$$

and the norm $\|x\|_{\ell^2(\mathbb{N})} := \langle x, x \rangle_{\ell^2(\mathbb{N})}^{1/2}$. For $s > 0$ we denote

$$\ell^{2,s}(\mathbb{N}) := \{x \in \ell^2(\mathbb{N}) : \sum_{m=0}^{\infty} (1+m^2)^s |x(m)|^2 < \infty\} \quad (1.7)$$

(c) The canonical basis of $\ell^2(\mathbb{N})$ is denoted $\{e_n\}_{n \in \mathbb{N}}$ (i.e. $e_n(m) = \delta_{n,m}$ for $n, m \in \mathbb{N}$).

(d) The annihilation and creation operators, \hat{a} and \hat{a}^\dagger , are the linear maps $\ell^{2,1/2}(\mathbb{N}) \rightarrow \ell^2(\mathbb{N})$ satisfying

$$\hat{a}^\dagger e_n = \sqrt{n+1} e_{n+1} \text{ for } n \in \mathbb{N} \quad (1.8)$$

$$\hat{a} e_0 = 0 \text{ and } \hat{a} e_n = \sqrt{n} e_{n-1} \text{ for } n \in \mathbb{N} \setminus \{0\}. \quad (1.9)$$

(e) Using $(1, 0) \in \mathbb{C}^2$ and $(0, 1) \in \mathbb{C}^2$ as the canonical basis of the Euclidean space \mathbb{C}^2 , we denote by σ_x, σ_z, I_2 , the linear operators in \mathbb{C}^2 defined by the matrices

$$\sigma_x := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad I_2 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (1.10)$$

Definition 1.2. (a) The two-level system (TLS) Hamiltonian is the linear map in \mathbb{C}^2 defined by the matrix

$$H_{\text{TLS}} = \frac{1}{2} \begin{pmatrix} \Delta & 0 \\ 0 & -\Delta \end{pmatrix} = \frac{1}{2} \Delta \sigma_z \quad (1.11)$$

where Δ is a real parameter.

(b) The Hamiltonian of the single-mode radiation is the linear map $H_{\text{rad}} : \ell^{2,1}(\mathbb{N}) \rightarrow \ell^2(\mathbb{N})$ defined by the formula

$$H_{\text{rad}} e_n = \hat{a}^\dagger \hat{a} e_n = n e_n \text{ for } n \in \mathbb{N}. \quad (1.12)$$

(c) The *one-photon quantum Rabi Hamiltonian* is defined as the linear map $\mathbf{H}_{\text{Rabi}}^{(1)} : \mathbb{C}^2 \otimes \ell^{2,1}(\mathbb{N}) \rightarrow \mathbb{C}^2 \otimes \ell^2(\mathbb{N})$ given by

$$\mathbf{H}_{\text{Rabi}}^{(1)} = I_2 \otimes H_{\text{rad}} + H_{\text{TLS}} \otimes I_{\ell^2(\mathbb{N})} + g \sigma_x \otimes (\hat{a} + \hat{a}^\dagger), \quad (1.13)$$

where $g > 0$ is the coupling constant.

(d) The *two-photon quantum Rabi Hamiltonian* is defined as the linear map $\mathbf{H}_{\text{Rabi}}^{(2)} : \mathbb{C}^2 \otimes \ell^{2,1}(\mathbb{N}) \rightarrow \mathbb{C}^2 \otimes \ell^2(\mathbb{N})$ given by

$$\mathbf{H}_{\text{Rabi}}^{(2)} = I_2 \otimes H_{\text{rad}} + H_{\text{TLS}} \otimes I_{\ell^2(\mathbb{N})} + g \sigma_x \otimes (\hat{a}^2 + (\hat{a}^\dagger)^2), \quad (1.14)$$

where $g > 0$ is the coupling constant.

1.3. Main result. In what follows, we assume $0 < g < 1/2$ and introduce

$$\beta := \sqrt{1 - 4g^2}, \quad (1.15)$$

$$\alpha := \arctan \left(\sqrt{\frac{1-2g}{1+2g}} \right). \quad (1.16)$$

Let $\mathbf{H}_{0,\text{Rabi}}^{(2)}$ denote the operator given by (1.14) with $\Delta = 0$, i.e.

$$\mathbf{H}_{0,\text{Rabi}}^{(2)} = I_2 \otimes H_{\text{rad}} + g \sigma_x \otimes (\hat{a}^2 + (\hat{a}^\dagger)^2). \quad (1.17)$$

If $0 < g < 1/2$, then the spectrum of $\mathbf{H}_{0,\text{Rabi}}^{(2)}$ is explicitly known (see [11], [12]): it is composed of the sequence of eigenvalues

$$E_m^0 = m\beta + (\beta - 1)/2, \quad m = 0, 1, 2, \dots \quad (1.18)$$

and each eigenvalue E_m^0 is of multiplicity 2. Thus $0 < g < 1/2$ ensures the fact that $\mathbf{H}_{0,\text{Rabi}}^{(2)}$ is self-adjoint and has compact resolvent. Since $\mathbf{H}_{\text{Rabi}}^{(2)} - \mathbf{H}_{0,\text{Rabi}}^{(2)}$ is bounded, the operator $\mathbf{H}_{\text{Rabi}}^{(2)}$ is self-adjoint and has compact resolvent if $0 < g < 1/2$. The explicit values of eigenvalues of $\mathbf{H}_{\text{Rabi}}^{(2)}$ are not known when $\Delta \neq 0$, but we can describe their asymptotic behaviour. Our main result is the following

Theorem 1.3. *Assume that $0 < g < 1/2$. Then one can find $\{v_m^+\}_{m \in \mathbb{N}} \cup \{v_m^-\}_{m \in \mathbb{N}}$, an orthonormal basis of $\mathbb{C}^2 \otimes \ell^2(\mathbb{N})$, such that*

$$\mathbf{H}_{\text{Rabi}}^{(2)} v_m^\pm = E_m^\pm v_m^\pm, \quad m = 0, 1, 2, \dots$$

and the eigenvalue sequences $(E_m^+)_{m \in \mathbb{N}}$, $(E_m^-)_{m \in \mathbb{N}}$, satisfy the large m estimates

$$E_m^\pm = m\beta + (\beta - 1)/2 \pm \mathfrak{r}_m + O(m^{-1} \ln m) \quad (1.19)$$

with \mathfrak{r}_m given by the formula

$$\mathfrak{r}_m = \begin{cases} \frac{\Delta}{2} \sqrt{\frac{\beta}{\pi g m}} \cos((2m+1)\alpha) & \text{if } m \text{ is even} \\ \frac{\Delta}{2} \sqrt{\frac{\beta}{\pi g m}} \sin((2m+1)\alpha) & \text{if } m \text{ is odd,} \end{cases} \quad (1.20)$$

where β is given by (1.15) and α is given by (1.16).

Remarks. (a) One has $E_m^\pm - E_m^0 = O(m^{-1/2})$ in spite of the fact that $\mathbf{H}_{\text{Rabi}}^{(2)} - \mathbf{H}_{0, \text{Rabi}}^{(2)}$ is not compact. A similar fact was established in [30] for the one-photon Rabi model.

(b) Following [25], one can prove that the spectrum of $\mathbf{H}_{\text{Rabi}}^{(2)}$ is not discrete if $g \geq \frac{1}{2}$.

1.4. Overview of earlier results. The distribution of large eigenvalues of the QRM has been a subject of many numerical investigations. Concerning the earliest physical considerations, we refer to Schmutz [26]. It is well known (see e.g [28]) that the one-photon quantum Rabi Hamiltonian can be expressed as a direct sum of two Jacobi operators, i.e. operators defined by infinite tridiagonal matrices. Similarly, the two-photon quantum Rabi Hamiltonian can be expressed as a direct sum of four Jacobi operators (see [6]). A mathematical study of large eigenvalues of Jacobi matrices was initiated by J. Janas and S. Naboko in the paper [19], which contains fundamental ideas of the method of approximate diagonalizations.

The question of the behaviour of large eigenvalues of $\mathbf{H}_{\text{Rabi}}^{(1)}$, was first posed by E. A. Tur [28]-[29] and it was considered by A. Boutet de Monvel, S. Naboko and L. O. Silva in [1]-[2]. Due to the difficulty of the problem, the authors give a solution for a simpler class of operators ("modified Jaynes-Cummings models"). The two-term asymptotic formula for large eigenvalues of the one-photon quantum Rabi model, was proved by E. A. (Tur) Yanovich in [30] (see also [29]). The three-term asymptotic formula (1.2) was proved in [3]-[4]. We remark that the formula (1.2) allows one to recover the values of parameters g and Δ from the spectrum (see [5]). The two-term asymptotic formula (1.4) for large eigenvalues of the two-photon quantum Rabi model, was proved in [6]. The result of Theorem 1.3 and the main ideas of its proof were described in [7]. The same result was also described in [17].

1.5. Organization of the paper. Our approach is based on an analysis of operators acting in $\ell^2(\mathbb{Z})$. These operators are introduced in Section 2.1 and the corresponding three-term asymptotic formula is given in Theorem 2.3. In Theorem 2.3(a) we consider the operators \tilde{J}_γ^0 , which are special case of operators considered by Edward [10]. In Section 2.2 we describe explicit expressions of their eigenvalues and eigenvectors by means of the discrete Fourier transform. In Theorem 2.3(b) we consider the operators

\tilde{J}_γ^δ , which are viewed as perturbations of \tilde{J}_γ^0 . In Section 3 we show that the assertion of Theorem 2.3(b) follows from a GRWA result given in Proposition 3.1. The proof of Proposition 3.1 begins in Section 4 and is completed in Section 5. In Section 6 we show how to deduce Theorem 1.3 from Theorem 2.3(b). More precisely, we follow the following scheme:

- we prove Proposition 3.1, using Lemma 4.3 and Kato-Temple estimate (Theorem 5.2)
- we prove Theorem 2.3, using Lemma 2.6, Proposition 3.1 and an asymptotic formula from [6]
- we prove Theorem 1.3, using Theorem 2.3, Lemma 6.2 and Lemma 6.3.

2. AN AUXILIARY PROBLEM IN $\ell^2(\mathbb{Z})$

2.1. Behaviour of large eigenvalues for auxiliary operators \tilde{J}_γ^δ .

Notation 2.1. (a) We denote by $\mathcal{B}(\mathcal{V})$ the algebra of bounded operators on the Banach space \mathcal{V} and $\|\cdot\|_{\mathcal{B}(\mathcal{V})}$ is the corresponding operator norm.

(b) We denote by $\ell^2(\mathbb{Z})$ the complex Hilbert space of square-summable sequences $x : \mathbb{Z} \rightarrow \mathbb{C}$ equipped with the scalar product

$$\langle x, y \rangle_{\ell^2(\mathbb{Z})} = \sum_{k \in \mathbb{Z}} \overline{x(k)} y(k) \quad (2.1)$$

and the norm $\|x\|_{\ell^2(\mathbb{Z})} := \langle x, x \rangle_{\ell^2(\mathbb{Z})}^{1/2}$. For $s > 0$ we denote

$$\ell^{2,s}(\mathbb{Z}) := \{x \in \ell^2(\mathbb{Z}) : \|x\|_{\ell^{2,s}(\mathbb{Z})} < \infty\}$$

where

$$\|x\|_{\ell^{2,s}(\mathbb{Z})} := \left(\sum_{k \in \mathbb{Z}} (1 + k^2)^s |x(k)|^2 \right)^{1/2} \quad (2.2)$$

(c) The canonical basis of $\ell^2(\mathbb{Z})$ is denoted $\{\tilde{e}_j\}_{j \in \mathbb{Z}}$ (i.e. $\tilde{e}_j(k) = \delta_{j,k}$ for $j, k \in \mathbb{Z}$) and $\ell_{\text{fin}}^2(\mathbb{Z})$ denotes the set of finite linear combinations of vectors belonging to $\{\tilde{e}_j\}_{j \in \mathbb{Z}}$.

(d) If $(\tilde{d}_j)_{j \in \mathbb{Z}}$ is real valued, then $\tilde{D} := \text{diag}(\tilde{d}_j)_{j \in \mathbb{Z}}$ is the self-adjoint operator in $\ell^2(\mathbb{Z})$ satisfying

$$\tilde{D} \tilde{e}_j = \tilde{d}_j \tilde{e}_j \text{ for every } j \in \mathbb{Z}.$$

We denote

$$\Lambda := \text{diag}(j)_{j \in \mathbb{Z}}. \quad (2.3)$$

(e) We denote by S the shift defined in $\ell^2(\mathbb{Z})$ by the formula $(Sx)(j) = x(j-1)$.

Definition 2.2. In what follows, γ , δ and g are fixed real numbers.

(a) We define \tilde{J}_γ^0 as the linear map $\ell^{2,1}(\mathbb{Z}) \rightarrow \ell^2(\mathbb{Z})$ given by

$$\tilde{J}_\gamma^0 := \Lambda + g(S(\Lambda + \gamma) + (\Lambda + \gamma)S^{-1}) = \Lambda + g(S(\Lambda + \gamma) + \text{h.c.}) \quad (2.4)$$

(b) We define \tilde{J}_γ^δ as the linear map $\ell^{2,1}(\mathbb{Z}) \rightarrow \ell^2(\mathbb{Z})$ given by the formula

$$\tilde{J}_\gamma^\delta := \tilde{J}_\gamma^0 + D_\delta, \quad (2.5)$$

where

$$D_\delta := \text{diag}(\delta(-1)^j)_{j \in \mathbb{Z}}. \quad (2.6)$$

Using the above definition, we find that the action of \tilde{J}_γ^δ can be represented by the following tridiagonal, $\mathbb{Z} \times \mathbb{Z}$ matrix

$$\begin{pmatrix} \ddots & & & & & & & & & & \\ & -2 + \delta & g(-2 + \gamma) & 0 & 0 & 0 & & & & & \\ & g(-2 + \gamma) & -1 - \delta & g(-1 + \gamma) & 0 & 0 & & & & & \\ & 0 & g(-1 + \gamma) & \delta & g\gamma & 0 & & & & & \\ & 0 & 0 & g\gamma & 1 - \delta & g(1 + \gamma) & & & & & \\ & 0 & 0 & 0 & g(1 + \gamma) & 2 + \delta & & & & & \\ & & & & & & \ddots & & & & \end{pmatrix}$$

whose diagonal entries $\{\tilde{d}_\delta(j)\}_{j \in \mathbb{Z}}$ are given by

$$\tilde{d}_\delta(j) := j + \delta(-1)^j, \quad (2.7)$$

off-diagonal entries $\{\tilde{b}_\gamma(j)\}_{j \in \mathbb{Z}}$ are given by

$$\tilde{b}_\gamma(j) := g(j + \gamma) \quad (2.8)$$

and $\tilde{J}_\gamma^\delta \tilde{e}_j = \tilde{d}_\delta(j) \tilde{e}_j + \tilde{b}_\gamma(j) \tilde{e}_{j+1} + \tilde{b}_\gamma(j-1) \tilde{e}_{j-1}$ for every $j \in \mathbb{Z}$.

Theorem 2.3. *Let \tilde{J}_γ^0 be given by (2.4) and \tilde{J}_γ^δ by (2.5)-(2.6). If $0 < g < 1/2$ then*

(a) *the spectrum of \tilde{J}_γ^0 is composed of a non-decreasing sequence of eigenvalues $\{d_{\gamma,j}\}_{j \in \mathbb{Z}}$ of the form*

$$d_{\gamma,j} := \beta j + \left(\gamma - \frac{1}{2}\right)(\beta - 1), \quad (2.9)$$

where β is given by (1.15).

(b) *the spectrum of \tilde{J}_γ^δ is composed of a non-decreasing sequence of eigenvalues $\{\lambda_j(\tilde{J}_\gamma^\delta)\}_{j \in \mathbb{Z}}$ which can be labeled so that*

$$\lambda_j(\tilde{J}_\gamma^\delta) = d_{\gamma,j} + \mathfrak{r}_\gamma^\delta(j) + O(j^{-1} \ln j) \text{ as } j \rightarrow \infty \quad (2.10)$$

where $d_{\gamma,j}$ is given by (2.9),

$$\mathfrak{r}_\gamma^\delta(j) = \delta \left(\frac{\beta}{2\pi g j}\right)^{1/2} \cos(4\alpha j + \hat{\theta}_\gamma), \quad (2.11)$$

β is given by (1.15), α by (1.16) and

$$\hat{\theta}_\gamma = \left(\gamma - \frac{1}{2}\right)(4\alpha - \pi) + \frac{\pi}{4}. \quad (2.12)$$

Proof. (a) This result was proved in [10]. We describe a simplified proof in Section 2.2.

(b) See Section 3.2. \square

2.2. Diagonalization of \tilde{J}_γ^0 . In what follows, $\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$ is identified with $]-\pi, \pi]$ and $L^2(\mathbb{T})$ denotes the Hilbert space of Lebesgue square integrable functions $]-\pi, \pi] \rightarrow \mathbb{C}$ equipped with the scalar product

$$\langle f, g \rangle_{L^2(\mathbb{T})} := \int_{-\pi}^{\pi} \overline{f(\theta)} g(\theta) \frac{d\theta}{2\pi} \quad (2.13)$$

and the norm $\|f\|_{L^2(\mathbb{T})} = \langle f, f \rangle_{L^2(\mathbb{T})}^{1/2}$. We let $\mathcal{F}_\mathbb{T}$ denote the isometric isomorphism $L^2(\mathbb{T}) \rightarrow \ell^2(\mathbb{Z})$ given by

$$(\mathcal{F}_\mathbb{T} f)(j) = \int_{-\pi}^{\pi} f(\theta) e^{-ij\theta} \frac{d\theta}{2\pi} \quad (2.14)$$

and consider the operator

$$L_\gamma^0 := \mathcal{F}_\mathbb{T}^{-1} \tilde{J}_\gamma^0 \mathcal{F}_\mathbb{T} = -i \frac{d}{d\theta} + g \left(e^{i\theta} \left(-i \frac{d}{d\theta} + \gamma \right) + \text{h.c.} \right)$$

Similarly as in [10] and [6], we observe that the assumption $0 < g < 1/2$ ensures the fact that L_γ^0 is the first order linear elliptic differential operator,

$$L_\gamma^0 = \frac{1}{2} \left((1 + 2g \cos(\theta)) \left(-i \frac{d}{d\theta} \right) + \text{h.c.} \right) + (2\gamma - 1)g \cos(\theta)$$

and we introduce

$$\Phi(\theta) := \int_0^\theta \frac{\beta d\theta'}{1 + 2g \cos(\theta')}. \quad (2.15)$$

An easy calculation allows one to express the right-hand side of (2.15),

$$\Phi(\theta) = 2 \arctan \left(\sqrt{\frac{1-2g}{1+2g}} \tan \left(\frac{\theta}{2} \right) \right) \text{ if } -\pi < \theta < \pi. \quad (2.16)$$

Moreover $\Phi(\pi) = \pi$ and Φ is odd, i.e. $\Phi(-\theta) = -\Phi(\theta)$. Using the change of variable $\eta = \Phi(\theta)$, we can define the unitary operator acting in $L^2(\mathbb{T})$ according to the formula

$$(U_\Phi f)(\theta) = \Phi'(\theta)^{1/2} f(\Phi(\theta)) \quad (2.17)$$

and the direct computation (see [6]) gives

$$U_\Phi^{-1} L_\gamma^0 U_\Phi = \beta \left(-i \frac{d}{d\eta} + q_\gamma(\eta) \right) \quad (2.18)$$

with

$$q_\gamma(\eta) := \beta^{-1} (2\gamma - 1)g \cos(\Phi^{-1}(\eta)). \quad (2.19)$$

In the following, q_γ is given by (2.19) and

$$\tilde{q}_\gamma(\eta) := \int_0^\eta q_\gamma(\eta') d\eta'. \quad (2.20)$$

We claim that

$$\tilde{q}_\gamma(\Phi(\theta)) = \left(\gamma - \frac{1}{2} \right) (\theta - \beta^{-1} \Phi(\theta)). \quad (2.21)$$

Indeed,

$$\begin{aligned} \frac{d}{d\theta} \left(\tilde{q}_\gamma(\Phi(\theta)) \right) &= q_\gamma(\Phi(\theta)) \Phi'(\theta) = \frac{(2\gamma - 1)g \cos(\theta)}{\beta} \frac{\beta}{1 + 2g \cos(\theta)} = \\ &= \frac{2\gamma - 1}{2} \left(1 - \frac{1}{1 + 2g \cos(\theta)} \right) = \left(\gamma - \frac{1}{2} \right) (1 - \beta^{-1} \Phi'(\theta)) \end{aligned}$$

implies $\tilde{q}_\gamma(\Phi(\theta)) = (\gamma - \frac{1}{2})(\theta - \beta^{-1} \Phi(\theta)) + C_0$ and $C_0 = 0$ holds due to $\Phi(0) = 0$ and $\tilde{q}_\gamma(\Phi(0)) = \tilde{q}_\gamma(0) = 0$.

Using $\Phi(\pm\pi) = \pm\pi$ in (2.21), we compute

$$\langle q_\gamma \rangle := \frac{\tilde{q}_\gamma(\pi) - \tilde{q}_\gamma(-\pi)}{2\pi} = \left(\gamma - \frac{1}{2} \right) (1 - \beta^{-1}) \quad (2.22)$$

and remark that $\eta \rightarrow \langle q_\gamma \rangle \eta - \tilde{q}_\gamma(\eta)$ is a smooth 2π -periodic function. Then we define $(f_{q_\gamma, j})_{j \in \mathbb{Z}}$ to be the orthonormal basis in $L^2(\mathbb{T})$ given by

$$f_{q_\gamma, j}(\eta) = e^{ij\eta} e^{i(\langle q_\gamma \rangle \eta - \tilde{q}_\gamma(\eta))} \quad (2.23)$$

and (see [7], [10]), for every $j \in \mathbb{Z}$, we get

$$\beta \left(-i \frac{d}{d\eta} + q_\gamma \right) f_{q_\gamma, j} = d_{\gamma, j} f_{q_\gamma, j}, \quad (2.24)$$

where $d_{\gamma, j} = \beta(j + \langle q_\gamma \rangle) = \beta j + (\gamma - 1/2)(\beta - 1)$ is given (2.9). Combining (2.24) with (2.18), we obtain

Corollary 2.4. *Let $\{f_{\gamma, j}^0\}_{j \in \mathbb{Z}}$ be the orthonormal basis of $L^2(\mathbb{T})$ given by*

$$f_{\gamma, j}^0(\theta) := (U_\Phi f_{q_\gamma, j})(\theta) = \Phi'(\theta)^{1/2} e^{ij\Phi(\theta)} e^{i(\langle q_\gamma \rangle \Phi(\theta) - \bar{q}_\gamma(\Phi(\theta)))}. \quad (2.25)$$

Then

$$L_\gamma^0 f_{\gamma, j}^0 = d_{\gamma, j} f_{\gamma, j}^0 \quad (2.26)$$

holds with $d_{\gamma, j}$ given (2.9).

It is clear that the assertion of Theorem 2.3(a) follows from Corollary 2.4. Indeed, using (2.26) and $\tilde{J}_\gamma^0 = \mathcal{F}_\mathbb{T} L_\gamma^0 \mathcal{F}_\mathbb{T}^{-1}$, we get

$$\tilde{J}_\gamma^0 u_{\gamma, j}^0 = d_{\gamma, j} u_{\gamma, j}^0 \text{ with } u_{\gamma, j}^0 := \mathcal{F}_\mathbb{T} f_{\gamma, j}^0. \quad (2.27)$$

2.3. An auxiliary estimate.

Notation 2.5. (a) For $f \in L^2(\mathbb{T})$ we write $T_\pi f := f \circ \tau_\pi$ with $\tau_\pi : \mathbb{T} \rightarrow \mathbb{T}$ given by

$$\tau_\pi(\theta + 2\pi\mathbb{Z}) := \theta + \pi + 2\pi\mathbb{Z} \quad (2.28)$$

(b) For $j, n \in \mathbb{Z}$, we denote

$$V_\gamma^\delta(j, n) := \delta \langle f_{\gamma, j}^0, T_\pi f_{\gamma, n}^0 \rangle_{L^2(\mathbb{T})}, \quad (2.29)$$

where $f_{\gamma, j}^0$ is given by (2.25).

Lemma 2.6. *Let $\hat{c} > 0$ be small enough. Then there exists $\hat{C} > 0$ such that*

$$\sup_{\{k \in \mathbb{Z}: |k| \leq \hat{c}|n|\}} |V_\gamma^\delta(n+k, n)| \leq \hat{C}|n|^{-1/2} \quad (2.30)$$

holds for every $n \in \mathbb{Z} \setminus \{0\}$.

Proof. Let us assume that $n \neq 0$. By definition (2.25), $f_{\gamma, n}^0 = e^{in\Phi} p_\gamma$ holds with

$$p_\gamma(\theta) := e^{i\langle q_\gamma \rangle \Phi(\theta) - i\bar{q}_\gamma(\Phi(\theta))} \beta^{1/2} (1 + 2g \cos \theta)^{-1/2} \quad (2.31)$$

and we can express

$$V_\gamma^\delta(n+k, n) = \delta \int_{-\pi}^{\pi} e^{in\Psi_{n, k}(\theta)} p_\gamma(\theta) \overline{(T_\pi p_\gamma)(\theta)} \frac{d\theta}{2\pi} \quad (2.32)$$

with

$$\Psi_{n, k} = \Phi - T_\pi \Phi + \frac{k}{n} \Phi. \quad (2.33)$$

Denote $\Psi := \Phi - T_\pi \Phi$. Then

$$\Psi'(\theta) = \frac{\beta}{1 + 2g \cos \theta} - \frac{\beta}{1 - 2g \cos \theta} = 0 \Leftrightarrow \cos \theta = 0, \quad (2.34)$$

i.e. Ψ has two critical points: $-\frac{\pi}{2}$ and $\frac{\pi}{2}$. Since

$$\Psi''(\theta) = \left(\frac{2g\beta}{(1 + 2g \cos \theta)^2} + \frac{2g\beta}{(1 - 2g \cos \theta)^2} \right) \sin \theta, \quad (2.35)$$

we obtain $\Psi''(\pm\frac{\pi}{2}) = \pm 4g\beta \neq 0$ and choosing $c_0 > 0$ small enough we can ensure

$$|\Psi''(\theta)| \geq c_0 \text{ for } \theta \in \mathcal{I}_{c_0} := [-\frac{\pi}{2} - c_0, -\frac{\pi}{2} + c_0] \cup [\frac{\pi}{2} - c_0, \frac{\pi}{2} + c_0].$$

However $\Psi_{n,k} = \Psi + \frac{k}{n}\Phi$ and denoting $C_0 = \sup |\Phi''|$ we can estimate

$$|\Psi''_{n,k}(\theta)| \geq |\Psi''(\theta)| - C_0 \left| \frac{k}{n} \right|$$

and

$$\left| \frac{k}{n} \right| \leq \frac{c_0}{2C_0} \implies |\Psi''_{n,k}(\theta)| \geq \frac{c_0}{2} \text{ for } \theta \in \mathcal{I}_{c_0}. \quad (2.36)$$

Due to van der Corput Lemma (see Lemma 7.1(b)), there exists $C > 0$ such that

$$\left| \int_{\mathcal{I}_{c_0}} e^{in\Phi_{n,k}(\theta)} p_\gamma(\theta) \overline{(T_\pi p_\gamma)(\theta)} \frac{d\theta}{2\pi} \right| \leq C|n|^{-1/2}. \quad (2.37)$$

Moreover

$$c_1 := \inf\{|\Psi'(\theta)| : \theta \in [-\pi, \pi] \setminus \mathcal{I}_{c_0}\} > 0$$

and denoting $C_1 = \sup |\Phi'|$ we can estimate

$$|\Psi'_{n,k}(\theta)| \geq |\Psi'(\theta)| - C_1 \left| \frac{k}{n} \right|,$$

hence

$$\left| \frac{k}{n} \right| \leq \frac{c_1}{2C_1} \implies |\Psi'_{n,k}(\theta)| \geq \frac{c_1}{2} \text{ for } \theta \in [-\pi, \pi] \setminus \mathcal{I}_{c_0}. \quad (2.38)$$

Due to the non-stationary phase estimate (see Lemma 7.1(a)), there exists a constant $C' > 0$ such that

$$\left| \int_{[-\pi, \pi] \setminus \mathcal{I}_{c_0}} e^{in\Phi_{n,k}(\theta)} p_\gamma(\theta) \overline{(T_\pi p_\gamma)(\theta)} \frac{d\theta}{2\pi} \right| \leq C'|n|^{-1}. \quad (2.39)$$

Combining (2.37), (2.39) and (2.32), we complete the proof. \square

3. THE GRWA APPROACH

3.1. Statement of the result. The idea of GRWA consists in using the diagonal entries of a perturbation as the first correction for eigenvalues of a perturbed diagonal matrix (see [18] and [14]). Our proof of Theorem 2.3 is based on the following variant of the GRWA :

Proposition 3.1. *Let us fix $\beta_0 \in \mathbb{R}$, $\beta > 0$, $\hat{C} > 0$ and $\hat{c} > 0$. Let $J : \ell^{2,1}(\mathbb{Z}) \rightarrow \ell^2(\mathbb{Z})$ be given by the formula*

$$J = \beta\Lambda + \beta_0 + V, \quad (3.1)$$

where $\Lambda = \text{diag}(j)_{j \in \mathbb{Z}}$ and V is a bounded self-adjoint operator in $\ell^2(\mathbb{Z})$ satisfying

$$\sup_{\{k \in \mathbb{Z} : |k| \leq \hat{c}|n|\}} |\langle \tilde{e}_{n+k}, V\tilde{e}_k \rangle_{\ell^2(\mathbb{Z})}| \leq \hat{C}|n|^{-1/2} \quad (3.2)$$

for every $n \in \mathbb{Z} \setminus \{0\}$. Assume that the spectrum of J is composed of a non-decreasing sequence of eigenvalues $\{\lambda_j(J)\}_{j \in \mathbb{Z}}$ such that

$$\lambda_j(J) = j\beta + \beta_0 + o(1) \text{ as } j \rightarrow \infty. \quad (3.3)$$

Then one has

$$\lambda_j(J) = j\beta + \beta_0 + \langle \tilde{e}_j, V\tilde{e}_j \rangle_{\ell^2(\mathbb{Z})} + O(j^{-1} \ln j) \text{ as } j \rightarrow \infty \quad (3.4)$$

Proof. See Section 5. \square

3.2. Proof of Theorem 2.3(b). In what follows, we describe how to deduce the assertion of Theorem 2.3(b) from Proposition 3.1. We begin by introducing \tilde{U}_γ , the unitary operator in $\ell^2(\mathbb{Z})$ satisfying $\tilde{U}_\gamma \tilde{e}_j = u_{\gamma,j}^0$, where $\{\tilde{e}_j\}_{j \in \mathbb{Z}}$ is the canonical basis of $\ell^2(\mathbb{Z})$ and $\{u_{\gamma,j}^0\}_{j \in \mathbb{Z}}$ is the basis introduced in (2.27). We claim that the hypotheses of Proposition 3.1 are satisfied if

$$J = \tilde{U}_\gamma^{-1} \tilde{J}_\gamma^\delta \tilde{U}_\gamma, \quad (3.5)$$

where J_γ^δ is as in Definition 2.2. Indeed, if β is given by (1.15) and

$$\beta_0 = \left(\gamma - \frac{1}{2}\right)(\beta - 1), \quad (3.6)$$

then $\tilde{U}_\gamma^{-1} \tilde{J}_\gamma^\delta \tilde{U}_\gamma = \beta \Lambda + \beta_0 + V_\gamma^\delta$ holds with

$$V_\gamma^\delta := \tilde{U}_\gamma^{-1} D_\delta \tilde{U}_\gamma, \quad (3.7)$$

where $D_\delta = \text{diag}(\delta(-1)^j)_{j \in \mathbb{Z}}$ (see (2.6)). Therefore

$$\langle \tilde{e}_j, V_\gamma^\delta \tilde{e}_k \rangle_{\ell^2(\mathbb{Z})} = \langle u_{j,\gamma}^0, D_\delta u_{j,\gamma}^0 \rangle_{\ell^2(\mathbb{Z})} = \langle f_{j,\gamma}^0, \mathcal{F}_\mathbb{T}^{-1} D_\delta \mathcal{F}_\mathbb{T} f_{j,\gamma}^0 \rangle_{L^2(\mathbb{T})}, \quad (3.8)$$

where we used $u_{j,\gamma}^0 = \mathcal{F}_\mathbb{T} f_{j,\gamma}^0$ and the isometry $\mathcal{F}_\mathbb{T}$. Since $\mathcal{F}_\mathbb{T}^{-1} D_\delta \mathcal{F}_\mathbb{T} = \delta T_\pi$ holds with T_π as in Notation 2.5, (3.8) gives

$$\langle \tilde{e}_j, V_\gamma^\delta \tilde{e}_k \rangle_{\ell^2(\mathbb{Z})} = V_\gamma^\delta(j, k) \quad (3.9)$$

with $V_\gamma^\delta(\cdot, \cdot)$ expressed by (2.29) and due to Lemma 2.6, the estimate (3.2) holds if $V = V_\gamma^\delta$ is given by (3.7). Moreover, the result of [6, Proposition 5.1] ensures the fact that the spectrum of \tilde{J}_γ^δ is composed of the non-decreasing sequence of eigenvalues $\{\lambda_j(\tilde{J}_\gamma^\delta)\}_{j \in \mathbb{Z}}$, which can be labeled so that one has the estimate

$$\lambda_j(\tilde{J}_\gamma^\delta) = j\beta + \beta_0 + O(j^{-1/3}) \text{ as } j \rightarrow \infty, \quad (3.10)$$

i.e. (3.3) holds if J is given by (3.5). Thus all the hypotheses of Proposition 3.1 hold and choosing $\{\lambda_j(\tilde{J}_\gamma^\delta)\}_{j \in \mathbb{Z}}$ to be the sequence of eigenvalues of \tilde{J}_γ^δ such that $\lambda_j(\tilde{J}_\gamma^\delta) = \lambda_j(J)$ for every $j \in \mathbb{Z}$, we obtain

$$\lambda_j(\tilde{J}_\gamma^\delta) = j\beta + \beta_0 + \langle \tilde{e}_j, V_\gamma^\delta \tilde{e}_j \rangle_{\ell^2(\mathbb{Z})} + O(j^{-1} \ln j) \text{ when } j \rightarrow \infty \quad (3.11)$$

due to (3.4). The assertion of Theorem 2.3(b) follows from (3.11) and

$$\langle \tilde{e}_j, V_\gamma^\delta \tilde{e}_j \rangle_{\ell^2(\mathbb{Z})} = \mathfrak{r}_\gamma^\delta(j) + O(j^{-1}) \text{ when } j \rightarrow \infty. \quad (3.12)$$

where $\mathfrak{r}_\gamma^\delta$ is given by (2.11)-(2.12). The proof of (3.12) is given in Section 3.3.

3.3. Proof of (3.12). Due to (2.32),

$$V_\gamma^\delta(n, n) = \delta \int_{-\pi}^{\pi} e^{in\Psi(\theta)} h_\gamma(\theta) \frac{d\theta}{2\pi}, \quad (3.13)$$

where $\Psi := \Phi - T_\pi \Phi$ and $h_\gamma(\theta) = p_\gamma(\theta) \overline{(T_\pi p_\gamma)(\theta)}$ with p_γ given by (2.31). Since $\pm \frac{\pi}{2}$ are non-degenerated critical points of Ψ (see (2.34), (2.35)), the stationary phase formula gives

$$V_\gamma^\delta(n, n) = \delta \sum_{\nu=\pm 1} \frac{h_\gamma(\nu\pi/2) e^{in\Psi(\nu\pi/2) + i\nu\pi/4}}{\sqrt{2\pi n |\Psi''(\nu\pi/2)|}} + O(n^{-1}). \quad (3.14)$$

Since Φ is odd, we get

$$\Psi(\pm\pi/2) = \Phi(\pm\pi/2) - \Phi(\mp\pi/2) = 2\Phi(\pm\pi/2) = \pm 2\Phi(\pi/2) = \pm 4\alpha \quad (3.15)$$

where (2.16) was used to write $\Phi(\pi/2) = 2\alpha$ with α given by (1.16). By definition (2.31), one has $p_\gamma(\pi/2) = \sqrt{\beta} e^{i\tilde{\theta}_\gamma}$ with

$$\tilde{\theta}_\gamma := \langle q_\gamma \rangle \Phi(\pi/2) - \tilde{q}_\gamma(\Phi(\pi/2)) \quad (3.16)$$

and $\tilde{q}_\gamma \circ \Phi$ is odd, hence $p_\gamma(-\pi/2) = \sqrt{\beta} e^{-i\tilde{\theta}_\gamma}$ and $h_\gamma(\pi/2) = \beta e^{2i\tilde{\theta}_\gamma}$. Moreover, one has $h_\gamma(-\pi/2) = \overline{h_\gamma(\pi/2)}$, hence

$$h_\gamma(\pm\pi/2) = \beta e^{\pm 2i\tilde{\theta}_\gamma} \quad (3.17)$$

and we get two conjugated terms corresponding to $\nu = \pm 1$ in (3.14). Therefore, using (3.17), (3.15) and $|\Psi''(\pm\pi/2)| = 4g\beta$ in (3.14), we get

$$\begin{aligned} V_\gamma^\delta(n, n) &= \delta \frac{2\operatorname{Re}(\beta e^{2i\tilde{\theta}_\gamma} e^{i4n\alpha + i\pi/4})}{\sqrt{2\pi n} \cdot 4g\beta} + O(n^{-1}) \\ &= \delta \frac{\sqrt{\beta} \cos(4n\alpha + 2\tilde{\theta}_\gamma + \pi/4)}{\sqrt{2\pi n} g} + O(n^{-1}). \end{aligned}$$

To complete the proof of (3.12), it remains to check that $\hat{\theta}_\gamma := 2\tilde{\theta}_\gamma + \pi/4$ is given by (2.12). However, (2.22) gives

$$\langle q_\gamma \rangle \Phi(\pi/2) = \left(\gamma - \frac{1}{2}\right) (\Phi(\pi/2) - \beta^{-1} \Phi(\pi/2)), \quad (3.18)$$

(2.21) gives

$$-\tilde{q}_\gamma(\Phi(\pi/2)) = \left(\gamma - \frac{1}{2}\right) (\beta^{-1} \Phi(\pi/2) - \pi/2), \quad (3.19)$$

and summing up (3.18), (3.19), we find that (3.16) gives

$$\tilde{\theta}_\gamma = \left(\gamma - \frac{1}{2}\right) (\Phi(\pi/2) - \pi/2) = \left(\gamma - \frac{1}{2}\right) (2\alpha - \pi/2) \quad (3.20)$$

and $\hat{\theta}_\gamma = 2\tilde{\theta}_\gamma + \frac{\pi}{4} = (\gamma - \frac{1}{2})(4\alpha - \pi) + \frac{\pi}{4}$ as claimed in (2.12).

4. AUXILIARY OPERATORS Q_n

4.1. Definition of Q_n . Our aim is to introduce a sequence of auxiliary self-adjoint operators $\{Q_n\}_{n \in \mathbb{N}^*}$ which will be used in Section 5 to estimate $\lambda_n(J)$ thanks to an analysis of the operator

$$J'_n := e^{-iQ_n} J e^{iQ_n}. \quad (4.1)$$

Definition 4.1. Let V be a self-adjoint bounded operator in $\ell^2(\mathbb{Z})$ and denote

$$V(j, k) := (V\tilde{e}_k)(j) = \langle \tilde{e}_j, V\tilde{e}_k \rangle_{\ell^2(\mathbb{Z})} \text{ for } (j, k) \in \mathbb{Z}^2.$$

For $n \in \mathbb{N}^*$ define $(Q_n(j, k))_{(j, k) \in \mathbb{Z}^2}$ by

$$Q_n(j, k) = \begin{cases} i \frac{V(j, n)}{\beta(j-n)} & \text{when } j \neq n \text{ and } k = n \\ i \frac{V(n, k)}{\beta(n-k)} & \text{when } j = n \text{ and } k \neq n \\ 0 & \text{otherwise} \end{cases} \quad (4.2)$$

Then

$$Q_n(j, k) \neq 0 \Rightarrow (\text{either } (j \neq n \text{ and } k = n) \text{ or } (j = n \text{ and } k \neq n)) \quad (4.3)$$

and the relation

$$(Q_n \tilde{e}_k)(j) = \langle \tilde{e}_j, Q_n \tilde{e}_k \rangle_{\ell^2(\mathbb{Z})} = Q_n(j, k) \text{ for } (j, k) \in \mathbb{Z}^2$$

defines in $\ell^2(\mathbb{Z})$ a self-adjoint Hilbert-Schmidt operator Q_n due to

$$\sum_{(j,k) \in \mathbb{Z}^2} |Q_n(j, k)|^2 \leq \frac{2}{\beta^2} \sum_{j \in \mathbb{Z}} |V(j, n)|^2 = \frac{2}{\beta^2} \|V \tilde{e}_n\|_{\ell^2(\mathbb{Z})}^2 \leq \frac{2}{\beta^2} \|V\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}^2$$

Notation 4.2. If $n \in \mathbb{Z}$ then $\hat{\Pi}_n \in \mathcal{B}(\ell^2(\mathbb{Z}))$ (respectively $\tilde{\Pi}_n \in \mathcal{B}(\ell^2(\mathbb{Z}))$) denotes the orthogonal projection on $\mathbb{C}\tilde{e}_n$ (respectively on $(\mathbb{C}\tilde{e}_n)^\perp$), i.e.

$$\hat{\Pi}_n x := \langle \tilde{e}_n, x \rangle_{\ell^2(\mathbb{Z})} \tilde{e}_n = x(n) \tilde{e}_n \quad \text{and} \quad \tilde{\Pi}_n := I - \hat{\Pi}_n.$$

4.2. Properties of Q_n . We observe that (4.3) implies

$$\hat{\Pi}_n Q_n \hat{\Pi}_n = 0 = \tilde{\Pi}_n Q_n \tilde{\Pi}_n \quad (4.4)$$

and

$$Q_n = \tilde{\Pi}_n Q_n \hat{\Pi}_n + \hat{\Pi}_n Q_n \tilde{\Pi}_n. \quad (4.5)$$

More properties of Q_n are given in

Lemma 4.3. *Assume that V satisfies the assumption of Proposition 3.1, i.e. V is a bounded self-adjoint operator in $\ell^2(\mathbb{Z})$ satisfying*

$$|j - n| \leq \hat{c}|n| \Rightarrow |V(j, n)| \leq \hat{C}|n|^{-1/2}, \quad (4.6)$$

where $\hat{C}, \hat{c} > 0$ are independent of n . Let $Q_n \in \mathcal{B}(\ell^2(\mathbb{Z}))$ be as in Definition 4.1. Then

(a) one has large n estimate

$$\|Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} = O(n^{-1/2}) \quad (4.7)$$

(b) $\ell^{2,1}(\mathbb{Z})$ is an invariant subspace of Q_n and for every $x \in \ell^{2,1}(\mathbb{Z})$,

$$i\beta[Q_n, \Lambda]x = i\beta(Q_n \Lambda - \Lambda Q_n)x = V_n x \quad (4.8)$$

holds with

$$V_n := \tilde{\Pi}_n V \hat{\Pi}_n + \hat{\Pi}_n V \tilde{\Pi}_n. \quad (4.9)$$

(c) $\ell^{2,1}(\mathbb{Z})$ is an invariant subspace of e^{itQ_n} and $t \rightarrow e^{itQ_n}x$ is of class $C^\infty(\mathbb{R}; \ell^{2,1}(\mathbb{Z}))$ if $x \in \ell^{2,1}(\mathbb{Z})$.

Proof. (a) By definition of Q_n ,

$$\|Q_n \tilde{e}_n\|_{\ell^2(\mathbb{Z})}^2 = \sum_{j \in \mathbb{Z} \setminus \{n\}} |Q_n(j, n)|^2 = \sum_{j \in \mathbb{Z} \setminus \{n\}} \frac{|V(j, n)|^2}{\beta^2(j-n)^2} \leq \beta^{-2}(\mathcal{M}_n + \mathcal{M}'_n)$$

holds with

$$\mathcal{M}_n := \sum_{\{j \in \mathbb{Z}: 0 < |j-n| \leq \hat{c}n\}} \frac{|V(j, n)|^2}{(j-n)^2} \leq \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{\hat{C}^2 n^{-1}}{m^2} = C_1 n^{-1}$$

due to (4.6) and

$$\mathcal{M}'_n := \sum_{\{j \in \mathbb{Z}: |j-n| > \hat{c}n\}} \frac{|V(j, n)|^2}{(j-n)^2} \leq \sum_{j \in \mathbb{Z}} \frac{|V(j, n)|^2}{\hat{c}^2 n^2} = \frac{\|V \tilde{e}_n\|_{\ell^2(\mathbb{Z})}^2}{\hat{c}^2 n^2} \leq C_2 n^{-2}.$$

This shows that $\|Q_n \widehat{\Pi}_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} = O(n^{-1/2})$. Moreover, due to (4.5),

$$\|Q_n \widetilde{\Pi}_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} = \|\widehat{\Pi}_n Q_n \widetilde{\Pi}_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq \|\widehat{\Pi}_n Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} = \|Q_n \widehat{\Pi}_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}$$

is $O(n^{-1/2})$, which completes the proof of (4.7).

(b) Let V_n be given by (4.9). Then one has

$$V_n(j, k) := (V_n \tilde{e}_k)(j) = \langle \tilde{e}_j, V_n \tilde{e}_k \rangle_{\ell^2(\mathbb{Z})} = \begin{cases} V(j, n) & \text{when } j \neq n \text{ and } k = n \\ V(n, k) & \text{when } j = n \text{ and } k \neq n \\ 0 & \text{otherwise} \end{cases} \quad (4.10)$$

and combining (4.10) with (4.2) we obtain

$$i\beta(j - k)Q_n(j, k) = -V_n(j, k) \text{ for every } j, k \in \mathbb{Z}. \quad (4.11)$$

However, (4.11) implies

$$i\beta(j + \frac{1}{2})(Q_n \tilde{e}_k)(j) = i\beta(Q_n(k + \frac{1}{2})\tilde{e}_k)(j) - (V_n \tilde{e}_k)(j) \quad (4.12)$$

and ensures $Q_n \tilde{e}_k \in \ell^{2,1}(\mathbb{Z})$. Moreover, (4.12) ensures the equality

$$i\beta(\Lambda + \frac{1}{2})Q_n x = i\beta Q_n(\Lambda + \frac{1}{2})x - V_n x \quad (4.13)$$

for $x \in \ell_{\text{fin}}^2(\mathbb{Z})$. If $y \in \ell_{\text{fin}}^2(\mathbb{Z})$, then using (4.13) with $x = (\Lambda + \frac{1}{2})^{-1}y$, we obtain

$$i\beta(\Lambda + \frac{1}{2})Q_n(\Lambda + \frac{1}{2})^{-1}y = i\beta Q'_n y \quad (4.14)$$

with $Q'_n \in \mathcal{B}(\ell^2(\mathbb{Z}))$ given by

$$Q'_n := Q_n - (i\beta)^{-1}V_n(\Lambda + \frac{1}{2})^{-1}. \quad (4.15)$$

Due to (4.14), for every $y \in \ell_{\text{fin}}^2(\mathbb{Z})$ one has

$$i\beta Q_n(\Lambda + \frac{1}{2})^{-1}y = i\beta(\Lambda + \frac{1}{2})^{-1}Q'_n y \quad (4.16)$$

and by continuity, (4.16) holds for every $y \in \ell^2(\mathbb{Z})$. Consider now $x \in \ell^{2,1}(\mathbb{Z})$ and take $y = (\Lambda + \frac{1}{2})x$ in (4.16). This gives

$$i\beta Q_n x = i\beta(\Lambda + \frac{1}{2})^{-1}Q'_n(\Lambda + \frac{1}{2})x \in \ell^{2,1}(\mathbb{Z}) \quad (4.17)$$

and

$$i\beta(\Lambda + \frac{1}{2})Q_n x = i\beta Q'_n(\Lambda + \frac{1}{2})x = i\beta Q_n(\Lambda + \frac{1}{2})x - V_n x, \quad (4.18)$$

implying (4.8) for every $x \in \ell^{2,1}(\mathbb{Z})$.

(c) We first observe that (4.16) holds for every $y \in \ell^2(\mathbb{Z})$ and implies (4.14) for every $y \in \ell^2(\mathbb{Z})$. Since (4.14) implies $(Q'_n)^m y = (\Lambda + \frac{1}{2})(Q_n)^m(\Lambda + \frac{1}{2})^{-1}y$ for every $m \in \mathbb{N}$ and $y \in \ell^2(\mathbb{Z})$, we have

$$\begin{aligned} e^{itQ_n}(\Lambda + \frac{1}{2})^{-1}y &= \lim_{N \rightarrow \infty} \sum_{m=0}^N \frac{(it)^m}{m!} (Q_n)^m (\Lambda + \frac{1}{2})^{-1}y = \\ &= \lim_{N \rightarrow \infty} \sum_{m=0}^N \frac{(it)^m}{m!} (\Lambda + \frac{1}{2})^{-1} (Q'_n)^m y = (\Lambda + \frac{1}{2})^{-1} e^{itQ'_n} y \in \ell^{2,1}(\mathbb{Z}) \end{aligned}$$

if $y \in \ell^2(\mathbb{Z})$ and setting $x = (\Lambda + \frac{1}{2})^{-1}y$, we find that $t \rightarrow (\Lambda + \frac{1}{2})e^{itQ_n}x = e^{itQ'_n}y$ is $C^\infty(\mathbb{R}; \ell^2(\mathbb{Z}))$ for every $x \in \ell^{2,1}(\mathbb{Z})$. \square

5. PROOF OF PROPOSITION 3.1

5.1. Taylor's expansion formula. We will investigate $J'_n = e^{-iQ_n} J e^{iQ_n}$, using a special version of the Taylor's expansion formula. Assume that B and $Q \in \mathcal{B}(\ell^2(\mathbb{Z}))$ and denote

$$F_{tQ}(B) := e^{-itQ} B e^{itQ} \text{ for } t \in \mathbb{R}, \quad (5.1)$$

$$\text{ad}_{iQ}^1(B) := [B, iQ] = i(BQ - QB)$$

and

$$\text{ad}_{iQ}^{m+1}(B) := [\text{ad}_{iQ}^m(B), iQ] \text{ for } m \in \mathbb{N}^*.$$

Then

$$\frac{d^m}{dt^m} F_{tQ}(B) = e^{-itQ} \text{ad}_{iQ}^m(B) e^{itQ} = F_{tQ}(\text{ad}_{iQ}^m(B)) \quad (5.2)$$

and the Taylor formula gives

$$F_{tQ}(B) = B + \sum_{m=1}^{N-1} \frac{t^m}{m!} \text{ad}_{iQ}^m(B) + \mathcal{R}_Q^{t,N}(B) \quad (5.3)$$

with

$$\mathcal{R}_Q^{t,N}(B) := \frac{t^N}{(N-1)!} \int_0^1 F_{stQ}(\text{ad}_{iQ}^N(B)) (1-s)^{N-1} ds. \quad (5.4)$$

We can also consider the case when B is an unbounded symmetric operator in $\ell^2(\mathbb{Z})$, defined on a dense domain $\mathcal{D}(B)$. Suppose that $\mathcal{D}(B)$ is an invariant subspace for Q and e^{itQ} for every $t \in \mathbb{R}$. If $t \rightarrow B e^{itQ} x$ is $C^1(\mathbb{R}, \ell^2(\mathbb{Z}))$ for every $x \in \mathcal{D}(B)$, then

$$\frac{d}{dt} \langle e^{itQ} x, B e^{itQ} y \rangle_{\ell^2(\mathbb{Z})} = \langle B e^{itQ} x, iQ e^{itQ} y \rangle_{\ell^2(\mathbb{Z})} + \langle iQ e^{itQ} x, B e^{itQ} y \rangle_{\ell^2(\mathbb{Z})}$$

holds for every $x, y \in \mathcal{D}(B)$. If the form $(x, y) \rightarrow \langle Bx, iQy \rangle_{\ell^2(\mathbb{Z})} + \langle iQx, By \rangle_{\ell^2(\mathbb{Z})}$ can be extended from $\mathcal{D}(B) \times \mathcal{D}(B)$ to a bounded form on $\ell^2(\mathbb{Z}) \times \ell^2(\mathbb{Z})$, then we can introduce $[B, iQ] \in \mathcal{B}(\ell^2(\mathbb{Z}))$ defined by this form and we can write

$$\frac{d}{dt} F_{tQ}(B) = e^{-itQ} [B, iQ] e^{itQ} = F_{tQ}([B, iQ]),$$

hence (5.3)-(5.4) still hold for every $N \in \mathbb{N}^*$.

5.2. Step 1 of the proof of Proposition 3.1.

Notation 5.1. If L_n, L'_n are linear maps defined on a dense subspace of the Banach space \mathcal{V} and $L_n - L'_n$ can be extended to a bounded operator on \mathcal{V} , then $\|L_n - L'_n\|_{\mathcal{B}(\mathcal{V})}$ denotes the norm of this extension and the notation $L_n = L'_n + O(\eta_n)$ means that choosing $n_0 \in \mathbb{N}$ and $C > 0$ large enough, we ensure $\|L_n - L'_n\|_{\mathcal{B}(\mathcal{V})} \leq C\eta_n$ for all $n \geq n_0$. If $x_n, x'_n \in \mathcal{V}$, then we write $x_n = y_n + O(\eta_n)$ if and only if we can find $C > 0$, $n_0 \in \mathbb{N}$, such that $\|x_n - x'_n\|_{\mathcal{V}} \leq C\eta_n$ holds for all $n \geq n_0$.

In what follows, V satisfies the hypotheses of Proposition 3.1 and without any loss of generality we assume $\beta_0 = 0$, i.e. $J = \beta\Lambda + V$. We consider

$$J'_n = e^{-iQ_n} (\beta\Lambda + V) e^{iQ_n} \quad (5.5)$$

and claim the large n estimate

$$J'_n = \beta\Lambda + V - V_n + [V - \frac{1}{2}V_n, iQ_n] + O(n^{-1}), \quad (5.6)$$

where $V_n = \widetilde{\Pi}_n V \widehat{\Pi}_n + \widehat{\Pi}_n V \widetilde{\Pi}_n$ is given by (4.9).

We first use (5.3)-(5.4) with $N = 2$ and write

$$e^{-iQ_n} V e^{iQ_n} = V + [V, iQ_n] + \mathcal{R}_{Q_n}^{1,2}(V), \quad (5.7)$$

$$\|\mathcal{R}_{Q_n}^{1,2}(V)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq \|\text{ad}_{iQ_n}^2(V)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq C \|Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}^2. \quad (5.8)$$

We also use (5.3)-(5.4) with $N = 3$ and write

$$e^{-iQ_n} \beta \Lambda e^{iQ_n} = \beta \Lambda + [\beta \Lambda, iQ_n] + \frac{1}{2} [[\beta \Lambda, iQ_n], iQ_n] + \mathcal{R}_{Q_n}^{1,3}(\beta \Lambda), \quad (5.9)$$

$$\|\mathcal{R}_{Q_n}^{1,3}(\beta \Lambda)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq \|\text{ad}_{iQ_n}^3(\beta \Lambda)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}. \quad (5.10)$$

Using $[\beta \Lambda, iQ_n] = -V_n$ (see Lemma 4.3b) in (5.9)-(5.10), we obtain

$$e^{-iQ_n} \beta \Lambda e^{iQ_n} = \beta \Lambda - V_n - [\frac{1}{2} V_n, iQ_n] + \mathcal{R}_{Q_n}^{1,3}(\beta \Lambda), \quad (5.11)$$

$$\|\mathcal{R}_{Q_n}^{1,3}(\beta \Lambda)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq \|\text{ad}_{iQ_n}^2(V_n)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq C \|Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}^2 \quad (5.12)$$

and summing up (5.11), (5.7), we obtain

$$e^{-iQ_n} J e^{iQ_n} = \beta \Lambda + V - V_n + [V - \frac{1}{2} V_n, iQ_n] + \mathcal{R}_{Q_n}^{1,2}(V) + \mathcal{R}_{Q_n}^{1,3}(\beta \Lambda), \quad (5.13)$$

$$\|\mathcal{R}_{Q_n}^{1,2}(V) + \mathcal{R}_{Q_n}^{1,3}(\beta \Lambda)\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq C \|Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}^2. \quad (5.14)$$

To complete the proof, we observe that $\|Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))}^2 = O(n^{-1})$ follows from (4.7).

5.3. Step 2 of the proof of Proposition 3.1. We denote

$$W_n := [V - \frac{1}{2} V_n, iQ_n] \quad (5.15)$$

and we claim that the following estimate

$$W_n(n, n) := (W_n \tilde{e}_n)(n) = \langle \tilde{e}_n, W_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} = O(n^{-1} \ln n) \quad (5.16)$$

holds when $n \rightarrow \infty$.

Indeed, $W_n = [V'_n, iQ_n]$ holds with $V'_n := V - \frac{1}{2} V_n$ and

$$|V'_n(j, k)| \leq 2|V(j, k)| \text{ for all } j, k \in \mathbb{Z}, \quad (5.17)$$

hence we can estimate

$$\begin{aligned} |(V'_n Q_n)(n, n)| &\leq \sum_{j \in \mathbb{Z}} 2|V(n, j) Q_n(j, n)| \leq \\ &\leq \sum_{j \in \mathbb{Z} \setminus \{n\}} \frac{2|V(j, n)|^2}{\beta |j - n|} \leq 2\beta^{-1} (\mathcal{M}_n + \mathcal{M}'_n) \end{aligned}$$

with

$$\mathcal{M}_n := \sum_{\{j \in \mathbb{Z}: 0 < |j - n| \leq \hat{c}n\}} \frac{|V(j, n)|^2}{|j - n|} \leq \sum_{\{m \in \mathbb{Z}: 0 < |m| \leq \hat{c}n\}} \frac{\hat{C}^2 n^{-1}}{|m|} \leq C_1 n^{-1} \ln n$$

due to (4.6) and

$$\mathcal{M}'_n := \sum_{\{j \in \mathbb{Z}: |j - n| > \hat{c}n\}} \frac{|V(j, n)|^2}{|j - n|} \leq \sum_{j \in \mathbb{Z}} \frac{|V(j, n)|^2}{\hat{c}n} = \frac{\|V \tilde{e}_n\|_{\ell^2(\mathbb{Z})}^2}{\hat{c}n} \leq C_2 n^{-1}.$$

5.4. **Step 3 of the proof of Proposition 3.1.** Let J'_n be as above and denote

$$d'_n := \beta n + V(n, n). \quad (5.18)$$

We claim that the following estimates

$$\langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} = d'_n + O(n^{-1} \ln n) \quad (5.19)$$

$$(J'_n - \langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})}) \tilde{e}_n = O(n^{-1/2}) \quad (5.20)$$

hold when $n \rightarrow \infty$.

Indeed, writing

$$\begin{aligned} \beta\Lambda + V - V_n &= \beta\Lambda + V - \tilde{\Pi}_n V \hat{\Pi}_n - \hat{\Pi}_n V \tilde{\Pi}_n \\ &= \beta\Lambda + \hat{\Pi}_n V \hat{\Pi}_n + \tilde{\Pi}_n V \tilde{\Pi}_n = \beta\Lambda + V(n, n) \hat{\Pi}_n + \tilde{\Pi}_n V \tilde{\Pi}_n \end{aligned}$$

in (5.6), we find

$$J'_n \tilde{e}_n = (\beta\Lambda + V(n, n) \hat{\Pi}_n + \tilde{\Pi}_n V \tilde{\Pi}_n + W_n) \tilde{e}_n + O(n^{-1}), \quad (5.21)$$

where $W_n = [V - \frac{1}{2}V_n, iQ_n]$ is given by (5.15). However (5.21) implies

$$J'_n \tilde{e}_n = d'_n \tilde{e}_n + W_n \tilde{e}_n + O(n^{-1}), \quad (5.22)$$

hence

$$\langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} = d'_n + \langle \tilde{e}_n, W_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} + O(n^{-1}) \quad (5.23)$$

and (5.19) follows from (5.16).

In order to prove (5.20), we observe that (5.19) implies

$$(J'_n - \langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})}) \tilde{e}_n = (J'_n - d'_n) \tilde{e}_n + O(n^{-1} \ln n) \quad (5.24)$$

and (5.22) ensures

$$(J'_n - d'_n) \tilde{e}_n = W_n \tilde{e}_n + O(n^{-1}). \quad (5.25)$$

Thus (5.20) follows from (5.24), (5.25) and

$$\|W_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} \leq C \|Q_n\|_{\mathcal{B}(\ell^2(\mathbb{Z}))} = O(n^{-1/2}).$$

5.5. **End of the proof of Proposition 3.1.** In what follows, $\sigma(L)$ denotes the spectrum of the operator L . We assume that the hypotheses of Proposition 3.1 hold and $\beta_0 = 0$. Then (3.3) ensures $\lambda_n(J) = \beta n + o(1)$ as $n \rightarrow \infty$ and

$$\sigma(J) \cap [\beta(n - \frac{1}{2}), \beta(n + \frac{1}{2})] = \{\lambda_n(J)\} \text{ for } n \geq n_0 \quad (5.26)$$

if $n_0 \in \mathbb{N}$ is chosen large enough. Let $\{\lambda_j(J'_n)\}_{j \in \mathbb{Z}}$ be the sequence of eigenvalues of J'_n labeled so that $\lambda_j(J'_n) = \lambda_j(J)$ for every $j \in \mathbb{Z}$. Then (5.26) gives

$$\sigma(J'_n) \cap [\beta(n - \frac{1}{2}), \beta(n + \frac{1}{2})] = \{\lambda_n(J'_n)\} \text{ for } n \geq n_0 \quad (5.27)$$

and we will use $L = \tilde{J}'_n$, $x = e_n$, $d' = \beta(n - \frac{1}{2})$ and $d'' = \beta(n + \frac{1}{2})$ with $n \geq n_0$ in

Theorem 5.2. (Kato-Temple) *Assume that the operator L is self-adjoint in the Hilbert space \mathcal{H} and has exactly one eigenvalue λ in the interval $[d', d'']$. If x is an element of the domain of L such that $\eta := \langle x, Lx \rangle_{\mathcal{H}}$ belongs to $]d', d''[$, then*

$$\eta - \frac{\|(L - \eta)x\|^2}{\eta - d'} \leq \lambda \leq \eta + \frac{\|(L - \eta)x\|^2}{d'' - \eta}. \quad (5.28)$$

Proof. See [16]. □

We claim that the assumptions of Theorem 5.2 are satisfied for $n \geq n_0$ if n_0 is chosen large enough. Indeed, (5.27) holds and

$$\eta := \langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} \in [\beta(n - \frac{1}{4}), \beta(n + \frac{1}{4})] \text{ for } n \geq n_0 \quad (5.29)$$

holds due to (5.19) and $V(n, n) = O(n^{-1/2})$ as $n \rightarrow \infty$. Since (5.29) implies the inequality $\min\{d'' - \eta, \eta - d'\} \geq \frac{1}{4}\beta$, the estimate (5.28) gives

$$|\lambda_n(J'_n) - \langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})}| \leq 4\beta^{-1} \| (J'_n \tilde{e}_n - \langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} \tilde{e}_n) \|^2 = O(n^{-1}), \quad (5.30)$$

where the last estimate is due to (5.20). The estimate (5.30) implies

$$\lambda_n(J'_n) = \langle \tilde{e}_n, J'_n \tilde{e}_n \rangle_{\ell^2(\mathbb{Z})} + O(n^{-1}) \quad (5.31)$$

and using (5.19) in the right hand side of (5.31), we obtain

$$\lambda_n(J) = \lambda_n(J'_n) = d'_n + O(n^{-1} \ln n),$$

completing the proof of Proposition 3.1.

6. PROOF OF THEOREM 1.3

6.1. Step 1 of the proof of Theorem 1.3. As before $0 < g < \frac{1}{2}$ and $\{e_m\}_{m \in \mathbb{N}}$ is the canonical basis of $\ell^2(\mathbb{N})$.

Definition 6.1. For $\nu = \pm 1$, $\mu = 0, 1$, we define $\hat{J}_{\nu, \mu}$ as the linear map $\hat{J}_{\nu, \mu} : \ell^{2,1}(\mathbb{N}) \rightarrow \ell^2(\mathbb{N})$ given by

$$\begin{cases} \hat{J}_{\nu, \mu} e_m = \hat{d}_{\nu, \mu}(m) e_m + \hat{b}_\mu(m) e_{m+1} + \hat{b}_\mu(m-1) e_{m-1} \text{ for } m \in \mathbb{N}^*, \\ \hat{J}_{\nu, \mu} e_0 = \hat{d}_{\nu, \mu}(0) e_0 + \hat{b}_\mu(0) e_1 \end{cases} \quad (6.1)$$

where

$$\begin{cases} \hat{d}_{\nu, \mu}(m) := 2m + \mu + (-1)^m \nu \Delta / 2 \\ \hat{b}_\mu(m) := g \sqrt{(2m+1+\mu)(2m+2+\mu)}. \end{cases} \quad (6.2)$$

Lemma 6.2. The operator $\mathbf{H}_{\text{Rabi}}^{(2)}$ is similar to the direct sum

$$\hat{J}_{1,0} \oplus \hat{J}_{-1,0} \oplus \hat{J}_{1,1} \oplus \hat{J}_{-1,1} \quad (6.3)$$

Proof. See Section 2.2 in [6]. \square

6.2. Step 2 of the proof of Theorem 1.3. Let \tilde{J}_γ^δ , \tilde{d}_δ , \tilde{b}_γ be as in Definition 2.2. If $\hat{d}_{\nu, \mu}$ and \hat{b}_μ are given by (6.2), then it is easy to check the following relations

$$\begin{cases} \hat{d}_{\pm 1, 0}(m) = 2\tilde{d}_{\pm \Delta/4}(m) \\ \hat{b}_0(m) = 2\tilde{b}_{3/4}(m) + O(m^{-1}) \text{ when } m \rightarrow \infty \end{cases} \quad (6.4)$$

$$\begin{cases} \hat{d}_{\pm 1, 1}(m) = 2\tilde{d}_{\pm \Delta/4}(m) + 1 \\ \hat{b}_1(m) = 2\tilde{b}_{5/4}(m) + O(m^{-1}) \text{ when } m \rightarrow \infty \end{cases} \quad (6.5)$$

The above properties allow us to prove

Lemma 6.3. *Let $0 < g < \frac{1}{2}$ and let $\{\lambda_n(\hat{J}_{\nu,\mu})\}_{n \in \mathbb{N}}$ denote the non-decreasing sequence of eigenvalues of $\hat{J}_{\nu,\mu}$. If $\{\lambda_j(\tilde{J}_\gamma^\delta)\}_{j \in \mathbb{Z}}$ is the sequence of eigenvalues of \tilde{J}_γ^δ satisfying (2.10), then one has*

$$\lambda_n(\hat{J}_{\pm 1,0}) = 2\lambda_n(\tilde{J}_{3/4}^{\pm \Delta/4}) + O(n^{-1}) \text{ as } n \rightarrow \infty \quad (6.6)$$

and

$$\lambda_n(\hat{J}_{\pm 1,1}) = 2\lambda_n(\tilde{J}_{5/4}^{\pm \Delta/4}) + 1 + O(n^{-1}) \text{ as } n \rightarrow \infty. \quad (6.7)$$

Proof. We identify $\ell^2(\mathbb{N})$ with $\{x \in \ell^2(\mathbb{Z}) : x(k) = 0 \text{ for } k \in \mathbb{Z} \setminus \mathbb{N}\}$. Let $\Pi_{\geq 0}$ be the orthogonal projection $\ell^2(\mathbb{Z}) \rightarrow \ell^2(\mathbb{N})$ and let $\hat{J}_\gamma^\delta := \Pi_{\geq 0} \tilde{J}_\gamma^\delta |_{\ell^2(\mathbb{N})}$. Identifying \tilde{e}_m with e_m when $m \in \mathbb{N}$, we get

$$\begin{cases} \hat{J}_\gamma^\delta e_m = \hat{d}_\delta(m)e_m + \hat{b}_\gamma(m)e_{m+1} + \hat{b}_\gamma(m-1)e_{m-1} \text{ for } m \in \mathbb{N}^*, \\ \hat{J}_\gamma^\delta e_0 = \hat{d}_\delta(0)e_0 + \hat{b}_\gamma(0)e_1 \end{cases} \quad (6.8)$$

We observe that (6.4)-(6.5) imply

$$2\hat{J}_{3/4}^{\pm \Delta/4} - \hat{J}_{\pm 1,0} \in \mathcal{B}(\ell^2(\mathbb{N}), \ell^{2,1}(\mathbb{N})), \quad (6.9)$$

$$2\hat{J}_{5/4}^{\pm \Delta/4} + 1 - \hat{J}_{\pm 1,1} \in \mathcal{B}(\ell^2(\mathbb{N}), \ell^{2,1}(\mathbb{N})) \quad (6.10)$$

and using the result of Rozenblum [24, Theorem 1.1] as indicated in the proof of (3.6) in [6, Section 3.2], we find that (6.9)-(6.10) ensure

$$2\lambda_n(\hat{J}_{3/4}^{\pm \Delta/4}) = \lambda_n(2\hat{J}_{3/4}^{\pm \Delta/4}) = \lambda_n(\hat{J}_{\pm 1,0}) + O(n^{-1}) \text{ as } n \rightarrow \infty, \quad (6.11)$$

$$2\lambda_n(\hat{J}_{5/4}^{\pm \Delta/4}) + 1 = \lambda_n(2\hat{J}_{5/4}^{\pm \Delta/4} + 1) = \lambda_n(\hat{J}_{\pm 1,1}) + O(n^{-1}) \text{ as } n \rightarrow \infty, \quad (6.12)$$

where $\{\lambda_n(\hat{J}_\gamma^\delta)\}_{n \in \mathbb{N}}$ is the non-decreasing sequence of eigenvalues of \hat{J}_γ^δ .

Let $\{\lambda_j(\tilde{J}_\gamma^\delta)\}_{j \in \mathbb{Z}}$ be the sequence of eigenvalues of \tilde{J}_γ^δ satisfying (2.10). We complete the proof, combining (6.11)-(6.12) and the estimate

$$\lambda_n(\hat{J}_\gamma^\delta) - \lambda_n(\tilde{J}_\gamma^\delta) = O(n^{-N}) \text{ as } n \rightarrow \infty, \quad (6.13)$$

which holds for every $N \geq 1$. The estimate (6.13) is proved in Section 6.1 of [6]. \square

6.3. End of the proof of Theorem 1.3. Using the asymptotic estimates (2.10) in the right hand side of (6.6) and (6.7), we obtain

$$\lambda_n(\hat{J}_{\pm 1,0}) = 2 \left(n\beta + \frac{1}{4}(\beta - 1) + \mathfrak{r}_{3/4}^{\pm \Delta/4}(n) \right) + O(n^{-1} \ln n), \quad (6.14)$$

$$\lambda_n(\hat{J}_{\pm 1,1}) = 2 \left(n\beta + \frac{3}{4}(\beta - 1) + \mathfrak{r}_{5/4}^{\pm \Delta/4}(n) \right) + 1 + O(n^{-1} \ln n). \quad (6.15)$$

Due to (2.12),

$$\begin{aligned} \hat{\theta}_{3/4} &= \frac{1}{4}(4\alpha - \pi) + \frac{\pi}{4} = \alpha, \\ \hat{\theta}_{5/4} &= \frac{3}{4}(4\alpha - \pi) + \frac{\pi}{4} = 3\alpha - \frac{\pi}{2} \end{aligned}$$

and (6.14)-(6.15) give

$$\lambda_n(\hat{J}_{\pm 1,0}) = 2n\beta + \frac{1}{2}(\beta - 1) \pm \frac{\Delta}{2} \left(\frac{\beta}{2\pi gn} \right)^{1/2} \cos((4n+1)\alpha) + O(n^{-1} \ln n),$$

$$\lambda_n(\hat{J}_{\pm 1,1}) = (2n+1)\beta + \frac{1}{2}(\beta - 1) \pm \frac{\Delta}{2} \left(\frac{\beta}{2\pi gn} \right)^{1/2} \sin((4n+3)\alpha) + O(n^{-1} \ln n)$$

Due to Lemma 6.2, $\sigma(\mathbf{H}_{\text{Rabi}}^{(2)}) = \sigma(\hat{J}_{1,0}) \cup \sigma(\hat{J}_{-1,0}) \cup \sigma(\hat{J}_{1,1}) \cup \sigma(\hat{J}_{-1,1})$ and we complete the proof, taking

$$\begin{cases} E_{2n}^{\pm} = \lambda_n(\hat{J}_{\pm 1,0}) \\ E_{2n+1}^{\pm} = \lambda_n(\hat{J}_{\pm 1,1}) \end{cases} \quad (6.16)$$

for $n \in \mathbb{N}$.

7. APPENDIX: ESTIMATES OF OSCILLATORY INTEGRALS

Lemma 7.1. *Let us fix real numbers θ_0, θ_1 and $c_0 > 0$. Assume that $\{\Psi_{\tau}\}_{\tau \in \mathcal{T}}$ is a family of functions $[\theta_0, \theta_1] \rightarrow \mathbb{R}$, bounded in $C^2([\theta_0, \theta_1])$, i.e. $\sup_{\theta_0 \leq \theta \leq \theta_1} |\Psi_{\tau}^{(k)}(\theta)| < \infty$ for $k = 0, 1, 2$. For $\lambda \in \mathbb{R}$ and $h \in C^2([\theta_0, \theta_1])$ we denote*

$$M_{\tau}(\lambda) := \int_{\theta_0}^{\theta_1} e^{i\lambda\Psi_{\tau}(\theta)} h(\theta) d\theta. \quad (7.1)$$

(a) *If $|\Psi'_{\tau}(\theta)| \geq c_0 > 0$ holds for every $\theta \in [\theta_0, \theta_1]$ and $\tau \in \mathcal{T}$, then there is a constant C such that the estimate*

$$|M_{\tau}(\lambda)| \leq C|\lambda|^{-1} \quad (7.2)$$

holds for all $\tau \in \mathcal{T}$ and $\lambda \in \mathbb{R} \setminus \{0\}$.

(b) *If $|\Psi''_{\tau}(\theta)| \geq c_0 > 0$ holds for every $\theta \in [\theta_0, \theta_1]$ and $\tau \in \mathcal{T}$, then there is a constant C such that the estimate*

$$|M_{\tau}(\lambda)| \leq C|\lambda|^{-1/2} \quad (7.3)$$

holds for all $\tau \in \mathcal{T}$ and $\lambda \in \mathbb{R} \setminus \{0\}$.

Proof. (a) The non-stationary phase estimate (7.2) follows by using the standard integration by parts.

(b) The estimate (7.3) follows from van der Corput Lemma (see [27, Section VIII.1.2, p. 354]). \square

REFERENCES

- [1] A. Boutet de Monvel, S. Naboko, and L. O. Silva, *Eigenvalue asymptotics of a modified Jaynes–Cummings model with periodic modulations*, *Comptes Rendus, Mathématique, Acad. Sci. Paris* **338** (2004), no. 1, 103–107.
- [2] ———, *The asymptotic behaviour of eigenvalues of a modified Jaynes–Cummings model*, *Asymptot. Anal.* **47** (2006), no. 3–4, 291–315.
- [3] A. Boutet de Monvel and L. Zielinski, *Asymptotic behaviour of large eigenvalues for Jaynes–Cummings type models*, *J. Spectr. Theory*, vol. 7, 2017, pp. 559–631, available at [arXiv:1503.05841](https://arxiv.org/abs/1503.05841).
- [4] ———, *Oscillatory Behavior of Large Eigenvalues in Quantum Rabi Models*, *International Mathematics Research Notices*, vol. 7, 2021, pp. 5155–5213, available at [arXiv:1711.03366](https://arxiv.org/abs/1711.03366).
- [5] A. Boutet de Monvel and L. Zielinski, *On the spectrum of the quantum Rabi Models*, *Analysis as a Tool in Mathematical Physics : In Memory of Boris Pavlov. Operator Theory: Advances and Applications*, vol. 276, Birkhäuser, 2020.
- [6] ———, *Asymptotic behavior of large eigenvalues of the two-photon Rabi model*, *Schrödinger Operators, Spectral Theory and Number Theory*, vol. 348, Springer Proceedings in Mathematics & Statistics, 2021, pp. 89–115, available at <https://hal.science/hal-04147636>.
- [7] ———, *Asymptotic formula for large eigenvalues of the two-photon Rabi model*, *Comptes Rendus, Mathématique*, 2023, pp. 1761–1766, available at <https://doi.org/10.5802/crmath.515>.
- [8] D. Braak, Q.-H. Chen, M. T. Batchelor, and E. Solano, *Semi-classical and quantum Rabi models: in celebration of 80 years*, *J. of Physics A* **49** (2016), 300301.
- [9] L. Duan, Y.-F. Xie, D. Braak, and Q.-H. Chen, *Two-photon Rabi model: analytic solutions and spectral collapse*, *Journal of Physics A: Mathematical and Theoretical* **49** (2016), no. 46, 464002.
- [10] J. Edward, *Spectra of Jacobi matrices, differential equations on the circle, and the $su(1, 1)$ Lie algebra*, *SIAM J. Math. Anal.* **24** (2006), no. 3, 824–831.

- [11] C. Emary and R. F. Bishop, *Bogoliubov transformations and exact isolated solutions for simple nonadiabatic Hamiltonians*, Journal of Mathematical Physics **43** (2002), no. 8, 3916–3926.
- [12] ———, *Exact isolated solutions for the two-photon Rabi Hamiltonian*, J. Physics A: Math. Gen **35** (2002), no. 39, 8231–8241.
- [13] I. D. Feranchuk, L. I. Komarov, and A. P. Ulyanenko, *Two-level system in a one-mode quantum field: numerical solution on the basis of the operator method*, J. Phys. A: Math. Gen. (1996), 4035–4047.
- [14] I. Feranchuk, A. Ivanov, V.-H. Le, and A. Ulyanenko, *Non-perturbative Description of Quantum Systems*, Lecture Notes in Physics **894** (Springer International Publishing, 2015).
- [15] C. Gerry, *Two-photon Jaynes-Cummings model interacting with the squeezed vacuum*, Phys. Rev. A **37** (1988), no. 7, 2683–2686.
- [16] M. E. Harrell, *Generalizations of Temple’s Inequality*, Proceedings of the American Mathematical Society **69** (1978), no. 2, 271–276.
- [17] E. A. Ianovich, *Eigenvalues asymptotics of unbounded operators. Two-photon quantum Rabi model* (December 2023), arXiv:2312.05646.
- [18] E. K. Irish, *Generalized Rotating-Wave Approximation for Arbitrarily Large Coupling*, Physical Review Letters **99** (2007), no. 17.
- [19] J. Janas and S. Naboko, *Infinite Jacobi matrices with unbounded entries: asymptotics of eigenvalues and the transformation operator approach*, SIAM J. Math. Anal. **36** (2004), no. 2, 643–658 (electronic).
- [20] E. T. Jaynes and F. W. Cummings, *Comparison of quantum and semiclassical radiation theories with application to the beam maser*, Proc. IEEE **51** (1963), no. 1, 89–109.
- [21] X. Qiongtao, Z. Honghua, T. B. Murray, and L. Chaohong, *The quantum Rabi model: solution and dynamics*, J. Phys. A: Math. Theor. **50** (2017), 113001.
- [22] I. I. Rabi, *On the process of space quantization*, Phys. Rev. **49** (1936), 324.
- [23] I. I. Rabi, *Space quantization in a gyrating magnetic field*, Phys. Rev. **51** (1937), 652.
- [24] G. V. Rozenblum, *Near-similarity of operators and the spectral asymptotic behavior of pseudodifferential operators on the circle*, Tr. Mosk. Mat. Obs. **36** (1975), 59–84.
- [25] J. Sahbani, *Spectral theory of certain unbounded Jacobi matrices*, Journal of Mathematical Analysis and Applications **342** (2008), no. 1, 663–681.
- [26] M. Schmutz, *Two-level system coupled to a boson mode: the large n limit*, Journal of Physics A: Mathematical and General **19** (1986), no. 17, 3565–3577.
- [27] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy; Monographs in Harmonic Analysis, III.
- [28] È. A. Tur, *Jaynes-Cummings model: solution without rotating wave approximation*, Optics and Spectroscopy **89** (2000), no. 4, 574–588.
- [29] ———, *Jaynes-Cummings model without rotating wave approximation*, arXiv:math-ph/0211055v1 22 Nov 2002.
- [30] E. A. (Tur) Yanovich, *Asymptotics of eigenvalues of an energy operator in a problem of quantum physics*, J. Janas, P. Kurasov, A. Laptev, and S. Naboko (eds), Operator Methods in Mathematical Physics OTAMP 2010, Bedlewo, Oper. Theory Adv. Appl., vol. 227, Birkhäuser Verlag, Basel, 2013.

¹UNIVERSITÉ PARIS CITÉ, BÂTIMENT SOPHIE GERMAIN, BOÎTE COURRIER 7012, 75205 PARIS CEDEX 13, FRANCE, E-MAIL: ANNE.BOUTET-DE-MONVEL@IMJ-PRG.FR

²LABORATOIRE DE MATHÉMATIQUES PURES ET APPLIQUÉES JOSEPH LIOUVILLE UR 2597, UNIVERSITÉ DU LITTORAL CÔTE D’OPALE, F-62228 CALAIS, FRANCE, E-MAIL: LECH.ZIELINSKI@UNIV-LITTORAL.FR