



HAL
open science

Observation-only learning of 4DVarNet neural schemes for the reconstruction of sea surface turbidity dynamics from gappy satellite images

Clément Dorffer, Frédéric Jourdin, David Mouillot, Rodolphe Devillers, Ronan Fablet

► To cite this version:

Clément Dorffer, Frédéric Jourdin, David Mouillot, Rodolphe Devillers, Ronan Fablet. Observation-only learning of 4DVarNet neural schemes for the reconstruction of sea surface turbidity dynamics from gappy satellite images. EGU 2023: European Geophysical Union, European Geophysical Union, Apr 2023, Vienna, Austria. pp.1-1, 10.5194/egusphere-egu23-15977 . hal-04363493

HAL Id: hal-04363493

<https://hal.science/hal-04363493>

Submitted on 26 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Observation-only learning of 4DVarNet neural schemes for the reconstruction of sea surface turbidity dynamics from gappy satellite images

Clément Dorffer¹, Frédéric Jourdin², David Mouillot³, Rodolphe Devillers⁴, and Ronan Fablet¹

¹IMT Atlantique, UMR CNRS Lab-STICC, INRIA team Odyssey, Brest, France

²Service Hydrographique et Océanographique de la Marine (SHOM), 29603 Brest, France

³MARBEC, University of Montpellier, CNRS, IFREMER, IRD, Montpellier, France

⁴Espace-Dev (IRD-UM-UG-UR-UA-UNC), Station SEAS-OI, 97410 Saint-Pierre, La Réunion, France

Optical remote sensing is increasingly used to assess various sea surface biogeochemical parameters (e.g., Chl-a, turbidity). If today's systems offer better spatiotemporal coverage, the space-time sampling depends on both the satellite orbit and the cloud cover. The resulting sea surface observations generally present large proportions of missing data, making their completion challenging.

Here, we explore neural interpolation schemes as an approach for image gap filling, and their training from observation-only datasets with large missing data rates (with a mean of 65% of missing data and up to 100% for the worst days of the time-series), i.e., when no reference gap-free data are available to run a classic supervised learning approach. We propose and assess different strategies based on real or simulated missing data patterns to discard parts of the available data for learning. We combine these learning strategies with 4DVarNet schemes, which are state-of-the-art neural interpolation schemes backed on a variational data assimilation formulation. The approach was tested in a turbidity reconstruction context, using a multi-modal satellite dataset (CMEMS product: `oceancolour_med_bgc_l3_my_009_143`) at 1 km spatial resolution with daily images from year 2019 to 2021, off the French coast in the western Mediterranean Sea.

Our learned variational algorithm significantly outperforms state-of-the-art interpolation techniques, including optimal interpolation and DINEOF, with a 37% gain in RMSE reached in preliminary tests.