

First Detection of Hepatitis E Virus (Rocahepevirus ratti Genotype C1) in Synanthropic Norway Rats (Rattus norvegicus) in Romania

Daniela Porea, Cristian Raileanu, Luciana Alexandra Crivei, Vasilica Gotu,

Gheorghe Savuta, Nicole Pavio

▶ To cite this version:

Daniela Porea, Cristian Raileanu, Luciana Alexandra Crivei, Vasilica Gotu, Gheorghe Savuta, et al.. First Detection of Hepatitis E Virus (Rocahepevirus ratti Genotype C1) in Synanthropic Norway Rats (Rattus norvegicus) in Romania. 10th European Meeting on Viral Zoonoses, Sep 2023, Saint-Raphaël, France. hal-04365095

HAL Id: hal-04365095 https://hal.science/hal-04365095

Submitted on 27 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First Detection of Hepatitis E Virus (Rocahepevirus ratti Genotype C1) in Synanthropic Norway Rats (Rattus norvegicus) in Romania

Daniela Porea^{1,2}, Cristian Raileanu¹, Luciana Alexandra Crivei^{1,3}, Vasilica Gotu⁴, Gheorghe Savuta^{1,3} and Nicole Pavio⁵

- 1 Department of Public Health, Faculty of Veterinary Medicine, Iasi University of Life Sciences (IULS), 700490 Iasi, Romania;
- 2 Danube Delta National Institute for Research and Development (DDNI), 820112, Tulcea, Romania;
- 3 Regional Center of Advanced Research for Emerging Diseases, Zoonoses and Food Safety, IULS;
- 4 Faculty of Veterinary Medicine, University of Agronomical Sciences and Veterinary Medicine of Bucharest, 011464 Bucharest, Romania;
- 5 Agence Nationale de Sécurité Sanitaire de L'alimentation de L'environnement et du Travail (ANSES),
- Institut National de Recherche pour L'agriculture L'alimentation et L'environnement (INRAE),

École Nationale Vétérinaire d'Alfort (ENVA), UMR Virology, 94700 Maisons-Alfort, France.

Introduction

Hepatitis E virus (HEV) is an emerging zoonotic pathogen with different viral genera and species reported in a wide range of animals. Rodents, particularly rats, carry the specific genus rat HEV (Rocahepevirus genus, genotype C1) and are exposed occasionally to HEV-3 (Paslahepevirus genus, genotype 3), a zoonotic genotype identified in humans and widely distributed in domestic and feral pigs.

Results

Nine rat liver samples were identified as being positive for rat HEV RNA (17.3%). Positive rats originated from each analyzed county and from four of the seven sites studied, indicating a broad geographical distribution of rat HEV in this species in the study area. All samples tested from other animal species, within the same environment, were negative for HEV RNA (Table 2).



The detection of rat HEV in humans with acute or chronic hepatitis and the worldwide distribution of rat HEV in rodents suggest that rat HEV infection may be an emerging disease. The role of rats in the transmission of the zoonotic HEV-3 genotype, the most commonly detected genotype in humans worldwide, the main animal reservoir of which is domestic pigs, also needs to be clarified. In this study, the presence of HEV was investigated in synanthropic Norway rats (Rattus Norvegicus) from Eastern Romania, in areas where the presence of HEV-3 was previously reported in pigs, wild boars and humans.

Materials & Methods

Using methods capable of detecting different HEV species, the presence of HEV RNA was investigated in 69 liver samples collected from 52 rats and other animal species, originated in different counties in Eastern Romania (Figure 1).



Species	Type of Samples	County						
Wild		Suceava	Iasi	Bacau	Tulcea	Vrancea		
Rattus norvegicus	_ Liver _	3/16 (3/10 ⁴)	1/6 (1/1 ⁵)	2/16 (2/11 ²)	3/14 (3/12 1)	-		
Vulpes vulpes		-	-	-	-	0/9		
Mustela putorius		-	0/1	-	0/3	-		
Mus musculus		-	0/1	-	-	-		
Talpa europaea		0/1	-	-	-	-		
Myocastor coypus		-	-	-	0/1	-		
Ondatra zibethicus		-	-	-	0/1	-		
Domestic								
Sus scrofa domesticus	Pool faecal	-	-	-	0/10	-		

^{1, 2, 4, 5} Numbers corresponding to the different settings indicated in Table 1 (column 1).

High sequence identity (85–89% nt) was found with other European *Rocahepevirus*. The nucleotide identity between our isolates ranged from 85% to 91%, and phylogenetic analysis showed that they were clustered by county. These sequences clustered within the proposed subtype GI, but are more distant from those of Southern Europe such as Spain and may constitute a new European clade or subtype. The analysis of complete genomes would improve this comparison. As rat populations may have followed trade routes, the identification of other sequences from different countries would contribute to drawing a possible map of rat HEV virus evolution and spread.

Figure 1. Geographical distribution of the counties included in the study, with the representation of the analyzed species.

Rats were trapped at various locations near human settlements and animal farms (Table 1). In 2017, 10 pool fecal samples were collected from two indoor pig farms (Sus scrofa domesticus) where rats were trapped in Tulcea and Bacau counties.

Table 1. Data on the samples from rats

NIa		Number of Kats/Location/Counties				
INO.	Location/Site	Suceava	lasi	Bacau	Tulcea	
1	Intensive system pig farm	-	-	-	12	
2	Semi-intensive system pig farm	5	-	11	_	
3	Extensive system pig farm	1	-	5	-	
4	Farm with different animal species, forestry site	10	_	_	-	
5	Cattle farm, metropolitan area	-	1	-	_	
6	Urban settlements, near garbage cans	-	5	-	-	
7	Rural settlements, population's household	-	-	-	2	
Total		16	6	16	14	

. • . •



For HEV RNA detection, three different assays were used:

. A TaqMan real-time RT-PCR assay (Jothikumar et al., 2006; Barnaud et al., 2012), and a RT-nested PCR specific for HEV genotypes 1-4 (Cooper et al, 2005; Bouquet et al., 2011),

. A nested broad-spectrum RT-PCR (NBS-RT-PCR) for hepeviruses, including rat HEV (Johne et al., 2010).

Figure 2. Phylogenetic tree based on the alignment of the generated sequences with the best hits to our sequences in GenBank and with reference sequences for the members of the subfamily Othohepevirinae (the rat HEV strains obtained in our study are highlighted by black diamonds).

Conclusions

The first detection and molecular characterization of rat HEV in Romania represent the novelty of the present study. .The results of the molecular investigations regarding the presence of hepatitis E virus in rats allowed the classification of the isolated strains into the rat HEV

genotype and indicated the presence of a new source of human infections in the study area.

.To better assess the role of rats in the epidemiology of HEV infection in humans, it is recommended to screen for rat HEV in humans with a suspicion of hepatitis. (Porea et al 2023, Viruses)