



# Another rational solutions to the Gardner equation

Pierre Gaillard

## ► To cite this version:

| Pierre Gaillard. Another rational solutions to the Gardner equation. 2024. hal-04369040

HAL Id: hal-04369040

<https://hal.science/hal-04369040>

Preprint submitted on 2 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Another rational solutions to the Gardner equation

Pierre Gaillard,

Université de Bourgogne-Franche Comté,  
Institut de mathématiques de Bourgogne,  
9 avenue Alain Savary BP 47870  
21078 Dijon Cedex, France :  
E-mail : Pierre.Gaillard@u-bourgogne.fr

## Abstract

Rational solutions to the Gardner (G) equation are constructed in terms of a quotient of determinants involving certain particular polynomials. This gives a very efficient method to construct solutions to this equation. We construct very easily explicit expressions of these rational solutions for the first orders for  $n = 1$  until 8.

**Key Words :** Gardner equation, rational solutions.

**PACS numbers :**

33Q55, 37K10, 47.10A-, 47.35.Fg, 47.54.Bd

## 1 Introduction

We consider the Gardner equation in the following normalization

$$u_t + 6u(u - 2)u_x + u_{xxx} + 6u_x = 0, \quad (1)$$

where the subscripts  $x$  and  $t$  stand for partial derivatives.

This equation was introduced by Gardner [1] in 1968. He considered this equation as an auxiliary mathematical tool in the derivation of the infinite set of local conservation laws of the Korteweg de Vries equation. This equation is used to describe nonlinear wave effects in several physical contexts: for example, in plasma physics [2, 3], fluid flows [4], quantum fluid dynamics [5], in ocean and atmosphere [6].

A lot of research has been done to solve this equation. We can quote many methods as the Hirota method [8], the series expansion method [9], the mapping method [10] or the method of leading-order analysis [11].

Here, we used particular polynomials to construct rational solutions to the Gardner equation. We obtain rational solutions as a quotient of polynomials in  $x$  and  $t$ . We give explicit solutions for the first orders.

## 2 Rational solutions to the Gardner equation

We consider the following polynomials  $p_n(x, t)$  defined by

$$\begin{cases} p_n(x, t) = \sum_{k=0}^n \frac{x^k}{k!} \frac{t^{\left(\frac{n-k}{3}\right)}}{\left(\frac{n-k}{3}\right)!} \left(1 - \left[\frac{1}{2}(n-k+1-3\left[\frac{n-k}{3}\right])\right]\right), & k \geq 0, \\ p_n(x, t) = 0, & n < 0, \end{cases} \quad (2)$$

with  $[x]$  denoting the largest integer less or equal to  $x$ .

We denote  $A_n(x, t)$  the determinant defined by

$$A_n(x, t) = \det(p_{n+1-2i+j}(x, t))_{\{1 \leq i \leq n, 1 \leq j \leq n\}} \quad (3)$$

With these notations we have the following result

**Theorem 2.1** *The function  $v_n(x, t)$  defined by*

$$v_n(x, t) = 1 - \partial_X \left( \ln \frac{A_{n+1}(X, T)}{A_n(X, T)} \right)_{|X=ix, T=4it} \quad (4)$$

*is a rational to the Gardner equation (1)*

$$u_t + 6u(u-2)u_x + u_{xxx} + 6u_x = 0,$$

## 3 Explicit construction of rational solutions to the Gardner equation for the first orders

The solutions we present in the following are singular. There is therefore no interest in constructing figures in the plan  $(x, t)$  of the coordinates.

### 3.1 First order rational solution

The function  $v_1(x, t)$  defined by

$$v_1(x, t) = -\frac{-x^4 - 12xt - 2ix^3 + 12it}{x(x^3 + 12t)} \quad (5)$$

is a rational to the Gardner equation (1).

### 3.2 Second order rational solution

The function  $v_2(x, t)$  defined by

$$v_2(x, t) = \frac{-i(-72ix^6t + 8640it^3 - ix^9 + 72x^5t + 4320x^2t^2 + 3x^8)}{(-60tx^3 + 720t^2 - x^6)(x^3 + 12t)} \quad (6)$$

is a rational to the Gardner equation (1).

### 3.3 Third order rational solution

The function  $v_3(x, t)$  defined by

$$v_3(x, t) = \frac{n(x, t)}{d(x, t)} \quad (7)$$

with

$$n(x, t) = -x^{16} - 4ix^{15} - 240tx^{13} - 600ix^{12}t - 10080t^2x^{10} - 36000ix^9t^2 - 172800t^3x^7 + 2419200it^3x^6 - 18144000t^4x^4 + 36288000it^4x^3 + 217728000t^5x + 217728000it^5$$

and

$$d(x, t) = (x^9 + 180tx^6 + 302400t^3)(-x^6 - 60tx^3 + 720t^2)x$$

is a rational to the Gardner equation (1).

### 3.4 Rational solution of order four

The function  $v_4(x, t)$  defined by

$$v_4(x, t) = \frac{n(x, t)}{d(x, t)} \quad (8)$$

with

$$\begin{aligned} n(x, t) = & -i(-ix^{25} + 5x^{24} - 600ix^{22}t + 2280tx^{21} - 100800ix^{19}t^2 + 352800t^2x^{18} \\ & - 6955200ix^{16}t^3 + 4838400t^3x^{15} - 254016000ix^{13}t^4 + 2794176000t^4x^{12} \\ & + 39626496000it^5x^{10} + 259096320000t^5x^9 - 365783040000ix^7t^6 + 5120962560000t^6x^6 + \\ & 76814438400000it^7x^4 - 153628876800000t^7x^3 + 460886630400000it^8x \\ & + 460886630400000t^8) \end{aligned}$$

and

$$d(x, t) = (-x^{15} - 420tx^{12} - 25200t^2x^9 - 2116800t^3x^6 + 254016000t^4x^3 + 1524096000t^5)(x^9 + 180tx^6 + 302400t^3)x$$

is a rational to the Gardner equation (1).

### 3.5 Rational solution of order five

The function  $v_5(x, t)$  defined by

$$v_5(x, t) = \frac{n(x, t)}{d(x, t)} \quad (9)$$

with

$$\begin{aligned}
n(x, t) = & -i(ix^{36} - 6x^{35} + 1260itx^{33} - 6300tx^{32} + 544320it^2x^{30} - 2419200t^2x^{29} + \\
& 110073600it^3x^{27} - 381024000t^3x^{26} + 11430720000it^4x^{24} - 50295168000t^4x^{23} + \\
& 329204736000it^5x^{21} + 192036096000t^5x^{20} + 119062379520000it^6x^{18} + 1371137725440000t^6x^{17} + \\
& 7189831434240000it^7x^{15} + 129970029772800000t^7x^{14} + 1393721170329600000it^8x^{12} - \\
& 116143430860800000t^8x^{11} - 59620294508544000000it^9x^9 + 319394434867200000000t^9x^8 - \\
& 4292661204615168000000it^{10}x^6 + 2146330602307584000000t^{10}x^5 \\
& + 128779836138455040000000t^{11}x^2 + 128779836138455040000000it^{12})
\end{aligned}$$

and

$$d(x, t) = (-x^{21} - 840tx^{18} - 166320t^2x^{15} - 16934400t^3x^{12} + 1397088000t^4x^9 - \\
352066176000t^5x^6 - 14082647040000t^6x^3 + 84495882240000t^7)(-x^{15} - 420tx^{12} - \\
25200t^2x^9 - 2116800t^3x^6 + 254016000t^4x^3 + 1524096000t^5)$$

is a rational to the Gardner equation (1).

### 3.6 Rational solution of order six

The function  $v_6(x, t)$  defined by

$$v_6(x, t) = \frac{n(x, t)}{d(x, t)} \quad (10)$$

with

$$\begin{aligned}
n(x, t) = & -i(ix^{49} - 7x^{48} + 2352itx^{46} - 14448tx^{45} + 2116800it^2x^{43} - 11733120t^2x^{42} + \\
& 979655040it^3x^{40} - 4792435200t^3x^{39} + 259756761600it^4x^{37} - 1190674598400t^4x^{36} + \\
& 40391592192000it^5x^{34} - 148315878144000t^5x^{33} + 4855696699392000it^6x^{31} - \\
& 8843902341120000t^6x^{30} + 61513002270720000it^7x^{28} - 7215948343296000000t^7x^{27} - \\
& 86555891849011200000ix^{25}t^8 - 2329982402356224000000t^8x^{24} \\
& - 29124513867423744000000ix^{22}t^9 - 196587700520008089600000t^9x^{21} \\
& + 125918062002044928000000it^{10}x^{19} - 19426796003252994048000000t^{10}x^{18} - \\
& 26702499023308652544000000ix^{16}t^{11} + 573430854357312602112000000t^{11}x^{15} - \\
& 480928298059060346880000000000ix^{13}t^{12} + 89978213950266259537920000000t^{12}x^{12} - \\
& 13953068593089999903129600000000ix^{10}t^{13} + 6295896804199146297753600000000t^{13}x^9 - \\
& 12251474862225365768601600000000ix^{7}t^{14} - 171520648071155120760422400000000t^{14}x^6 - \\
& 857603240355775603802112000000000ix^{4}t^{15} - 1715206480711551207604224000000000t^{15}x^3 + \\
& 5145619442134653622812672000000000it^{16}x - 5145619442134653622812672000000000t^{16})
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & (-x^{27} - 1512tx^{24} - 680400t^2x^{21} - 139708800t^3x^{18} - 5029516800t^4x^{15} - \\
& 3168595584000t^5x^{12} + 604145558016000t^6x^9 + 108746200442880000t^7x^6 \\
& + 60897872248012800000t^9)(-x^{21} - 840tx^{18} - 166320t^2x^{15} - 16934400t^3x^{12} + \\
& 1397088000t^4x^9 - 352066176000t^5x^6 - 14082647040000t^6x^3 + 84495882240000t^7)x
\end{aligned}$$

is a rational to the Gardner equation (1).

### 3.7 Rational solution of order seven

The function  $v_7(x, t)$  defined by

$$v_7(x, t) = \frac{n(x, t)}{d(x, t)} \quad (11)$$

with

$$\begin{aligned} n(x, t) = & x^{64} + 8ix^{63} + 4032tx^{61} + 29232itx^{60} + 6652800t^2x^{58} + 44331840it^2x^{57} + \\ & 6008325120t^3x^{55} + 36696844800it^3x^{54} + 3329540121600t^4x^{52} + 18820451865600it^4x^{51} + \\ & 1197306651340800t^5x^{49} + 6162707171174400it^5x^{48} + 293930755795353600t^6x^{46} + \\ & 1373247357176217600it^6x^{45} + 46643754897653760000t^7x^{43} + 2604421310052925440000it^7x^{42} + \\ & 5156738132927496192000t^8x^{40} + 7389254129539645440000it^8x^{39} \\ & + 1123297792337944903680000t^9x^{37} - 28613498613600193413120000it^9x^{36} \\ & + 621308836469927647641600000t^{10}x^{34} - 9639939765218166551347200000it^{10}x^{33} + \\ & 96034709039084017916313600000t^{11}x^{31} - 1768604112551541393063936000000it^{11}x^{30} + \\ & 12747529948908855776772096000000t^{12}x^{28} - 103264051547071763510722560000000it^{12}x^{27} + \\ & 3002223914988325881595822080000000t^{13}x^{25} - 3265018050783059977332326400000000it^{13}x^{24} - \\ & 498353242970741203369407283200000000t^{14}x^{22} \\ & - 2066613083889788928871869972480000000it^{14}x^{21} \\ & - 104928442540121429745671444889600000000t^{15}x^{19} \\ & - 464599008554226302534480717414400000000it^{15}x^{18} \\ & - 5587991432946635075558051099443200000000t^{16}x^{16} \\ & + 1181438340409670179520687741337600000000it^{16}x^{15} \\ & - 2113873981271113908649122909388800000000000t^{17}x^{13} \\ & - 853337549281549667333619616579584000000000it^{17}x^{12} \\ & + 12515617389462728454226421043167232000000000t^{18}x^{10} \\ & - 774271245280321336575024352670515200000000000it^{18}x^9 \\ & - 763664789865522414156188402633932800000000000t^{19}x^7 \\ & - 1069130705811731379818663763687505920000000000it^{19}x^6 \\ & + 80184802935879853486399782276562944000000000000t^{20}x^4 \\ & + 160369605871759706972799564553125888000000000000it^{20}x^3 \\ & + 320739211743519413945599129106251776000000000000t^{21}x \\ & - 320739211743519413945599129106251776000000000000it^{21} \end{aligned}$$

and

$$\begin{aligned} d(x, t) = & (-x^{36} - 2520tx^{33} - 2162160t^2x^{30} - 884822400t^3x^{27} - 163459296000t^4x^{24} - \\ & 30207277900800t^5x^{21} + 4833164464128000t^6x^{18} - 891718843631616000t^7x^{15} - \\ & 831255956185374720000t^8x^{12} - 34833582925863321600000t^9x^9 \\ & - 2194515724329389260800000t^{10}x^6 + 131670943459763355648000000t^{11}x^3 + \\ & 526683773839053422592000000t^{12})(-x^{27} - 1512tx^{24} - 680400t^2x^{21} - 139708800t^3x^{18} - \\ & 5029516800t^4x^{15} - 3168595584000t^5x^{12} + 604145558016000t^6x^9 + 108746200442880000t^7x^6 + \\ & 60897872248012800000t^9)x \end{aligned}$$

is a rational to the Gardner equation (1).

### 3.8 Rational solution of order eight

The function  $v_8(x, t)$  defined by

$$v_8(x, t) = \frac{n(x, t)}{d(x, t)} \quad (12)$$

with

$$\begin{aligned} n(x, t) = & -i(-ix^{81} + 9x^{80} - 6480itx^{78} + 54000tx^{77} - 17962560it^2x^{75} + 139708800t^2x^{74} - \\ & 284407200000it^3x^{72} + 206689190400t^3x^{71} - 28894574016000it^4x^{69} + 196729549632000t^4x^{68} - \\ & 20027877904742400it^5x^{66} + 127548405221529600t^5x^{65} - 9845156013428736000it^6x^{63} + \\ & 58650450772193280000t^6x^{62} - 3496233842718769152000it^7x^{60} + 19742220213202206720000t^7x^{59} - \\ & 917522577057504337920000it^8x^{57} + 45842043443808400588800000t^8x^{56} \\ & - 186047506160225456947200000it^9x^{54} + 705970885405282482585600000t^9x^{53} - \\ & 259331566184437918806835200000it^{10}x^{51} + 3793368402563618404682956800000t^{10}x^{50} + \\ & 6395652736671085473890304000000it^{11}x^{48} + 297650067947402651243642880000000t^{11}x^{47} + \\ & 5207428862069036677958467584000000it^{12}x^{45} + 129905635334268499051254644736000000t^{12}x^{44} + \\ & 1702586892784983334684867952640000000it^{13}x^{42} \\ & + 296206395405993443048000197754880000000t^{13}x^{41} \\ & + 396994691532759366938073508085760000000it^{14}x^{39} \\ & + 42323605252136969901466129426022400000000t^{14}x^{38} \\ & - 4349155069043839525357532208955392000000it^{15}x^{36} \\ & + 583120649740812817456115098224427008000000t^{15}x^{35} \\ & - 202466214174573975549688586510008320000000it^{16}x^{33} \\ & + 28159381522332939098979530493552230400000000t^{16}x^{32} \\ & + 5518612694466199647008064688942649180160000000it^{17}x^{30} \\ & - 31710925844235863073697671356573181542400000000t^{17}x^{29} \\ & + 1253795797863202772049566358144828624076800000000it^{18}x^{27} \\ & - 3779312133537331858253773128621109542912000000000t^{18}x^{26} \\ & + 127205858675790678540997355173101824901120000000000it^{19}x^{24} \\ & - 454698326162344677609910772685294048116736000000000it^{19}x^{23} \\ & + 1234128242496442531104836094950737862197248000000000it^{20}x^{21} \\ & - 28547550061967276275702055938476943173746688000000000it^{20}x^{20} \\ & + 446177931156674484955283193242186295080386560000000000it^{21}x^{18} \\ & + 3908572835159623913132810908575137703584071680000000000it^{21}x^{17} \\ & + 13662139312759241693421665342506948154118635520000000000it^{22}x^{15} \\ & + 370790999888692634519688589966263523867203993600000000000it^{22}x^{14} \\ & + 2648353159088714543647892051009039180644535500800000000000it^{23}x^{12} \\ & - 1471307310604841413137717806116132878135853056000000000000it^{23}x^{11} \\ & - 424839985937147958043516016516033368561727569920000000000000it^{24}x^9 \\ & + 333802846093473395605619727262597646727071662080000000000000it^{24}x^8 \\ & - 3058847898747465297913315318915440253644438503424000000000000it^{25}x^6 \\ & + 1019615966249155099304438439638480084548146167808000000000000it^{25}x^5 \\ & + 61176957974949305958266306378308805072888770068480000000000000it^{26}x^2 \\ & + 407846386499662039721775375539203381925846712320000000000000it^{27}) \end{aligned}$$

and

$$d(x, t) = (x^{45} + 3960tx^{42} + 5821200t^2x^{39} + 4324320000t^3x^{36} + 1743565824000t^4x^{33} +$$

$$\begin{aligned}
& 455986052121600 t^5 x^{30} + 33630769396224000 t^6 x^{27} + 18378107874846720000 t^7 x^{24} + \\
& 3172344159319695360000 t^8 x^{21} - 6261336530923932057600000 t^9 x^{18} \\
& - 1088291697919577411420160000 t^{10} x^{15} - 98937546915666185433907200000 t^{11} x^{12} + \\
& 512147301681095481284608000000 t^{12} x^9 - 645305600118180390641860608000000 t^{13} x^6 - \\
& 19359168003545411719255818240000000 t^{14} x^3 + 77436672014181646877023272960000000 t^{15}) \\
& (-x^{36} - 2520 t x^{33} - 2162160 t^2 x^{30} - 884822400 t^3 x^{27} - 163459296000 t^4 x^{24} - \\
& 30207277900800 t^5 x^{21} + 4833164464128000 t^6 x^{18} - 891718843631616000 t^7 x^{15} - \\
& 831255956185374720000 t^8 x^{12} - 34833582925863321600000 t^9 x^9 \\
& - 2194515724329389260800000 t^{10} x^6 + 131670943459763355648000000 t^{11} x^3 + \\
& 526683773839053422592000000 t^{12})
\end{aligned}$$

is a rational to the Gardner equation (1).

### 3.9 Rational solution of order nine

The function  $v_9(x, t)$  defined by

$$v_9(x, t) = \frac{n(x, t)}{d(x, t)} \quad (13)$$

with

$$\begin{aligned}
n(x, t) = & -i(-ix^{100} + 10x^{99} - 9900ix^{97}t + 93060tx^{96} - 43243200ix^{94}t^2 + 383961600t^2x^{93} - \\
& 111198700800ix^{91}t^3 + 934226092800t^3x^{90} - 189254729664000ix^{88}t^4 + 1506596547456000t^4x^{87} - \\
& 227051874137088000ix^{85}t^5 + 1713219123103488000t^5x^{84} - 1996525579342694400000ix^{82}t^6 + \\
& 1428145956200291328000t^6x^{81} - 1319595993742189363200000ix^{79}t^7 \\
& + 895581632952719032320000t^7x^{78} - 667603033565633490124800000ix^{76}t^8 \\
& + 428022377616371217039360000t^8x^{75} - 26201771314573067845632000000ix^{73}t^9 + \\
& 158789776023912561028300800000t^9x^{72} - 79820511350954286201569280000000ix^{70}t^{10} + \\
& 48867778365532304849672601600000t^{10}x^{69} - 1845794510242697463355146240000000ix^{67}t^{11} + \\
& 10500563590228982713863477657600000t^{11}x^{66} - 4369386748577714555583890718720000000ix^{64}t^{12} - \\
& 3509125889298278795399059734528000000t^{12}x^{63} \\
& - 2297677505784692695552786044026880000000ix^{61}t^{13} \\
& - 61533493572432454507107017860055040000000t^{13}x^{60} \\
& - 1449815926554684377327575785642393600000000ix^{58}t^{14} \\
& - 3665293540829092709903367569229742080000000t^{14}x^{57} \\
& - 65471826215870206689389630766270382080000000ix^{55}t^{15} \\
& - 13042434125415100832957182218387377356800000000t^{15}x^{54} \\
& - 159482048483172697860011723263699845120000000000ix^{52}t^{16} \\
& - 321470231876680131350631334233741176340480000000t^{16}x^{51} \\
& - 32634096279242127691173671142756597104640000000000ix^{49}t^{17} \\
& - 502266661319331641970710360305345033666560000000000t^{17}x^{48} \\
& - 11090341091536021381954548300183335243612160000000000ix^{46}t^{18} \\
& - 1648734229738630070957192326433181511188480000000000t^{18}x^{45} \\
& + 27768030541570534401038610181385410880471040000000000it^{19}x^{43} \\
& - 1744391947969980802122124052001220299533058048000000000t^{19}x^{42} \\
& + 176353619079396164221683485553881739201481801728000000000it^{20}x^{40} \\
& - 984738559457980653135258698718009747572885815296000000000t^{20}x^{39}
\end{aligned}$$

and

$$\begin{aligned}
d(x,t) = & (-x^{54} - 5940tx^{51} - 13899600t^2x^{48} - 17254036800t^3x^{45} - 12586365792000t^4x^{42} - \\
& 5851581261926400t^5x^{39} - 1623511727405568000t^6x^{36} - 400077271429355520000t^7x^{33} - \\
& 28551097433877258240000t^8x^{30} + 50926698237612176179200000t^9x^{27} \\
& - 42318665025272343181393920000t^{10}x^{24} - 24094202601815176923316224000000t^{11}x^{21} - \\
& 4866679734224610446090698752000000t^{12}x^{18} - 89328732359216685504566132736000000t^{13}x^{15} - \\
& 56277101386306511867876663623680000000t^{14}x^{12} \\
& + 4292066932395643305123393545699328000000t^{15}x^9 \\
& + 772572047831215794922210838225879040000000t^{16}x^6 \\
& + 216320173392740422578219034703246131200000000t^{18})(x^{45} + 3960tx^{42} + 5821200t^2x^{39} + \\
& 4324320000t^3x^{36} + 1743565824000t^4x^{33} + 455986052121600t^5x^{30} + 33630769396224000t^6x^{27} + \\
& 18378107874846720000t^7x^{24} + 3172344159319695360000t^8x^{21} - 6261336530923932057600000t^9x^{18} - \\
& 1088291697919577411420160000t^{10}x^{15} - 98937546915666185433907200000t^{11}x^{12} + \\
& 5121473016810955481284608000000t^{12}x^9 - 645305600118180390641860608000000t^{13}x^6 - \\
& 19359168003545411719255818240000000t^{14}x^3 + 7743667201418164687702327296000000t^{15})x
\end{aligned}$$

is a rational to the Gardner equation (1).

### 3.10 Rational solution of order ten

The function  $v_{10}(x, t)$  defined by

$$v_{10}(x, t) = \frac{n(x, t)}{d(x, t)} \quad (14)$$

with

$$\begin{aligned} n(x, t) = & -i(-ix^{121} + 11x^{120} - 14520ix^{118}t + 151800tx^{117} - 95135040ix^{115}t^2 + 948261600t^2x^{114} - \\ & 375171825600ix^{112}t^3 + 3570850483200t^3x^{111} - 1001486566080000ix^{109}t^4 + 9111786780096000t^4x^{108} - \\ & 1928677978260633600ix^{106}t^5 + 16783754184791193600t^5x^{105} - 2788145214059132928000ix^{103}t^6 + \\ & 23214265448682147840000t^6x^{102} - 3107387740090288951296000ix^{100}t^7 + 24758895105773203415040000t^7x^{99} - \\ & 2720957040436266106675200000ix^{97}t^8 + 20739568173400178020638720000t^8x^{96} \\ & - 1897998004600345592443699200000ix^{94}t^9 + 13838512826996381618660966400000t^9x^{93} \\ & - 1065031427670194137372455075840000ix^{91}t^{10} + 7441832235420948554463447613440000t^{10}x^{90} - \\ & 484359826252919742513781250457600000ix^{88}t^{11} \\ & + 3216285223633520093028577443840000000t^{11}x^{87} \\ & - 180973477671370782253997564834611200000ix^{85}t^{12} \\ & + 1094007551086362182820948800962560000000t^{12}x^{84} \\ & - 55662624659826717571311231389663232000000ix^{82}t^{13} \\ & + 366409302919024594307060726369353728000000t^{13}x^{81} \\ & - 106292105088373711675989148044990873600000000ix^{79}t^{14} \\ & + 2877196120826052801365210508052502937600000000t^{14}x^{78} \\ & + 4280030807361509169541614389644146769920000000it^{15}x^{76} \\ & + 304538976982963877702699431050813489807360000000t^{15}x^{75} \\ & + 6885670109355945389014278608796839313408000000000it^{16}x^{73} \\ & + 228041011956866466611552032321683792592896000000000t^{16}x^{72} \\ & + 4453721520274708640987473009279000243077120000000000it^{17}x^{70} \\ & + 117762805364914466257694778298876385933918208000000000t^{17}x^{69} \\ & + 1928767694961190051291693412300125238030499840000000000it^{18}x^{67} \\ & + 42939910195222368442373794245841509282389950464000000000t^{18}x^{66} \\ & + 654735071078952292481291250564426986112984023040000000000it^{19}x^{64} \\ & + 11100640725265289005976333178685141575591650131968000000000t^{19}x^{63} \\ & + 11898388510029268300683286437262873928059593739468000000000it^{20}x^{61} \\ & + 2548058689427082387186572761563839262072167014072320000000000t^{20}x^{60} \\ & - 7153367620935353165234889584513031133610669475102720000000000ix^{58}t^{21} \\ & + 501342812165200991524948146064415704363708955523809280000000000t^{21}x^{57} \\ & + 1122098111161432183874791372551241736238612859961999360000000000it^{22}x^{55} \\ & - 5328063166643583956850588870472355563633395222917939200000000000t^{22}x^{54} \\ & + 18666731499151103582697435263135280572769467169281723596800000000000it^{23}x^{52} \\ & - 11190072052434997260774808915853970834697517783347888128000000000000t^{23}x^{51} \\ & + 8211604920987036524942436185081467249913411096884077920256000000000000it^{24}x^{49} \\ & - 472356267985166166429107110009412445346002202412648836890624000000000000t^{24}x^{48} \\ & + 2074576890338308833679718042344039076057947005326331519455723520000000000it^{25}x^{46} \\ & - 88712810537698606548709845510973812347610188041362448225350451200000000000t^{25}x^{45} \\ & + 310982628871002671709704390044044172790555050597378687653799526400000000000it^{26}x^{43} \\ & - 1310226464648804607175637470426054827417012493833765156242325504000000000000t^{26}x^{42} \end{aligned}$$



$$56277101386306511867876663623680000000 t^{14}x^{12} + 4292066932395643305123393545699328000000 t^{15}x^9 \\ + 772572047831215794922210838225879040000000 t^{16}x^6 + 216320173392740422578219034703246131200000000 t^{18})x$$

is a rational to the Gardner equation (1).

We could go on for greater orders but the expressions of the solutions become too long to be given in this text.

## 4 Conclusion

We have constructed rational solutions to the Gardner equation and we obtain an infinite hierarchy of families of rational solutions of this equation as a quotient of a polynomials in  $x$  and  $t$ .

At order  $k$  the numerator is a polynomial of degree  $(k+1)^2$  en  $x$  and of degree  $\left[\frac{(k+1)^2}{3}\right]$  in  $t$ , where as usual,  $[x]$  denotes the bigger integer less or equal to  $x$ .

The denominator is also a polynomial of degree  $(k+1)^2$  en  $x$  and of degree  $\left[\frac{(k+1)^2}{3}\right]$  in  $t$  at order  $k$ , with the same notation as previously.

It will relevant to study in details the structure of these polynomials.

## References

- [1] R.M. Miura, C.S. Gardner and M.D. Kruskal, Korteweg-deVries equation and generalizations. II. Existence of conservation laws and constants of motion,, J. Math. Phys., V. 9, 1204-1209, 1968
- [2] S. Watanabe, Ion Acoustic Soliton in Plasma with Negative Ion, J. Phys. Soc. Japan, V. 53, 950-956, 1984
- [3] M.S. Ruderman, T. Talipova, and E. Pelinovsky, Dynamics of modulationally unstable ion-acoustic wavepackets in plasmas with negative ions, J. Plasma Phys., V. 74, 639-656, 2008
- [4] R. Grimshaw, Environmental Stratified Flows, Topics in Environmental Fluid Mechanics, Kluwer, 2002
- [5] E. Demler and A. Maltsev, Semiclassical solitons in strongly correlated systems of ultracold bosonic atoms in optical lattices, Ann. Phys., V. 326, 1775-1805, 2011
- [6] R. Grimshaw, D. Pelinovsky, E. Pelinovsky, T. Talipova, Wave Group Dynamics in Weakly Nonlinear Long-Wave Models, Phys. D, V. 159, 35-37, 2001

- [7] M.H. Alford, T. Peacock, T.Y. Tang, The formation and fate of internal waves in the South China Sea, *Nature*, V 521, 65-69, 2015
- [8] M. Wadati, Wave propagation in nonlinear lattice III, *J. Phys. Soc. Jpn.*, V. 38, 681-686, 1975
- [9] M. Coffey On series expansions giving closed-form solutions of Korteweg-de Vries-like equations, *J. Appl. Math.*, V. 50, N. 6, 1580-1592, 1990
- [10] S.Y. Lou, L.L Chen, Solitary wave solutions and cnoidal wave solutions to the combined KdV and mKdV equation, *Math Meth. Appl. Sci.*, V. 17, 339-347, 1994
- [11] J Zhang, New Solitary Wave Solution of the Combined KdV and mKdV Equation, *Int. Jour. Of Theo. Phys.*, V. 37, 1541-1546, 1998
- [12] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. **161**, 2002
- [13] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. **68**, 77-90, 2004
- [14] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, 2009
- [15] P. Gaillard, V.B. Matveev, Wronskian and Casorati determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 404409-1-16, 2009
- [16] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. **185**, 247-258, 2010
- [17] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. **44**, 435204-1-15, 2011
- [18] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. **13**, N. 2, 71-153, 2012
- [19] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. **54**, 013504-1-32, 2013
- [20] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, 2013
- [21] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, 2013

- [22] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, 2013
- [23] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, 2013
- [24] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013
- [25] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013
- [26] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [27] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [28] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014
- [29] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [30] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [31] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [32] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [33] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, 2015
- [34] P. Gaillard, Other 2N-2 parameters solutions to the NLS equation and 2N+1 highest amplitude of the modulus of the N-th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [35] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, N. **5**, 346-364, 2015

- [36] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, Jour. Phys. : Conf. Ser., V. **633**, 012106-1-6, 2016
- [37] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, Journal of Theoretical and Applied Physics, V. **10**, 1-7, 2016
- [38] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, Commun. Theor. Phys, V. **65**, N. **2**, 136-144, 2016
- [39] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, Annals Of Physics, V. **367**, 1-5, 2016
- [40] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, Adv. Res., V. **10**, 83-89, 2016
- [41] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, Theor. And Math. Phys., V. **189**, N. **1**, 1440-1449, 2016
- [42] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, Jour. of Math. Phys., V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [43] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, Jour. Of Bas. And Appl. Res. Int., V. **21**, N. 3, 130-139, 2017
- [44] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on  $2N - 2$  parameters, Int. Jour. of Appl. Sci. And Math., V. **4**, N. 3, 60-70, 2017
- [45] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, New Hor. in Math. Phys., V. 1, N. 1, 26-31, 2017
- [46] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, Jour. Of Bas. And Appl. Res. Int., V. **21**, N. 2, 92-98, 2017
- [47] P. Gaillard,  $N$ -Order rational solutions to the Johnson equation depending on  $2N - 2$  parameters, Int. Jour. of Adv. Res. in Phys. Sci., V. **4**, N. 9, 19-37, 2017
- [48] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, Int. Jour. of Adv. Res. in Phys. Sci., V. **4**, N. 11, 24-30, 2017
- [49] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, Int. Jour. of Inn. In Sci. and Math., V. **6**, N. 1, 14-19, 2018

- [50] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. **196**, N. 2, 1174-1199, 2018
- [51] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. **2018**, 1-18, 2018
- [52] P. Gaillard, Families of Solutions of Order 5 to the Johnson Equation Depending on 8 Parameters, *NHIMP*, V. **2**, N. 4, 53-61, 2018
- [53] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. **1141**, 012102-1-10, 2018
- [54] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. , 109-112, 2019
- [55] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. **2**, N. 4, 1-4, 2019
- [56] P. Gaillard, Multi-parametric families of solutions of order N to the Boussinesq and KP equations and the degenerate rational case, *UJMA*, V. **3**, N. 2, 44-52, 2020
- [57] P. Gaillard, The mKdV equation and multi-parameters rational solutions, *Wave Motion*, V. **100**, 102667-1-9, 2021
- [58] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, *Jour. Of Geom. And Phys.*, V. **161**, 104059-1-12, 2021
- [59] P. Gaillard, 2N parameter solutions to the Burgers equation, *Jour. Of Appl. Nonlin. Dyn.*, V. **1**, N. **1**, 1-6, 2021
- [60] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. **3**, N. 3, 14-21, 2021
- [61] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. **11**, N. 4, 2102-2113, 2021
- [62] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. **17**, N. 6, 27-34, 2021
- [63] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. **107**, 102824-1-11, 2021
- [64] P. Gaillard, Multiparametric rational solutions of order N to the KPI equation and the explicit case of order 3, *Arch. Of cur. Res. Int.*, V. **21**, N . 6, 58-71, 2021

- [65] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. **108**, 102828-1-9, 2022
- [66] P. Gaillard, Rogue waves of the Lakshmanan Porsezian Daniel Equation Depending on Multi-parameters, *As. Jour. Of Adv. Res. And Rep.*, V 16, N. 3, 32-40, 2022
- [67] P. Gaillard, Quasi-rational and rational solutions to the defocusing NLS equation, *Jour. Of Math. Sci. And Mod. Res. And Rep.*, V 5, N. 1, 24-34, 2022
- [68] P. Gaillard, Rational solutions to the KdV equation in terms of particular polynomials, *Jour. Of Nonlin. Mod. and Ana.*, V. 4, N. 4, 615-626, 2022
- [69] P. Gaillard, Rational solutions to the Painlev II equation from particular polynomials, *Int. J. Of Appl. Nonlin. Sci.*, V. 3, N. 3, 189-196, 2022
- [70] P. Gaillard, Multi-lump solutions to the KPI equation with a zero degree of derivation, *Jour. Of Theor. And Appl. Phys.*, V. 16, N. 4, 1-8, 2022
- [71] P. Gaillard, Families of solutions to the KPI equation given by an extended Darboux transformation, *Part. Diff. Equ. And Appl.*, V. 3, N. 80, 1-14, 2022
- [72] P. Gaillard, Rogue waves of the Hirota equation in terms of quasi-rational solutions depending on multi-parameters, *WSEAS Trans. On Math.*, DOI: 10.37394/23206.2023.22.24, V. 22, 190-203, 2023
- [73] P. Gaillard, Different representations of the solutions to the cylindrical nonlinear Schrödinger equation, *Dis. Nonlin. And Compl.*, V. 12, N. 3, 539-553, 2023
- [74] P. Gaillard, Rational solutions to the KdV equation from Riemann theta functions, *Jour. Of Theor. And Appl. Phys.*, V. 17, N. 3, 1-5, 2023
- [75] P. Gaillard, On Hadamard's Coefficients for Two Dimensional Schrödinger Operators, *Int. Jour. Of Theor. Phys.*, V. 62, N. 205, 1-16, 2023
- [76] P. Gaillard, Rational solutions to the Burgers' equation from particular polynomials, *WSEAS Trans. On Math*, DOI: 10.37394/23206.2023.22.94, 867-873, 2023