



**HAL**  
open science

# Closure Results for Arbitrarily Partitionable Graphs

Julien Bensmail

► **To cite this version:**

Julien Bensmail. Closure Results for Arbitrarily Partitionable Graphs. *Opuscula Mathematica*, 2024, 44 (6), pp.773-788. hal-04370225v2

**HAL Id: hal-04370225**

**<https://hal.science/hal-04370225v2>**

Submitted on 29 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Closure Results for Arbitrarily Partitionable Graphs

Julien Bensmail<sup>a</sup>

<sup>a</sup>Université Côte d'Azur, CNRS, Inria, I3S, France

---

## Abstract

A well-known result of Bondy and Chvátal establishes that a graph of order  $n$  is Hamiltonian if and only if its  $n$ -closure (obtained through repeatedly adding an edge joining any two non-adjacent vertices with degree sum at least  $n$ ) also is. In this work, we investigate such closure results for arbitrarily partitionable graphs, a weakening of Hamiltonian graphs being those graphs that can be partitioned into arbitrarily many connected graphs of arbitrary orders. Among other results, we establish closure results for arbitrary partitions into connected graphs of order at most 3, for arbitrary partitions into connected graphs of order exactly any  $\lambda$ , and for the property of being arbitrarily partitionable in full.

*Keywords:* connected partition; arbitrarily partitionable graph; closure; traceability.

---

## 1. Introduction

Let  $n \geq 1$  be a positive integer. An  $n$ -graph  $G$  is a graph of order  $n$ , while an  $n$ -partition  $\pi = (\lambda_1, \dots, \lambda_p)$  is a partition of  $n$  (i.e.,  $\lambda_1 + \dots + \lambda_p = n$ ). A partition  $(S_1, \dots, S_p)$  of the vertex set  $V(G)$  of  $G$  is called a *realisation* of  $\pi$  in  $G$  if each part  $S_i$  is *connected* and has cardinality  $\lambda_i$ , that is, if, for every  $i \in \{1, \dots, p\}$ , the graph  $G[S_i]$  is a connected graph of order  $\lambda_i$ . We now say  $G$  is *arbitrarily partitionable* (AP) if every  $n$ -partition is realisable in  $G$ , or, in other words, if  $G$  can be partitioned into arbitrarily many connected graphs with arbitrary orders. All these notions have been introduced and considered independently by Barth, Baudon, and Puech in [1], and by Horňák and Woźniak in [13], in connection, in particular, with a practical network sharing problem. Since then, quite some interesting aspects of AP graphs have been introduced and investigated in literature, see e.g. the latest references [3, 4, 5, 6, 8, 10, 15] on the topic for further information.

One of the main sources of motivation behind the investigations in the current work, is the fact that APness can be perceived both as a strengthening of *perfect matchings* (sets of  $\lfloor n/2 \rfloor$  pairwise disjoint edges<sup>1</sup>) and a weakening of *Hamiltonian paths* (paths going through all  $n$  vertices exactly once). Indeed, note that an AP  $n$ -graph, depending on the parity of  $n$ , admits realisations of  $(1, 2, \dots, 2)$  or  $(2, \dots, 2)$ , which form perfect matchings; and that every path is obviously AP, from which we get that any traceable graph, i.e., any graph admitting a Hamiltonian path, is AP. From this, interesting questions on AP graphs arise when wondering how classical results on perfect matchings and Hamiltonian paths adapt to AP graphs. This line of research was initiated by Marczyk, who, in [16], proved:

**Theorem 1.1** (Marczyk [16]). *Let  $G$  be a connected  $n$ -graph in which the degree sum of any two non-adjacent vertices is at least  $n - 2$ . Then  $G$  is AP if and only if  $G$  admits a perfect matching.*

---

<sup>1</sup>For convenience, unless stated otherwise, the order of any graph is denoted by  $n$  throughout.

30 Clearly, Theorem 1.1 stands as a weakening, to APness, of the well-known sufficient  
 31 condition by Ore for graphs to admit Hamiltonian paths or cycles (stating that any con-  
 32 nected  $n$ -graph in which the degree sum parameter is at least  $n$  or  $n-1$  admits Hamiltonian  
 33 cycles or paths, respectively [18]). A remarkable fact also, is that the statement of Theo-  
 34 rem 1.1 involves the two notions (perfect matchings and Hamiltonian paths) between which  
 35 APness is comprised; although these two notions are rather distant in general, this shows  
 36 there are stronger connections between the two in the context of AP graphs.

37 Marczyk’s Theorem 1.1 opened the way to an interesting line of research on AP graphs,  
 38 being to investigate how well-known sufficient conditions for Hamiltonicity or traceability  
 39 weaken to APness. In the very line of Theorem 1.1, better results involving the degree sum  
 40 of pairs of non-adjacent vertices were established in [12, 17], and such results for triples of  
 41 pairwise non-adjacent vertices were considered in [3]. In [14], sufficient conditions in terms  
 42 of number of edges have also been established, while toughness properties of AP graphs  
 43 have been studied in [5]. Last, in [6], the authors considered several other classical concepts  
 44 and results for Hamiltonicity and traceability, such as forbidden induced patterns and the  
 45 square operation, and proved that some adapt to APness while some others do not.

46 In the current work, we pursue this line of research by considering yet another classical  
 47 aspect borrowed from the study of Hamiltonian and traceable graphs, being that of **graph**  
 48 **closures**. Recall that, for a graph  $G$  and some  $k \geq 1$ , the  $k$ -closure  $\text{clos}(G, k)$  of  $G$  is  
 49 obtained upon repeatedly adding an edge  $uv$  between two non-adjacent vertices  $u$  and  $v$   
 50 satisfying  $d(u) + d(v) \geq k$ . Equivalently, note that the  $k$ -closure  $\text{clos}(G, k)$  of  $G$  is obtained  
 51 through a  $k$ -closing sequence  $G_0, \dots, G_m$ , being a sequence of  $n$ -graphs where  $G_0 = G$  and  
 52  $G_m = \text{clos}(G, k)$ , and, for every  $i \in \{1, \dots, m\}$ , the graph  $G_i$  is obtained from  $G_{i-1}$  by  
 53 adding a single edge  $uv$  such that  $d_{G_{i-1}}(u) + d_{G_{i-1}}(v) \geq k$ .

54 Graph closures emerged in a seminal work [7] of Bondy and Chvátal, in which they pro-  
 55 vided the celebrated Bondy-Chvátal sufficient condition for Hamiltonicity and traceability:

56 **Theorem 1.2** (Bondy, Chvátal [7]). *An  $n$ -graph is Hamiltonian if and only if its  $n$ -closure*  
 57 *is Hamiltonian. Likewise, an  $n$ -graph is traceable if and only if its  $(n-1)$ -closure is traceable*

58 Among other remarkable aspects of interest, Bondy and Chvátal’s Theorem 1.2 is im-  
 59 portant in that it implies several other classical sufficient conditions for Hamiltonicity and  
 60 traceability, such as Dirac’s conditions [11] and Ore’s conditions [18]. It is also worth men-  
 61 tioning that the concept of graph closure does not restrict to Hamiltonian cycles and paths  
 62 only, and can also be employed to express sufficient conditions for graphs to admit other  
 63 types of objects. For more details on this point, and more generally on anything related  
 64 to graph closures, we refer the interested reader to the survey [9].

65 Our main intent in the current work is to initiate the study of graph closures in the  
 66 very context of AP graphs. In particular, as explained earlier when introducing Marczyk’s  
 67 Theorem 1.1, given any sufficient condition for Hamiltonicity and traceability, it is natural  
 68 to wonder whether it weakens to APness. Bondy and Chvátal’s Theorem 1.2 being one of  
 69 the most influential results on graph closures, it is thus legitimate to wonder how it adapts  
 70 (or not) to APness. As a first step, we consider  $(n-2)$ -closures in Section 2, showing  
 71 that, for APness, we cannot just weaken Theorem 1.2 to  $(n-2)$ -closures. Our arguments  
 72 lead us to considering arbitrary partitions into connected graphs of order at most 3 in  
 73 Section 3, for which we prove the tight result that it is necessary and sufficient to consider  
 74  $(n-1)$ -closures. Then, in Section 4, we consider weaker closures, and establish more general  
 75 results for partitions into connected graphs of order any fixed  $\lambda$ , and for the AP property  
 76 in full. We finish off in Section 5 with discussions for further work on the topic.

77 **2. On the APness of  $(n - 2)$ -closures**

78 Given how sufficient conditions on degree sums of pairs of non-adjacent vertices essen-  
 79 tially weaken from  $n$  for Hamiltonicity to  $n - 1$  for traceability (Ore's Theorem) and to  
 80  $n - 2$  for APness (Marczyk's Theorem 1.1), a first, legitimate question is whether Bondy  
 81 and Chvátal's Theorem 1.2 weakens the same way to APness, or, in other words, whether  
 82  $n$ -graphs are AP if and only if their  $(n - 2)$ -closure is. Likewise, just like how Bondy and  
 83 Chvátal's Theorem 1.2 implies Ore's Theorem, one can also wonder whether there is some  
 84 result on APness and  $(n - 2)$ -closures that would imply Marczyk's Theorem 1.1.

85 Previous studies on AP graphs have highlighted that, when realising an  $n$ -partition  
 86  $\pi$  in an  $n$ -graph, an important parameter to take into account is the *spectrum*  $\text{sp}(\pi)$   
 87 of  $\pi$ , which is the set of all values appearing in  $\pi$  (thus,  $\text{sp}(\pi)$  is essentially  $\pi$  with all  
 88 duplicates removed). Perhaps one of the most significant results on partition spectra is one  
 89 by Ravoux stating that, in graphs with large diameter, the AP property relies solely on the  
 90 realisability of partitions with small spectrum [20]. More generally speaking, as described  
 91 in [5], something we learn from previous works on AP graphs is that, 1) the smaller the  
 92 spectrum of a partition  $\pi$  is, and 2) the smaller the values of  $\pi$  are, the less chances there  
 93 are that  $\pi$  is realisable in a given graph. In other words, to establish that some graphs with  
 94 particular properties are not necessarily AP, very generally speaking one should consider  
 95 the realisability of partitions with small values and low variety of values.

96 From these thoughts, regarding the hopes we have exposed for  $(n - 2)$ -closures earlier,  
 97 we get to the matter of establishing whether, in general,  $n$ -partitions with small spectrum  
 98 are realisable if and only if they are in the  $(n - 2)$ -closure. As a starting point, it thus  
 99 makes sense to wonder about perfect matchings<sup>2</sup>, or, in other words, about realisations of  
 100 partitions with spectrum  $\{2\}$ . Below, we prove that, indeed, already for this particular  
 101 type of partitions we cannot weaken Bondy and Chvátal's Theorem 1.2 to value  $n - 2$  as  
 102 is. For transparency, let us mention that upcoming Theorem 2.3 is actually a particular  
 103 case of a previous result of Plummer and Saito [19]; still, we provide a straight, thorough  
 104 proof, as our arguments stand as a good introduction to the ones to be used later on.

105 To begin with, let us remind Tutte's condition for the existence of perfect matchings.

106 **Theorem 2.1** (Tutte [21]). *A graph  $G$  with even order has a perfect matching if and only*  
 107 *if for every  $S \subseteq V(G)$  the graph  $G - S$  has at most  $|S|$  connected components with odd order.*

108 We now proceed with our first result, Theorem 2.3. Before that, we just need to  
 109 introduce an auxiliary result that will show Theorem 2.3 is indeed the best we can hope  
 110 for. Recall that for any two positive integers  $p, q \geq 1$ , we denote by  $K_{p,q}$  the complete  
 111 bipartite graph with partition classes of cardinality  $p$  and  $q$ .

112 **Lemma 2.2.** *For any  $p \geq 1$ ,  $\text{clos}(K_{p,p+2}, n - 2)$  has perfect matchings while  $K_{p,p+2}$  has not.*

113 *Proof.* Set  $G = K_{p,p+2}$  for some  $p \geq 1$ , and denote by  $U$  the partition class of  $G$  with  
 114 cardinality  $p$ , and by  $V$  that with cardinality  $p + 2$ . Note that all vertices of  $U$  have degree  
 115  $|V| = p + 2$ , while all vertices of  $V$  have degree  $|U| = p$ . Set  $n = |V(G)| = 2p + 2$ .

116 First off, note that  $G - U$  consists of  $p + 2$  connected components, all of which have odd  
 117 order, 1. Meanwhile,  $|U| = p$ . Thus, by Tutte's Theorem,  $G$  admits no perfect matchings.

118 Now consider  $G' = \text{clos}(G, n - 2)$ . Note that, in  $G$ , any two non-adjacent vertices  $u$  and  
 119  $v$  either both belong to  $U$ , or both belong to  $V$ . Also, if  $u, v \in U$ , then  $d_G(u) + d_G(v) =$   
 120  $2(p + 2) = 2p + 4 = n + 2$ ; while, if  $u, v \in V$ , then  $d_G(u) + d_G(v) = 2p = n - 2$ . Thus, for any two

---

<sup>2</sup>From here on, perfect matchings are only considered for graphs with even order.

121 non-adjacent vertices  $u$  and  $v$  of  $G$ , we have  $d_G(u) + d_G(v) \geq n - 2$ , which implies that  $u$   
 122 and  $v$  are adjacent in  $G'$ . Thus  $G'$  is complete, and hence it admits perfect matchings.  $\square$

123 **Theorem 2.3.** *Every graph  $G$  with even order  $n \geq 2$  admits perfect matchings if and only*  
 124 *if  $\text{clos}(G, n - 1)$  does. Besides, value  $n - 1$  in the previous statement is best possible.*

125 *Proof.* Set  $G' = \text{clos}(G, n - 1)$ . Clearly, if  $G$  admits perfect matchings, then so does  $G'$ .  
 126 Thus, it remains to prove that if  $G'$  admits perfect matchings, then so does  $G$ . Consider  
 127 an  $(n - 1)$ -closing sequence  $G_0, \dots, G_m$  where  $G_0 = G$  and  $G_m = G'$ . It suffices to prove  
 128 that, for any  $i \in \{1, \dots, m\}$ , from any perfect matching of  $G_i$  we can deduce one of  $G_{i-1}$ .  
 129 Set thus  $H = G_{i-1}$  and  $H' = G_i$ , and let  $M$  be a perfect matching of  $H'$ .

130 If  $M$  is a perfect matching of  $H$ , then we are done. Otherwise, it means  $M$  contains  
 131  $uv$ , the unique edge of  $H'$  not in  $H$ . Then, by definition,  $d_H(u) + d_H(v) \geq n - 1$ . Now, for  
 132 every neighbour  $w \neq v$  of  $u$  in  $H$ , note that if  $uw' \in M$  for some  $w' \notin \{u, v, w\}$ , and we have  
 133  $vw' \in E(H)$ , then  $M \setminus \{uw, ww'\} \cup \{uw, vw'\}$  is a perfect matching of  $H$ . If there is no such  
 134 configuration in  $H$ , then it means that, for every edge  $uw \in E(H)$  such that  $uw' \in M$ , we  
 135 cannot have  $vw' \in E(H)$ . From this, we deduce that  $d_H(v) \leq n - 2 - d_H(u)$ , and thus that  
 136  $d_H(u) + d_H(v) \leq n - 2$ , contradicting that  $uv$  is an edge of  $H'$ . Thus we can always derive  
 137 a perfect matching of  $H$  from  $M$ , and we are done.

138 The last part of the statement follows *e.g.* from Lemma 2.2.  $\square$

139 Due to Theorem 2.3, it is not true that any  $n$ -graph is AP if and only if its  $(n - 2)$ -  
 140 closure is AP, which would have stood as a smooth and natural weakening of Bondy and  
 141 Chvátal's Theorem 1.2 to APness. The fact that perfect matchings are an obstruction to  
 142 such a result is actually not that surprising either, as, recall, Marczyk's Theorem 1.1, to  
 143 weaken Ore's Theorem from  $n - 1$  to  $n - 2$ , already had to exclude such partitions from the  
 144 equation. In the present case, however, things are a bit different, as we can actually adapt  
 145 Theorem 2.3 to partitions with spectrum  $\{3\}$ , see upcoming Theorem 2.5.

146 As earlier, we start off by introducing some construction that will show the next result  
 147 is best possible. In the present case, the construction actually holds for any size value  
 148  $\lambda \geq 2$ , although upcoming Theorem 2.5 deals only with the particular case where  $\lambda = 3$ .  
 149 For any three positive integers  $p, q, r \geq 1$ , we denote by  $T(p, q, r)$  the graph obtained from  
 150 the disjoint union of three cliques  $K_p, K_q$ , and  $K_r$  on  $p, q$ , and  $r$  vertices, respectively, by  
 151 adding a universal vertex  $v$ . Note that  $v$  is a cut-vertex of  $T(p, q, r)$ , whose removal results  
 152 in exactly three connected components, being  $K_p, K_q$ , and  $K_r$ .

153 **Theorem 2.4.** *For every  $\lambda \geq 2$  and  $p \geq 1$  with  $p \equiv \lambda - 1 \pmod{\lambda}$ ,  $\text{clos}(T(1, p, p), n - 2)$  has*  
 154 *realisations of  $(\lambda, \dots, \lambda)$  while  $T(1, p, p)$  has not.*

155 *Proof.* For any  $\lambda \geq 2$  and  $p \geq 1$  with  $p \equiv \lambda - 1 \pmod{\lambda}$ , set  $G = T(1, p, p)$  and  $n = |V(G)| =$   
 156  $2p + 2$ . Since  $G$  has a cut-vertex  $v$  whose removal results in three connected components,  
 157  $K, K',$  and  $K''$ , having order 1,  $p$ , and  $p$ , respectively, and  $p \equiv \lambda - 1 \pmod{\lambda}$ , it should be  
 158 clear that, in any realisation  $\mathcal{R}$  of  $(\lambda, \dots, \lambda)$  in  $G$ , there must be one part containing  $v$  and  
 159  $\lambda - 1$  vertices of  $K'$ , and similarly one part containing  $v$  and  $\lambda - 1$  vertices of  $K''$ , which is  
 160 impossible. Thus  $\mathcal{R}$  cannot exist, and  $G$  does not admit any realisation of  $(\lambda, \dots, \lambda)$ .

161 Let us now consider  $G' = \text{clos}(G, n - 2)$ . Note that, in  $G$ , all vertices of  $K'$  and  $K''$   
 162 have degree  $p = \frac{n-2}{2}$ , vertex  $v$  has degree  $n - 1$ , and the unique vertex of  $K$  has degree 1.  
 163 Thus, in  $G'$ , any two vertices of  $K'$  and  $K''$  are adjacent. This implies  $G'$  is traceable, from  
 164 which we get that  $G'$  is AP, and thus admits realisations of  $(\lambda, \dots, \lambda)$ , as claimed.  $\square$

165 **Theorem 2.5.** *Every graph  $G$  of order  $n \equiv 0 \pmod{3}$  at least 3 admits realisations of*  
 166  *$(3, \dots, 3)$  if and only if  $\text{clos}(G, n-1)$  does. Besides, value  $n-1$  in the previous state-*  
 167 *ment is best possible.*

168 *Proof.* Set  $G' = \text{clos}(G, n-1)$ . As in the proof of Theorem 2.3, we focus on an  $(n-1)$ -  
 169 closing sequence  $G_0, \dots, G_m$  where  $G_0 = G$  and  $G_m = G'$ , our goal being to prove that, for  
 170 any  $i \in \{1, \dots, m\}$ , any realisation  $\mathcal{R}$  of  $(3, \dots, 3)$  in some  $H' = G_i$  can be turned into one  
 171 of  $H = G_{i-1}$ . Assume thus  $\mathcal{R}$  does not hold as is in  $H$ ; thus, there is a part  $S = \{u, v, w\}$  of  
 172  $\mathcal{R}$  that contains the unique edge  $vw$  of  $H'$  not in  $H$ , thereby making  $H[S]$  not connected.  
 173 Then we can assume  $H[S]$  contains  $uv$ , while  $w$  is isolated in  $H[S]$ .

174 Let  $n_3, n_2$ , and  $n_1$  be the number of parts of  $\mathcal{R}$  different from  $S$  in which, in  $H$ , vertex  $v$   
 175 has exactly three, two, or one neighbour, respectively. Then  $d_H(v) = 1 + 3n_3 + 2n_2 + n_1$ . We  
 176 now analyse the number of possible edges incident to  $w$  going to some part  $S' = \{x, y, z\}$   
 177 of  $\mathcal{R}$  different from  $S$  in which  $v$  also has neighbours (all these adjacencies being in  $H$ ).

- 178 • If  $v$  is adjacent to all of  $x, y$ , and  $z$ , then we claim a realisation of  $(3, \dots, 3)$  in  $H$  can  
 179 be deduced if  $w$ , in  $H$ , is also adjacent to any vertex of  $S'$ . Indeed, assume, w.l.o.g.,  
 180 that  $wx$  is an edge of  $H$ . Since  $H[S']$  is connected,  $H'$  (and thus  $H$ ) necessarily  
 181 contains one of  $xy$  or  $xz$ . In the former case, replacing  $S$  and  $S'$  in  $\mathcal{R}$  with  $\{u, v, z\}$   
 182 and  $\{w, x, y\}$  results in a realisation of  $(3, \dots, 3)$  in  $H$ . In the latter case, we can  
 183 instead replace parts  $S$  and  $S'$  with  $\{u, v, y\}$  and  $\{w, x, z\}$ .
- 184 • If  $v$  is adjacent only to, say,  $x$  and  $y$  in  $S'$ , then we claim a realisation of  $(3, \dots, 3)$   
 185 in  $H$  can be deduced if  $w$ , in  $H$ , is also adjacent to any two vertices in  $S'$ . First off,  
 186 note that, for similar arguments as in the previous case, we would be done if these  
 187 at least two neighbours of  $w$  in  $S'$  include  $z$ . So suppose  $w$  is only adjacent to  $x$  and  
 188  $y$ . Since  $H'[S']$  is connected, we have that  $xz$  or  $yz$  lies in  $H'$  (and thus in  $H$ ). In  
 189 the former case, replace, in  $\mathcal{R}$ , parts  $S$  and  $S'$  with  $\{u, v, y\}$  and  $\{w, x, z\}$  to get a  
 190 realisation in  $H$ . In the latter case, replace  $S$  and  $S'$  with  $\{u, v, x\}$  and  $\{w, y, z\}$ .
- 191 • If  $v$  is adjacent only to, say,  $x$  in  $S'$ , then, for similar reasons as earlier, we claim a  
 192 realisation of  $(3, \dots, 3)$  in  $H$  can be obtained from  $\mathcal{R}$  if  $w$  is adjacent, in  $H$ , to all  
 193 three vertices of  $S'$ . This is because, in that case, replacing, in  $\mathcal{R}$ , parts  $S$  and  $S'$   
 194 with  $\{u, v, x\}$  and  $\{w, y, z\}$  yields a desired realisation.

Now, if none of the situations above occurs, then we deduce that

$$d_H(w) \leq n - 3 - 3n_3 - 2n_2 - n_1 = n - 2 - (1 + 3n_3 + 2n_2 + n_1) = n - 2 - d_H(v),$$

195 and hence  $d_H(w) \leq n - 2 - d_H(v)$ , and  $d_H(v) + d_H(w) \leq n - 2$ , a contradiction.

196 To conclude the proof, remark that *e.g.* Theorem 2.4 shows the second part of the  
 197 statement also holds true. □

198 Although the first part of the statement of Theorem 2.5 holds only for partitions con-  
 199 taining value  $\lambda = 3$  only, Theorem 2.4 implies that if the first part of Theorem 2.5 also held  
 200 for any  $\lambda \geq 4$ , then value  $n-1$  would be best possible. Anyhow, Theorem 2.5 shows that  
 201 Bondy and Chvátal's Theorem 1.2 does not weaken to APness by just considering threshold  
 202  $n-2$ , and this is not due to perfect matchings only, as Theorem 2.3 could indicate.

### 3. Connected partitions into parts of size at most 3

In previous Theorems 2.3 and 2.5, we were only concerned with connected partitions into parts all having the same size,  $\lambda$ , for very small values of  $\lambda$ , namely 2 and 3. To get a flavour of what it would take, with the same approach, to generalise our arguments to any partition, in the next result we consider a generalisation of Theorems 2.3 and 2.5 to partitions with spectrum included in  $\{1, 2, 3\}$ , or, in other words, to connected partitions into parts of size at most 3. As will be recalled right after the proof, such a result is not interesting only for generalisation purposes, but also because such partitions, in previous works on APness, have been proved to be of prime interest for certain graph classes.

**Theorem 3.1.** *For every  $n \geq 2$  and every  $n$ -partition  $\pi$  with spectrum included in  $\{1, 2, 3\}$ , every graph  $G$  of order  $n$  admits realisations of  $\pi$  if and only if  $\text{clos}(G, n-1)$  does. Besides, value  $n-1$  in the previous statement is best possible.*

*Proof.* The proof goes similarly as the proofs of Theorems 2.3 and 2.5. Since one of the two directions is obvious, we focus on proving the less straight direction. Set  $G' = \text{clos}(G, n-1)$ , and let  $\pi$  be an  $n$ -partition with spectrum included in  $\{1, 2, 3\}$ . We consider an  $(n-1)$ -closing sequence  $G_0, \dots, G_m$  where  $G_0 = G$  and  $G_m = G'$ . Again, it suffices to prove that, for any  $i \in \{1, \dots, m\}$ , any realisation of  $\pi$  in  $G_i$  yields one of  $G_{i-1}$ . So, we consider  $H = G_{i-1}$  and  $H' = G_i$ , and let  $\mathcal{R}$  be a realisation of  $\pi$  in  $H'$ . If  $\mathcal{R}$  does not hold directly in  $H$ , then it is because of a part  $S$  containing the unique edge of  $H'$  that does not belong to  $H$  (that is,  $H[S]$  is not connected while  $H'[S]$  is). Then,  $|S| \in \{2, 3\}$ .

To begin with, we assume  $S$  has cardinality 2, and set  $S = \{u, v\}$ . Then,  $uv \in E(H')$  and  $uv \notin E(H)$ . As in the proofs of Theorems 2.3 and 2.5, we analyse the possible neighbours of  $v$  in  $H$  w.r.t. those of  $u$ .

- If  $u$  has, in  $H$ , a neighbour in a part  $S'$  of cardinality 2 of  $\mathcal{R}$  different from  $S$ , and  $v$  is adjacent to the other vertex of  $S'$ , then, through the exact same arguments as in the proof of Theorem 2.3, we can deduce, from  $\mathcal{R}$ , a realisation of  $\pi$  in  $H$ .
- If  $u$  is adjacent, in  $H$ , to a vertex  $w$  such that  $S' = \{w\}$  is a part of  $\mathcal{R}$ , then note that, from  $\mathcal{R}$ , replacing  $S$  and  $S'$  with  $\{u, w\}$  and  $\{v\}$  results in a realisation in  $H$ .
- Assume last that  $u$  has, in  $H$ , neighbours in a part  $S' = \{w, x, y\}$  of cardinality 3.
  - If  $u$  is adjacent to all of  $w, x$ , and  $y$ , and  $v$  is also adjacent to some vertex of  $S'$ , then a realisation of  $\pi$  in  $H$  can be obtained from  $\mathcal{R}$  as follows. Assuming, w.l.o.g., that  $v$  is adjacent to  $w$ , then just replace parts  $S$  and  $S'$  with  $\{u, x, y\}$  and  $\{v, w\}$ . This indeed results in a desired realisation.
  - If  $u$  is adjacent, in  $H$ , to exactly two vertices in  $S'$ , say  $w$  and  $x$  w.l.o.g., then we claim a realisation of  $\pi$  in  $H$  can be deduced from  $\mathcal{R}$  in case  $v$  is also adjacent to two vertices of  $S'$  in  $H$ . We distinguish two cases.
    - \* Assume first that  $v$  is adjacent to  $y$ . In that case, it suffices to replace parts  $S$  and  $S'$  of  $\mathcal{R}$  with  $\{u, w, x\}$  and  $\{v, y\}$ .
    - \* Assume second that  $v$  is adjacent to  $w$  and  $x$  (and not to  $y$ ). Since  $H'[S']$  (and thus  $H[S']$ ) is connected, then one of  $w$  and  $x$  is adjacent to  $y$  in  $H'$  (and  $H$ ). Assuming w.l.o.g. that  $xy$  is an edge of  $H$ , we can here replace parts  $S$  and  $S'$  of  $\mathcal{R}$  with  $\{u, w\}$  and  $\{v, x, y\}$  to be done.
  - If  $u$  is adjacent, in  $H$ , to only one vertex of  $S'$ , say  $w$  w.l.o.g., and  $v$  is adjacent, in  $H$ , to all of  $w, x$ , and  $y$ , then we can obtain a desired realisation by replacing parts  $S$  and  $S'$  of  $\mathcal{R}$  with  $\{u, w\}$  and  $\{v, x, y\}$ .

Now denote by  $m_1$  and  $m_2$  the number of parts of size 2 of  $\mathcal{R}$  in which  $u$  has (in  $H$ ) exactly one or two neighbours, respectively, and by  $n_1$ ,  $n_2$ , and  $n_3$  the number of parts of size 3 in which  $u$  has (in  $H$ ) exactly one, two, or three neighbours. Then

$$d_H(u) = m_1 + 2m_2 + n_1 + 2n_2 + 3n_3.$$

Also, since none of the cases above apply, we have

$$d_H(v) \leq n - 2 - m_1 - 2m_2 - n_1 - 2n_2 - 3n_3 = n - 2 - d_H(u),$$

248 and thus  $d_H(u) + d_H(v) \leq n - 2$ , a contradiction.

249 Now suppose  $S = \{u, v, w\}$ , where  $H[S]$  contains the edge  $uv$  while  $w$  is isolated (because  $H'$  contains the edge  $vw$  while  $H$  does not contain it). For the same reasons as in  
250  
251 the proof of Theorem 2.5, we can assume that, in  $\mathcal{R}$ , there is no  $S' \neq S$  of size 3 in which:

- 252 •  $v$  has three neighbours and  $w$  has one neighbour;
- 253 •  $v$  has two neighbours and  $w$  has two neighbours;
- 254 •  $v$  has one neighbour and  $w$  has three neighbours.

255 We now lead a similar analysis regarding the possible neighbours of  $v$  in  $H$  in parts  $S' \neq S$   
256 of  $\mathcal{R}$  of cardinality 1 or 2.

- 257 • Assume first that  $v$  has, in  $H$ , a neighbour  $x$  in a part  $S' = \{x\}$  of size 1. Then a  
258 realisation of  $\pi$  in  $H$  is obtained upon replacing  $S$  and  $S'$  with  $\{u, v, x\}$  and  $\{w\}$ .
- 259 • Assume second that  $v$  has, in  $H$ , neighbours in a part  $S' = \{x, y\}$  of cardinality 2.  
260 Recall that  $xy$  is an edge of both  $H'$  and  $H$ .
  - 261 – Assume first  $v$  is adjacent to both  $x$  and  $y$  in  $H$ . If  $w$  is adjacent to any of  $x$  and  
262  $y$  in  $H$ , then a realisation of  $\pi$  in  $H$  is obtained from  $\mathcal{R}$  upon replacing parts  $S$   
263 and  $S'$  with either  $\{u, v, x\}$  and  $\{w, y\}$ , or  $\{u, v, y\}$  and  $\{w, x\}$ .
  - 264 – Assume second  $v$  is adjacent only to  $x$  in  $G$ . If  $w$  is adjacent to  $x$  and  $y$ , then a  
265 realisation is obtained when replacing parts  $S$  and  $S'$  with  $\{u, v, x\}$  and  $\{w, y\}$ .

Now set  $m_1$  and  $m_2$  the number of parts of cardinality 2 of  $\mathcal{R}$  in which  $v$  has (in  $H$ ) exactly one or two neighbours, respectively, and  $n_1$ ,  $n_2$ , and  $n_3$  the number of parts of cardinality 3 in which  $v$  has (in  $H$ ) exactly one, two, or three neighbours. Then

$$d_H(v) = 1 + m_1 + 2m_2 + n_1 + 2n_2 + 3n_3.$$

Meanwhile, by the arguments above, we have

$$d_H(w) \leq n - 3 - m_1 - 2m_2 - n_1 - 2n_2 - 3n_3 = n - 2 - d_H(v),$$

266 from which we deduce  $d_H(v) + d_H(w) \leq n - 2$ , a final contradiction.

267 The very last part of the statement follows *e.g.* from Lemma 2.2 and Theorem 2.4.  $\square$

268 Theorem 3.1 makes more particular sense in the context of graph classes for which the  
269 AP property is known to rely only on the realisability of partitions with spectrum included  
270 in  $\{1, 2, 3\}$ . Such matters relate to questions raised first by Barth and Fournier in [2] on  
271 the complexity of determining whether a graph is AP. Note that it is not clear whether  
272 this problem lies in NP or co-NP, since the APness of an  $n$ -graph relies on the realisability



273 of an exponential number of  $n$ -partitions, while the non-APness of an  $n$ -graph implies an  
 274 exponential number of partitions of the vertex set do not stand as realisations of particular  
 275  $n$ -partitions. Still, following the terminology from [4], it is believed that, perhaps, every  
 276  $n$ -graph  $G$  admits a *polynomial kernel of partitions*, i.e., a set  $K$  of a polynomial number  
 277 (function of  $n$ ) of  $n$ -partitions such that  $G$  is AP if and only if all partitions of  $K$  are  
 278 realisable in  $G$ . If this was true, then this would imply deciding APness lies in NP.

279 Polynomial kernels of partitions have been proved to exist for a few graph classes, such  
 280 as subdivided stars [1, 2], graphs with large diameter [20], superclasses of split graphs [6,  
 281 8, 15], and others [4, 6]. In particular, the polynomial kernel from [8] for split graphs  
 282 contains partitions with spectrum included in  $\{1, 2, 3\}$  only, which kernel also holds for  
 283 superclasses of split graphs [15], in particular for  $\{2K_2, C_4\}$ -free graphs. Thus, a direct,  
 284 general consequence of Theorem 3.1 and of these thoughts, is the following:

285 **Corollary 3.2.** *If  $\mathcal{G}$  is a class of graphs of order  $n \geq 2$  admitting  $n$ -partitions with spectrum*  
 286 *included in  $\{1, 2, 3\}$  as a polynomial kernel, then every  $G \in \mathcal{G}$  is AP if and only if  $\text{clos}(G, n -$   
 287  $1)$  is.*

288 Note that Corollary 3.2 would extend the same way to any  $n$ -graph class for which  
 289  $n$ -partitions with spectrum included in  $\{1, 2, 3\}$  stand as a polynomial kernel. Recall also  
 290 that building graph closures can be done in polynomial time (see e.g. [9]).

#### 291 4. More general results for weaker closures

292 In this section, we investigate how the proof arguments from the proofs of previous  
 293 Theorems 2.3, 2.5, and 3.1 could be generalised to other partitions with spectrum of size 1  
 294 (Subsection 4.1) and even to APness (Subsection 4.2), provided we consider weaker closures.

##### 295 4.1. Connected partitions into parts of size $\lambda$

296 Before proceeding with our main result here, let us establish the following useful lemma.

297 **Lemma 4.1.** *If  $G$  is an  $n$ -graph with two non-adjacent vertex  $u$  and  $v$  such that  $d(u) +$   
 298  $d(v) \geq n + c$ , then  $u$  and  $v$  have at least  $c + 2$  common neighbours. In particular,  $c \leq n - 4$ .*

*Proof.* Since  $u$  and  $v$  are not adjacent, we have  $N(u) \cup N(v) \subseteq V(G) \setminus \{u, v\}$ , which set  
 has cardinality  $n - 2$ . Now, if  $u$  and  $v$  had at most  $c + 1$  common neighbours only, then we  
 would have

$$d(u) + d(v) \leq 2(c + 1) + (n - 2) - (c + 1) = n + c - 1,$$

299 a contradiction.

300 The last part is because having  $d(u) + d(v) \geq 2n - 3$  would imply  $u$  and  $v$  have at least  
 301  $n - 1$  common neighbours, which is impossible in loopless graphs.  $\square$

302 We now essentially adapt the first part of the statement of Theorem 2.5 to partitions  
 303 with spectrum  $\{\lambda\}$  for any  $\lambda \geq 4$ . Recall that Theorem 2.4 is precisely about such partitions,  
 304 and thus indicates that, even in this context, we must at least consider  $(n - 1)$ -closures.

305 **Theorem 4.2.** *For every  $\lambda \geq 4$ , every graph  $G$  of order  $n \equiv 0 \pmod{\lambda}$  at least  $\lambda$  admits*  
 306 *realisations of  $(\lambda, \dots, \lambda)$  if and only if  $\text{clos}(G, 2n - (\frac{2n}{\lambda} + \lambda - 1))$  does.*

307 *Proof.* Set  $\alpha = 2n - (\frac{2n}{\lambda} + \lambda - 1)$ , and  $G' = \text{clos}(G, \alpha)$ . Consider an  $\alpha$ -closing sequence  
 308  $G_0, \dots, G_m$  where  $G_0 = G$  and  $G_m = G'$ . It again suffices to prove that any realisation  $\mathcal{R}$   
 309 of  $(\lambda, \dots, \lambda)$  of  $H' = G_i$  for any  $i \in \{1, \dots, m\}$  yields one of  $H = G_{i-1}$ . If  $\mathcal{R}$  does not hold

310 as is in  $H$ , then it is because a part  $S$  of  $\mathcal{R}$  is not connected in  $H$  (while it is in  $H'$ ). That  
 311 is,  $H'[S]$  contains the only edge  $uv$  that is present in  $H'$  but not in  $H$ .

Since  $H[S]$  is not connected, and  $H[S] = H'[S] - uv$ , vertices  $u$  and  $v$  cannot have any  
 common neighbour in  $S$ . We claim there is necessarily a part  $S' \neq S$  of  $\mathcal{R}$  in which  $u$  and  $v$   
 have at least  $\lambda - 1$  common neighbours. Indeed, note first that the value of  $\lambda$  ensures that  
 $n \geq 2\lambda$ , as, for  $n = \lambda$ , we have  $\alpha = \lambda - 1$ , while  $H$  in the present case is not connected and  
 thus  $d_H(u) + d_H(v) \leq n - 2 = \lambda - 2$ , a contradiction. Thus,  $|\mathcal{R}| \geq 2$ , and if  $u$  and  $v$  had at  
 most  $\lambda - 2$  common neighbours in all of the  $\frac{n}{\lambda} - 1 \geq 1$  parts of  $\mathcal{R}$  different from  $S$ , then  $u$   
 and  $v$  would have at most

$$\left(\frac{n}{\lambda} - 1\right)(\lambda - 2) = n - \left(\frac{2n}{\lambda} + \lambda - 2\right)$$

312 common neighbours, while, by Lemma 4.1, vertices  $u$  and  $v$  have at least  $n - (\frac{2n}{\lambda} + \lambda - 3)$   
 313 common neighbours in  $H$ , a contradiction.

314 Thus, let  $S' \neq S$  be any part of  $\mathcal{R}$  in which  $u$  and  $v$  have at least  $\lambda - 1$  common neighbours.  
 315 Let also  $S_u$  and  $S_v$  be the vertex sets of the exactly two connected components of  $H[S]$ ,  
 316 where  $S_u$  contains  $u$  while  $S_v$  contains  $v$ . In case  $u$  and  $v$  have exactly  $\lambda - 1$  common  
 317 neighbours in  $S'$ , then let also  $w$  be the unique vertex of  $S'$  not adjacent to both  $u$  and  $v$ ;  
 318 otherwise, if  $u$  and  $v$  are both adjacent to all vertices of  $S'$ , then let  $w$  be any vertex of  $S'$ .  
 319 In both cases, let  $w'$  be any neighbour of  $w$  in  $H[S']$ .

320 Since  $\lambda \geq 4$ , we must have  $\lambda - |S_u| \geq 2$  or  $\lambda - |S_v| \geq 2$ . Assume w.l.o.g. the former  
 321 inequality holds. Now replace  $S$  and  $S'$  in  $\mathcal{R}$  with both  $S_u \cup \{w, w'\} \cup X$  and  $S_v \cup Y$ , where  
 322  $X$  is any set of  $\lambda - (|S_u| + 2) \geq 0$  vertices of  $S' \setminus \{w, w'\}$ , and  $Y = S' \setminus \{w, w'\} \setminus X$ . Since  
 323  $ww'$  is an edge of  $H$ , and  $u$  and  $v$  are adjacent to all vertices in  $S' \setminus \{w\}$ , it can be observed  
 324 that this results in a realisation of  $(\lambda, \dots, \lambda)$  in  $H$ .  $\square$

325 Note that  $2n - (\frac{2n}{\lambda} + \lambda - 1)$  is a non-decreasing function of  $\lambda$ , so the most interesting  
 326 case, yielding the stronger closure, is for  $\lambda = 4$ , for which case Theorem 4.2 deals with  
 327  $(\frac{3}{2}n - 3)$ -closures. Recall that, regardless of the actual value of  $\lambda$ , in any case we cannot  
 328 hope to establish results involving closures stronger than  $(n - 1)$ -closures.

#### 329 4.2. APness

330 We here consider the AP property in full, that is, we wonder about generalisations of  
 331 Bondy and Chvátal's Theorem 1.2 to AP graphs. By the last part of Lemma 4.1, recall  
 332 that we must consider degree sum thresholds at most  $2n - 4$ . In what follows, we prove that  
 333 an  $n$ -graph  $G$  is indeed AP if and only if its  $(2n - 4)$ -closure is. This result being surely far  
 334 from optimal, we then improve it down to  $(2n - 5)$ -closures and even to  $(2n - 6)$ -closures,  
 335 for the sake mainly of showcasing why it might be tedious to go even lower.

336 **Theorem 4.3.** *Every graph  $G$  of order  $n \geq 3$  is AP if and only if  $\text{clos}(G, 2n - 4)$  is.*

337 *Proof.* Set  $G' = \text{clos}(G, 2n - 4)$ . Again, it suffices to prove that if the  $(2n - 4)$ -closure  
 338  $\text{clos}(G, 2n - 4)$  is AP, then  $G$  is also AP. Actually, we can consider a  $(2n - 4)$ -closing sequence  
 339  $G_0, \dots, G_m$  where  $G_0 = G$  and  $G_m = G'$ , and prove that, for  $H' = G_i$  and  $H = G_{i-1}$  for some  
 340  $i \in \{1, \dots, m\}$ , any realisation  $\mathcal{R}$  of any  $n$ -partition  $\pi$  in  $H'$  yields one of  $H$ . If  $\mathcal{R}$  holds as  
 341 is in  $H$ , then we are done. Otherwise, some part  $S$  of  $\mathcal{R}$  contains the only edge  $uv$  of  $H'$   
 342 not in  $H$ , and we have that  $H[S]$  is not connected.

343 Let us denote by  $S_u$  and  $S_v$  the subsets of vertices of  $S$  such that  $S_u$  contains the  
 344 vertices of  $H[S]$  belonging to the same connected component as  $u$ , and  $S_v$  contains those  
 345 belonging to the same connected component as  $v$ . Then,  $S = S_u \cup S_v$ , and we have  $u \in S_u$

346 and  $v \in S_v$ . By Lemma 4.1, vertices  $u$  and  $v$  have at least  $n - 2$  common neighbours in  $H$ ,  
347 and, actually, since  $u$  and  $v$  are not adjacent in  $H$ , we have  $N_H(u) = N_H(v) = V(H) \setminus \{u, v\}$ .  
348 In particular, since  $H[S]$  is not connected, it must be that  $|S_u| = |S_v| = 1$ , as otherwise  $u$   
349 and  $v$  would have, in  $H$ , a common neighbour in either  $S_u$  or  $S_v$ , making  $H[S]$  connected.  
350 Thus,  $|S| = 2$ , and since  $n - 2 \geq 1$  we have  $|\pi| \geq 2$ . Now, consider  $S' \neq S$  another part of  $\mathcal{R}$ .  
351 Then, as mentioned above, in  $H$ , vertices  $u$  and  $v$  are adjacent to all vertices of  $S'$ . It then  
352 suffices, to obtain a realisation of  $\pi$  in  $H$ , to start from  $\mathcal{R}$ , and, denoting by  $w$  any vertex  
353 of  $S'$ , to replace parts  $S$  and  $S'$  with, say,  $\{u, w\}$  and  $\{v\} \cup S' \setminus \{w\}$ .  $\square$

354 **Theorem 4.4.** *Every graph  $G$  of order  $n \geq 4$  is AP if and only if  $\text{clos}(G, 2n - 5)$  is.*

355 *Proof.* The proof goes similarly as that of Theorem 4.3 (from a  $(2n - 5)$ -closing sequence).  
356 This time, in  $H$  vertices  $u$  and  $v$  have at least  $n - 3$  common neighbours. Since  $H[S]$  is not  
357 connected, vertices  $u$  and  $v$  have no common neighbours in  $S$ , which implies  $2 \leq |S| \leq 3$ .

358 If  $|S| = 3$ , then we can assume, w.l.o.g., that  $S_u = \{u\}$  and  $S_v = \{v, w\}$ . Thus  $vw$  is  
359 an edge of  $H$ , and  $u$  and  $v$  admit all vertices of  $V(H) \setminus \{u, v, w\}$  as common neighbours.  
360 Then, again, we can just consider any part  $S' \neq S$  of  $\mathcal{R}$  (such exists since  $2n - 5 \geq 3$ ), and,  
361 denoting by  $x$  any vertex of  $S'$ , a realisation of  $\pi$  in  $H$  is obtained when starting from  $\mathcal{R}$ ,  
362 and replacing parts  $S$  and  $S'$  with  $\{v, w, x\}$  and  $\{u\} \cup S' \setminus \{x\}$ , respectively.

363 Now assume  $|S| = 2$ , i.e.,  $S$  contains  $u$  and  $v$  only. If  $N_H(u) = N_H(v) = V(H) \setminus \{u, v\}$ ,  
364 then a realisation of  $\pi$  in  $H$  can be obtained similarly as in the proof of Theorem 4.3.  
365 Hence, we can last assume  $u$  and  $v$  have only  $n - 3$  common neighbours in  $H$ . Let us thus  
366 denote by  $w$  the only vertex of  $V(H) \setminus \{u, v\}$  not adjacent to both  $u$  and  $v$  in  $H$ . Again,  
367 since  $2n - 5 \geq 3$ , we must have  $|\pi| \geq 2$ . If  $\mathcal{R}$  contains a part  $S' \neq S$  not containing  $w$ , then  
368 we can again reach our conclusion similarly as in the proof of Theorem 4.3. Otherwise,  
369  $|\pi| = 2$ , and  $\mathcal{R}$  thus contains only two parts,  $S$  and  $S'$ , where  $w \in S'$ . By arguments above,  
370  $u$  and  $v$  are adjacent to all vertices of  $S' \setminus \{w\}$ .

- 371 • If  $|S'| = 1$ , then  $n = 3$ , a case not covered by the statement.
- 372 • If  $|S'| = 2$ , then  $S' = \{w, x\}$  where  $x$  is a common neighbour of  $u$  and  $v$  in  $H$ . In that  
373 case,  $n = 4$ , and since  $d_H(u) + d_H(v) \geq 2n - 5 = 3$ , for  $uv$  to be an edge of  $H'$  it must  
374 be that  $uw$  is an edge of  $H$ , w.l.o.g. Then a realisation of  $\pi$  in  $H$  is obtained upon  
375 considering parts  $\{u, w\}$  and  $\{v, x\}$ .
- 376 • If  $|S'| \geq 3$ , then consider any vertex  $x$  of  $S' \setminus \{w\}$  such that  $H[S' \setminus \{x\}]$  is connected.  
377 Such an  $x$  can be obtained e.g. by considering a leaf different from  $w$  of a spanning  
378 tree of  $H[S']$ . Since  $|S'| \geq 3$ , recall that  $H[S' \setminus \{x\}]$  contains a vertex adjacent to  
379 both  $u$  and  $v$ . Then a realisation of  $\pi$  in  $H$  is obtained from  $\mathcal{R}$  when considering  
380 parts  $\{u, x\}$  and  $\{v\} \cup S' \setminus \{x\}$ .

381 This concludes the proof.  $\square$

382 **Theorem 4.5.** *Every  $n$ -graph  $G$  with  $n \geq 5$  is AP if and only if  $\text{clos}(G, 2n - 6)$  is.*

383 *Proof.* We follow the lines of the proofs of Theorems 4.3 and 4.4 (but this time considering  
384 a  $(2n - 6)$ -closing sequence). Here, in  $H$ , vertices  $u$  and  $v$  have at least  $n - 4$  common  
385 neighbours. Because  $H[S]$  is not connected, vertices  $u$  and  $v$  have no common neighbours  
386 in  $S$ , from which we deduce  $2 \leq |S| \leq 4$ . Also, keep in mind throughout that  $n \geq 5$ .

387 Similarly as in the proof of Theorem 4.4, if  $|S| = 4$ , then  $u$  and  $v$  are adjacent to all  
388 vertices of  $V(H) \setminus S$ . If, say,  $|S_u| = 1$  and  $|S_v| = 3$ , then, again, it suffices to consider any  
389 part  $S' \neq S$  of  $\mathcal{R}$  (it can be checked that such exists, as  $n \geq 5$  which implies  $2n - 6 \geq 4$ ), and,

390 denoting by  $w$  any vertex of  $S'$  (being a common neighbour of  $u$  and  $v$  in  $H$ ), a realisation  
 391 of  $\pi$  in  $H$  is obtained when replacing parts  $S$  and  $S'$  of  $\mathcal{R}$  with  $S_v \cup \{w\}$  and  $S' \cup \{u\} \setminus \{w\}$ .  
 392 Otherwise,  $|S_u| = |S_v| = 2$ ; we consider a few cases.

- 393 • If there is a part  $S' \neq S$  of  $\mathcal{R}$  with  $|S'| \geq 4$ , then a realisation of  $\pi$  in  $H$  is obtained  
 394 upon replacing  $S$  and  $S'$  with  $S_u \cup \{w, x\}$  (where  $w$  and  $x$  are any two vertices of  $S'$ )  
 395 and  $S_v \cup S' \setminus \{w, x\}$ .
- 396 • If there is a part  $S' \neq S$  with  $|S'| = 3$ , then, setting  $S' = \{w, x, y\}$ , it here suffices to  
 397 replace  $S$  and  $S'$  with, say,  $S_u \cup \{w, x\}$  and  $S_v \cup \{y\}$ .
- 398 • If there is a part  $S' \neq S$  with  $|S'| = 2$ , then, setting  $S' = \{w, x\}$ , it here suffices to  
 399 replace  $S$  and  $S'$  with, say,  $S_u$  and  $S_v \cup \{w, x\}$ .
- 400 • If there is a part  $S' \neq S$  with  $|S'| = 1$ , then, setting  $S' = \{w\}$ , it here suffices to replace  
 401  $S$  and  $S'$  with, say,  $S_u \setminus \{u\}$  and  $S_v \cup \{w, u\}$ .

402 Thus, when  $|S| = 4$ , a realisation of  $\pi$  in  $H$  can be deduced in all cases.

403 If  $|S| = 3$ , then set  $S = \{u, v, w\}$ , where  $S_u = \{u\}$  and  $S_v = \{v, w\}$  (thus  $vw$  is an edge  
 404 of  $H$ ). If  $u$  and  $v$  are, in  $H$ , both adjacent to all vertices of  $V(H) \setminus S$  (there are such,  
 405 since  $2n - 6 \geq 4$ ), then a realisation of  $\pi$  in  $H$  can be obtained similarly as in the proof of  
 406 Theorem 4.4. Thus, we can assume there is some  $x \in V(H) \setminus S$  such that all vertices of  
 407  $V(H) \setminus S \setminus \{x\}$  are common neighbours of  $u$  and  $v$ . Actually, again, if there is some part  
 408  $S' \neq S$  of  $\mathcal{R}$  not containing  $x$ , then a realisation can be obtained. So we get to assuming  
 409 that  $|\pi| = 2$ , and that, besides  $S$ , the only part  $S'$  of  $\mathcal{R}$  contains  $x$ .

- 410 • If  $|S'| = 1$ , then  $n = 4$ , a case not covered by the statement.
- 411 • If  $|S'| \geq 2$ , then  $S'$  contains a vertex  $y$  adjacent to  $x$  being a common neighbour of  $u$   
 412 and  $v$ . Furthermore,  $u$  and  $v$  are both adjacent to all vertices of  $S' \setminus \{x\}$ . Then the  
 413 two parts  $\{u, x, y\}$  and  $\{v, w\} \cup S' \setminus \{x, y\}$  form a realisation of  $\pi$  in  $H$ .

414 It now remains to consider when  $|S| = 2$ , that is, when  $S = \{u, v\}$ . In this case, there are  
 415 at most two vertices of  $V(H) \setminus \{u, v\}$  that are not common neighbours of  $u$  and  $v$  in  $H$ . We  
 416 can actually suppose that there exist exactly two such vertices  $w$  and  $x$  of  $V(H) \setminus \{u, v\}$   
 417 that are not common neighbours, as, if there is none or only one such vertex, then we can  
 418 proceed similarly as in the proof of Theorem 4.4. Likewise, if there is a part  $S' \neq S$  of  
 419  $\mathcal{R}$  that contains only common neighbours of  $u$  and  $v$  in  $H$ , then a realisation can also be  
 420 deduced. So, besides  $S$ , we can last assume either there is only one part  $S'$  (that is,  $|\pi| = 2$ ,  
 421 and  $S'$  contains both  $w$  and  $x$ ), or there are exactly two parts  $S_w$  and  $S_x$  (that is,  $|\pi| = 3$ ,  
 422 and  $S_w$  contains  $w$  while  $S_x$  contains  $x$ ).

- 423 • Assume first  $|\pi| = 2$ . Thus,  $\mathcal{R}$  contains a unique part  $S' \neq S$ , that contains both  $w$   
 424 and  $x$ . If there is a spanning tree of  $H[S']$  that contains a leaf  $y$  different from  $w$  and  
 425  $x$ , then recall that that leaf is adjacent to both  $u$  and  $v$  in  $H$ , and the two parts  $\{u, y\}$   
 426 and  $\{v\} \cup S' \setminus \{y\}$  then form a realisation of  $\pi$  in  $H$  (unless  $S' = \{w, x, y\}$ , in which  
 427 case necessarily  $wx$  is an edge, and we could instead consider the two parts  $\{u, v, y\}$   
 428 and  $\{w, x\}$ ). So we can now assume that any vertex of  $S' \setminus \{w, x\}$  is a cut-vertex of  
 429 any spanning tree of  $H[S']$ , which implies any spanning tree  $T$  of  $H[S']$  must be a  
 430 path with end-vertices  $w$  and  $x$ . In that case, a realisation of  $\pi$  in  $H$  is obtained upon  
 431 considering parts  $\{w, w'\}$  and  $\{u, v\} \cup S' \setminus \{w, w'\}$ , where  $w'$  is the unique neighbour  
 432 of  $w$  in  $T$ . Note indeed that if this was not the case, then we would have  $w' = x$ , thus  
 433  $|S'| = 2$  and  $n = 4$ , which peculiar case is not covered by the statement.

434 • Assume now  $|\pi| = 3$ . Thus,  $\mathcal{R}$  contains exactly two parts  $S_w$  and  $S_x$  different from  $S$ ,  
435 where  $S_w$  and  $S_x$  contain  $w$  and  $x$ , respectively. Also, in  $H$  vertices  $u$  and  $v$  are both  
436 adjacent to all vertices of  $S_w \setminus \{w\}$  and  $S_x \setminus \{x\}$ . Here, assume, say, that  $H[S_w]$   
437 contains a spanning tree with a leaf  $y$  different from  $w$ . Then, a realisation of  $\pi$  in  
438  $H$  is obtained upon replacing, in  $\mathcal{R}$ , parts  $S$  and  $S_w$  with  $\{u, y\}$  and  $\{v\} \cup S_w \setminus \{y\}$ .  
439 Now, if none of  $H[S_w]$  and  $H[S_x]$  admits such a spanning tree, then  $|S_w| = |S_x| = 1$   
440 and  $n = 4$ , which peculiar case is not covered by the statement.

441 We are thus done in all cases. □

## 442 5. Conclusions

443 In this work, inspired by Bondy and Chvátal's influential Theorem 1.2, we have in-  
444 vestigated closure results for connected partitions of graphs, in particular in the context  
445 of the AP property which has been regarded as a weakening of Hamiltonicity. Looking  
446 at Theorem 1.2, a natural first question (in view of Marczyk's Theorem 1.1) was whether  
447  $(n - 2)$ -closures are necessary and sufficient for the AP property. In Section 2, we proved  
448 that this is not true, because of several particular types of  $n$ -partitions. For such parti-  
449 tions, we proved, in particular throughout Section 3, that it is sufficient and necessary to  
450 consider  $(n - 1)$ -closures. In last Section 4, we then considered other  $n$ -partitions, includ-  
451 ing the whole set of  $n$ -partitions (and thus the AP property), and proved that it is here  
452 sufficient to consider weaker closures (namely  $(2n - c)$ -closures for some  $c$ ).

453 Let us mention that the fact that  $(n - 2)$ -closures do not suffice for APness (Theorems 2.3  
454 and 2.5) is not too surprising, given that Marczyk's Theorem 1.1 already had to exclude  
455 perfect matchings from the equation. Given all connections between the results involved,  
456 the fact that perfect matchings form an obstruction to APness and  $(n - 2)$ -closures can  
457 thus be regarded as a counterpart. What is more surprising is that perfect matchings, in  
458 our case, are not the only obstruction, recall Theorem 2.5.

459 Given the results we came up with, we thus have no obvious objection again the fact  
460 that, perhaps, in general  $(n - 1)$ -closures are necessary and sufficient.

461 **Question 5.1.** *Is it true that any graph is AP if and only if its  $(n - 1)$ -closure is?*

462 If true, a notable consequence of Question 5.1 is that it would imply Marczyk's Theo-  
463 rem 1.1, just like how Bondy and Chvátal's Theorem 1.2 implies Ore's Theorem. A way to  
464 progress towards Question 5.1 could be to first generalise Theorems 4.3 to 4.5 to  $(2n - c)$ -  
465 closures for any fixed  $c \geq 4$ . However, as highlighted by the proofs of Theorems 4.3 to 4.5,  
466 proceeding this way is not quite obvious. In particular, in our arguments, it is important  
467 that we keep control over the structure of  $H[S]$ , which is less and less obvious as  $c$  in-  
468 creases, but easy to do when  $c$  is very small (as in the cases we considered). Also, the fact  
469 that  $uv$  is an edge of  $H'$  but not of  $H$  is a very local property, which becomes less and  
470 less easy to exploit as the diameter of  $H[S]$  increases (which might be the case when  $c$   
471 is large). Altogether, an annoying point is that  $|S|$  being large implies  $u$  and  $v$  have lots  
472 of common neighbours in  $V(H) \setminus S$  but the structure of  $H[S]$  might get hard to handle,  
473 while  $|S|$  being small implies the structure of  $H[S]$  is easier to deal with but  $u$  and  $v$  have  
474 less common neighbours in  $V(H) \setminus S$ , which is a crucial point.

475 To summarise, while some of our arguments in the proofs of Theorems 4.3 to 4.5 could  
476 obviously be generalised to larger values of  $c$ , some others do not; thus, to go further, it  
477 would be crucial to come up with other arguments. For instance, it might be useful to  
478 exploit that if, for some  $c$ , the closure  $\text{clos}(G, 2n - c)$  is AP but  $H[S]$  has a very faulty  
479 structure, then we could deduce other  $n$ -partitions showing that  $\text{clos}(G, 2n - c)$  is not AP.

480 One has to take into account also that if  $c$  is too large w.r.t.  $n$ , then we might fall  
 481 into pathological cases, as better highlighted by Theorem 4.5. Note indeed that if we  
 482 consider as  $G$  the disjoint union of two edges, then  $n = 4$  and  $2n - 6 = 2$ , while, for any  
 483 two non-adjacent vertices  $u$  and  $v$  of  $G$  we have  $d_G(u) + d_G(v) = 2$ . Thus in that case  
 484  $\text{clos}(G, 2n - 6)$  is complete and thus AP, while  $G$  is obviously not AP (consider partitioning  
 485 it following  $(1, 3)$ ). So the statement of Theorem 4.5 is actually best possible (w.r.t.  $n$ ),  
 486 and, as considering larger values of  $c$ , we must make sure to focus on large enough values of  
 487  $n$  only, or add additional constraints (such as focusing at least on connected graphs only).

488 Other directions of interest include other results of the type of Theorem 3.1, to establish  
 489 other results of the sort of Corollary 3.2, for other polynomial kernels of partitions. As a  
 490 more general perspective, we wonder whether it could be worth studying the APness of  
 491 graphs w.r.t. the traceability or the Hamiltonicity of some of their closures, and *vice versa*.

## 492 References

- 493 [1] D. Barth, O. Baudon, J. Puech. Decomposable trees: a polynomial algorithm for  
 494 tripodes. *Discrete Applied Mathematics*, 119(3):205–216, 2002.
- 495 [2] D. Barth, H. Fournier. A degree bound on decomposable trees. *Discrete Mathematics*,  
 496 306(5):469–477, 2006.
- 497 [3] J. Bensmail. A  $\sigma_3$  condition for arbitrarily partitionable graphs. *Discussiones Mathe-*  
 498 *maticae Graph Theory*, 44(2):755–776, 2024.
- 499 [4] J. Bensmail. On three polynomial kernels of sequences for arbitrarily partitionable  
 500 graphs. *Discrete Applied Mathematics*, 202:19–29, 2016.
- 501 [5] J. Bensmail. Toughness Properties of Arbitrarily Partitionable Graphs. Preprint, avail-  
 502 able online at <https://hal.science/hal-04312057>. 2023.
- 503 [6] J. Bensmail, B. Li. More Aspects of Arbitrarily Partitionable Graphs. *Discussiones*  
 504 *Mathematicae Graph Theory*, 42(4):1237–1261, 2022.
- 505 [7] J.A. Bondy, V. Chvátal. A method in graph theory. *Discrete Mathematics*, 15(2):111–  
 506 135, 1976.
- 507 [8] H. Broersma, D. Kratsch, G.J. Woeginger. Fully decomposable split graphs. *European*  
 508 *Journal of Combinatorics*, 34(3):567–575, 2013.
- 509 [9] H. Broersma, Z. Ryjáček, I. Schiermeyer. Closure Concepts: A Survey. *Graphs and*  
 510 *Combinatorics*, 16:17–48, 2000.
- 511 [10] C. Buchanan, B. Du Preez, K.E. Perry, P. Rombach. Toughness of recursively parti-  
 512 tionable graphs. *Theory and Applications of Graphs*, 10(2), Article 4, 2023.
- 513 [11] G.A. Dirac. Some theorems on abstract graphs. *Proceedings of the London Mathemat-*  
 514 *ical Society*, 2:69–81, 1952.
- 515 [12] M. Horňák, A. Marczyk, I. Schiermeyer, M. Woźniak. Dense arbitrarily vertex decom-  
 516 posable graphs. *Graphs and Combinatorics*, 28:807–821, 2012.
- 517 [13] M. Horňák, M. Woźniak. On arbitrarily vertex decomposable trees. *Discrete Mathe-*  
 518 *matics*, 308(7):1268–1281, 2008.

- 519 [14] R. Kalinowski, M. Piłśniak, I. Schiermeyer, M. Woźniak. Dense arbitrarily partition-  
520 able graphs. *Discussiones Mathematicae Graph Theory*, 36:5–22, 2016.
- 521 [15] F. Liu, B. Wu, J. Meng. Arbitrarily partitionable  $\{2K_2, C_4\}$ -free graphs. *Discussiones*  
522 *Mathematicae Graph Theory*, 42:485–500, 2022.
- 523 [16] A. Marczyk. A note on arbitrarily vertex decomposable graphs. *Opuscula Mathemat-*  
524 *ica*, 26(1):109–118, 2006.
- 525 [17] A. Marczyk. An Ore-type condition for arbitrarily vertex decomposable graphs. *Dis-*  
526 *crete Mathematics*, 309:3588–3594, 2009.
- 527 [18] O. Ore. Note on hamilton circuits. *American Mathematical Monthly*, 67:55, 1960.
- 528 [19] M.D. Plummer, A. Saito. Closure and factor-critical graphs. *Discrete Mathematics*,  
529 215:171–179, 2000.
- 530 [20] R. Ravaux. Decomposing trees with large diameter. *Theoretical Computer Science*,  
531 411:3068–3072, 2010.
- 532 [21] W.T. Tutte. The factorization of locally finite graphs. *Canadian Journal of Mathe-*  
533 *matics*, 2:44–49, 1950.