

Existence and Uniqueness for the SQG Vortex-Wave System when the Vorticity is Constant near the Point-Vortex

Dimitri Cobb, Martin Donati, Ludovic Godard-Cadillac

To cite this version:

Dimitri Cobb, Martin Donati, Ludovic Godard-Cadillac. Existence and Uniqueness for the SQG Vortex-Wave System when the Vorticity is Constant near the Point-Vortex. 2024. hal-04370483

HAL Id: hal-04370483 <https://hal.science/hal-04370483>

Preprint submitted on 4 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence and Uniqueness for the SQG Vortex-Wave System when the Vorticity is Constant near the Point-Vortex

Dimitri Cobb,* Martin Donati[†]and Ludovic Godard-Cadillac[‡]

January 4, 2024

Abstract

This article studies the vortex-wave system for the Surface Quasi-Geostrophic equation with parameter $0 < s < 1$. We obtained local existence of classical solutions in $H⁴$ under the standard "plateau hypothesis", H^2 -stability of the solutions, and a blow-up criterion. In the sub-critical case $s > 1/2$ we established global existence of weak solutions. For the critical case $s = 1/2$, we introduced a weaker notion of solution (V-weak solutions) to give a meaning to the equation and prove global existence.

Introduction

The purpose of this work is to study the vortex-wave system associated to (SQG), the surface quasi-geostrophic equation. Equation (SQG) is a partial differential equation set on the plane that is widely used to model atmospheric flows outside the tropical region. It is given by

$$
\frac{\partial}{\partial t}\theta + v \cdot \nabla \theta = 0, \qquad \text{with} \quad v = -\nabla^{\perp}(-\Delta)^{-s}\theta,
$$
\n(SQG)

where $0 \lt s \lt 1$ and where \perp designates the counter-clockwise rotation of angle $\pi/2$ in the plane. In this equation, $\theta : [0, T) \times \mathbb{R}^2 \to \mathbb{R}$ is the potential temperature of the fluid, that will be referred throughout this article as the *active scalar*, and $v : [0, T) \times \mathbb{R}^2 \to \mathbb{R}^2$ is the velocity of the fluid. For the physical content of (SQG), we recommend the two reference books [31, 34]. The second equation in (SQG) is called the quasi-geostrophic Biot-Savart law in reference to the 2D Euler equation written in term of the vorticity ω :

$$
\frac{\partial}{\partial t}\omega + v \cdot \nabla \omega = 0, \qquad \text{with} \quad v = -\nabla^{\perp}(-\Delta)^{-1}\omega. \tag{Euler}
$$

The 2D Euler equation can be seen as the limit case in (SQG) where $s \to 1^-$. The similarity of these two equations gave raise to a wide range of studies trying to investigate whether the special solutions known for 2D Euler have their equivalent for (SQG). The most studied cases are traveling and co-rotating solutions, either from bifurcation arguments [1, 13, 21, 22] or variationnal arguments [6, 19, 20].

However close these two equations may seem, the theory for the Cauchy problem associated to the evolution equation is very different. Existence and uniqueness globally in time for the 2D Euler equation is well-known since the works by Youdovich if the initial vorticity lays in L^{∞} . In this case, the Euler Biot-Savart law gives that the velocity field is log-Lipschitz, so that the Osgood lemma applies (see [36, 37]). For the (SQG) equation, the Biot-Savart law is more singular and such an argument does not apply. The Cauchy theory for (SQG) remains to this day an outstanding problem. The global existence is only known for weak solutions [27], which are nonunique in some cases when the regularity is too low [5]. Numerical simulations [32] suggests that, unlike the 2D Euler equations, the L^{∞} setting does not give raise to global in time strong solutions. Existence and uniqueness on a short interval of time for (SQG) has been proved for the sub-critical and critical case ($s > 1/2$) by [9] and the super-critical case $(0 < s < 1/2)$ has been done by [8] in the H^k setting with $k \geq 4$. The study of the Cauchy problem for these equations is very active, trying to exhibit finite time singularity or to prove global existence in a well-chosen class of regularity (see for instance [11, 14, 23, 26] and references therein).

Nevertheless, one feature of the 2D Euler equation that is preserved by (SQG) is the point-vortex system. This consists in the assumption that at the initial time the active scalar is sharply concentrated around a finite number of points and therefore writes at the limit as a pondered sum of Dirac masses. This model should be understood as a mathematical formalization of the intuitive notion of "centers of whirlpools". These Dirac masses then evolve in time in virtue of the transport equations respectively (SQG) and (Euler). This system is well-posed

^{*}Universit¨at Bonn, Hausdorff Center for Mathematics (HCM), Endenicher Allee 62, D-53115 Bonn, Deutschland.

[†]Université de Lyon, ENS de Lyon, Unité de Mathématiques Pures et Appliquées, 46 allée d'Italie, 69364 Lyon Cedex 07, France. Université de Bordeaux, Bordeaux INP, Institut de Mathématiques de Bordeaux (IMB), UMR 5251, F-33400 Talence, France.

on a given interval of time as long as the distances between vortices remain positive (absence of collapse). Global well-posedness for almost every initial datum has been obtained for the Euler case in the plane by [29], and in bounded smooth domains by [12]. The extension of this result to the quasi-geostrophic point-vortex system is done by [15, 18].

One natural question then lays in the possibility to study a mixed system involving both a regular part for the active-scalar (driven by the PDE) coupled with a singular part made of point-vortices. This coupled system was introduced for the 2D Euler equation independently by [28, 33] and called the "vortex-wave system". Using a Lagrangian formulation of the vortex-wave system, the existence of solutions for this system was proved in presence of a single point-vortex. Uniqueness for this system has been achieved by [28] and [24] respectively for Lagrangian and Eulerian solutions, under the assumption that the point-vortex lays in a region where the vorticity is constant (plateau assumption). It is worth mentioning other works related to the vortex-wave system such as [4, 7, 10, 16, 17, 25, 30, 35].

The main goal of the present paper is to study the vortex-wave system for the Surface Quasi-Geostrophic equation. Our first result is the local existence and uniqueness of a classical solution for the SQG vortex-wave system under the same "plateau assumption" as in [24]. Unlike the Euler case, it is not clear whether this result can be made global since the velocity field is less regular due to the singularity of the Biot-Savart law kernel. We cannot obtain strong solution with θ being only in $L^1 \cap L^{\infty}$ so that we make a much stronger assumption on the initial datum: $\theta_0 \in H^4(\mathbb{R}^2)$. Such a regularity allows us to proceed to energy estimates relying on the commutator structure of the equation. While the energy estimates are comparable to the ones of [8], new difficulties arise due to the presence of the point-vortex. Passing to the limit in a regularized system gives the existence of the solution on a bounded interval of time. We were also able to obtain a uniqueness result with a proof that relies on the point-vortex staying in its initial "plateau", combined with a precise H^2 -stability estimate. We also provide a directional blow-up criterion for the case where the point-vortex reaches the boundary of its "plateau" in finite time. In our work, we mainly focused on the novelties induced by the presence of the point-vortex but we believe it is possible to improve the regularity using some of the refined techniques developed in the literature on SQG (see [23] or [26] for instance).

The second part of the article studies weak solutions (in $L^1 \cap L^{\infty}$) for the SQG vortex-wave system. Since the Youdovitch theory does not apply for SQG, we cannot expect global solutions in $L^1 \cap L^\infty$ to be unique with characteristics that are well-defined. In the case $s > 1/2$ the Biot-Savart law has good regularizing properties, and we can prove global existence. The critical case $s = 1/2$ is more difficult since it is no longer possible to obtain regularity on the velocity field. The Biot-Savart law is here a 0-order pseudo-differential operator. The first consequence is that one must relax the definition of the point-vortex system to apply the Peano theorem. Furthermore, the main difficulty lays in the definition of the velocity field generated by the point-vortex when acting on the smooth part of the active scalar. Indeed, the case $s = 1/2$ implies that the kernel of the Biot-Savart law is no longer L^1_{loc} . To overcome this problem, we introduce a weaker notion of solution, that we called V-weak solutions, adapted to the study of singularities localized on a set of small Hausdorff dimension. We show the existence of a global V-weak solution and study its properties. We show that this weaker notion of solutions shares with the standard notion of weak solutions several important properties. In particular, we prove that if the V-weak solution is \mathcal{C}^1 , then it is a classical solution.

1 Presentation of the Problem and Main Result

1.1 The Vortex-Wave System for SQG

We first define for $0 < s \leq 1$ the function:

$$
\forall x \in \mathbb{R}^2 \setminus \{0\}, \qquad K_s(x) := c_s \frac{x^{\perp}}{|x|^{4-2s}}, \qquad \text{with } c_s := \frac{(1-s)\Gamma(1-s)}{2^{2s-1}\pi\Gamma(s)},
$$
(1.1)

which is the kernel of the Biot-Savart operator $-\nabla^{\perp}(-\Delta)^{-s}$ associated to the Green's function of $(-\Delta)^s$. Remark that $c_s \to \frac{1}{2\pi} =: c_1$ as $s \to 1^-$. The case $s = 1$ corresponds to the case of (Euler) while $0 < s < 1$ corresponds to (SQG). The point-vortex system in the plane is given by the following equations. If z_1, \ldots, z_N are the positions of the point-vortices in the plane and a_1, \ldots, a_N their intensities, then the evolution of the ith vortex with $i = 1, \ldots, N$ is given by:

$$
\frac{\mathrm{d}}{\mathrm{d}t}z_i(t) = c_s \sum_{\substack{j=1 \ j \neq i}}^N a_j \frac{(z_i(t) - z_j(t))^{\perp}}{|z_i(t) - z_j(t)|^{4-2s}}.
$$
\n(1.2)

Indeed, if one plugs an active scalar of the form $\theta(0, x) = \sum_{j=1}^{N} a_j \delta_{x_j}$ in (SQG) or (Euler) then the associated velocity field writes, using (1.1):

$$
v(0,x) := c_s \sum_{j=1}^{N} a_j \frac{(x-z_j)^{\perp}}{|x-z_j|^{4-2s}}.
$$

Since the active scalar is simply transported by the velocity of the fluid, we indeed obtain (1.2) under the natural assumption that a point-vortex does not self-interact.

For this article, we mainly focus on the vortex-wave system in the case of a single vortex moving in time $t \mapsto z(t) \in \mathbb{R}^2$. The case of several point-vortices is similar as long as the distances between the point-vortices remain positive on the studied interval of time, see Section 7. To give a proper definition of the vortex-wave system, the authors in [28] suggest a Lagrangian point-of-view while in [24] they work with an Eulerian pointof-view. In this work we follow the second point-of-view.

We prove the following well-posedness result for the vortex-wave system:

Theorem 1.1. Let $0 < s < 1$. Consider $\theta_0 \in H^k$ an initial datum for some integer $k \geq 4$ and $z_0 \in \mathbb{R}^2$ an initial point-vortex position. Assume that θ_0 is constant on some neighborhood of z_0 (plateau assumption). Then the following assertions hold true.

(i) Local existence and uniqueness: there exists time $T > 0$ such that the PDE system

$$
\begin{cases}\n\partial_t \theta + (v + H) \cdot \nabla \theta = 0, \\
\frac{d}{dt} z(t) = v(t, z(t)), \\
v = -\nabla^{\perp} (-\Delta)^{-s} \theta = K_s \star \theta, \\
H(t, x) = K_s (x - z(t)),\n\end{cases}
$$
\n(1.3)

with initial datum respectively θ_0 and z_0 , has a unique solution $\theta \in L^{\infty}([0,T); H^k(\mathbb{R}^2)) \cap C^1([0,T) \times \mathbb{R}^2)$ and $z \in C^1([0, T); \mathbb{R}^2)$.

(ii) Short time H^2 -stability: let (θ_1, z_1) and (θ_2, z_2) be two solutions in $L^{\infty}([0, T], H^k(\mathbb{R}^2))$ of (1.4) s, with $\theta_1(0)$ and $\theta_2(0)$ locally constant around z_1 and z_2 respectively. Then there exists $\rho > 0$ and $T_0 > 0$ depending only on $\theta_1(0)$ and $\theta_2(0)$ such that if

$$
|z_2(0) - z_1(0)| < \rho,
$$

then, for every $t \in [0, T_0]$, and for every $\ell \in \mathbb{N}$ such that $2 \leq \ell \leq k - 2$,

$$
\|\theta_2(t,\cdot)-\theta_1(t,\cdot)\|_{H^{\ell}}+|z_2(t)-z_1(t)|\lesssim \|\theta_2(0,\cdot)-\theta_1(0,\cdot)\|_{H^{\ell}}+|z_2(0)-z_1(0)|.
$$

(iii) Blow-up criterion: let (θ, z) be a solution of (1.4) on an interval $[0, T^*)$ with $T^* < \infty$. Let

$$
R(t) := \sup \{ r \ge 0 \,, \, \nabla \theta(t) \equiv 0 \text{ on } B(z(t), r) \}
$$
\n
$$
(1.4)
$$

and recall that the plateau hypothesis states that $R(0) > 0$. Assume that $R(t) \longrightarrow_{t \to T^*} 0$. Then

$$
\int_0^{T^*} \|\theta(t)\|_{H^{3-2s}} \mathrm{d}t = +\infty.
$$

Let us formulate a few remarks about Theorem 1.1.

In (1.4), the equation $v = -\nabla^{\perp}(-\Delta)^{-s}\theta$ is well-defined as a convolution with K_s only when $s > 1/2$. Yet it is not always clear whether this convolution is well-defined. When $s \leq 1/2$ we use Fourier multipliers to give a meaning to the velocity v (see Proposition 2.5 hereafter). Note also that for a fixed time $t \in [0, T)$, the product $H\theta$ is a well-defined and measurable function taking finite values for every $x \neq z(t)$. It is also well-defined on all \mathbb{R}^2 when seen as a principal value distribution, because $\theta(t) \in H^k(\mathbb{R}^2) \subset H^4(\mathbb{R}^2)$ and $H(t) \in \dot{B}^{2s-1}_{1,\infty}(\mathbb{R}^2) \subset H^{-2}(\mathbb{R}^2)$. The convolution against the Biot-savart kernel K_s is taken with respect to the space variable $x \in \mathbb{R}^2$. As a convention throughout this article, the gradient operators ∇ and ∇^{\perp} , convolutions and fractional Laplace operators $(-\Delta)^s$ and $(-\Delta)^{-s}$ are always taken with respect to the space variable $x \in \mathbb{R}^2$.

Concerning the regularity of strong solutions, in this work we limit ourselves to $H⁴$ solutions for simplicity reasons and focus on what is new in presence of a point-vortex. However, we believe that both existence and uniqueness results can be extended to lower levels of regularity on θ , according to [23] and [26]. For the stability result, the hypothesis $2 \leq \ell \leq k-2$ is always satisfied at least by $\ell = 2$ since $k \geq 4$. In the case $s \geq 1/2$ we also obtain a L^2 -stability result (see Section 4 for rigorous statement and proof). This L^2 -stability fails in the case $s < 1/2$ for which we need at least to control the H^2 norm of θ .

The system (1.4) can blow-up in two different ways: either the active scalar could lose regularity on its own as in the usual SQG equations, or the point-vortex could collide with the boundary of the constant part of the active scalar, making the system singular. Theorem 1.1-(*iii*) investigates this second type of blow-up. In Section 5, we actually obtain a sharper and directional criterion, which is the following Theorem 1.2, from which we deduce Theorem 1.1-*(iii)*. More precisely, we prove:

Theorem 1.2. Let (θ, z) be a strong solution of (1.4) as given by Theorem 1.1 on an interval $[0, T^*)$, with $T^* < \infty$. Let

$$
N(t) := \max_{\{x \in \mathbb{R}^2, |x - z(t)| = R(t)\}} \frac{-\big(x - z(t)\big) \cdot \big(v(x) - v(z(t))\big)}{R^2(t)(1 - \ln R(t))}.
$$

Then,

$$
R(t) \underset{t \to T^*}{\longrightarrow} 0 \qquad \Longleftrightarrow \qquad \int_0^{T^*} N(t) \mathrm{d}t = +\infty \, .
$$

The quantity $t \mapsto N(t)$ is a weaker version of the log-Lipschitz norm of v that only takes into account the capacity of v to bring fluid particles from the outside of $B(z(t), R(t))$ towards the point-vortex.

On the other hand, in the sub-critical case (namely $s > 1/2$) we have no difficulty in proving the existence of global weak solutions as follows.

Theorem 1.3 (Global weak solutions if $s > 1/2$). We assume that $s > 1/2$. Let $\theta_0 \in L^1 \cap L^\infty(\mathbb{R}^2)$ and $z_0 \in \mathbb{R}^2$. Then there exists $\theta \in L^{\infty}(\mathbb{R}_+; L^1 \cap L^{\infty})$ and $z \in C^0(\mathbb{R}_+; \mathbb{R}^2)$ such that (θ, z) is a weak solution of the vortex-wave system with initial datum (θ_0, z_0) . More precisely, for any test function $\psi \in \mathcal{D}([0, T) \times \mathbb{R}^2)$, we have

$$
\int_0^T \int_{\mathbb{R}^2} \theta(t, x) \left(\frac{\partial \psi}{\partial t}(t, x) + (v + H) \cdot \nabla \psi(t, x) \right) dx dt + \int_{\mathbb{R}^2} \theta_0(x) \psi(0, x) dx = 0,
$$
\n(1.5)

where v and H are given by (1.4) and where z is a solution of the integral equation

$$
z(t) = z_0 + \int_0^t v(\tau, z(\tau)) \, \mathrm{d}\tau. \tag{1.6}
$$

Moreover, this solution satisfies $\|\theta(t, \cdot)\|_{L^p} \le \|\theta_0\|_{L^p}$ for all $p \in [1, +\infty]$. Finally, the function θ is continuous in time with values in $L^1 \cap L^\infty(\mathbb{R}^2)$ equipped with the weak- \star topology.

1.2 Global Existence of Weak Solutions in the Critical Case

From the result given by Theorem 1.3, one naturally wonders what happens in the critical case where $s = 1/2$.

1.2.1 Duhamel Formula for the Quasi-Point-Vortex System

One important difference in the case $s = 1/2$ is that the function K_s ceases to be locally L^1 . Moreover, the convolution with K_s is now a Fourier multiplier of order 0 since such convolution correspond to the differential operator $-\nabla^{\perp}(-\Delta)^s$. As a consequence, there is no hope for the velocity $v := K_s \star \theta$ to be a continuous function with θ being only in $L^1 \cap L^\infty$. To give a meaning to the vortex-wave system when $s = 1/2$, we have to weaken the notion of point-vortex to obtain the regularity needed to define the flow. For that purpose, we rewrite the equation on the evolution of the point-vortex (1.22) as follows:

$$
z(t) = z_0 + \int_0^t \int_{\mathbb{R}^2} v(t', x) \, \delta_0(x - z(t')) \, dx \, dt', \tag{1.7}
$$

where δ_0 is the centered Dirac mass. In the view of such a formulation we introduce the notion of quasi pointvortex, which is a relaxation of the equation above:

$$
z(t) = z_0 + \int_0^t \int_{\mathbb{R}^2} v(t', x) \, \chi(x - z(t')) \, dx \, dt', \tag{1.8}
$$

where χ is a fixed L^1 function such that $\int \chi = 1$. This function χ must be understood as an approximation of the Dirac mass. In other words, the computation of the velocity for the point-vortex is not the simple evaluation of the velocity field v at one point z (which is ill-defined) but the computation of an average of v on a "small" neighborhood of z. The main weakness of this approach is that the solution depends on the choice of $\chi \in L^1$ and it is unclear whether one can pass to the limit $\chi \to \delta_0$. Yet this relaxation (1.14) is the most simple and reasonable way to give a meaning to (1.13).

1.2.2 A Remark on the Structure of the Non-Linearity

Another problem arising in the critical case $s = 1/2$ is to give a meaning for almost every $t \ge 0$ to the following integrals appearing in (1.11):

$$
\int_{\mathbb{R}^2} \theta(t, x) v(t, x) \cdot \nabla \psi(t, x) dx \quad \text{and} \quad \int_{\mathbb{R}^2} \theta(t, x) H(t, x) \cdot \nabla \psi(t, x) dx.
$$
\n(1.9)

Concerning the first one, we simply follow the idea developed in [27] which consists in using the commutator structure of this term to regain some regularity on θ . This idea is reminiscent from previous works on SQG.

Lemma 1.4 (Commutator formulation). Let $\theta \in C^1$ and $\varphi \in \mathcal{D}(\mathbb{R}^2)$. We have:

$$
\int_{\mathbb{R}^2} \theta(x) \, K_s \star \theta(x) \cdot \nabla \varphi(x) \, dx = \frac{1}{2} \int_{\mathbb{R}^2} K_s \star \theta(x) \cdot \left[(-\Delta)^s, \nabla \varphi \right] (-\Delta)^{-s} \theta(x) \, dx,\tag{1.10}
$$

where $[A, B] := AB - BA$ is the commutator between A and B.

This lemma is proved later on (Section 2.2). It is proved in [27], using this commutator formulation, that this integral makes sense in L^p for $4/3 < p < 2$ (see Lemma 6.1 hereafter).

1.2.3 A Weaker Notion of Weak Solution

The remaining difficulty is the second integral in (1.15), which is more difficult to treat since $H = -\nabla^{\perp}(-\Delta)^{s}\delta_{z}$, which more singular than $v := -\nabla^{\perp}(-\Delta)^s \theta$. One possible idea would have been to regularize $\delta_0(x - z(t))$ in the same spirit than (1.14). Nevertheless, such an approach would have made loose the information on the deformation of θ in the neighborhood of the singularity of a point-vortex.

To pass-by this difficulty, we choose to introduce a weaker notion of solution that is weaker than the standard notion of weak solution. We first recall that, in a general formulation, the standard notion of weak solution writes:

$$
u \in \mathcal{D}'(\Omega; \mathbb{R}^d)
$$
 is a weak solution iff: $\forall \phi \in \mathcal{D}(\Omega; \mathbb{R}^d)$, $\mathcal{F}(u, \phi) = 0$ (1.11)

where $\mathcal{F}: \mathcal{D}' \times \mathcal{D} \to \mathbb{R}$ is the function that represent the weak formulation of the studied equation (the function $u \mapsto \mathcal{F}(u, \phi)$ may be well-defined only on a subspace of \mathcal{D}' .

Definition 1.5 (*V*-weak solutions). Let Ω be a domain and $V \subseteq \mathcal{D}(\Omega;\mathbb{R}^d)$. Then

 $u \in \mathcal{D}'(\Omega;\mathbb{R}^d)$ is a V-weak solution to (1.17) iff: $\forall \phi \in V$, $\mathcal{F}(u,\phi) = 0$.

In other words, we are not able to have the weak formulation (1.17) for all test functions $\phi \in \mathcal{D}(\Omega;\mathbb{R}^d)$ but only for functions in a subset V. To have this notion of V-weak solution be relevant, this set V must be the biggest possible. Whenever we manipulate it, we systematically study how big the set $V \subseteq \mathcal{D}(\Omega;\mathbb{R}^d)$ can be to ensure that we remain close to the standard notion of weak solution. Note that when u and $\mathcal F$ are smooth enough, the two notion coincide whenever V is dense in $\mathcal{D}(\Omega;\mathbb{R}^d)$. To measure how much the set V coincide with $\mathcal{D}(\Omega;\mathbb{R}^d)$, we introduce the notion of Γ -coincidences for sets of functions:

Definition 1.6 (Γ-coincidence). Let V and W two sets of functions $\Omega \to \mathbb{R}^d$ and let $\Gamma \subseteq \Omega$. We say that the two sets V and W are Γ-coincident if they coincide except on Γ. More precisely, for all $\Omega_0 \subseteq \Omega$ such that the distance between Ω_0 and Γ is positive:

$$
\forall\,f\in V,\ \, \exists\;g\in W,\quad \left.f\right|_{_{\Omega_0}}\equiv g_{\big|_{_{\Omega_0}}}\qquad\text{and symmetrically}:\qquad\forall\;g\in W,\;\;\exists\;f\in V,\quad \left.f\right|_{_{\Omega_0}}\equiv g_{\big|_{_{\Omega_0}}}.
$$

This notion of coincidence is adapted to the study of singularities (choosing Γ being the set where the singularity is supported). This definition can be interpreted intuitively as follows: as soon as we lay at positive distance from the singularity supported on Γ then the notion of V-weak solution coincides with the standard notion of weak solution (taking $W = \mathcal{D}(\Omega; \mathbb{R}^d)$ in the definition above).

To better sustain the relevancy of this notion of V -weak solution, but also to give a better intuition, we mention the following easy property:

Property 1.7. let $u \in \mathcal{D}'(\Omega;\mathbb{R}^d)$ be a V-weak solution of some equation (see definition 1.5), with V being Γ -coincident with $\mathcal{D}(\Omega : \mathbb{R}^d)$ for some $\Gamma \subseteq \Omega$.

Then u is a solution in the sense of distributions in $\Omega \setminus \Gamma$:

 $u \in \mathcal{D}'(\Omega \setminus \Gamma;\mathbb{R}^d)$ is such that: $\forall \phi \in \mathcal{D}(\Omega \setminus \Gamma;\mathbb{R}^d)$, $\mathcal{F}(u,\phi) = 0$.

In other words, we have the following hierarchy :

solution in $\mathcal{D}'(\Omega) \implies V$ -weak solution (Γ-coincident) \implies solution in \mathcal{D}' $\prime'(\Omega \setminus \Gamma).$

Note also that $\mathcal{D}(\Omega \setminus \Gamma)$ is Γ-coincident with $\mathcal{D}(\Omega)$. Nevertheless, in practice the set V should be chosen larger so that the least possible information is lost in the neighborhood of the singularity.

1.2.4 Global Existence of V-Weak Solutions in the Critical Case

With this weaker notion of solution at hand and this commutator formulation, we are able to prove the following theorem:

Theorem 1.8 (Global V-weak solutions to the *quasi* vortex-wave if $s = 1/2$). Assume that $s = 1/2$. Let $\theta_0 : \mathbb{R}^2 \to \mathbb{R}, \text{ let } z_0 \in \mathbb{R}^2 \text{ and let } T > 0.$ Let $\chi \in L^1(\mathbb{R}^2)$ such that $\int \chi = 1$. Then there exists $V \subseteq \mathcal{D}([0, T) \times \mathbb{R}^2)$ such that:

(i) There exists $\theta \in L^{\infty}([0,T); L^{1} \cap L^{\infty}(\mathbb{R}^{2}))$ and $z \in C^{0,1}([0,T); \mathbb{R}^{2})$ a V-weak solution to the vortex-wave system in the sense that for all test functions $\psi \in V$:

$$
\int_0^T \int_{\mathbb{R}^2} \left(\frac{\partial \psi}{\partial t}(t, x) + v(t, x) \cdot \left[(-\Delta)^s, \nabla \psi(t, x) \right] (-\Delta)^{-s} \right. \\
\left. + H(t, x) \cdot \nabla \psi(t, x) \right) \theta(t, x) dx dt + \int_{\mathbb{R}^2} \omega_0(x) \psi(0, x) dx = 0,
$$
\n(1.12)

where v and H are given by (1.4), and we have for all $t \in [0, T)$,

$$
z(t) = z_0 + \int_0^t \int_{\mathbb{R}^2} v(t', x) \chi(x - z(t')) \, dx \, dt'.
$$
 (1.13)

(ii) The set V is such that the product $H \cdot \nabla \psi$ lays in $L^1(\mathbb{R}^2)$. Moreover, the set V is dense in $L^p(\mathbb{R}^2)$ for all $p \in [1, +\infty]$, dense in $W^{1,p}(\mathbb{R}^2)$ for all $p < +\infty$.

(iii) The set V is Γ -coincident with $\mathcal{D}([0,T) \times \mathbb{R}^2)$ for Γ the trajectory of the point-vortex:

$$
\Gamma \, := \, \bigcup_{t \in [0,T)} \big\{ (t,z(t)) \big\} \quad \subseteq \, [0,T) \times \mathbb{R}^2.
$$

(iv) If moreover a solution θ of (1.21) has regularity \mathcal{C}^1 on $[0,T) \times \mathbb{R}^2$ and is constant on a neighborhood of $z(t)$ for all $t \in [0,T)$, then (θ, z) is a classical solution on $[0,T)$ to the the quasi vortex-wave with $s = 1/2$.

Point *(iv)* of this theorem is a main and standard property for any weak solution of any equation. This property can be seen as one of the main interest of the notion of weak solution because the notion of weak solution makes sense as long as we can reconstruct a classical solution whenever the weak solution turns out to be smooth enough. What we prove at Point (iv) of this theorem is that this important property is also satisfied by our notion of V -weak solutions. The fact that the weak and V -weak solutions share this property is the main argument in favor of this new weaker notion of solution.

Concerning Point (iii) of the theorem, this property of Γ-coincidence means, from an intuitive point-of-view, that the solution is a distribution "outside" the singularity (meaning at positive distance from the point-vortex) and is not defined "inside" the singularity.

2 Functional Analysis and Regularization

2.1 Fourier Analysis and Besov spaces

We first define a very convenient tool for Fourier analysis in a context of non-linear equations which is the Littlewood-Paley analysis. For a more detailed presentation, see [3]. Recall that the Fourier transform of $f:\mathbb{R}^2\to\mathbb{R}$ is defined by

$$
\widehat{f}(\xi) = \mathcal{F}f(\xi) := \int_{\mathbb{R}^2} f(x)e^{-ix\cdot\xi} dx
$$

Definition 2.1 (Littlewood-Paley decomposition). We consider a nonnegative function $\phi \in \mathcal{D}(\mathbb{R}^2)$ that is radial and such that $\phi(\xi) = 1$ if $|\xi| \le 1/2$ and $\phi(\xi) = 0$ if $|\xi| \ge 1$. We define $\psi(\xi) := \phi(\xi/2) - \phi(\xi)$, and we call this function the Fourier cut-off function.

We then define the j-th homogeneous dyadic block of the Littlewood-Paley decomposition for a given tempered distribution $f \in \mathcal{S}'(\mathbb{R}^2)$ by

$$
\dot{\Delta}_j f := \mathcal{F}^{-1}\bigg(\psi\Big(\frac{\xi}{2^j}\Big)\,\mathcal{F}(f)\bigg).
$$

We also define the non-homogenous dyadic blocks $(\triangle_j)_{j\geq -1}$ in the following way: when $j\geq 0$, we set $\triangle_j = \dot{\triangle}_j$, and for $j = -1$, we note

$$
\Delta_{-1}f = \mathcal{F}^{-1}\left(\phi(\xi)\mathcal{F}(f)\right). \tag{2.1}
$$

With the cut-off function ϕ , we define a low frequencies cutting operator by setting, for $j \in \mathbb{Z}$

$$
S_j f := \mathcal{F}[\phi(2^{-j}\xi)\mathcal{F}(f)].\tag{2.2}
$$

We also define a *high frequencies cutting operator* by:

$$
\mathcal{H}_j f := (1 - \mathcal{S}_j) f = \sum_{k=j}^{+\infty} \dot{\Delta}_k f.
$$

With the Littlewood-Paley decomposition, it is possible to define the Besov spaces:

Definition 2.2 (Inhomogeneous Besov spaces). For $s \in \mathbb{R}$, and $p, q \in [1, +\infty]$, the distribution $f \in \mathcal{S}'$ belongs to the inhomogeneous Besov space $B_{p,q}^s$ if and only if the following Besov norm is finite:

$$
||f||_{B_{p,q}^s} := ||\Delta_{-1}f||_{L^p} + \bigg(\sum_{j=0}^{+\infty} 2^{jqs} ||\Delta_j f||_{L^p}^q \bigg)^{\frac{1}{q}}.
$$

With this norm, the space $B_{p,q}^s$ is a Banach space.

Definition 2.3 (Homogeneous Besov spaces). For $s \in \mathbb{R}$, and $p, q \in [1, +\infty]$, the distribution $f \in \mathcal{S}'$ belongs to the homogeneous Besov space $\dot{B}_{p,q}^s$ if and only if the following Besov semi-norm is finite:

$$
\|f\|_{\dot{B}^{s}_{p,q}}\;:=\bigg(\sum_{j=-\infty}^{+\infty}2^{jqs}\,\|\dot{\bigtriangleup}_jf\|_{L^{p}}^{q}\bigg)^{\frac{1}{q}}.
$$

It is standard to prove that $B_{2,2}^s = H^s$ for $s \in \mathbb{R}$ and that $\dot{B}_{2,2}^s = \dot{H}^s$ in the case $s < 1$. One of the most useful properties of homogeneous spaces is their scaling invariance: if $f \in \dot{B}_{p,r}^s$ and $f_\lambda(x) = f(\lambda x)$, then we have $||f_\lambda||_{\dot{B}_{p,r}^s} \approx \lambda^{s-2/p} ||f||_{\dot{B}_{p,r}^s}.$

We will also need a Bernstein inequality: if a dyadic block Δf belongs to L^p , then it belongs to all L^q spaces for $q \geq p$, but the price to pay is a power of 2^j , which is roughly the regularity scale of $\Delta_j f$. This is formalized in the following Lemma.

Lemma 2.4 (see Lemma 2.1, p. 52, in [3]). Consider $1 \le p \le q \le +\infty$, $j \in \mathbb{Z}$, and $f \in \mathcal{S}'$ such that $\Delta_j f \in L^p$. Then the following inequality holds:

$$
\|\dot{\triangle}_j f\|_{L^q} \le C(d) 2^{jd\left(\frac{1}{p} - \frac{1}{q}\right)} \|\dot{\triangle}_j f\|_{L^p}.
$$

We are able to give a meaning to the convolution with the kernel $K_s: x \mapsto x^{\perp}/|x|^{4-2s}$ for all value of s provided that the function θ is H^1 :

Proposition 2.5. Consider $s \in (0,1)$. Then the operator $\nabla^{\perp}(-\Delta)^{-s}$ defines a bounded Fourier multiplier $H^{1-2s} \longrightarrow L^{\infty} + L^2$. In other words, the function

$$
\widehat{g}(\xi) = i\xi^{\perp}|\xi|^{-2s}\widehat{\theta}(\xi)
$$

is the Fourier transform a function $g = \Delta_{-1}g + (\text{Id} - \Delta_{-1})g$ with

$$
\|\triangle_{-1}g\|_{L^{\infty}} + \big\|(\mathrm{Id} - \triangle_{-1})g\big\|_{L^2} \lesssim \|\theta\|_{H^{1-2s}}.
$$

Proof. We split the function q into two parts, a low frequency one and a high frequency one: we have, by making use of the small frequencies cutting operator of (2.3),

$$
\begin{aligned}\n\widehat{g}(\xi) &= \mathcal{F}[\triangle_{-1}g](\xi) + \mathcal{F}[(\mathrm{Id} - \triangle_{-1})g](\xi) \\
&= \phi(\xi)i\xi^{\perp}|\xi|^{-2s}\widehat{\theta}(\xi) + (1 - \phi(\xi))i\xi^{\perp}|\xi|^{-2s}\widehat{\theta}(\xi) := \widehat{g_1}(\xi) + \widehat{g_1}(\xi),\n\end{aligned}
$$

where $\phi \in \mathcal{D}$ is the Littlewood-Paley function from Definition 2.1. Let us focus on the low frequency part, which will give a L^{∞} function: we have

$$
||g_1||_{L^{\infty}} \lesssim ||\widehat{g_1}||_{L^1} \lesssim \int \phi(\xi)|\xi|^{1-2s}|\widehat{\theta}(\xi)|d\xi.
$$

Now, the Cauchy-Schwarz inequality provides, as ϕ is supported in a ball $B(0, 1)$,

$$
\int \phi(\xi)|\xi|^{1-2s}|\widehat{\theta}(\xi)|\mathrm{d}\xi \lesssim \|\widehat{\theta}\|_{L^2} \left(\int_0^1 r^{2(1-2s)} r \mathrm{d}r\right)^{1/2} \lesssim \|\theta\|_{L^2}.
$$

On the other hand, the high frequency term reads

$$
\widehat{g_2}(\xi) = (1 - \phi(\xi)) i\xi^{\perp} |\xi|^{-2s} \widehat{\theta}(\xi) = (1 - \phi(\xi)) \frac{i\xi^{\perp} |\xi|^{-2s}}{1 + |\xi|^{1-2s}} (1 + |\xi|^{1-2s}) \widehat{\theta}(\xi),
$$

so that we do indeed have $||g_2||_{L^2} \le ||\theta||_{H^{1-2s}}$.

In the particular case where $s = 1/2$, it is possible to give a meaning to the convolution with $K_{\frac{1}{2}}$ when θ is only in L^1 or L^2 (which will be usefull for the study of weak solutions:

Lemma 2.6. The following estimates hold:

$$
\|K_{\frac12}\star \theta\|_{L^1}\lesssim \|\theta\|_{L^1},\qquad and \qquad \|K_{\frac12}\star \theta\|_{L^2}\lesssim \|\theta\|_{L^2}
$$

Proof. The Fourier Multiplyer associated to the convolution with $K_{\frac{1}{2}}$ is the symbol $\xi^{\perp}/|\xi|$, which is bounded by 1. This gives the L^1 estimate above using Directly the definition of the Fourier transform. The L^2 estimate is obtained similarly, using the Fourier-Plancherel formula. \Box

2.2 Commutator formulation

To start-with, we prove the commutator formulation stated by Lemma 1.4:

Proof. On the one hand we simply write

$$
I := \int_{\mathbb{R}^2} \theta(x) \, K_s \star \theta(x) \cdot \nabla \varphi(x) \, dx = \int_{\mathbb{R}^2} K_s \star \theta(x) \cdot \nabla \varphi(x) \, (-\Delta)^s (-\Delta)^{-s} \theta(x) \, dx \tag{2.3}
$$

On the other hand, recalling that $K_s \star = \nabla^{\perp}(-\Delta)^{-s}$, an integration by part gives,

$$
I = -\int_{\mathbb{R}^2} (-\Delta)^{-s} \theta(x) \nabla^{\perp} \cdot (\theta \nabla \varphi)(x) dx = -\int_{\mathbb{R}^2} (-\Delta)^{-s} \theta(x) \nabla^{\perp} \theta(x) \cdot \nabla \varphi(x) dx,
$$

where for the second equality we used the Schwarz theorem that states $\nabla^{\perp} \cdot \nabla \phi = 0$. We now remark that $(-\Delta)^{-s}$ is a convolution operator. Writting $1 = (-\Delta)^{-s}(-\Delta)^s$, The Fubini theorem then gives:

$$
I = -\int_{\mathbb{R}^2} \nabla^{\perp} \theta(x) \cdot (-\Delta)^{-s} (-\Delta)^s \Big((-\Delta)^{-s} \theta(x) \nabla \varphi \Big) (x) dx = -\int_{\mathbb{R}^2} K_s \star \theta(x) \cdot (-\Delta)^s \Big((-\Delta)^{-s} \theta \nabla \varphi \Big) (x) dx \tag{2.4}
$$

Summing (2.16) and (2.18) gives (1.16).

From this reformulation of the non-linear term as a commutator, we have the following properties that are classical for the analysis of SQG (see given references for proofs). We make widely use of these results throughout the article:

Lemma 2.7 (lemma 2.2 in [27]). For $s \ge 1/2$, define the singular integral operator $J := [(-\Delta)^s, \nabla \phi]$. For $1 < p < +\infty$, there exists $C_p \geq 0$ such that

$$
||J(f)||_{L^p} \leq C_p ||\nabla^2 \phi||_{L^\infty} ||f||_{L^p}.
$$

Lemma 2.8 (See Proposition 2.1 in [8]). Consider $s \in \mathbb{R}$, $j \in \{1,2\}$ and the operator $A_j = \partial_j(-\Delta)^{-s}$. For any $\epsilon > 0$ the bound

$$
\left\|[A_j,g]f\right\|_{L^2} \leq C(s)\|f\|_{L^2}\left\|(-\widehat{\Delta)^{(1-2s)/2}g}\right\|_{L^1} + C(s)\|(-\Delta)^{-s}f\|_{L^2}\left\|(-\widehat{\Delta)^{1/2}g}\right\|_{L^1}.
$$

holds as long as all the norms above make sense (and are finite).

2.3 Regularization of the Kernel

In this paragraph, we define the regularization $K_{s,\epsilon}$ of the kernel K_s that we will be using throughout the article, as well as the approximate system that will serve to construct solutions.

First of all, we fix a smooth cut-off function $\chi \in C^{\infty}$ such that $0 \leq \chi \leq 1$ and

$$
\chi(x) = 1
$$
 for $|x| \le \frac{1}{2}$, $\chi(-x) = \chi(x)$ and $\text{supp}(\chi) \subset B(0, 1)$.

For any radius $\epsilon > 0$, we also define a scaled version of this cut-off

$$
\chi_{\epsilon}(x) = \chi\left(\frac{x}{\epsilon}\right).
$$

This allows to truncate smoothly the kernel K_s , and so get rid of the singularity. We define the truncated kernel $K_{s,\epsilon}$ by

$$
K_{s,\epsilon}(x) := (1 - \chi_{\epsilon}(x)) K_s(x),
$$

so that $K_{s,\epsilon}(x) = K_s(x)$ for all $|x| \geq \epsilon$. We also have the inequality

$$
|K_{s,\varepsilon}(x)| \le |K_s(x)| = \frac{c_s}{|x|^{3-2s}}.\tag{2.5}
$$

In particular, the convergence $K_{s,\epsilon} \longrightarrow K_s$ holds uniformly locally in $\mathbb{R}^2 \setminus \{0\}$ as the truncation parameter goes to zero $\epsilon \to 0^+$ (see also Lemma 3.7). Also note that $K_{s,\epsilon}$ is a smooth odd function $K_{s,\epsilon}(x) = -K_{s,\epsilon}(-x)$.

With the help of the kernel $K_{s,\epsilon}$, we may define an approximate system, for which the existence of solutions is easier. More precisely, we consider the following set of PDEs

$$
\begin{cases} \partial_t \theta_{\epsilon} + \text{div} \left((v_{\epsilon} + \mathcal{H}_{\epsilon}) \theta_{\epsilon} \right) = 0 \\ \frac{d}{dt} z_{\epsilon}(t) = v_{\epsilon} \left(t, z_{\epsilon}(t) \right) \\ v_{\epsilon} = K_{s,\epsilon} \star \theta_{\epsilon} \\ \mathcal{H}_{\epsilon}(t, x) = K_{s,\epsilon} \left(x - z_{\epsilon}(t) \right), \end{cases} \tag{2.6}
$$

which we equip with the initial datum

$$
\theta_{\epsilon}(0) = \theta_{0,\epsilon} := \chi_{1/\epsilon} \cdot (S_{j_{\epsilon}} \theta_0),\tag{2.7}
$$

where S_j is the Littlewood-Paley approximation operator (2.4) and $j_{\epsilon} \geq 1/\epsilon$. In this way, the initial datum is smooth and compactly supported.

Since the velocity field v_{ε} is divergence-free $(\nabla \cdot \nabla^{\perp} = 0$ by the Stokes theorem), the Liouville theorem [2] states that for all measurable set $X \subseteq \mathbb{R}^2$:

$$
\mathcal{L}^2(X) = \mathcal{L}^2(S^t_{\varepsilon}X),
$$

where \mathcal{L}^2 is the Lebesgue measure on \mathbb{R}^2 and $t \geq 0 \mapsto S^t$ is the flow of the equation. The active scalar being transported by the flow, we deduce the following conservation property for all $p \in [1, +\infty]$:

$$
\forall t > 0, \qquad \|\theta_{\varepsilon}(t,.)\|_{L^p} = \|\theta_0\|_{L^p}.
$$
\n(2.8)

Before examining the existence of classical solutions, we first show that the approximate system (2.24) has global (approximate) solutions.

Proposition 2.9. Consider any $\varepsilon > 0$ and the initial data $(\theta_{0,\varepsilon}, z_0)$ defined by (2.25). Then the approximate system has a global smooth solution $(\theta_{\epsilon}, z_{\epsilon})$.

Proof. Since the approximate system (2.24) is fully non-linear PDE system, it is not immediately obvious that it has global solutions. To construct these, we resort to a second approximation: define for all $N \geq 1$ the operator E_N by its Fourier transform

$$
\forall f \in L^2, \qquad \widehat{E_N f}(\xi) := \mathbb{1}_{|\xi| \le N} \widehat{f}(\xi)
$$

and consider the system of equations

$$
\begin{cases}\n\partial_t \theta_N + E_N \operatorname{div} \left((v_N + \mathcal{H}_N) \theta_N \right) = 0 \\
\frac{d}{dt} z_N(t) = v_N(t, z_N(t)) \\
v_N = K_{s,\varepsilon} \star \theta_N \\
\mathcal{H}_N(t, x) = K_{s,\varepsilon} (x - z_N(t)).\n\end{cases}
$$
\n(2.9)

We associate to (2.29) the initial data

$$
(\theta_N(0), z_0) := (E_N \theta_{0,\varepsilon}, z_0)
$$

with the perspective of using the Cauchy-Lipschitz theorem to construct a solution (θ_N, z_N) . With that in mind, we define the Banach space

$$
X_N := \Big\{ (\theta, z) \in L^2 \times \mathbb{R}^2, \quad \text{supp}(\widehat{f}) \subset B(0, N) \Big\},
$$

which we endow with the norm

$$
\big\|(\theta,z)\big\|_{X_N}:=\|\theta\|_{L^2}+|z|.
$$

In order to apply the Cauchy-Lipschitz theorem, we must check that the map

$$
F: \left(\begin{array}{c} \theta \\ z \end{array}\right) \longmapsto \left(\begin{array}{c} E_N \operatorname{div} \left(\left(\theta \star K_{s,\varepsilon} + K_{s,\varepsilon} (x-z) \right) . \theta \right) \\ K_{s,\varepsilon} \star \theta(t,z) \end{array}\right)
$$

is locally Lipschitz on $X_N \longrightarrow X_N$. We therefore consider two elements (θ_1, z_1) and (θ_2, z_2) of X_N and study the difference $||F(\theta_2, z_2) - F(\theta_1, z_1)||_{X_N}$. We have

$$
||F(\theta_2, z_2) - F(\theta_1, z_1)||_{X_N} \lesssim N ||(\theta_2 - \theta_1) \star K_{s,\varepsilon} \cdot \theta_2||_{L^2} + N ||\theta_1 \star K_{s,\varepsilon} \cdot (\theta_2 - \theta_1)||_{L^2} + N ||(K_{s,\varepsilon}(x - z_2) - K_{s,\varepsilon}(x - z_1)).\theta_2||_{L^2} + N ||K_{s,\varepsilon}(x - z_1).(\theta_2 - \theta_1)||_{L^2} + |K_{s,\varepsilon} \star \theta_2(t, z_2) - K_{s,\varepsilon} \star \theta_2(t, z_1)| + |K_{s,\varepsilon} \star \theta_2(t, z_1) - K_{s,\varepsilon} \star \theta_1(t, z_1)| := \sum_{k=1}^6 M_k.
$$

We start by noting that the approximate kernel $K_{s,\varepsilon}$ is a L^2 function for any value of $\varepsilon > 0$. Therefore, we bound the first two terms M_1 and M_2 by

$$
M_1 + M_2 \lesssim N \| (\theta_2 - \theta_1) \star K_{s,\varepsilon} \|_{L^\infty} \| \theta_2 \|_{L^2} + N \| \theta_1 \star K_{s,\varepsilon} \|_{L^\infty} \| \theta_2 - \theta_1 \|_{L^2}
$$

$$
\lesssim 2N \| K_{s,\varepsilon} \|_{L^2} \| \theta_2 \|_{L^2} \| \theta_2 - \theta_1 \|_{L^2}.
$$

For the next two terms, we note that the kernel $K_{s,\varepsilon}$ also is a bounded Lipschitz function. This implies the following bounds for M_3 and M_4 ,

$$
M_3+M_4\lesssim N\|\nabla K_{s,\varepsilon}\|_{L^\infty}|z_2-z_1|\|\theta_2\|_{L^2}+N\|K_{s,\varepsilon}\|_{L^\infty}\|\theta_2-\theta_1\|_{L^2}.
$$

Finally, for the last two terms, M_5 and M_6 , we also use the fat that $K_{s,\varepsilon} \in W^{1,\infty}$ to write

$$
\begin{aligned} M_5 + M_6 &\lesssim \|K_{s,\varepsilon} \star \theta_2\|_{L^\infty} |z_2 - z_1| + \|K_{s,\varepsilon} \star (\theta_2 - \theta_1)\|_{L^\infty} \\ &\lesssim \|\nabla K_{s,\varepsilon}\|_{L^2} \|\theta_2\|_{L^2} |z_2 - z_1| + \|K_{s,\varepsilon}\|_{L^2} \|\theta_2 - \theta_1\|_{L^2} .\end{aligned}
$$

Putting together these three estimates, we see that

$$
\big\|F(\theta_2,z_2)-F(\theta_1,z_1)\big\|_{X_N}\lesssim C(s,\varepsilon,N)\|\theta_2\|_{L^2}\big(\|\theta_2-\theta_1\|_{L^2}+|z_2-z_1|\big),
$$

so that the map is indeed $X_N \longrightarrow X_N$ Lipschitz. Accordingly, the Cauchy-Lipschitz theorem provides the existence of a (unique) maximal solution

$$
(\theta_N,z_N): (T_N^-,T_N^+) \longrightarrow X_N.
$$

We now prove that this solutions is global, or, since we only are interested in non-negative times, that T_N^+ = + ∞ . For this, we remark that the conservation of L^2 norms also applies to solutions of (2.29), since the approximate kernel $K_{s,\varepsilon}$ is divergence free: this follows from the relation

$$
\operatorname{div}(K_{s,\varepsilon}) = -\nabla \chi_{\varepsilon} \cdot K_s + (1 - \chi_{\varepsilon}) \operatorname{div}(K_s)
$$

in which the first term $\nabla \chi_{\varepsilon} \cdot K_s$ is zero because χ_{ε} is, by choice, a radial function, whereas $K_s(x)$ is colinear to x^{\perp} . Consequently, the vector field $v_N + \mathcal{H}_N$ is divergence-free and

$$
\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int |\theta_N(t)|^2 \mathrm{d}x = 0,
$$

and this implies that there is conservation of L^2 norms $\|\theta_N(t)\|_{L^2} = \|\theta_N(0)\|_{L^2} \le \|\theta_0\|_{L^2}$. Furthermore, the approximate velocity v_N also satisfies the bound

$$
||v_N||_{W^{1,\infty}} \lesssim ||K_{s,\varepsilon}||_{H^1} ||\theta_N||_{L^2}.
$$

Using this in the differential equation satisfied by the particle trajectory, we see that

$$
\left|\frac{\mathrm{d}}{\mathrm{d}t}z_N\right| \lesssim \|K_{s,\varepsilon}\|_{H^1} \|\theta_N\|_{L^2}|z_N| \leq C(s,\varepsilon) \|\theta_0\|_{L^2}|z_N|.
$$

Grönwall's inequality then gives the global estimate

$$
|z_N(t)| \le (||\theta_0||_{L^2} + |z_0|)e^{C(s,\varepsilon)t||\theta_0||_{L^2}}.
$$
\n(2.10)

Note that the upper bound in this inequality is independent on N . In particular, we deduce that the approximate solutions (θ_N, z_N) remain bounded in the space X_N , so that they cannot blow-up in finite time and $T_N^+ = +\infty$.

Now, in order to construct the solution $(\theta_{\varepsilon}, z_{\varepsilon})$, we prepare to take the limit $N \to +\infty$. From estimate (2.44), we already know that the sequence $(\theta_N, z_N)_N$ is uniformly bounded in $L^\infty(\mathbb{L}^2) \times L^\infty_{\text{loc}}$ with respect to the approximation parameter $N \geq 1$. Consequently, since the kernel $K_{s,\varepsilon}$ lies in H^{ℓ} for any $\ell \in \mathbb{R}$, we see that, for any fixed time $T > 0$,

$$
\|\partial_t \theta_N\|_{L_T^{\infty}(H^{-1})} \le \|v_N + \mathcal{H}_N\|_{L^{\infty}} \|\theta_N\|_{L^2}
$$

\n
$$
\le (\|K_{s,\varepsilon}\|_{L^{\infty}} + \|K_{s,\varepsilon}\|_{L^2} \|\theta_N\|_{L^2}) \|\theta_N\|_{L^2}
$$

\n
$$
\le C(s,\varepsilon) (\|\theta_0\|_{L^2}^2 + \|\theta_0\|_{L^2}). \tag{2.11}
$$

The sequence (θ_N) is then bounded in the space $W^{1,\infty}_{\text{loc}}(H^{-1})$. We deduce the strong convergence of the sequence (up to an extraction),

$$
\theta_N \underset{N \to +\infty}{\longrightarrow} \theta_{\varepsilon} \quad \text{in } C_T^0(H_{\text{loc}}^{-1-\delta})
$$

for any $\delta > 0$. In particular, the regularity of the approximate kernel $K_{s,\varepsilon} \in H^{\ell}$ yields the convergence

$$
v_N \underset{N \to +\infty}{\longrightarrow} v_{\varepsilon} \qquad \text{in } C_T^0(H_{\text{loc}}^{\ell'}) \tag{2.12}
$$

for any $\ell' \in \mathbb{R}$, from which we deduce convergence of the product

$$
\theta_N v_N \underset{N \to +\infty}{\longrightarrow} \theta_\varepsilon v_\varepsilon \quad \text{in } C_T^0(H_{\text{loc}}^{-1-\delta}).
$$

In the same way, by using the differential equation $\frac{d}{dt}z_N = v_N(t, z_N)$, we see that the approximate particle trajectories are bounded in the space $W^{1,\infty}(\mathbb{R}_+;\mathbb{R}^2)$, and Ascoli's theorem then shows that the sequence (z_N) is uniformly convergent (up to an extraction). Associating this with the convergence (2.47) provides, for any $t \in [0, T)$,

$$
\begin{aligned} \left| v_N(t, z_N) - v_\varepsilon(t, z_\varepsilon) \right| &\leq \left| v_N(t, z_N) - v_N(t, z_\varepsilon) \right| + \left| v_N(t, z_\varepsilon) + v_\varepsilon(t, z_\varepsilon) \right| \\ &\leq \left\| z_N - z_\varepsilon \right\|_\infty \| v_N \|_{H^3} + \| v_N - v_\varepsilon \|_{H^2} \\ &\xrightarrow[N \to +\infty]{} 0 \,. \end{aligned}
$$

The inequality above proves that the functions $v_N(t, z_N)$ converge uniformly to $v_{\varepsilon}(t, z_{\varepsilon})$. In exactly the same way, we see that the functions $\mathcal{H}_N(t,x) - K_{s,\varepsilon}(x-z_N(t))$ converge to $\mathcal{H}_\varepsilon(t,x) = K_{s,\varepsilon}(x-z_\varepsilon(t))$.

Overall, we see that the functions $(\theta_{\varepsilon}, v_{\varepsilon}, \mathcal{H}_{\varepsilon})$ solve the approximate system (2.24) with initial datum $\theta_{0,\varepsilon}$. Note that because $v_{\varepsilon} = K_{s,\varepsilon} * \theta_{\varepsilon}$ is a smooth function, the C^{∞} regularity of the initial datum is propagated by the flow.

 \Box

3 Local Existence of Classical Solutions

The section is devoted to the proof of existence of solutions as stated in Theorem 1.1- (i) . More precisely, we prove the following.

Theorem 3.1. Consider $k \geq 4$ and a set of vortex-wave initial data $(\theta_0, z_0) \in H^k(\mathbb{R}^2) \times \mathbb{R}^2$ such that θ_0 is constant (equal to, say a fixed $\beta \in \mathbb{R}$) on a ball $B(z_0, R_0)$ centered at the point vortex and of positive radius $R_0 > 0.$

Then there exists a time $T > 0$ such that the vortex-wave system (1.4) has a solution $(\theta, z) \in L^{\infty}([0, T); H^k(\mathbb{R}^2)) \times$ $C^1([0,T);\mathbb{R}^2)$. In addition, the time T may be chosen such that

$$
T \geq \frac{C}{E^{6+3k-6s} + E^{1/2}},
$$

where $E = ||\theta_0||_{H^k}^2 + \frac{1}{R_0}$.

3.1 A Priori Estimates

This first paragraph is devoted to the establishment of a priori estimates for our system: assuming that the solution is regular, we will derive high regularity estimates on some time interval. These are summarized in the following proposition.

Proposition 3.2. Consider an initial datum (θ_0, z_0) and a solution (θ, z) that are very regular, both in terms of smoothness and integrability. Furthermore, assume that there is a $R_0 > 0$ such that θ_0 is constant, say equal to a fixed $\beta \in \mathbb{R}$, on the ball $B(z_0, R_0)$.

For every time, we set $R(t) \geq 0$ to be the largest radius $r \geq 0$ such that the function $\theta(t)$ is constant on the ball $B(z(t), R(t))$, see (1.7). Then, for every integer $k \geq 4$, there exists a time $T > 0$ such that the bound

$$
\left\|\theta(t)\right\|_{H^k}^2 \leq C\left(\|\theta_0\|_{H^k}^2 + \frac{1}{R(t)}\right)
$$

holds for every $0 \le t \le T$. Moreover, the time T satisfies the inequality

$$
T \ge \frac{C}{E^{6+3k-6s} + E^{1/2}}.
$$

Adopt the notation of Proposition 3.2. The proof is based on an energy method: consider a multi-index $\alpha \in \mathbb{N}^2$ whose length is $k := |\alpha| = \alpha_1 + \alpha_2$. By integrating the transport equation against $\partial^{2\alpha}\theta$, we obtain an energy estimate involving the \dot{H}^k norm of θ , namely

$$
\frac{1}{2} \frac{d}{dt} \int |\partial^{\alpha} \theta|^{2} dx = - \int \partial^{\alpha} \theta \cdot \partial^{\alpha} \operatorname{div}(\theta \cdot \nabla^{\perp} (-\Delta)^{-s} \theta) dx - \int \partial^{\alpha} \theta \cdot \partial^{\alpha} \operatorname{div}(\theta H)
$$
\n
$$
= J_{1} + J_{2}.
$$
\n(3.1)

3.1.1 Estimates for the First Integral

We will first start by focusing on the integral J_1 , involving a trilinear product of derivatives of θ , and turn afterwards our attention to J_2 . This is the purpose of the following lemma.

Lemma 3.3. With the notation defined above, the inequality

$$
|J_1(t)| \le C(s,k) ||\theta(t)||_{H^k}^3
$$
\n(3.2)

holds for all times $t > 0$.

Proof. We distribute the derivatives in the product $\theta.\nabla^{\perp}(-\Delta)^{-s}\theta$ by means of the binomial formula:

$$
\int \partial^{\alpha} \theta \cdot \partial^{\alpha} \operatorname{div}(\theta \cdot \nabla^{\perp}(-\Delta)^{-s}\theta) dx = \sum_{\gamma \leq \alpha} \begin{pmatrix} \alpha \\ \gamma \end{pmatrix} \int \partial^{\alpha} \theta \cdot \operatorname{div}(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \nabla^{\perp}(-\Delta)^{-s}\theta) dx,
$$

where the binomial coefficient is defined by

$$
\left(\begin{array}{c} \alpha \\ \gamma \end{array}\right) := \frac{\gamma!}{\alpha!(\alpha-\gamma)!} = \frac{\gamma_1!\gamma_2!}{\alpha_1!\alpha_2!(\alpha_1-\gamma_1)!(\alpha_2-\gamma_2)!}.
$$

We distinguish between four cases in the sum. Firstly, all derivatives could be of order lower than k , so that we may easily bound the integral by using only $\|\theta\|_{H^k}$. More precisely, assume that $2 \le |\gamma| \le k-1$, so that

$$
\left| \int \partial^{\alpha} \theta \cdot \mathrm{div} \left(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \nabla^{\perp} (-\Delta)^{-s} \theta \right) \mathrm{d}x \right| \leq \| \partial^{\alpha} \theta \|_{L^{2}} \| \partial^{\alpha - \gamma} \nabla^{\perp} (-\Delta)^{-s} \theta \|_{L^{\infty}} \| \nabla \partial^{\gamma} \theta \|_{L^{2}} \leq \| \theta \|_{H^{k}} \| \partial^{\alpha - \gamma} \nabla^{\perp} (-\Delta)^{-s} \theta \|_{L^{\infty}} \| \theta \|_{H^{k}}
$$

In order to bound the L^{∞} norm in the previous inequality, we make use of the Sobolev embedding $H^{1+\varepsilon} \subset L^{\infty}$, which holds for any $\varepsilon > 0$. Because the exponent $s > 0$ is positive, the operator $\nabla^{\perp}(-\Delta)^{-s}$ is of order smaller than 1, so that

$$
\|\partial^{\alpha-\gamma}\nabla^{\perp}(-\Delta)^{-s}\theta\|_{L^{\infty}} \le \|\partial^{\alpha-\gamma}\nabla^{\perp}(-\Delta)^{-s}\theta\|_{H^{1+\varepsilon}} \le \|\theta\|_{H^{\sigma}},
$$

where the index $\sigma = (k-2) + (1-s) + (1+\varepsilon)$ is smaller than $\sigma \leq k$ if ε is chosen small enough. Therefore we have the bound

$$
\left| \int \partial^{\alpha} \theta \cdot \text{div} \left(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \nabla^{\perp} (-\Delta)^{-s} \theta \right) \text{d}x \right| \leq \| \theta \|_{H^{k}}^{3} \leq \| \theta \|_{H^{k}}^{3}.
$$
\n(3.3)

Secondly, when we have $|\gamma|=1$, similar computations lead to

$$
\left| \int \partial^{\alpha} \theta \cdot \text{div} \left(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \nabla^{\perp} (-\Delta)^{-s} \theta \right) dx \right| \leq \| \partial^{\alpha} \theta \|_{L^{2}} \| \partial^{\gamma} \nabla \theta \|_{L^{\infty}} \| \partial^{\alpha - \gamma} \nabla^{\perp} (-\Delta)^{-s} \theta \|_{L^{2}} \leq \| \theta \|_{H^{k}} \| \partial^{\gamma} \nabla \theta \|_{H^{1+\varepsilon}} \| \theta \|_{H^{k-s/2}} \leq \| \theta \|_{H^{k}}^{3}
$$
\n(3.4)

provided that the exponent k be greater than $k \geq 4 > 3 + \varepsilon$.

The third case of interest is when $\gamma = \alpha$, so that the integral cancels because the velocity field $v = \nabla^{\perp}(-\Delta)^{-s}\theta$ is divergence free:

$$
\int \partial^{\alpha} \theta \cdot \operatorname{div} \left(\partial^{\alpha} \theta \cdot \nabla^{\perp} (-\Delta)^{-s} \theta \right) dx = 0.
$$
 (3.5)

Finally, we are left with the case where $\gamma = 0$, so the full weight of the derivatives bears on the velocity field $v = \nabla^{\perp}(-\Delta)^{-s}\theta$. In order to circumvent the loss of derivatives induced by the operator $A = \nabla^{\perp}(-\Delta)^{-s}$, we take advantage of the commutator structure in the equation: by noting $A = (A_1, A_2) = (-\partial_2(-\Delta)^{-s}, \partial_1(-\Delta)^{-s}),$ we define two skew-symmetric operators A_1 and A_2 so that

$$
\int \partial^{\alpha} \theta \cdot \text{div} \left(\theta \cdot \partial^{\alpha} \nabla^{\perp} (-\Delta)^{-s} \theta \right) dx = \int \partial^{\alpha} \theta \cdot \partial_{j} \theta \cdot \partial^{\alpha} A_{j} \theta dx
$$

=
$$
- \int A_{j} (\partial^{\alpha} \theta \cdot \partial_{j} \theta) \cdot \partial^{\alpha} \theta dx
$$

=
$$
\frac{1}{2} \int \partial^{\alpha} \theta \cdot [\partial_{j} \theta, A_{j}] \partial^{\alpha} \theta dx,
$$

where there is an implicit sum on the repeated index $j \in \{1, 2\}$. Using Proposition 2.8 with $\sigma = -s$ to bound the commutator appearing in the last integral, we get, for any $\varepsilon > 0$,

$$
\left| \int \partial^{\alpha} \theta \cdot \text{div} \left(\theta \cdot \partial^{\alpha} \nabla^{\perp} (-\Delta)^{-s} \theta \right) \text{d}x \right| \lesssim \| \partial^{\alpha} \theta \|_{L^{2}} \left\| \left[\partial_{j} \theta, A_{j} \right] \partial^{\alpha} \theta \right\|_{L^{2}} \lesssim \| \partial^{\alpha} \theta \|_{L^{2}}^{2} \int |\xi|^{2-2s} |\widehat{\theta}(\xi)| \text{d}\xi + \| \partial^{\alpha} \theta \|_{L^{2}} \| \partial^{\alpha} \theta \|_{\dot{H}^{-2s}} \int |\xi|^{2} |\widehat{\theta}(\xi)| \text{d}\xi.
$$

In order to bound the integrals above, note that the function $\xi \mapsto |\xi|^{2-2s}$ is bounded on a neighborhood of $\xi = 0$. As a consequence, for every $\varepsilon > 0$, we may use the Cauchy-Schwarz inequality to write

$$
\int (|\xi|^{2-2s} + |\xi|^2)|\widehat{\theta}(\xi)|d\xi \le \int \frac{|\xi|^2}{1 + |\xi|^{3+\varepsilon}} (1 + |\xi|)^{3+\varepsilon} |\widehat{\theta}(\xi)|d\xi
$$

$$
\le C(\varepsilon) \|\theta\|_{H^{3+\varepsilon}}.
$$

Furthermore, the since $k = |\alpha| \ge 4 > 2s$, the homogeneous norm $||\theta||_{\dot{H}^{2s}}$ is bounded by $||\theta||_{H^k}$. We deduce

$$
\left| \int \partial^{\alpha} \theta \, \mathrm{div} \left(\theta \cdot \partial^{\alpha} \nabla^{\perp} (-\Delta)^{-s} \theta \right) \mathrm{d} x \right| \lesssim \|\theta\|_{H^{k}}^{3}.
$$
 (3.6)

Putting together the bounds (3.10), (3.11), (3.12) and (3.16), we then get the following estimate for the second summand in (3.4):

$$
\left| \int \partial^{\alpha} \theta . \partial^{\alpha} \operatorname{div} (\theta . \nabla^{\perp} (-\Delta)^{-s} \theta) \, dx \right| \lesssim \| \theta \|_{H^{k}}^{3}.
$$

3.1.2 Radius Estimates

Now, we take care of the last integral J_2 in (3.4) involving the influence of the point vortex through $H(t, x)$. This will create no particular problem: the point vortex is assumed to be in a region where θ is constant, and this will allow us to ignore any contribution of $H(t, x)$ near the singular point $z(t)$. In order to implement this argument, for any time, we need to prove that the θ remains constant near $z(t)$. We prove the following lemma.

Lemma 3.4. Let (θ, z) be a strong solution on $[0, T)$ to (1.4) (in the sense of Thrm 1.1) and assume that

$$
\theta_0 \equiv \beta \qquad on \ B(z(0), R_0),
$$

for some $\beta \in \mathbb{R}$. Recall that R, defined in relation (1.7), is given by

$$
R(t) = \sup \{ r \ge 0 , \ \theta(t) \equiv \beta \text{ on } B(z(t), r) \}.
$$

Then for every $t \in [0, T)$,

$$
R(t) \ge R(0) \exp\left(-\int_0^t \|\nabla v(\tau,\cdot)\|_{L^\infty} d\tau\right).
$$

In addition, we also have the inequality

$$
\frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{1}{R(t)}\right) \lesssim \frac{\|\nabla v\|_{L^{\infty}}}{R(t)} \lesssim \frac{\|\theta\|_{H^3}}{R(t)}.\tag{3.7}
$$

Proof. Let $x \in \mathbb{R}^2$ such that $\theta(t, x) \neq \beta$. Let $\tau \mapsto X(\tau)$ be the trajectory of the fluid particle passing in x at time t. Then

$$
\frac{d}{d\tau}|X(\tau) - z(\tau)|\Big|_{\tau=t} = \frac{x - z(t)}{|x - z(t)|} \cdot (v(x, t) + H(x, t) - v(z(t), t))
$$
\n
$$
= \frac{x - z(t)}{|x - z(t)|} \cdot (v(x, t) - v(z(t), t))
$$
\n
$$
\geq -|x - z(t)||\nabla v||_{L^{\infty}}
$$
\n
$$
\geq -R(t)||\nabla v||_{L^{\infty}}
$$

where we used the fact that $(x - z(t)) \cdot H(x, t) = 0$. By taking the infimum on every such point x, we have that

$$
\frac{\mathrm{d}}{\mathrm{d}t}R(t) \ge -R(t)\|\nabla v\|_{L^{\infty}},\tag{3.8}
$$

namely that

$$
R(t) \ge R(0) \exp\left(-\int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau\right).
$$

Finally, inequality (3.21) follows from (3.23) and the Sobolev embeddings $H^k \subset H^3 \subset W^{1,\infty}$. \Box

3.1.3 Estimates for the Second Integral

We are now in a favorable position to bound the integral J_2 . Our estimates are contained in the lemma immediately below.

Lemma 3.5. With the notation of Proposition 3.2, the inequality

$$
\left| \int \partial^{\alpha} \theta . \partial^{\alpha} \operatorname{div}(\theta . H) \, \mathrm{d}x \right| \lesssim \|\theta\|_{H^{k}}^{2} \left(\frac{1}{R(t)^{4-2s}} + \frac{1}{R(t)^{2+k-2s}} \right) \tag{3.9}
$$

holds for all times $t \geq 0$.

Proof. We now define a smooth truncation of the singular contribution of the point vortex $H(t, x)$. Let $\chi \in \mathcal{D}$ be the radial and smooth cut-off function defined in Subsection 2.3, and set, for any time such that $R(t) > 0$,

$$
\widetilde{H}_R(t,x) = \left(1 - \chi_{R(t)}(x - z(t))\right)H(t,x), \quad \text{where } \chi_R(y) := \chi\left(\frac{y}{R}\right).
$$

Then, because θ is constant on the ball $B(z(t), R(t))$, and so $\nabla \theta = 0$ on that area, we have

$$
\operatorname{div}(\theta H) = H \cdot \nabla \theta = \widetilde{H}_R \cdot \nabla \theta = \operatorname{div}(\theta \widetilde{H}_R),
$$

which leads to expressing the last integral in the energy estimate (3.4) as

$$
\int \partial^{\alpha} \theta \cdot \partial^{\alpha} \operatorname{div}(\theta \widetilde{H}_{R}) \, dx = \sum_{|\gamma| \leq |\alpha|} {\alpha \choose \gamma} \int \partial^{\alpha} \theta \cdot \operatorname{div}(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \widetilde{H}_{R}) \, dx.
$$

In estimating the integrals above, there are three cases. Firstly, when $\gamma = \alpha$, we have

$$
\int \partial^{\alpha} \theta . \nabla \partial^{\alpha} \theta \cdot \widetilde{H}_R \,dx = 0
$$

owing to the fact that the function \widetilde{H}_R is divergence free. Indeed, by writing the divergence of \widetilde{H}_R , we see that

$$
\operatorname{div} \tilde{H}_R(x) = (1 - \chi_R(x - z)) \operatorname{div} H(x) - \nabla \chi_R(x - z) \cdot K_s(x - z).
$$

The first term must be zero because H is smooth and divergence free on the support of $1 - \chi_R(x - z)$. The second is also zero because the vector $K_s(x-z)$ is colinear to $(x-z)^{\perp}$ while the radial symmetry of the cut-off function χ implies that $\nabla \chi_R(x-z)$ is colinear to $x-z$. Secondly, we explore the case where $|\gamma| \leq k-3$. We then have, thanks to the Sobolev embedding $H^2\subset L^\infty,$

$$
\left| \int \partial^{\alpha} \theta \cdot \operatorname{div} \left(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \widetilde{H}_R \right) \mathrm{d}x \right| \leq \| \partial^{\alpha} \theta \|_{L^2} \| \partial^{\alpha - \gamma} \widetilde{H}_R \|_{L^2} \| \nabla \partial^{\gamma} \theta \|_{L^{\infty}} \n\lesssim \frac{\| \theta \|_{H^k}^2}{R(t)^{2 + k - |\gamma| - 2s}} \n\lesssim \| \theta \|_{H^k}^2 \left(\frac{1}{R(t)^{5 - 2s}} + \frac{1}{R(t)^{2 + k - 2s}} \right).
$$

Lastly, in the case where $k-2 \le |\gamma| \le k-1$, so that $1 \le |\alpha - \gamma| \le 2$, we instead evaluate the L^{∞} norm of $\partial^{\gamma}H_R$ to get

$$
\left| \int \partial^{\alpha} \theta \cdot \operatorname{div} \left(\partial^{\gamma} \theta \cdot \partial^{\alpha - \gamma} \widetilde{H}_R \right) \mathrm{d}x \right| \leq \| \partial^{\alpha} \theta \|_{L^2} \| \partial^{\alpha - \gamma} \widetilde{H}_R \|_{L^{\infty}} \| \nabla \partial^{\gamma} \theta \|_{L^2}
$$

$$
\lesssim \frac{\| \theta \|_{H^k}^2}{R(t)^{3 + k - |\gamma| - 2s}}
$$

$$
\lesssim \| \theta \|_{H^k}^2 \left(\frac{1}{R(t)^{4 - 2s}} + \frac{1}{R(t)^{5 - 2s}} \right).
$$

Note that because $k \geq 4$, the exponent $4 - 2s$ in this last upper bound is smaller than the exponent $2 + k - 2s$ obtained above. Overall, we have the estimate

$$
\left| \int \partial^{\alpha} \theta . \partial^{\alpha} \operatorname{div}(\theta. \widetilde{H}_R) \, dx \right| \lesssim \|\theta\|_{H^k}^2 \left(\frac{1}{R(t)^{4-2s}} + \frac{1}{R(t)^{2+k-2s}} \right).
$$

3.1.4 Proof of the A Priori Estimates

Finally, with the help of the lemmas above, we can prove Proposition 3.2.

Proof (of Proposition 3.2). Plugging estimates (3.5) from Lemma 3.3 and (3.25) from Lemma 3.5 along with (3.21) into inequality (3.4) , we obtain a differential inequality:

$$
\frac{\mathrm{d}}{\mathrm{d}t} \left(\|\theta\|_{H^k}^2 + \frac{1}{R(t)} \right) \lesssim \|\theta\|_{H^k}^3 + \frac{\|\theta\|_{H^3}}{R(t)} + \|\theta\|_{H^k}^2 \left(\frac{1}{R(t)^{4-2s}} + \frac{1}{R(t)^{2+k-2s}} \right)
$$

$$
\lesssim \|\theta\|_{H^k}^3 + \frac{1}{R(t)^{3/2}} + \frac{1}{R(t)^{6+3k-6s}}
$$

$$
\lesssim \left(\|\theta\|_{H^k}^2 + \frac{1}{R(t)} \right)^{3/2} + \left(\|\theta\|_{H^k}^2 + \frac{1}{R(t)} \right)^{6+3k-6s}.
$$

Define $E(t) := ||\theta||_{H^k}^2 + R(t)^{-1}$ and set, for notational convenience, $q + 1 := 6 + 3k - 6s$. Then, by integrating the inequality above, we obtain, for $T \geq 0$

$$
E(T) \le E(0) + C \int_0^T \left(E(t)^{3/2} + E(t)^{q+1} \right) dt.
$$

Let $[0, T^*]$ be the time interval on which the integral above stays smaller than the initial data:

$$
T^* := \sup \left\{ T > 0, \quad \int_0^T \left(E(t)^{3/2} + E(t)^{q+1} \right) dt \le E(0) \right\}.
$$

Then, as long as $0 \leq T \leq T^*$, the energy $E(t)$ satisfies the inequality

$$
E(T) \le CE(0),\tag{3.10}
$$

from which we infer

$$
\int_0^T \left(E(t)^{3/2} + E(t)^{q_2+1} \right) dt \leq CTE(0)^{3/2} + CTE(0)^{q+1},
$$

and consequently

$$
T^* \ge \frac{C}{E(0)^q + E(0)^{1/2}}.
$$

3.2 Construction of a solution

This Subsection is dedicated to the construction of a local solution of (1.4). The proof relies on the regularized dynamics defined in Subsection 2.3, whose solutions will naturally satisfy the a priori estimates of Proposition 3.2.

3.2.1 Step 1: Approximate Solutions and Uniform Bounds

Lemma 3.6. Assume that the initial data (θ_0, z_0) are as in Proposition 3.1. Then for all $\varepsilon > 0$, the approximate system (2.24) equipped with the initial data ($\theta_{0,\varepsilon}$, z₀) has a global solution (θ_{ε} , z_e). In addition, this solution satisfies the bounds of Proposition 3.2, namely if $R_{\varepsilon}(t)$ is such that

$$
\theta_{\varepsilon}(t,x) = \beta \qquad \text{for } |x - z_{\varepsilon}(t)| \le R_{\varepsilon}(t),
$$

then we have a time $T_{\varepsilon} > 0$ such that the inequalities

$$
\|\theta_{\varepsilon}(t)\|_{H^k} + \frac{1}{R_{\varepsilon}(t)} \lesssim \|\theta_0\|_{H^k} + \frac{1}{R_0}
$$
\n(3.11)

hold for $t \in [0, T_{\varepsilon}].$ In addition T_{ε} satisfies the lower bound $T_{\varepsilon} \geq C^{-1}T^*$ independently of the approximation parameter $\varepsilon > 0$.

Proof. The only non-trivial point is the existence of a global approximate solution, which is given by Proposition 2.9. The exact same computations leading to Proposition 3.2 show that they also follow the a priori estimates. \Box

3.2.2 Step 2: Convergence of the Regularized Kernels

Before showing that the approximate solutions $(\theta_{\varepsilon}, z_{\varepsilon})$ converge when $\varepsilon \to 0^+$, we must study the regularized kernel $K_{s,\varepsilon}$. We already know that $K_{s,\varepsilon} \longrightarrow K_s$ locally uniformly in $\mathbb{R}^2 \setminus \{0\}$, however, we will need a more precise norm convergence. This is the purpose of the following lemma.

Lemma 3.7. The kernels $K_{s,\epsilon}$ are uniformly bounded in the homogeneous Besov space $\dot{B}^{2s-1}_{1,\infty}$. In addition, for any $\sigma < 2s - 1$, we have the convergence

$$
K_{s,\epsilon} \longrightarrow K_s \qquad \text{in } \dot{B}^{\sigma}_{1,\infty}.
$$
\n
$$
(3.12)
$$

Proof. As seen from equation (1.1), the kernel K_s is a homogeneous function of degree $2s - 3$. A simple scaling argument (see Proposition 2.21, p. 65 in [3]) shows that $K_s \in \dot{B}_{1,\infty}^{2s-1}$. Now, from (3.51), we see that the difference $K_s - K_{s,\epsilon}$ reads

$$
K_s(x) - K_{s,\epsilon}(x) = \chi \left(\frac{x}{\epsilon}\right) \frac{x^{\perp}}{|x|^{4-2s}}
$$

=
$$
\left(\frac{1}{\epsilon}\right)^{3-2s} \chi \left(\frac{x}{\epsilon}\right) \frac{(x/\epsilon)^{\perp}}{|x/\epsilon|^{4-2s}}
$$

=
$$
\left(\frac{1}{\epsilon}\right)^{3-2s} (\chi K_s) \left(\frac{x}{\epsilon}\right).
$$

This shows that the difference $K_s - K_{s,\epsilon}$ is a scaled version of the function χK_s . Therefore, for $\sigma \in \mathbb{R}$, we have, by the scaling propertied of homogeneous Besov spaces,

$$
\left\|K_s - K_{s,\epsilon}\right\|_{\dot{B}_{1,\infty}^{\sigma}} \lesssim \left(\frac{1}{\epsilon}\right)^{\sigma-(2s-1)} \left\|\chi K_s\right\|_{\dot{B}_{1,\infty}^{\sigma}}.
$$

In particular, the convergence (3.42) holds as soon as $\sigma < 2s - 1$, provided that we indeed have $\chi K_s \in \dot{B}^{\sigma}_{1,\infty}$. This follows from the fact that homogeneous and non-homogeneous Besov spaces coincide for functions supported in a given compact subset of \mathbb{R}^2 (see Proposition 2.93, p. 108, in [3]). \Box

3.2.3 Step 3: Time Compactness and Convergence of the Temperatures

We start by showing strong convergence of the approximate (potential) temperatures (θ_{ε}) , as an application of Ascoli's theorem.

Lemma 3.8. Let $(\theta_{\varepsilon}, z_{\varepsilon})$ be the approximate solution defined in Lemma 3.6. Then, for any $\ell' < k$, we have the strong convergence (up to an extraction)

$$
\theta_{\epsilon} \longrightarrow \theta \qquad in \ L^{\infty}([0, T); H_{\text{loc}}^{\ell}). \tag{3.13}
$$

Proof. While the uniform bounds of Lemma 3.6 provide space compactness on the sequence $(\theta_{\epsilon})_{\epsilon>0}$, we will also need compactness with respect to the time variable: by using the evolution equation for θ_{ε} , we see that

$$
\partial_t \theta = - v_\varepsilon \cdot \nabla \theta_\varepsilon - \chi_{R_\varepsilon} H_\varepsilon \cdot \nabla \theta_\varepsilon.
$$

Because θ_{ε} is constant (equal to β) on a neighborhood of $z_{\varepsilon}(t)$, the products in the right-hand side are well-defined and we have, for all times $t \in [0, T)$,

$$
\|\partial_t \theta_{\varepsilon}\|_{H^{k-1}} \lesssim \|v_{\varepsilon}\|_{H^{k-1}} \|\nabla \theta_{\varepsilon}\|_{H^{k-1}} + \|\chi_{R_{\varepsilon}} H_{\varepsilon}\|_{H^{k-1}} \|\nabla \theta_{\varepsilon}\|_{H^{k-1}}
$$

\n
$$
\lesssim \|\theta_{\varepsilon}\|_{H^k}^2 + \frac{1}{R_{\varepsilon}(t)^{k+1-2s}} \|\theta_{\varepsilon}\|_{H^k}
$$

\n
$$
\lesssim \|\theta_0\|_{H^k}^2 + \frac{1}{R_0^{k+1-2s}} \|\theta_0\|_{H^k}.
$$
\n(3.14)

In particular, we see that the sequence $(\theta_{\varepsilon})_{\varepsilon>0}$ is uniformly bounded in the space $W^{1,\infty}([0,T); H^{k-1})$. While this already provides both time and space compactness, it will be convenient to use an interpolation argument to get convergence in a stronger space topology. Consider $\ell \in (k-1,k)$ and $\sigma \in (0,1)$ such that $\ell = (k-1)\sigma + k(1-\sigma)$. Then, for $t_1, t_2 \in [0, T)$, we have

$$
\|\theta(t_1)-\theta(t_2)\|_{H^{\ell}} \lesssim \|\theta(t_1)-\theta(t_2)\|_{H^{k-1}}^{\sigma}\|\theta(t_1)-\theta(t_2)\|_{H^k}^{1-\sigma} \lesssim 2^{1-\sigma}|t_2-t_1|^{\sigma}\|\theta\|_{W^{1,\infty}(H^{k-1})}^{\sigma}\|\theta\|_{L^{\infty}(H^k)}^{1-\sigma}.
$$

Consequently, the sequence $(\theta_{\epsilon})_{\epsilon>0}$ is bounded in the space $C^{0,\sigma}([0,T); H^{\ell})$, and we deduce the existence of a θ with a strong convergence, up to taking a subsequence $(\epsilon_n)_n$,

$$
\theta_{\epsilon} \longrightarrow \theta \qquad \text{in } L^{\infty}([0,T); H_{\text{loc}}^{\ell'}),
$$

for any $\ell' < \ell < k$. Note that in reality, because $\theta_{\epsilon} \in C^{0}([0, T); H^{\ell'}_{\text{loc}})$, the convergence above is in that space, so that the limit θ is continuous with respect to the time variable. \Box

3.2.4 Step 4: Convergence of the Velocity Fields

In the same manner, we show the convergence of the approximate velocity fields (v_{ε}) . Here, there is a slight twist because $v_{\varepsilon} = K_{s,\varepsilon} \star \theta_{\varepsilon}$ is a convolution by a kernel depending on the approximation parameter $\varepsilon > 0$. This means that we must also use the convergence properties of the sequence $(K_{s,\varepsilon})$ from Lemma 3.7.

Lemma 3.9. Let (v_{ε}) the approximate velocities defined in Lemma 3.6. Then we have pointwise convergence (up to an extraction)

$$
v_{\epsilon} \longrightarrow v \qquad a.e.
$$

Furthermore, the limit v is given by $v = K_s \star \theta$.

Proof. Recall from Subsection 2.3 that the approximate kernel $K_{s,\varepsilon}$ is given by

$$
K_{s,\varepsilon}(x) = \left(1 - \chi(\varepsilon^{-1}x)\right)K(x). \tag{3.15}
$$

and that Lemma 3.7 shows that it converges to K_{ε} in the space $\dot{B}^{\sigma}_{1,\infty}$ for any $\sigma < 2s - 1$. We write the difference $v_{\varepsilon} - v$ as a sum of two terms

$$
v_{\varepsilon} - v = (K_{s,\varepsilon} - K_s) \star \theta_{\varepsilon} + K_s \star (\theta_{\varepsilon} - \theta).
$$

To handle the first of these terms, we use the Littlewood-Paley decomposition: remembering that $\dot{\Delta}_j \dot{\Delta}_m = 0$ as soon as $|j - m| \geq 2$, we have

$$
\begin{aligned} \left\| (K_{s,\varepsilon} - K_s) \star \theta_{\varepsilon} \right\|_{L^{\infty}} &\leq \sum_{j \in \mathbb{Z}} \left\| \dot{\bigtriangleup}_j (K_{s,\varepsilon} - K_s) \star \theta_{\varepsilon} \right\|_{L^{\infty}} \\ &\lesssim \sum_{|j-m| \leq 1} \left\| \dot{\bigtriangleup}_j (K_s - K_{s,\varepsilon}) \star \dot{\bigtriangleup}_m \theta_{\varepsilon} \right\|_{L^{\infty}} \\ &\lesssim \sum_{|j-m| \leq 1} \left\| \dot{\bigtriangleup}_j (K_s - K_{s,\varepsilon}) \right\|_{L^1} \left\| \dot{\bigtriangleup}_m \theta_{\varepsilon} \right\|_{L^{\infty}} \end{aligned}
$$

Let us focus on the series with negative indices $j < 0$. The Bernstein-type inequality of Lemma 2.4 applied to $\Delta_m \theta_{\epsilon}$ and $\Delta_m (\theta_{\epsilon} - \theta)$ provides, for any $\sigma < 2s - 1$,

$$
\sum_{j<0, |j-m|\leq 1} \|\dot{\Delta}_j (K_s - K_{s,\epsilon})\|_{L^1} \|\dot{\Delta}_m \theta_{\epsilon}\|_{L^{\infty}}
$$
\n
$$
\lesssim \sum_{j<0, |j-m|\leq 1} 2^{-j\sigma} \|K_s - K_{s,\epsilon}\|_{\dot{B}^{\sigma}_{1,\infty}} 2^j \|\dot{\Delta}_m \theta_{\epsilon}\|_{L^2}
$$
\n
$$
\lesssim \|K_s - K_{s,\epsilon}\|_{\dot{B}^{\sigma}_{1,\infty}} \|\theta_{\epsilon}\|_{L^2} \sum_{j<0} 2^{j(1-\sigma)}.
$$

By remembering that $\sigma < 2s - 1 < 1$, we see that the series above is convergent. Now, we have to deal with nonnegative indices $j \geq 0$. The Bernstein type inequality of Lemma 2.4 once again yields

$$
\sum_{j\geq 0, |j-m|\leq 1} \|\dot{\Delta}_j (K_s - K_{s,\epsilon})\|_{L^1} \|\dot{\Delta}_m \theta_{\epsilon}\|_{L^{\infty}}
$$
\n
$$
\lesssim \sum_{j\geq 0, |j-m|\leq 1} 2^{-j\sigma} \|K_s - K_{s,\epsilon}\|_{\dot{B}_{1,\infty}^{\sigma}} 2^j \|\dot{\Delta}_m \theta_{\epsilon}\|_{L^2}
$$
\n
$$
\lesssim \sum_{j\geq 0, |j-m|\leq 1} 2^{-j(\sigma+1)} \|K_s - K_{s,\epsilon}\|_{\dot{B}_{1,\infty}^{\sigma}} \|\theta_{\epsilon}\|_{H^2}
$$
\n
$$
\lesssim \|K_s - K_{s,\epsilon}\|_{\dot{B}_{1,\infty}^{\sigma}} \|\theta_{\epsilon}\|_{H^2} \sum_{j\geq 0} 2^{-j(\sigma+1)}.
$$

Because $\sigma < 2s - 1 < 1$, the sum above converges. As a consequence, we have shown that

$$
\left\|(K_{s,\varepsilon}-K_s)\star \theta_\varepsilon\right\|_{L^\infty}\lesssim \|K_{s,\varepsilon}-K_s\|_{\dot{B}_{1,\infty}^\sigma}\|\theta_\varepsilon\|_{H^2}\underset{\varepsilon\to 0^+}{\longrightarrow}0\,.
$$

We now focus on the second term $K_s \star (\theta_\varepsilon - \theta)$. The issue is that we do not have norm convergence of the sequence (θ_{ε}) in the space H^2 , as Lemma 3.8 only provides H^2_{loc} convergence. This means that we cannot show L^{∞} convergence of the sequence $(K_s * \theta_{\varepsilon})$. Instead, we will prove weak convergence. We first note that by repeating the computations above, we see that, for any $f \in \mathcal{S}$,

$$
||K_s \star f||_{L^{\infty}} \lesssim ||K_s||_{\dot{B}^{\sigma}_{1,\infty}} ||f||_{H^2}.
$$
\n(3.16)

This means that, for any $x \in \mathbb{R}^2$, the function $y \mapsto K_s(x-y)$ defines an element of the space H^{-2} . Furthermore, the weak- (\star) convergence

$$
\theta_{\varepsilon} \stackrel{*}{\rightharpoonup} \theta \qquad \text{in } L_T^{\infty}(H^2),
$$

which follows from the fact that $k \geq 4$, implies that

$$
K_s \star \theta_{\varepsilon}(x) = \langle K_s(x - \cdot), \theta_{\varepsilon} \rangle_{H^{-2} \times H^2}
$$

\n
$$
\longrightarrow_{\varepsilon \to 0^+} \langle K_s(x - \cdot), \theta \rangle_{H^{-2} \times H^2} = K_s \star \theta(x).
$$

Overall, we have proved that

$$
v_{\varepsilon} = K_{s,\varepsilon} \star \theta_{\varepsilon} \xrightarrow[\varepsilon \to 0^+]{} v = -\nabla^{\perp} (-\Delta)^{-s} \theta
$$
 a.e.

 \Box

3.2.5 Step 5: Convergence of the Vortex Trajectories

We wish to secure pointwise convergence of the vortex trajectories ODE $z_{\epsilon} = v(t, z_{\epsilon})$.

Lemma 3.10. Consider the sequence of functions (z_{ϵ}) defined in Lemma 3.6 and the limit velocity from Lemma 3.9. Then we have the convergence (up to an extraction)

$$
z_{\epsilon} \longrightarrow z(t) \qquad in \ C^{1,\alpha}([0,T))
$$

and the limit z is a solution of the ODE $z' = v(t, z)$.

Proof. Recall from (3.47) that the sequence $(\theta_{\epsilon})_{\epsilon>0}$ is bounded in $W^{1,\infty}([0,T); H^3)$, so that the sequence $(v_{\epsilon})_{\epsilon>0}$ is bounded in the spaces

$$
W^{1,\infty}([0,T);H^2) \subset W^{1,\infty}([0,T);C^{0,\alpha}) \subset C^{0,\alpha}([0,T)\times\mathbb{R}^2)
$$

for any Hölder exponent $0 < \alpha < 1$. The trajectory derivatives $z'_{\epsilon}(t) = v_{\epsilon}(t, z_{\epsilon}(t))$ must therefore also be bounded in $C^{0,\alpha}([0,T))$. This implies the convergence of the sequence $(z_{\epsilon})_{\epsilon>0}$ to

$$
z_{\epsilon} \longrightarrow z(t) \qquad \text{in } C^{1,\alpha}([0,T)), \tag{3.17}
$$

for some $z \in C^{1,\alpha}([0,T))$. We deduce

$$
\left| v_{\varepsilon}(t, z(t_{\varepsilon})) - v(t, z(t)) \right| \leq \left| v_{\varepsilon}(t, z(t_{\varepsilon})) - v_{\varepsilon}(t, z(t)) \right| + \left| v_{\varepsilon}(t, z(t_{\varepsilon})) - v(t, z(t)) \right|
$$

\n
$$
\leq \left| v_{\varepsilon} \right| \left| w_{1,\infty} \right| v(t, z(t_{\varepsilon})) - v(t, z(t)) \right| + \left| v_{\varepsilon}(t, z(t)) - v(t, z(t)) \right|
$$

\n
$$
\longrightarrow_{\varepsilon \to 0^{+}} 0.
$$

The convergence above flows from Lemma 3.9. We deduce that the particle trajectory solves the limit ODE, namely

$$
z'(t) = v(t, z(t)).
$$
\n(3.18)

 \Box

3.2.6 Step 6: End of the Proof of Existence

In this paragraph, we finish the proof of Theorem 3.1. It is mostly a matter of assembling the convergence results from the lemmas above.

Proof (of Theorem 3.1). We begin by studying the convergence of the non-linear terms $v_{\varepsilon} \cdot \nabla \theta_{\varepsilon}$ and $\mathcal{H}_{\varepsilon} \cdot \nabla \theta_{\varepsilon}$ appearing in the first equation of (2.24). Thanks to Lemmas 3.9 and 3.8, we write

$$
v_{\epsilon} \cdot \nabla \theta_{\epsilon} \longrightarrow v \cdot \nabla \theta \qquad \text{a.e.}. \tag{3.19}
$$

On the other hand, the product $\mathcal{H}_{\varepsilon} \cdot \nabla \theta_{\varepsilon}$ requires more attention because the functions

$$
\mathcal{H}_{\epsilon}(t,x) = K_{s,\epsilon}(x - z_{\epsilon}(t))
$$

possess a double dependency on ϵ , in the kernel $K_{s,\epsilon}$, and in the argument $x-z_{\epsilon}(t)$. However, as the functions θ_{ϵ} are constant around the singularity of \mathcal{H}_{ϵ} , we only need to care about the values of $\mathcal{H}_{\epsilon}(t,x)$ for $|x-z_{\epsilon}(t)| > R_{\epsilon}(t)$. Let us define the target function

$$
H(t, x) := K_s(x - z(t)).
$$

From the uniform convergence (3.63), we infer that the approximate and target point vortices are close for $\epsilon > 0$ small enough:

$$
|z_{\epsilon}(t) - z(t)| \le \frac{1}{4} \min \left\{ R_{\epsilon}(t), R(t) \right\} \quad \text{for } t \in [0, T).
$$

Thanks to the uniform bound (3.41) in Lemma 3.6, we deduce that for such ϵ , both derivatives $\nabla \theta_{\epsilon}$ and $\nabla \theta$ vanish on a ball $B(z(t), R_*)$ for some constant radius $R_* > 0$ depending only on the initial data. Outside of this ball, we get convergence

$$
||K_{s,\epsilon}(x-z_{\epsilon}(t))-K_{s}(x-z(t))||_{L^{\infty}({}^{c}B(z(t),R_{*}))}\longrightarrow 0 \text{ uniformly on } [0,T),
$$

from which we infer, together with (3.45),

$$
\mathcal{H}_{\epsilon} \cdot \nabla \theta_{\epsilon} \longrightarrow H \cdot \nabla \theta \qquad \text{in } L^{\infty}([0, T); L^{2}).
$$
\n(3.20)

Let us summarize what we have obtained so far: putting together all convergences (3.45) , (3.66) , (3.71) and the fact that $z(t)$ solves the limit ODE (3.65), we see that (z, θ) is indeed a solution of the Vortex-Wave system (1.4). It only remains to show that (z, θ) is a solution associated to the initial datum (z_0, θ_0) . The fact that $z(0) = z_0$ is a consequence of the uniform convergence (3.63) and the equality $z_\epsilon(0) = z_0$. Concerning the initial value for the scalar unknown, it is a consequence of the time continuity of θ (see the comments immediately following (3.45)) and the strong convergence of the initial data

$$
\theta_{0,\epsilon} \longrightarrow \theta_0 \quad \text{in } L^2.
$$

These remarks end the proof of Theorem 3.1.

3.3 Time Regularity

In this paragraph, we check that the solution constructed in Theorem 3.1 is indeed a classical solution of the PDE problem, *i.e.* that all derivatives are well-defined in the sense of $C¹$ functions (with respect to time and space). Together with the existence of strong solutions, this completes the proof of the existence part of Theorem 1.1.

Proposition 3.11. Let $k \geq 4$ be an integer and $\theta_0 \in H^k$ an initial datum. Then the solution $(\theta, z) \in$ $L^{\infty}([0,T); H^k) \times C^{k+s-1-\varepsilon}([0,T); \mathbb{R}^2)$ constructed in Theorem 3.1 is a classical solution (it is C^1 in both time ans space). More precisely, we have the following regularity: for any $\ell < k - 1$,

$$
\theta \in C^1([0,T);H^{\ell}).
$$

Proof. Be repeating the computations in (3.47), we already know that the solution satisfies $\theta \in W^{1,\infty}([0,T); H^{k-1})$ and $z \in C^1([0,T]; \mathbb{R}^2)$, but to show that the function θ has C^1 regularity in both time and space, we study the second derivative $\partial_t \theta$, which reads

$$
\partial_t^2 \theta = -\partial_t v \cdot \nabla \theta - v \cdot \nabla \partial_t \theta - \partial_t H \cdot \nabla \theta - H \cdot \nabla \partial_t \theta.
$$
\n(3.21)

On the one hand, from $\theta \in W^{1,\infty}([0,T); H^{k-1})$, we deduce that $\partial_t v \in L^{\infty}([0,T); H^{k-2})$, so that

$$
\|\partial_t v \cdot \nabla \theta\|_{H^{k-2}} \lesssim \|\partial_t v\|_{H^{k-2}} \|\nabla \theta\|_{H^{k-2}} < +\infty.
$$
\n(3.22)

Similarly, we have

$$
||v \cdot \nabla \partial_t \theta||_{H^{k-2}} \lesssim ||v||_{H^{k-2}} ||\partial_t \theta||_{H^{k-1}} < +\infty.
$$
\n(3.23)

On the other hand, we look at the last two terms in the righthandside of (3.74) , which involve $H(t, x)$. Because the the function θ is constant, equal to β , on the ball $B(z(t), R(t))$, the derivatives $\nabla \theta$ and $\partial_t \nabla \theta$ must vanish on that ball. Therefore, we have

$$
\partial_t H \cdot \nabla \theta + H \cdot \nabla \partial_t \theta = \chi_R \partial_t H \cdot \nabla \theta + \chi_R H \cdot \nabla \partial_t \theta,
$$

which is a considerable help, as the function $\chi_R H$ is C^{∞} spacewise smooth at any given time. Let us compute the derivative $\partial_t H$. We have

$$
\partial_t H(t,x) = -\nabla K_s(x - z(t)) \cdot v(t, z(t)),
$$

so that

$$
\|\chi_R \partial_t H \cdot \nabla \theta\|_{H^{k-1}} \lesssim \|\nabla \theta\|_{H^{k-1}} \|\chi_R \partial_t H\|_{H^{k-1}} \le \|\chi_{R(t)}(x) \nabla K_s(x - z(t)) \cdot v(t, z(t))\|_{H^{k-1}} \|\theta\|_{H^k}
$$

$$
\lesssim \frac{1}{R(t)^{k+2-2s}} \|\theta\|_{H^k}^2 < +\infty.
$$
 (3.24)

In the same way, we also have

$$
\|\chi_R H \cdot \nabla \partial_t \theta\|_{H^{k-2}} \lesssim \|\chi_R H\|_{H^{k-2}} \|\partial_t \theta\|_{H^{k-1}} \lesssim \frac{1}{R(t)^{k+1-2s}} \|\partial_t \theta\|_{H^{k-2}} < +\infty.
$$
\n(3.25)

By putting together (3.75), (3.76), (3.79) and (3.80), we find that

$$
\partial_t^2 \theta \in L^\infty([0,T); H^{k-2}).
$$

This shows that θ has C^1 regularity with respect to time in the H^{k-2} topology. However, we are not quite finished yet: if we want to show that θ is C^1 with respect to both time and space, then the case $k = 4$ requires a bit more attention, as $H^2 \not\subset C^1$. This issue is resolved by interpolating between the spaces $W^{1,\infty}([0,T); H^{k-1})$ and $W^{2,\infty}([0,T); H^{k-2})$. For any $\sigma \in [0,1]$, we set

$$
\ell = (k-2)\sigma + (k-1)(1-\sigma).
$$

Then, for any times $t_1, t_2 \geq 0$, we may write,

$$
\|\partial_t \theta(t_2) - \partial_t(t_1)\|_{H^{\ell}} \le \|\partial_t \theta(t_2) - \partial_t(t_1)\|_{K^{k-2}}^{\sigma}\|\partial_t \theta(t_2) - \partial_t \theta(t_1)\|_{H^{k}}^{\sigma-1}
$$

$$
\le 2^{1-\sigma}|t_2 - t_1|^{\sigma}\|\partial_t^2 \theta\|_{W^{1,\infty}(H^{k-2})}^{\sigma}\|\partial_t \theta\|_{L^{\infty}(H^{k-1})}^{1-\sigma}
$$

Consequently, we deduce Hölder regularity on the time derivative

$$
\partial_t \theta \in C^{0,\sigma}([0,T);H^{\ell}),
$$

which provides the space-time C^1 regularity we were looking for: when $0 < \sigma < 1$, the space above is a subspace of $C^0([0,T); C^1 \cap W^{1,\infty})$. \Box

4 Uniqueness and stability of classical solutions

In this section, we conclude the proof of Theorem 1.1- (i) and Theorem 1.1- (ii) by proving the following proposition.

Proposition 4.1. Let $T > 0$, $k \geq 4$ and (θ_1, z_1) and (θ_2, z_2) be two solutions in $L^{\infty}([0, T], H^k)$ of (1.4), as constructed in Section 3, such that there exists β_1 , β_2 , $R_1 > 0$ and $R_2 > 0$ such that

$$
\theta_i \equiv \beta_i \quad on \ B(z_i(0), R_i).
$$

Then there exists $\rho > 0$ depending only on $\|\theta_1(0, \cdot)\|_{H^k}$, $\|\theta_2(0, \cdot)\|_{H^k}$, R_1 , R_2 and T such that if

$$
|z_2(0) - z_1(0)| < \rho,\tag{4.1}
$$

then, there exists a time $T_0 \in [0, T)$ such that for every $t \in [0, T_0]$ and for every $\ell \in \mathbb{N}$ such that $2 \leq \ell \leq k - 2$,

$$
\|\theta_2(t,\cdot)-\theta_1(t,\cdot)\|_{H^{\ell}}+|z_2(t)-z_1(t)|\lesssim \|\theta_2(0,\cdot)-\theta_1(0,\cdot)\|_{H^{\ell}}+|z_2(0)-z_1(0)|,
$$

where the omitted multiplicative constant depends only on $\|\theta_1(0, \cdot)\|_{H^k}$, $\|\theta_2(0, \cdot)\|_{H^k}$, R_1 , R_2 and T.

Theorem 1.1- (ii) is a simple reformulation of Proposition 4.1 and uniqueness of classical solutions of (1.4) as stated in Theorem 1.1-(i) comes from applying Proposition 4.1 to two solutions with same initial datum (θ_0 , z_0) with θ_0 constant in a neighborhood of z_0 as we will detail in Section 4.4.

Moreover in this section we obtain en route the L^2 -stability when $s \geq 1/2$ as follows.

Proposition 4.2. If in addition to the hypotheses of Proposition 4.1 we assume that $s \geq 1/2$, then

$$
\|\theta_2(t,\cdot)-\theta_1(t,\cdot)\|_{L^2}+|z_2(t)-z_1(t)|\lesssim \|\theta_2(0,\cdot)-\theta_1(0,\cdot)\|_{L^2}+|z_2(0)-z_1(0)|.
$$

The plan of the proof of Proposition 4.1 is the following. We start by looking for a convenient time T_0 such that both point-vortices remain in the constant part of each θ_i . Then we estimate the time derivative of $|z_1(t) - z_2(t)|$ and $||\theta_2(t, \cdot) - \theta_1(t, \cdot)||_{H^1}$, then conclude by a Gronwall's type argument.

In the computations, we denote by $\delta\theta = \theta_2 - \theta_1$, $\delta v = v_2 - v_1$ and $\delta z = z_2 - z_1$ the difference functions.

4.1 Existence of ρ and T_0

We start by proving the following lemma concerning the existence of constants ρ and T_0 in Proposition 4.1.

Lemma 4.3. Let (θ_1, z_1) and (θ_2, z_2) chosen as in Proposition 4.1. Then there exists a choice of ρ and T_0 such that if (4.2) is satisfied, then for every $t \in [0,T_0]$, for every $i, j \in \{1,2\}$, θ_i remains constant in the disk $B(z_i(t), \rho).$

Proof. For both $i \in \{1,2\}$, recall that $\|\nabla v_i(t, \cdot)\|_{L^\infty} \leq C \|\theta_i(t, \cdot)\|_{H^3}$, and from relation (3.37), that $\|\theta_i(t, \cdot)\|_{H^3}$ is bounded in time on [0, T] by a constant depending only on $\|\theta_i(0)\|_{H^k}$, R_i and T. Therefore, there exists a constant $\rho = \rho(||\theta_1(0)||_{H^k}, ||\theta_2(0)||_{H^k}, R_1, R_2, T)$ such that for every $i \in \{1,2\}$ and for every $t \in [0,T]$,

$$
R_i \exp\left(-\int_0^t \|\nabla v_i(\tau,\cdot)\|_{L^\infty} d\tau\right) > 4\rho.
$$

By Lemma 3.4 applied successively to θ_1 and θ_2 , we infer that for every $t \in [0, T]$,

$$
\theta_i(t, \cdot) \equiv \beta_i \quad \text{on } B(z_i(t), 4\rho). \tag{4.2}
$$

There remains to prove that $\theta_1 \equiv \beta_1$ on $B(z_2(t), \rho)$.

We recall that the definition of the regularized kernel $K_{s,\varepsilon}$ and its properties from Subsection 2.3. We now establish that for every $\varepsilon \in (0, 4\rho]$, we have that

$$
(K_{s,\rho} \star \theta_i)(z_i) = K_{s,\varepsilon} \star \theta_i(z_i). \tag{4.3}
$$

Indeed, by relation (4.6),

$$
K_{s,\varepsilon} \star \theta_i(z_i) = \int_{\mathbb{R}^2} K_{s,\varepsilon}(z_i - y) \theta_i(y, t) dy
$$

=
$$
\int_{\mathbb{R}^2 \setminus B(z_i, 4\rho)} K_{s,\varepsilon}(z_i - y) \theta_i(y, t) dy + \beta_i \int_{B(z_i, 4\rho)} K_{s,\varepsilon}(z_i - y) dy.
$$

By skew-symmetry, we have that

$$
\int_{B(0,4\rho)} K_{s,\varepsilon}(y) dy = \int_{B(0,4\rho)} K_{s,\rho}(y) dy = 0.
$$
\n(4.4)

Then, we observe that for every $\varepsilon \in (0, 4\rho)$, $K_{s,\varepsilon} \equiv K_{s,\rho}$ on $\mathbb{R}^2 \setminus B(z_i, 4\rho)$, which proves (4.7) .

Using the same arguments than the ones used to obtain (3.57), we have that (see also Lemma 3.7), we have that

$$
||K_{s,\epsilon} \star \theta_i - v||_{L^{\infty}} \lesssim ||K_{s,\epsilon} - K_s||_{\dot{B}^{\sigma}_{1,\infty}} ||\theta_i||_{H^2} \longrightarrow 0 \quad \text{as } \epsilon \to 0^+,
$$

with $\sigma < 2s - 1$. Since the functions $K_{s,\epsilon} \star \theta$ are continuous, the convergence $K_{s,\epsilon} \star \theta_i \longrightarrow v$ holds pointwise. Consequently, we have:

$$
(K_{s,\varepsilon} \star \theta_i(t,\cdot))(z_i(t)) = v_i(t,z_i(t)).
$$
\n(4.5)

Gathering relations (4.7) and (4.11), we obtain that

$$
\frac{\mathrm{d}}{\mathrm{d}t}z_i(t) = v_i(t, z_i(t)) = (K_{s,\rho} \star \theta_i)(z_i). \tag{4.6}
$$

Observing that for every $t \in [0, T_0]$,

 $||(K_{s,\rho} \star \theta_i)(t,\cdot)||_{L^{\infty}} \leq ||K_{s,\rho}||_{L^{\infty}} ||\theta_i(t,\cdot)||_{L^1} = ||K_{s,\rho}||_{L^{\infty}} ||\theta_i(0,\cdot)||_{L^1},$

by taking

$$
T_0 < \rho \left(\|K_{s,\rho}\|_{L^\infty} \max_{i \in \{1,2\}} \|\theta_i(0,\cdot)\|_{L^1} \right)^{-1},
$$

we deduce from relation (4.12) that for each $i \in \{1,2\}$ and every $t \in [0, T_0]$ that

$$
|z_i(t) - z_i(0)| \le \rho.
$$

Adding then the assumption (4.2) that $|z_1(0) - z_2(0)| < \rho$, we obtain for every $t \in [0, T_0]$ that

$$
|z_2(t) - z_1(t)| \le |z_2(t) - z_2(0)| + |z_2(0) - z_1(0)| + |z_1(t) - z_1(0)| < 3\rho. \tag{4.7}
$$

Gathering relation (4.6) and (4.16), we obtain that for every $t \in [0, T_0]$, for every $i, j \in \{1, 2\}$,

$$
\theta_i(t, \cdot) \equiv \beta_i \quad \text{on } B(z_j(t), \rho).
$$

 \Box

4.2 Estimate of the distance between the point-vortices

We then obtain the following estimate on the evolution of the distance between the two point-vortices.

Lemma 4.4. Let (θ_1, z_1) and (θ_2, z_2) chosen as in Proposition 4.1, let ρ and T_0 be given by Lemma 4.3 and assume that (4.2) holds true. Then, for every $t \in [0, T_0]$, we have that

$$
\frac{\mathrm{d}}{\mathrm{d}t}|z_1(t)-z_2(t)|\lesssim |z_2(t)-z_1(t)|+\|\theta_2-\theta_1\|_{L^2}.
$$

We recall that the omitted constant may only depends on $\|\theta_1(0)\|_{H^k}$, $\|\theta_2(0)\|_{H^k}$, R_1, R_2, T .

Proof. We compute

$$
\frac{d}{dt}|z_1(t) - z_2(t)| \le |v_1(t, z_1(t)) - v_2(t, z_2(t))|
$$

\n
$$
\le |v_1(t, z_1(t)) - v_1(t, z_2(t))| + |\delta v(t, z_2(t))|.
$$

From one hand we have that

$$
|v_1(t,z_1(t))-v_1(t,z_2(t))| \leq \|\nabla v_1\|_{L^\infty} |\delta z| \lesssim \|\theta_1\|_{H^3} |\delta z| \lesssim |\delta z|,
$$

by recalling that $||\theta_1||_{H^3}$ is bounded on [0, T] by hypothesis. From the other hand, by Lemma 4.3, $\delta\theta$ is constant on $B(z_2(t), \rho)$ for every $t \in [0, T_0]$ so we have that

$$
\delta v(t, z_2(t)) = (K_s \star \delta \theta(t, \cdot))(z_2(t)) = (K_{s, \rho} \star \delta \theta(t, \cdot))(z_2(t)).
$$

By the Cauchy Schwartz inequality, we obtain

$$
|\delta v(t, z_2(t))| \leq \|K_{s,\rho}\|_{L^2} \|\delta\theta\|_{L^2} \lesssim \|\delta\theta\|_{L^2},
$$

which concludes the proof of Lemma 4.4.

4.3 Estimate on the H^{ℓ} norm of $\theta_2 - \theta_1$

We now turn to the estimate of $\delta\theta$.

Lemma 4.5. With the same notations as in Lemma 4.4, for every $t \in [0, T_0]$ and every $\ell \in \mathbb{N}$ such that $2 \leq \ell \leq k-2$, we have that

$$
\frac{\mathrm{d}}{\mathrm{d}t} || \theta_2 - \theta_1 ||_{H^{\ell}} \lesssim || \theta_2 - \theta_1 ||_{H^{\ell}} + |z_2 - z_1|.
$$

Proof. Let

$$
H_{i,\rho}(t,x) := K_{s,\rho}(x - z_i(t))
$$

and

$$
u_i := v_i + H_{i,\rho}
$$

for $i \in \{1,2\}$. We drop the dependence in ρ in the notation u_i since it's not relevant for our computations. One should simply remember that the u_i are smooth flows on [0, T_0].

By Lemma 4.3, we have that $\nabla \theta_i \equiv 0$ on $B(z_i(t), \rho)$, so that on the whole plane the equality

$$
H_i \cdot \nabla \theta_i = H_{i,\rho} \cdot \nabla \theta_i
$$

is satisfied. The active scalar θ_i is not only transported by $v_i + H_i$ but also by u_i . Consequently, by substracting the two transport equations on θ_1 and θ_2 respectively by the flows u_1 and u_2 , we have that

$$
\partial_t \delta \theta + u_1 \cdot \nabla \delta \theta + \delta u \cdot \nabla \theta_2 = 0, \tag{4.8}
$$

with the notation $\delta u = u_2 - u_1$. Let us also denote $\delta H_\rho = H_{2,\rho} - H_{1,\rho}$. We multiply relation (4.27) by $\delta \theta$ and integrate to obtain that

$$
\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t} \|\delta\theta\|_{L^2}^2 + \int_{\mathbb{R}^2} \delta v \cdot \nabla \theta_2 \,\delta\theta + \int_{\mathbb{R}^2} \delta H_\rho \cdot \nabla \theta_2 \,\delta\theta = 0. \tag{4.9}
$$

We start with the last term of this equation. Since δH_{ρ} is smooth, then

$$
\left| \int \delta H_{\rho} \cdot \nabla \theta_2 \, \delta \theta \right| \leq \| \delta H_{\rho} \|_{L^{\infty}} \| \nabla \theta_2 \|_{L^2} \| \delta \theta \|_{L^2}.
$$

More precisely, from Lemma 4.3, for every $t \in [0, T_0]$ we know that $H_{i,\rho} \cdot \nabla \theta_2 = 0$ in $B(z_i(t), \rho)$ for both $i \in \{1, 2\}$, so that

$$
\|\delta H_{\rho}\|_{L^{\infty}} = \sup_{x \in \mathbb{R}^2} |K_{s,\rho}(x - z_2(t)) - K_{s,\rho}(x - z_1(t))| \leq \|\nabla K_{s,\rho}\|_{L^{\infty}} |z_2(t) - z_1(t)|.
$$

Recalling then that $\|\theta_2\|_{H^4}$ is bounded on $[0,T_0]$ by a constant depending only on $\|\theta_2(0)\|_{H^4}$, R_2 and T, we obtain that

$$
\left| \int \delta H_{\rho} \cdot \nabla \theta_2 \, \delta \theta \right| \lesssim |\delta z| \|\delta \theta\|_{L^2}.
$$

In order to continue our estimates, in the middle term in relation (4.28), one has to control a norm of δv in terms of $\delta\theta$. In general, norms of δv cannot be controlled by $\|\delta\theta\|_{L^2}$ only so we use Proposition 2.5 to get that

$$
\left| \int \delta v \cdot \nabla \theta_2 \, \delta \theta \right| \leq \left| \int \Delta_{-1} \delta v \cdot \nabla \theta_2 \delta \theta \right| + \left| \int (I - \Delta_{-1}) \delta v \cdot \nabla \theta_2 \delta \theta \right|
$$

\n
$$
\leq \|\Delta_{-1} \delta v\|_{L^{\infty}} \|\nabla \theta_2\|_{L^2} \|\delta \theta\|_{L^2} + \|(I - \Delta_{-1} \delta v)\|_{L^2} \|\nabla \theta_2\|_{L^{\infty}} \|\delta \theta\|_{L^2}
$$

\n
$$
\lesssim \|\theta_2\|_{H^3} \|\delta \theta\|_{H^{1-2s}} \|\delta \theta\|_{L^2}.
$$

In the case $s \geq 1/2$, we obtain that

$$
\frac{\mathrm{d}}{\mathrm{d}t} \|\delta\theta\|_{L^2} \lesssim |\delta z| + \|\delta\theta\|_{L^2},\tag{4.10}
$$

from which we will obtain Proposition 4.2. We now continue our computations to prove H^{ℓ} stability in the general case $0 < s < 1$. Yet we proved in general that

$$
\frac{\mathrm{d}}{\mathrm{d}t} \|\delta\theta\|_{L^2}^2 \lesssim \left(|\delta z| + \|\delta\theta\|_{H^{1-2s}} \right) \|\delta\theta\|_{L^2}.\tag{4.11}
$$

Let $\ell \in \mathbb{N}$ such that $2s \leq \ell \leq k-2$. Similarly to what we did in Section 3.1, for any $\alpha \in \mathbb{N}^2$ such that $|\alpha| = \ell$, we have from applying ∂^{α} to relation (4.27), multiplying by $\partial^{\alpha}\delta\theta$ and integrating that

$$
\frac{1}{2} \frac{d}{dt} ||\partial^{\alpha} \delta \theta||_{L^{2}}^{2} = -\int \partial^{\alpha} (u_{1} \cdot \nabla \delta \theta) \partial^{\alpha} \delta \theta = \int \partial^{\alpha} (\delta u \cdot \nabla \theta_{2}) \partial^{\alpha} \delta \theta
$$
\n
$$
:= -\sum_{\gamma \leq \alpha} {\alpha \choose \gamma} (I_{\gamma} + J_{\gamma} + L_{\gamma})
$$
\n(4.12)

where, using the Leibniz rule, for every $(0,0) \leq \gamma \leq \alpha$,

$$
\begin{aligned} I_{\gamma} &= \int \partial^{\gamma} u_1 \cdot \partial^{\alpha - \gamma} \nabla \delta \theta \partial^{\alpha} \delta \theta, \\ J_{\gamma} &= \int \partial^{\gamma} \delta v \cdot \partial^{\alpha - \gamma} \nabla \theta_2 \partial^{\alpha} \delta \theta, \\ L_{\gamma} &= \int \partial^{\gamma} \delta H_{\rho} \cdot \partial^{\alpha - \gamma} \nabla \theta_2 \partial^{\alpha} \delta \theta. \end{aligned}
$$

Estimating the terms I_{γ}

We first observe that by integration by parts, $I_{0,0} = 0$ since $\nabla \cdot u_1 = 0$. For $\gamma \neq (0,0)$, we have that $|\alpha - \gamma| \leq \ell - 1$ so

$$
|I_{\gamma}| \leq \|\partial^{\gamma} u_1\|_{L^{\infty}} \|\partial^{\alpha-\gamma} \nabla \delta \theta\|_{L^2} \|\partial^{\alpha} \delta \theta\|_{L^2} \leq \|\partial^{\gamma} u_1\|_{L^{\infty}} \|\delta \theta\|_{H^{\ell}}^2.
$$

Using that $u_1 = v_1 + H_{1,\rho}$,

$$
\|\partial^\gamma u_1\|_{L^\infty} \le \|\partial^\gamma v_1\|_{L^\infty} + \|\partial^\gamma H_{1,\rho}\|_{L^\infty}.
$$

The term $\|\partial^{\gamma}H_{1,\rho}\|_{L^{\infty}} < +\infty$ is a constant that depends only on γ and ρ (with our notations, $\|\partial^{\gamma}H_{1,\rho}\|_{L^{\infty}} \lesssim 1$). We then use Proposition 2.5 to get that there exists $\varepsilon > 0$ small enough such that

$$
\|\partial^{\gamma}v_{1}\|_{L^{\infty}} \leq \|\triangle_{-1}\partial^{\gamma}v_{1}\|_{L^{\infty}} + \|(\mathrm{Id}-\triangle_{-1})\partial^{\gamma}v_{1}\|_{L^{\infty}}
$$

\n
$$
\lesssim \|\theta_{1}\|_{H^{k}} + \|(\mathrm{Id}-\triangle_{-1})\partial^{\gamma}v_{1}\|_{H^{1+\varepsilon}}
$$

\n
$$
\lesssim \|\theta_{1}\|_{H^{k}} + \|\partial^{\gamma}\theta_{1}\|_{H^{1+\varepsilon+1-2s}}
$$

\n
$$
\lesssim \|\theta_{1}\|_{H^{k}} + \|\theta_{1}\|_{H^{\ell+2-2s+\varepsilon}}
$$

\n
$$
\lesssim \|\theta_{1}\|_{H^{k}},
$$

where the last inequality comes from the hypothesis that $\ell + 2 \leq k$. In conclusion,

$$
\|\partial^{\gamma}v_{1}\|_{L^{\infty}} \lesssim \|\theta_{1}\|_{H^{k}} \lesssim \|\theta_{1}(0)\|_{H^{k}} \lesssim 1,
$$

$$
|I_{\gamma}| \lesssim \|\delta\theta\|_{H^{\ell}}^{2}.
$$
 (4.13)

Estimating the terms J_{γ}

so that for every $\gamma \leq \alpha$,

We start by the case $\gamma \neq \alpha$. We have that

$$
J_{\gamma} = \int (\Delta_{-1} \partial^{\gamma} \delta v) \cdot \partial^{\alpha - \gamma} \nabla \theta_2 \partial^{\alpha} \delta \theta + \int (\mathrm{Id} - \Delta_{-1}) \partial^{\gamma} \delta v \cdot \partial^{\alpha - \gamma} \nabla \theta_2 \partial^{\alpha} \delta \theta
$$

The first term satisfies, using Proposition 2.5, that

$$
\left| \int \left(\Delta_{-1} \partial^{\gamma} \delta v \right) \cdot \partial^{\alpha - \gamma} \nabla \theta_2 \partial^{\alpha} \delta \theta \right| \leq \|\Delta_{-1} \partial^{\gamma} \delta v\|_{L^{\infty}} \|\partial^{\alpha - \gamma} \nabla \theta_2\|_{L^2} \|\delta \theta\|_{H^{\ell}} \leq \|\partial^{\gamma} \delta \theta\|_{L^2} \|\theta_2\|_{H^{\ell+1}} \|\delta \theta\|_{H^{\ell}} \leq \|\delta \theta\|_{H^{\ell}}^2,
$$

where we used that $\ell \geq 2 > 2 - 2s$. For the second term, if $\gamma \neq (0,0)$ we have for $\varepsilon \in (0,1)$ that

$$
\left| \int \left(\mathrm{Id} - \triangle_{-1} \right) \partial^{\gamma} \delta v \cdot \partial^{\alpha - \gamma} \nabla \theta_2 \partial^{\alpha} \delta \theta \right| \leq \| (\mathrm{Id} - \triangle_{-1}) \partial^{\gamma} \delta v \|_{L^2} \| \partial^{\alpha - \gamma} \nabla \theta_2 \|_{L^{\infty}} \| \partial^{\alpha} \delta \theta \|_{L^2}
$$

$$
\lesssim \| \partial^{\gamma} \delta \theta \|_{H^{1-2s}} \| \partial^{\alpha - \gamma} \nabla \theta_2 \|_{H^{1+\varepsilon}} \| \delta \theta \|_{H^{\ell}}
$$

$$
\lesssim \| \delta \theta \|_{H^{1-2s+\ell-1}} \| \theta_2 \|_{H^{\ell-1+1+\varepsilon}} \| \delta \theta \|_{H^{\ell}}
$$

$$
\lesssim \| \delta \theta \|_{H^{\ell}}^2,
$$

where we used that $\gamma \neq \alpha$ so that $|\gamma| \leq \ell - 1$. In the case $\gamma = (0, 0)$, we have that

$$
\left| \int \left(\mathrm{Id} - \triangle_{-1} \right) \delta v \cdot \partial^{\alpha} \nabla \theta_2 \partial^{\alpha} \delta \theta \right| \leq \| (\mathrm{Id} - \triangle_{-1}) \delta v \|_{L^{\infty}} \| \partial^{\alpha} \nabla \theta_2 \|_{L^2} \| \partial^{\alpha} \delta \theta \|_{L^2}
$$

$$
\lesssim \| (\mathrm{Id} - \triangle_{-1}) \delta v \|_{H^{1+\varepsilon}} \| \theta_2 \|_{H^{\ell+1}} \| \delta \theta \|_{H^{\ell}}
$$

$$
\lesssim \| \delta \theta \|_{H^{1+\varepsilon+1-2s}} \| \delta \theta \|_{H^{\ell}}
$$

$$
\lesssim \| \delta \theta \|_{H^{\ell}}^2.
$$

We now estimate $|J_{\alpha}|$. As in Section 3.1, let us denote by $A = (A_1, A_2) = \left(-\partial_2(-\Delta)^{-s}, \partial_1(-\Delta)^{-s}\right)$, which is a (formally) skew-symmetric operator. We have that

$$
J_{\alpha} = -\sum_{j=1}^{2} \int \partial^{\alpha} A_{j}(\delta \theta) \partial_{j} \theta_{2} \partial^{\alpha} \delta \theta
$$

$$
= -\sum_{j=1}^{2} \int \partial^{\alpha} \delta \theta A_{j}(\partial_{j} \theta_{2} \partial^{\alpha} \delta \theta)
$$

and by summing both of these last relations we obtain that

$$
J_{\alpha} = \frac{1}{2} \int \partial^{\alpha} \delta \theta \big[A_j, \delta_j \theta_2 \big] (\partial^{\alpha} \delta \theta).
$$

We now use Lemma 2.8 to get that

$$
|J_\alpha| \lesssim \left(\|\partial^\alpha \delta \theta \|_{L^2} \left\| \mathcal{F} \left[(-\Delta)^{(1-2s)/2} \delta_j \theta_2 \right] \right\|_{L^1} + \| (-\Delta)^s \partial^\alpha \delta \theta \|_{L^2} \left\| \mathcal{F} \left[(-\Delta)^{1/2} \delta_j \theta_2 \right] \right\|_{L^1} \right) \|\partial^\alpha \delta \theta \|_{L^2}.
$$

We now compute each of these terms. Using the Cauchy-Schwarz inequality, we have for any $\varepsilon > 0$ that

$$
\|\mathcal{F}\left[(-\Delta)^{1/2}\delta_j\theta_2\right]\|_{L^1} \leq \int |\xi|^2 |\widehat{\theta}_2(\xi)| d\xi
$$

=
$$
\int \frac{|\xi|^2}{(1+|\xi|)^{3+\varepsilon}} (1+|\xi|)^{3+\varepsilon} |\widehat{\theta}_2(\xi)| d\xi
$$

$$
\leq \left(\int \frac{|\xi|^4}{(1+|\xi|)^{6+2\varepsilon}} d\xi\right)^{1/2} \|\theta_2\|_{H^{3+\varepsilon}}
$$

$$
\lesssim 1
$$

and

$$
\left\| \mathcal{F} \left[(-\Delta)^{(1-2s)/2} \delta_j \theta_2 \right] \right\|_{L^1} \leq \int |\xi|^{2-2s} |\widehat{\theta}_2(\xi)| \, d\xi
$$

$$
\leq \int (1+|\xi|)^2 |\widehat{\theta}_2(\xi)| \, d\xi
$$

$$
= \int \frac{(1+|\xi|)^2}{(1+|\xi|)^{3+\varepsilon}} (1+|\xi|)^{3+\varepsilon} |\widehat{\theta}_2(\xi)| \, d\xi
$$

$$
\leq \left(\int \frac{(1+|\xi|)^4}{(1+|\xi|)^{6+2\varepsilon}} \, d\xi \right)^{1/2} \|\theta_2\|_{H^{3+\varepsilon}}
$$

$$
\lesssim 1.
$$

We compute now the final term using the fact that $\ell - 2s \geq 0$,

$$
\begin{aligned} \|(-\Delta)^s \partial^\alpha \delta \theta \|_{L^2} &= \left(\int \left(|\xi|^{-2s} |\xi|^\ell |\widehat{\delta \theta}(\xi)| \right)^2 \mathrm{d}\xi \right)^{1/2} \\ &\le \left(\int (1 + |\xi|^\ell)^2 |\widehat{\delta \theta}(\xi)|^2 \mathrm{d}\xi \right)^{1/2} \\ &= \|\delta \theta \|_{H^\ell} .\end{aligned}
$$

In conclusion, we have that

and thus for every $\gamma \leq \alpha$,

$$
|J_{\gamma}| \lesssim \|\delta\theta\|_{H^{\ell}}^2. \tag{4.14}
$$

 $|J_{\alpha}| \lesssim \|\delta\theta\|_{H^{\ell}}^2,$

Estimating the terms L_{γ}

Similarly to the L^2 estimate, we have that

$$
\begin{aligned} |L_\gamma|&\leq \|\partial^\gamma\delta H_\rho\|_{L^\infty}\|\partial^{\alpha-\gamma}\nabla\theta_2\|_{L^2}\|\partial^\alpha\delta\theta\|_{L^2}\\ &\lesssim \|\partial^\gamma\delta H_\rho\|_{L^\infty}\|\delta\theta\|_{H^\ell}. \end{aligned}
$$

Observing that $\partial^{\gamma} \delta H_{\rho}$ is a smooth map, we have that

$$
\|\partial^{\gamma}\delta H_{\rho}\|_{L^{\infty}} \le \|\nabla\partial^{\gamma} K_{s,\rho}\|_{L^{\infty}}|\delta z| \lesssim |\delta z|
$$

and thus in the end,

$$
|L_{\gamma}| \lesssim |\delta z| \|\delta \theta\|_{H^{\ell}}.\tag{4.15}
$$

Conclusion

Adding relations (4.34) and (4.35) summed over every $|\alpha| = \ell$, and gathering estimates (4.41), (4.53) and (4.56), we obtain that

$$
\frac{\mathrm{d}}{\mathrm{d}t} \|\delta\theta\|_{H^{\ell}}^2 \lesssim \left(|\delta z| + \|\delta\theta\|_{H^{1-2s}} \right) + |\delta z| \|\delta\theta\|_{H^{\ell}} + \|\delta\theta\|_{H^{\ell}}^2,
$$

which yields that

$$
\frac{\mathrm{d}}{\mathrm{d}t} \|\delta\theta\|_{H^{\ell}} \lesssim |\delta z| + \|\delta\theta\|_{H^{\ell}},
$$

which concludes the proof of Lemma 4.5.

4.4 Conclusion of the proof of Proposition 4.1 and uniqueness in Theorem 1.1

We now conclude the proof of Proposition 4.1.

Gathering the results of Lemma 4.3, Lemma 4.4 and Lemma 4.5, we obtain that there exists a time T_0 such that for every $t \in [0, T_0]$,

$$
\frac{\mathrm{d}}{\mathrm{d}t} \big(\|\delta\theta\|_{H^{\ell}} + |\delta z| \big) \leq C \big(\|\delta\theta\|_{H_{\ell}} + |\delta z| \big),
$$

where T_0 and the constant C depend only on $\|\theta_1(0,\cdot)\|_{H^k}$, $\|\theta_2(0,\cdot)\|_{H^k}$, R_1 , R_2 and T.

By the classical Gronwall's Lemma, we infer that

$$
\|\delta\theta\|_{H^{\ell}} + |\delta z| \leq \left(\|\delta\theta(0,\cdot)\|_{H^{\ell}} + |\delta z(0)| \right) e^{Ct}.
$$

This concludes the proof of Proposition 4.1, and hence of Theorem 1.1- (ii) .

Please note that in the case $s \geq 1/2$ replacing the result of Lemma 4.5 by relation (4.33), then by the same argument using Gronwall's Lemma, we prove Proposition 4.2.

We now prove the uniqueness of strong solutions to the vortex wave system (1.4) as stated in Theorem 1.1- (i) , and whose existence has been proved in Section 3. Let (θ_1, z_1) and (θ_2, z_2) be two solutions with same initial datum (θ_0, z_0) , with

$$
\theta_0 \equiv \beta \quad \text{ on } B(z_0, R_0).
$$

Assume that both solutions are defined on $[0, T)$. We then define \overline{T} the supremum of times such that the two solutions coincide. If $\bar{T} < T$, starting from time \bar{T} , those solutions clearly satisfy the hypotheses of Proposition 4.1 and satisfy the existence of a time $\overline{T}_0 \in (\overline{T}, T)$ such that for every $t \in [\overline{T}, T_0]$,

$$
\|\delta\theta(t)\|_{H^1} + |\delta z(t)| \lesssim \|\delta\theta(\bar{T})\|_{H^1} + |\delta z(\bar{T})| = 0,
$$

which is absurd by definition of \overline{T} . This proves that $\overline{T} = T$ and therefore that maximal strong solutions are unique. \Box

5 Blow-up criterion

A first obstacle to obtain global existence of strong solutions to the vortex-wave system is that the active scalar, whose self-interaction is governed by the SQG equations, can blow-up on his own and lose regularity. Although we believe that a proof that this blow-up can actually happen is not known, there are numerical evidences suggesting it [32]. Moreover, a second type of blow-up, specific to the vortex-wave system, may happen. Indeed, the proof of existence of solutions requires that the active scalar should be locally constant around the point-vortex. If the distance to the non-constant part $t \mapsto R(t)$ goes to 0 in finite time, the equations become singular, resulting in a loss of regularity. In this article we investigate this second type of blow-up.

5.1 The case where $R(t)$ vanishes : proof of Theorem 1.2

For any $f \in C^0(\mathbb{R}^2)$, $z \in \mathbb{R}^2$ and $R \in (0,1)$, we introduce the quantity $\mathcal{N}(f, z, R) \in (-\infty; +\infty]$ given by

$$
\mathcal{N}(f, z, R) := \max_{\{x \in \mathbb{R}^2, |x - z| = R\}} \frac{-(x - z) \cdot (f(x) - f(z))}{R^2 (1 - \ln R)},
$$

which evaluates the behavior of the radial component of $f(x) - f(z)$ near z. This quantity is a directional Log-Lipschitz norm. In particular, it satisfies that

$$
|\mathcal{N}(f,z,R)| \le ||f||_{LL} := \sup_{0 < |x-y| < 1} \frac{|f(x) - f(y)|}{|x-y|(1-\ln|x-y|)}.
$$

We prove the following blow-up criterion.

Proposition 5.1 (Criterion for the shrink of the constant part). Let (θ, z) be a strong solution of (1.4) on $[0, T^*)$ with $T^* < \infty$, such that

$$
\theta_0 \equiv \beta \qquad on \ B(z(0), R_0),
$$

for some $\beta \in \mathbb{R}$. We recall the definition of $t \mapsto R(t)$ given in relation (1.7). Then,

$$
R(t) \underset{t \to T^*}{\longrightarrow} 0 \iff \int_0^{T^*} \mathcal{N}(v(t,\cdot), z(t), R(t)) \mathrm{d}t = +\infty.
$$

Proof. We start similarly as in the proof of Lemma 3.4. Let $x \in \mathbb{R}^2$ such that $\theta(t, x) \neq \beta$. Let $\tau \mapsto X(\tau)$ be the trajectory of the fluid particle passing in x at time t . Then

$$
\frac{d}{d\tau}|X(\tau) - z(\tau)|\Big|_{\tau=t} = \frac{x - z(t)}{|x - z(t)|} \cdot (v(x, t) + H(x, t) - v(z(t), t))
$$
\n
$$
= \frac{x - z(t)}{|x - z(t)|} \cdot (v(x, t) - v(z(t), t))
$$
\n
$$
\geq \min_{\{y \in \mathbb{R}^2, |y - z(t)| = R(t)\}} \frac{y - z(t)}{|y - z(t)|} \cdot (v(y, t) - v(z(t), t))
$$
\n
$$
= -\max_{\{y \in \mathbb{R}^2, |y - z(t)| = R(t)\}} -\frac{y - z(t)}{|y - z(t)|} \cdot (v(y, t) - v(z(t), t))
$$

since $(x - z(t)) \cdot H(x, t) = 0$. Since $v \in C^1([0, T^*))$, the quantity $\mathcal{N}(v(t, \cdot), z(t), R(t))$ is well defined and finite by assuming without lost of generality that $R(0) < 1$. We then obtain that R satisfies the differential inequality

$$
\frac{\mathrm{d}}{\mathrm{d}t}R(t) \ge -\mathcal{N}\left(v(t,\cdot),z(t),R(t)\right)R(t)(1-\log(R(t)),
$$

from which we infer, as a particular case of Osgood's uniqueness theorem that for every $t \in [0, T^*)$,

$$
R(t) \ge \exp\left(-\exp\left(\int_0^t \mathcal{N}\big(v(\tau,\cdot),z(\tau),R(\tau)\big)\mathrm{d}\tau + \ln\big(1-\ln(R(0))\big)\right) + 1\right).
$$

and thus

$$
R(t) \underset{t \to T^*}{\longrightarrow} 0 \Longleftrightarrow \int_0^{T^*} \mathcal{N}(v(t,\cdot), z(t), R(t)) \mathrm{d}t = +\infty.
$$

Theorem 1.2 is a simple reformulation of Proposition 5.1 with $N(t) := \mathcal{N}(v(t, \cdot), z(t), R(t))$.

5.2 Proof of Theorem 1.1- (iii) and remarks

From Proposition 5.1, we obtain immediately Theorem 1.1- (iii) from the observation that

$$
N(t) \leq ||v(t)||_{LL} \lesssim ||\theta(t)||_{H^{3-2s}}.
$$

Let us mention that a naive approach to obtain a blow-up criterion could be to conclude directly from Lemma 3.4 that ∗

$$
R(t) \underset{t \to T^*}{\longrightarrow} 0 \implies \int_0^{T^*} \|\nabla v(t, \cdot)\|_{L^\infty} dt = +\infty.
$$

From this, we only infer that for every $\varepsilon > 0$,

$$
\int_0^{T^*} \|\theta(t,\cdot)\|_{H^{3-2s+\varepsilon}} \mathrm{d}t = +\infty,
$$

which is a weaker result than Theorem 1.1-(iii). Moreover, the quantity $t \mapsto N(t)$ is much more precise than $\|\nabla v(t)\|_{L^{\infty}}$ or even than $\|v(t)\|_{LL}$. In particular, let us observe that since $(x-z(t)) \cdot H(t,x) = 0$, the point-vortex seems to play no role into the possible vanishing of R. This is not a rigorous assertion at all since θ is affected by the velocity field H generated by the point-vortex. This make us think that should such a blow-up exist, then it might come from the same mechanism than the blow-up of the SQG equations alone.

6 Global Existence of weak solutions

6.1 Proof of Theorem 1.3

Let $\varepsilon > 0$. To start-with, we consider the functions θ_{ε} , z_{ε} , v_{ε} and H_{ε} global solution to the dynamic associated to the regularized kernel $K_{s,\varepsilon}$ (see Section 2.3 for details). Using the preservation of the L^p norms by the flow stated by (2.27), we have that $(\theta_{\varepsilon})_{\varepsilon>0}$ is bounded in $L^{\infty}(\mathbb{R}_+ \times \mathbb{R}^2)$. Therefore, by the Banach-Alaoglu theorem, we have θ_{ε} that converges weakly- \star in $L^{\infty}(\mathbb{R}_{+} \times \mathbb{R}^{2})$ up to an omitted extraction.

The right-hand side of (2.23) is (up to a multiplicative constant) the Green function of the fractional Laplace operator $(-\Delta)^{\sigma}$ with $\sigma = s - 1/2$. Note that we have $\sigma > 0$ by hypothesis. Therefore, using properties of homogeneous Fourier multipliers on Besov spaces (see [3, §2.3]), we have $v(t,.) \in B^{\sigma}_{p,p}(\mathbb{R}^2)$. More precisely, there exists a constant $C > 0$ such that

$$
\forall t > 0, \qquad \|\mathbf{v}_{\varepsilon}(t,.)\|_{B^{\sigma}_{p,p}} \leq C_T \|\theta_0\|_{L^p}
$$
\n(6.1)

for all $p \in [2, +\infty)$. By Sobolev inequalites, we have for p large enough that $B^{\sigma}_{p,p}(\mathbb{R}^2)$ is continuously embedded in the space of Hölder continuous fonctions $\mathcal{C}^{0,\sigma'}(\mathbb{R}^2) =: B_{\infty,\infty}^{\sigma'}$ with $0 < \sigma' < \sigma$. With (2.23) , we have that $K_{s,\varepsilon}$ converges strongly in $L^1_{loc}(\mathbb{R}^2)$ by the Lebesgue dominated convergence theorem. This gives as $\varepsilon \to 0$:

$$
v_{\varepsilon} := K_{s,\varepsilon} \star \theta_{\varepsilon} \longrightarrow v := K_s \star \theta, \quad \text{strongly in } L^1_{\text{loc}} \cap L^{\infty}(\mathbb{R}^+ \times \mathbb{R}^2).
$$

Thus, for any $\psi \in \mathcal{D}([0,T) \times \mathbb{R}^2)$ with $T > 0$ fixed,

$$
\int_0^T \int_{\mathbb{R}^2} \theta_{\varepsilon}(t,x) \, v_{\varepsilon}(t,x) \cdot \nabla \psi(t,x) \, dx \, dt \longrightarrow \int_0^T \int_{\mathbb{R}^2} \theta(t,x) \, v(t,x) \cdot \nabla \psi(t,x) \, dx \, dt. \tag{6.2}
$$

The weak convergence of θ_{ε} as $\varepsilon \to 0$ implies also

$$
\int_0^T \int_{\mathbb{R}^2} \theta_{\varepsilon}(t,x) \frac{\partial \psi}{\partial t}(t,x) dx dt \longrightarrow \int_0^T \int_{\mathbb{R}^2} \theta(t,x) \frac{\partial \psi}{\partial t}(t,x) dx dt.
$$
 (6.3)

Similarly as before we have $v \in C^{0,\sigma'}$. We can then define $t \mapsto z(t)$ using (1.22). The uniform convergence of v_{ε} towards v implies the uniform convergence of z_{ε} towards z. We observe that, since v is bounded, we have $z \in C^{0,1}([0,T);\mathbb{R}^2)$. This is enough to get the convergence:

$$
H_{\varepsilon}(t,x) := K_{s,\varepsilon}(x - z_{\varepsilon}(t)) \longrightarrow H(t,x) := K_{s}(x - z(t)), \quad \text{strongly in } L^{1}_{\text{loc}}([0,T) \times \mathbb{R}^{2}).
$$

Thus,

$$
\int_0^T \int_{\mathbb{R}^2} \theta_{\varepsilon}(t,x) \, H_{\varepsilon}(t,x) \cdot \nabla \psi(t,x) \, dx \, dt \longrightarrow \int_0^T \int_{\mathbb{R}^2} \theta(t,x) \, H(t,x) \cdot \nabla \psi(t,x) \, dx \, dt \tag{6.4}
$$

Since $t \mapsto z(t)$ satisfies (1.22) then with (6.3)(6.4)(6.6) we have the existence of (θ, z) weak solution to the SQG vortex-wave system $(1.11)(1.12)$.

Finally, since the weak convergence decreases the L^p norms, we have or all $p \in [1, +\infty]$ that $\|\theta(t,.)\|_{L^p} \le$ $\|\theta_{\varepsilon}(t,.)\|_{L^p} = \|\theta_{\varepsilon}(t,.)\|_{L^p}$, where the last equality is given by (2.27).

The only point that remains to be proved is the continuity in time of $t \mapsto \theta(t,.)$ with values in $L^1 \cap L^\infty(\mathbb{R}^2)$ endowed with the weak- \star topology. For $t_0, t \in \mathbb{R}_+$ and for φ a test function on \mathbb{R}^2 :

$$
\left| \langle \theta(t,.) - \theta(t_0,.), \varphi \rangle \right| \leq \left| \langle \theta(t,.) - \theta_{\varepsilon}(t,.), \varphi \rangle \right| + \left| \langle \theta_{\varepsilon}(t,.) - \theta_{\varepsilon}(t_0,.), \varphi \rangle \right| + \left| \langle \theta_{\varepsilon}(t_0,.) - \theta(t_0,.), \varphi \rangle \right|.
$$

We now remark that the estimate (2.45) still holds independently on N using the same reasoning from the evolution equation. As a consequence, we have:

$$
\left| \langle \theta_{\varepsilon}(t_0,.) - \theta_{\varepsilon}(t,.),\varphi \rangle \right| \ = \ \left| \int_{t_0}^t \int_{\mathbb{R}^2} \frac{\partial \theta_{\varepsilon}}{\partial t}(s,x) \, \varphi(x) \, dx \, ds \right| \leq (t-t_0) \|\varphi\|_{H^1} \Big(\|K_{s,\varepsilon}\|_{L^\infty} + \|K_{s,\varepsilon}\|_{L^2} \|\theta_{\varepsilon}\|_{L^2} \Big) \|\theta_{\varepsilon}\|_{L^2}.
$$

For fixed values of ε , this term vanishes as $t \to t_0$. We also have that $(\theta(t,.) - \theta_{\varepsilon}(t,.) , \varphi)$ vanishes as $\varepsilon \to 0$ uniformly with respect to $t \in [0, T]$ as a consequence of (6.4). Thus, $\langle \theta(t,.) - \theta(t_0,.) , \varphi \rangle \to 0$ as $t \to t_0$

6.2 Proof of Theorem 1.8

For the proof of the global existence of V-weak solutions to the vortex-wave system when $s = 1/2$, ie: Theorem 1.8- (i) et (ii) , we follow closely the construction by Marchand [27]. This allows us to make use of the technical lemmas previously in [27] and ease the reading of this proof and focus on what is new. Indeed, the main novelties of this proof lay in the convergence of the term of interaction between the ODE part and the PDE part.

6.2.1 Step 1: regularization of the system

To start-with, we consider the regularization of the kernel $K_{\frac{1}{2},\varepsilon}$ introduced at (2.23). By Proposition 2.9, the vortex-wave system associated to this regularized kernel admits a global smooth solution $(\theta_{\varepsilon}, z_{\varepsilon})$ with the respective associated velocity fields v_{ε} and H_{ε} .

Since the total velocity field $(v_{\varepsilon} + H_{\varepsilon})$ is divergence-free, the flow preserves all the L^p norm of functions. As a consequence, by the Banach-Alaoglu theorem, we have weak convergence of θ_{ε} towards some function $\theta \in L^{\infty}(\mathbb{R}_+; L^1 \cap L^{\infty}(\mathbb{R}^2))$ (up to an omitted extraction of subsequence) as $\varepsilon \to 0$. By property of the weak convergence:

$$
\forall t > 0, \qquad \|\theta(t,.)\|_{L^p} \le \liminf_{\varepsilon \to 0} \|\theta_{\varepsilon}(t,.)\|_{L^p} \le \limsup_{\varepsilon \to 0} \|\theta_{\varepsilon}(0,.)\|_{L^p} = \|\theta(0,.)\|_{L^p}.
$$
 (6.5)

We also have for all $T > 0$ and $\psi \in \mathcal{D}([0, T) \times \mathbb{R}^2)$,

$$
\int_0^T \int_{\mathbb{R}^2} \frac{\partial \psi}{\partial t}(t, x) \, \theta_{\varepsilon}(t, x) \, dx \, dt \longrightarrow \int_0^T \int_{\mathbb{R}^2} \frac{\partial \psi}{\partial t}(t, x) \, \theta(t, x) \, dx \, dt.
$$

Similarly as (6.1), a simple triangular inequality give the continuity in time with value in $L^{\infty} \cap L^{1}$ endowed with the weak- \star topology.

To conclude the proof of Theorem 1.8-(i), we study the convergence of the other terms as $\varepsilon \to 0$.

6.2.2 Step 2: convergence of the non-linear term

We now study the non-linear term using the commutator formulation (1.16) . We introduce the following notation for the associated bilinear form:

$$
\mathcal{I}_{\psi}(\theta,\vartheta) := \int_{\mathbb{R}^2} K_{\frac{1}{2}} \star \theta \cdot \left[(-\Delta)^{\frac{1}{2}}, \nabla \psi \right] (-\Delta)^{-\frac{1}{2}} \vartheta(x) dx
$$

To study the convergence of this term, we make use of the following technical lemma which are proved in [27]. First, we use the following estimate on the high-frequencies self interactions:

Lemma 6.1 (Proposition 2.3 in [27]). Let $\frac{4}{3} < p < 2$. For $\theta \in L^p$, the distribution $K_s \star \theta \cdot \theta$ is well-defined using (1.16) as a consequence of the following estimate on the high-frequencies:

$$
\left|\mathcal{I}_{\psi}\big(H_j\theta, H_j\theta\big)\right| \leq C 2^{-j\left(3-\frac{4}{p}\right)} \|\nabla^2 \psi\|_{L^\infty} \|H_j\theta\|_{L^p}^2,
$$

The proof of this result relies on Littlewood-Paley decomposition, the commutator estimates given by Lemma 2.7 and Sobolev embeddings. Note that a consequence of this lemma is that $\mathcal{I}(\theta,\theta)$ is well-defined for all $\theta \in L^p$ with $4/3 < p < 2$.

We also need the following compactness result for the low frequencies of θ :

Lemma 6.2 (Lemma 9.2 in [27]). Let $(\theta_n)_{n \in \mathbb{N}}$ be a sequence bounded in $L^{\infty}(\mathbb{R}_+; L^p(\mathbb{R}^2))$ for $1 < p < +\infty$. Then for every $j \in \mathbb{Z}$, the sequences $(S_j \theta_n)_{n \in \mathbb{N}}$ and $(K_{\frac{1}{2}} \star (S_j \theta_n))_{n \in \mathbb{N}}$ are pre-compact in $L^{\frac{p}{p-1}}(\mathbb{R}_+ \times \mathbb{R}^2)$.

The proof of this lemma uses the smoothing properties of the operators (S_i) and a compact embedding argument.

Lemma 6.3. Let θ_{ε} be a solution in the regularized vortex-wave system (2.24). We have convergence of the non-linear term: $\mathcal{I}_{\psi}(\theta_{\varepsilon}, \theta_{\varepsilon}) \longrightarrow \mathcal{I}_{\psi}(\theta, \theta)$.

Proof. we decompose the studied non-linear term between low and high frequencies (we fix $j \in \mathbb{Z}$):

$$
\mathcal{I}_{\psi}(\theta_{\varepsilon}, \theta_{\varepsilon}) = \mathcal{I}_{\psi}(S_j \theta_{\varepsilon}, S_j \theta_{\varepsilon}) + \mathcal{I}_{\psi}(S_j \theta_{\varepsilon}, H_j \theta_{\varepsilon}) + \mathcal{I}_{\psi}(H_j \theta_{\varepsilon}, S_j \theta_{\varepsilon}) + \mathcal{I}_{\psi}(H_j \theta_{\varepsilon}, H_j \theta_{\varepsilon}).
$$
\n(6.6)

It is a direct consequence of Lemma 6.2 that the three first integrals appearing in the decomposition (6.13) are converging as $\varepsilon \to 0$ towards their expected limit quantity:

$$
\mathcal{I}_{\psi}(\mathbf{S}_{j}\theta_{\varepsilon},\mathbf{S}_{j}\theta_{\varepsilon}) \longrightarrow \mathcal{I}(\mathbf{S}_{j}\theta,\mathbf{S}_{j}\theta),\mathcal{I}_{\psi}(\mathbf{S}_{j}\theta_{\varepsilon},\mathbf{H}_{j}\theta_{\varepsilon}) \longrightarrow \mathcal{I}(\mathbf{S}_{j}\theta,\mathbf{H}_{j}\theta),\mathcal{I}_{\psi}(\mathbf{H}_{j}\theta_{\varepsilon},\mathbf{S}_{j}\theta_{\varepsilon}) \longrightarrow \mathcal{I}(\mathbf{H}_{j}\theta,\mathbf{S}_{j}\theta).
$$

Concerning the remaining term (ie: the high-frequency self interaction), we simply write, using Lemma 6.1 and Equation (6.9):

$$
\left| \mathcal{I}_{\psi}(\mathbf{H}_{j} \theta_{\varepsilon}, \mathbf{H}_{j} \theta_{\varepsilon}) \right| + \left| \mathcal{I}_{\psi}(\mathbf{H}_{j} \theta, \mathbf{H}_{j} \theta) \right| \leq C_{0} 2^{-j}
$$

where the constant C_0 above depends on the test function ψ and on the initial datum θ_0 . We conclude

$$
\limsup_{\varepsilon \to 0} \left| \mathcal{I}_{\psi}(\theta_{\varepsilon}, \theta_{\varepsilon}) - \mathcal{I}_{\psi}(\theta, \theta) \right| \leq C_0 2^{-j}
$$

Since the estimate above holds for all $j \in \mathbb{Z}$, we infer as $\varepsilon \to 0$:

$$
\mathcal{I}_{\psi}(\theta_{\varepsilon}, \theta_{\varepsilon}) \longrightarrow \mathcal{I}_{\psi}(\theta, \theta). \tag{6.7}
$$

,

 \Box

From the convergence (6.17), we can conclude that we have the expected convergence of the non-linear term:

$$
\int_{\mathbb{R}^2} K_{\frac{1}{2},\varepsilon} \star \theta_{\varepsilon}(x) \cdot \nabla \psi(x) \, \theta_{\varepsilon}(x) \, dx \longrightarrow \mathcal{I}_{\psi}(\theta,\theta).
$$

6.2.3 Step 3: convergence of the ODE

Lemma 6.4. Let z_{ε} be the solution to the ODE in the regularized vortex-wave system (2.24). As $\varepsilon \to 0$, it converges towards $z \in C^{0,1}(\mathbb{R}_+;\mathbb{R}^2)$ solution to

$$
z(t) = z_0 + \int_0^t \int_{\mathbb{R}^2} v(t, x) \, \chi(x - z(t)) \, \mathrm{d}x. \tag{6.8}
$$

Proof. To start with, we recall that

$$
\frac{\mathrm{d}z_{\varepsilon}}{\mathrm{d}t}(t) = \int_{\mathbb{R}^2} v_{\varepsilon}(t,.) \,\chi\big(x-z_{\varepsilon}(t)\big) \,\mathrm{d}x.
$$

The kernel $\nabla^{\perp}(-\Delta)^{-\frac{1}{2}}$ is associated to the Fourier multiplyer $\xi^{\perp}/|\xi|$, which is a bounded Fourier multiplyer (see Lemma 2.6). On the other hand, since θ_{ε} converges weakly towards θ then the Fourier transform $\widehat{\theta_{\varepsilon}}$ converges weakly towards θ . Therefore, we get that $v_{\varepsilon}(t,.) := K_{\frac{1}{2}} \star \theta_{\varepsilon}(t,.)$ converges weakly towards $v(t,.) := K_{\frac{1}{2}} \star \theta(t,.)$

On the other hand, it is a general property of functions in $W^{1,p}$ that

$$
\left\| \chi(.-z_2) - \chi(.-z_1) \right\|_{L^p} \leq \|\nabla \chi\|_{L^p} |z_2 - z_1|.
$$

Therefore, fixing a $t > 0$, we have for any sequence ζ_{ε} converging towards some $\zeta \in \mathbb{R}^2$,

$$
\int_{\mathbb{R}^2} v_{\varepsilon}(t,.) \, \chi(x - \zeta_{\varepsilon}) \, dx \longrightarrow \int_{\mathbb{R}^2} v(t,.) \, \chi(x - \zeta) \, dx
$$

Then, by a bootstrap argument, we have convergence of dz_ε/dt towards $\int_{\mathbb{R}^2} v_\varepsilon(t,x) \chi(x-z_\varepsilon(t)) dx$ for every fixed $t > 0$ and thus we have local uniform convergence of z_{ε} towards a z that satisfy (6.19). \Box

6.2.4 Convergence of the singular part of the velocity field

To study the convergence of H_{ε} , the singular part of the velocity field (ie the velocity field generated by the point-vortex), it is necessary to exploit the fact that we only study V -weak solutions. In this parts and in the next part of the proof, we manipulate this notion of V-weak solutions. The objective is to show that this notion permits to obtain a non-trivial object in the presence of a localized singularity (in our case it is the point-vortex) even if the singularity is non-integrable.

It is now time to introduce the set V we are going to work with:

Lemma 6.5. Let $H_{\varepsilon}, z_{\varepsilon}$ defined by (2.24) and let $t \mapsto z(t)$ the limit of z_{ε} as $\varepsilon \to 0$ (see Lemma 6.4. Let $H(t,x) := K_{\frac{1}{2}}(x-z(t)).$ Let $\delta > 0$, we define

$$
V_{\delta} := \left\{ \psi \in \mathcal{D}(\mathbb{R}_{+} \times \mathbb{R}^{2}) : \forall t > 0, \quad \forall x \in \mathcal{B}(z(t), \delta), \quad \nabla \psi(t, x) = 0. \right\}.
$$

Then, for all $\psi \in V_{\delta}$, we have $\nabla \psi \cdot H \in L^1$ and for all $t > 0$:

$$
\int_{\mathbb{R}^2} \nabla \psi(t,x) \cdot H_{\varepsilon}(t,x) \,dx \longrightarrow \int_{\mathbb{R}^2} \nabla \psi(t,x) \cdot H(t,x) \,dx.
$$

Proof. The idea is to choose V so that the most singular part (that lays in a moving point) vanishes. We recall the definition of the regularization kernel from Subsection 2.3 where we introduced the functions

 $H_{\varepsilon}(t,x) := K_{\frac{1}{2},\varepsilon}(x - z_{\varepsilon}(t)), \quad \text{and} \quad H_{\varepsilon}(t,x) := K_{\frac{1}{2},\varepsilon}(x - z(t))$

To start with, we have for all $\psi \in V_{\delta}$, on any fixed interval of time $[0, T)$,

$$
\forall (t, x) \in [0, T) \times \mathbb{R}^2, \qquad \widetilde{H}_{\varepsilon}(t, x) \cdot \nabla \psi(t, x) = H(t, x) \cdot \nabla \psi(t, x)
$$

provided that $\varepsilon \leq \delta$. Indeed, the two functions $\widetilde{H}_{\varepsilon}(t, \cdot)$ and $H(t, \cdot)$ coincide outside $\mathcal{B}(z(t), \delta)$ while $\nabla \psi(t, \cdot)$ vanishes inside this ball. On the other hand, if we assume that ε is so small that for all $t \in [0, T]$, we have $|z_{\varepsilon}(t) - z(t)| \leq \delta/2$ and $\varepsilon \leq \delta/2$, then, using again $\nabla \psi(t,.) \equiv 0$ on $\mathcal{B}(z(t),\delta)$,

$$
\int_{\mathbb{R}^2} \nabla \psi(t, x) \cdot \left(H_{\varepsilon}(t, x) - \widetilde{H}_{\varepsilon}(t, x) \right) dx
$$
\n
$$
= \int_{\mathbb{R}^2 \setminus \mathcal{B}(z(t), \delta)} \nabla \psi(t, x) \cdot \left(K_{\frac{1}{2}, \varepsilon}(x - z_{\varepsilon}(t)) - K_{\frac{1}{2}, \varepsilon}(x - z(t)) \right) dx
$$
\n
$$
= \int_{\mathbb{R}^2 \setminus \left(\mathcal{B}(z(t), \frac{\delta}{2}) \cup \mathcal{B}(z_{\varepsilon}(t), \frac{\delta}{2}) \right)} \nabla \psi(t, x) \cdot \left(K_{\frac{1}{2}, \varepsilon}(x - z_{\varepsilon}(t)) - K_{\frac{1}{2}, \varepsilon}(x - z(t)) \right) dx
$$
\n
$$
= \int_{\mathbb{R}^2} \nabla \psi(t, x) \cdot \left(K_{\frac{1}{2}, \frac{\delta}{2}}(x - z_{\varepsilon}(t)) - K_{\frac{1}{2}, \frac{\delta}{2}}(x - z(t)) \right) dx.
$$

Thus,

$$
\left| \int_{\mathbb{R}^2} \nabla \psi(t,x) \cdot \left(H_{\varepsilon}(t,x) - \widetilde{H}_{\varepsilon}(t,x) \right) dx \right| \leq \|\nabla \psi(t,.)\|_{L^2} \left\| \nabla K_{\frac{1}{2},\frac{\delta}{2}}(t,.) \right\|_{L^2} |z_{\varepsilon}(t) - z(t)|.
$$

This implies the announced convergence.

Gathering all the convergences established before, we can conclude the proof of Theorem $1.8-(i)$ by setting $V := \bigcup_{\delta > 0} V_{\delta}.$

6.2.5 Conclusion of the proof of Theorem 1.8

Concerning Theorem 1.8-(*ii*), we already have for all $\psi \in V$ that $H \cdot \nabla \psi \in L^1$. For the *Γ*-coincident property, we note that $V \subseteq \mathcal{D}(\mathbb{R}_+ \times \mathbb{R}^2)$. Conversely, let $\Omega_0 \subseteq \mathbb{R}_+ \times \mathbb{R}^2$ such that $\delta_0 := \inf_{x \in \Omega_0} \inf_{y \in \Gamma} |x - y| > 0$. For a fixed $\varphi \in \mathcal{D}(\mathbb{R}_+ \times \mathbb{R}^2)$, it is possible to construct a $\psi \in V_{\delta_0/2}$ such that $\varphi \equiv \psi$ on Ω_0 by the Urysohn's lemma. More precisely, let $\chi : \mathbb{R}_+ \to \mathbb{R}_+$ monotonous, \mathcal{C}^{∞} and such that $\chi(s) = 1$ if $s \le \delta_0/2$ and $\chi(s) = 0$ if $s \ge \delta_0$. We define

$$
\psi_{\delta_0}(t,x) := \chi(\|x - z(t)\|) \int_{\mathcal{B}\left(z(t), \frac{\delta_0}{2}\right)} \varphi(t,x) \, \mathrm{d}x + \left(1 - \chi(\|x - z(t)\|)\right) \varphi(t,x),\tag{6.9}
$$

 \Box

where the barred integral denotes the mean value. It is direct to check that $\psi_{\delta_0} \in V_{\delta_0} \subset V$ and that $\psi_{\delta_0} \equiv \varphi$ in Ω_0 .

To prove Theorem 1.8-(iii), we assume that a V-weak solution (θ, z) has regularity \mathcal{C}^1 on an interval of time $[0, T)$ with $T \in (0, +\infty]$ and that $\theta(t,.)$ is constant on a neighborhood of $z(t)$ for all $t \in [0, T)$. We prove here that, with such regularity, this V -weak solution is actually a solution in the classical sense.

To start with, by hypothesis, there exists $\delta_0 > 0$ such that $\nabla \theta(t,.) \equiv 0$ on the ball $\mathcal{B}(z(t), \delta)$ for all $t \in [0, T)$. Let $0 < \delta < \delta_0$, let $\varphi \in \mathcal{D}([0,T) \times \mathbb{R}^2)$ and let ψ_{δ} defined by (6.29). The fact that θ is constant on $\mathcal{B}(z(t),\delta)$ implies

$$
\forall t \in [0, T), \quad \forall x \in \mathbb{R}^2, \qquad \theta(t, x) \nabla \varphi(t, x) = \theta(t, x) \nabla \psi_{\delta}(t, x). \tag{6.10}
$$

On the other hand, since $\psi_{\delta} \in V$, we use that θ is a V-weak solution to write

$$
\int_0^T \int_{\mathbb{R}^2} \left(\frac{\partial \psi_\delta}{\partial t}(t, x) + v(t, x) \cdot \left[(-\Delta)^s, \nabla \psi_\delta(t, x) \right] (-\Delta)^{-s} \n+ H(t, x) \cdot \nabla \psi_\delta(t, x) \right) \theta(t, x) dx dt + \int_{\mathbb{R}^2} \omega_0(x) \psi_\delta(0, x) dx = 0,
$$

Using Lemma 1.4, since θ is \mathcal{C}^1 , gives

$$
\int_0^T \int_{\mathbb{R}^2} \left(\frac{\partial \psi_\delta}{\partial t}(t,x) + \left(v(t,x) + H(t,x) \right) \cdot \nabla \psi_\delta(t,x) \right) \theta(t,x) \, dx \, dt + \int_{\mathbb{R}^2} \omega_0(x) \, \psi_\delta(0,x) \, dx = 0,
$$

Using (6.30), we can replace $\nabla \psi_{\delta}$ by $\nabla \varphi$ in the equality above. It then becomes possible to pass to the limit $\delta \rightarrow 0$ and get

$$
\int_0^T \int_{\mathbb{R}^2} \left(\frac{\partial \varphi}{\partial t}(t, x) + \left(v(t, x) + H(t, x) \right) \cdot \nabla \varphi(t, x) \right) \theta(t, x) dx dt + \int_{\mathbb{R}^2} \omega_0(x) \varphi(0, x) dx = 0,
$$

Since θ is a \mathcal{C}^1 function and since φ is any function in $\mathcal{D}([0,T)\times\mathbb{R}^2)$, we conclude that θ satisfy point-wise

$$
\frac{\partial \theta}{\partial t}(t,x) + \left(v(t,x) + H(t,x)\right) \cdot \nabla \theta(t,x) = 0, \quad \text{and} \quad \theta(0,x) = \omega_0(x).
$$

Similar manipulations give that $t \mapsto z(t)$ satisfy the quasi-point-vortex equation point-wise.

7 The case of several vortices

In the previous sections, we studied for the sake of clarity the case of a single point-vortex interacting with the background. Naturally, the results of this paper can be extended to the case of several point-vortices at least in non degenerate cases. Although this is not the main topic of this paper and we will not provide a proof of this claim, let us discuss briefly how to define and deal with several point-vortices.

Consider several point-vortices $(z_i)_{1\leq i\leq N}$ of intensities $a_i \in \mathbb{R}^*$. The vortex-wave system then writes

$$
\begin{cases}\nv(t,x) = K_s \star \theta(t,x), \\
H_i(t,x) = K_s(x-z_i(t)), \quad \forall i \in \{1,\ldots,N\}, \\
\frac{\partial \theta}{\partial t} + \nabla \cdot \left[\left(v + \sum_{i=1}^N a_i H_i \right) \theta \right] = 0, \\
\frac{dz_i}{dt}(t) = v(z_i(t),t) + \sum_{j \neq i} a_j H_j(z_i(t),t), \quad \forall i \in \{1,\ldots,N\}.\n\end{cases}
$$
\n(7.1)

 \Box

A common method in the study of point-vortices used to reduce this problem to the case of a single pointvortex is to consider the influence of other point-vortices as an exterior field. In that case we consider the equations

$$
\begin{cases}\nv(t,x) = K_s \star \theta(t,x), \\
H(t,x) = K_s(x - z(t)), \\
\frac{\partial \theta}{\partial t} + \nabla \cdot \left[\left(v + aH + F \right) \theta \right] = 0, \\
\frac{dz}{dt}(t) = v(z_i(t), t) + F(z_i(t), t),\n\end{cases}
$$

with F being a bounded, Lipschitz, divergence free map. It is then not difficult to reproduce the arguments of local existence and uniqueness of strong solutions for this new system. The remaining part consists in proving that the exterior field F defined as the influence of the other point-vortices retains its properties locally in time. This is the case if the active scalar is locally constant around the point-vortices at time 0 and if the point-vortices remain far from each other.

The main difficulty arising when considering several point-vortices is that there exists a new type of blow-up: collapses of point-vortices. Assume that

$$
\forall i \in \{1, \ldots, N\}, \quad \theta_0 \equiv \beta_i \quad \text{ on } B(z_i(0), R_i(0)),
$$

for some constants β_i and $R_i(0) > 0$ and let

$$
R_i(t) := \sup \{ r \ge 0, \ \theta(t) \equiv \beta_i \text{ on } B(z_i(t), r) \}.
$$

If for some $T > 0$,

$$
\lim_{t \to T} |z_i(t) - z_j(t)| = 0,\t\t(7.2)
$$

which is what we call a collision of the point-vortices z_i and z_j , then if $\beta_i \neq \beta_j$ this implies that

$$
\lim_{t \to T} R_i(t) = 0 \quad \text{and } \lim_{t \to T} R_j(t) = 0,
$$

which is a more complicated version of the type of blow up that we already discussed in Section 5. However if $\beta_i = \beta_i$ it could be possible that (7.5) holds true, while having that $R_i(t) > m$ and $R_i(t) > m$, with $m > 0$, for every $t \in [0, T)$. This is a new type of blow-up, which is a classical point-vortex collapse in an exterior field that this time represents the interaction with the background.

Both in the context of the Euler or SQG point-vortex system, if the intensities a_i have all the same sign, then collapses can not happen and the solution is always global in time. For the Euler vortex-wave system, this remains true: there exists a global in time strong solution when the initial vorticity is locally constant around the point-vortices, see [28]. For the general SQG vortex-wave system this may not be true anymore because of the possible regularity blow-up of the background. Indeed, we prove that a collapse of all positive point-vortices may happen if at least one of the R_i goes to 0, hence if a blow-up as described in Section 5 happens.

Proposition 7.1. Let $(\theta, (z_i)_i)$ be a classical (in the sense of Theorem 1.1-(i)) solution to the vortex-wave system (7.1) with N points, on at least $[0, T^*)$. If all the intensities have the same sign, then for a collision of point-vortices to happen at time T^* one must have that

$$
\min_{1 \le i \le N} R_i(t) \underset{t \to T^*}{\longrightarrow} 0.
$$

Proof. Assume without loss of generality that $a_i > 0$ for every $i \in \{1, ..., N\}$. Let

$$
H(t) = \sum_{i \neq j} a_i a_j \frac{1}{|z_i(t) - z_j(t)|^{2-2s}}
$$

and

$$
I(t) = \sum_{i=1}^{N} a_i z_i^2(t).
$$

Then

$$
\frac{d}{dt}I(t) = 2C_s \sum_{i=1}^{N} a_i z_i \cdot \left(\sum_{j \neq i} a_j \frac{(z_i(t) - z_j(t))^{1}}{|z_i(t) - z_j(t)|^{4-2s}} + v(t, z_i(t)) \right)
$$
\n
$$
= 2C_s \sum_{i \neq j} a_i a_j \frac{z_i(t) \cdot z_j^{1}(t)}{|z_i(t) - z_j(t)|^{4-2s}} + 2C_s \sum_{i=1}^{N} a_i z_i(t) \cdot v(t, z_i(t))
$$
\n
$$
= 2C_s \sum_{i=1}^{N} a_i z_i(t) \cdot v(t, z_i(t))
$$
\n
$$
\leq C(I(t) + 1)V(t),
$$

were we used that $z_i(t) \leq z_i^2(t) + 1$. Therefore, there exists a constant C such that

$$
\forall t \in [0, T^*), \quad I(t) \le C \exp\left(C \int_0^t V(t') \mathrm{d}t'\right),
$$

and thus

$$
\forall t \in [0, T^*), \quad \forall i \in \{1, \dots, N\}, \quad |z_i(t)| \le C\sqrt{I(t)} \le C \exp\left(C \int_0^t V(t') \mathrm{d}t'\right). \tag{7.3}
$$

We then have that

$$
\frac{d}{dt}H(t) = C_s \sum_{i=1}^{N} v(t, z_i(t)) \cdot \sum_{j \neq i} a_j \frac{(z_i(t) - z_j(t))^{1}}{|z_i(t) - z_j(t)|^{4-2s}} \n\le CV(t) \exp\left(C \int_0^t V(t')dt'\right) \sum_{i \neq j} a_i a_j \frac{1}{|z_i(t) - z_j(t)|^{2-2s}} \n\le CV(t) \exp\left(C \int_0^t V(t')dt'\right) H(t)
$$

where we used relation (7.12). Therefore,

$$
\forall t \in [0, T^*), \quad H(t) \le C \exp\left(C \int_0^t V(\tau) \exp\left(C \int_0^{\tau} V(t') dt'\right) d\tau\right).
$$

We now observe that

$$
|V(t)| \le \frac{C}{\min_{1 \le i \le N} (R_i(t))^{3-2s}}
$$

since, as in relation (4.11) and (4.9) , we have that

$$
v(t, z_i(t)) = K_{s, R_i(t)} \star \theta(t, \cdot)(z_i(t))
$$

=
$$
\int_{|x-z_i(t)| \ge R_i(t)} K_s(x-z_i(t))\theta(t, x) dx
$$

$$
\le ||\theta(t)||_{L^1} ||K_s||_{L^{\infty}(\mathbb{R}^2 \setminus B(z_i(t), R_i(t)))}
$$

$$
\le \frac{C_s}{(R_i(t))^{3-2s}}.
$$

If a collision of point-vortices occurs at time T^* , then $H(t) \longrightarrow_{t \to T^*} +\infty$. Gathering the previous relations, $\min_{1 \leq i \leq N} R_i(t) \underset{t \to T^*}{\longrightarrow} 0.$ \Box

Acknowledgments.

Martin Donati's work is supported by the Simons fundation project on Wave Turbulence. Martin Donati wishes to thank the Institut Fourier and Université Grenoble Alpes, where he is invited to conduct his research.

References

- [1] W. Ao, J. Dávila, M. Del Pino, M. Musso, and J. Wei. Travelling and Rotating Solutions to the Generalized Inviscid Surface Quasi-Geostrophic Equation. Trans. Am. Math. Soc., 374(9):6665–6689, 2021.
- [2] V.I. Arnold. Mathematical Methods of Classical Mechanics. Springer-Verlag, New-York, 1989.
- [3] H. Bahouri, J.-Y. Chemin, and R. Danchin. Fourier Analysis and Nonlinear Partial Differential Equations. Springer Berlin, Heidelberg, 2011.
- [4] Bjorland. The vortex-wave equation with a single vortex as the limit of the Euler equation. Comm. Math. Phys., 305:131–151, 2011.
- [5] T. Buckmaster, V. Shkoller, and V. Vicol. Nonuniqueness of weak solutions to the SQG equation. Comm. Pure Appl. Math., 72(9):1809–1874, 2019.
- [6] D. Cao, G. Qin, W. Zhan, and C. Zou. On the global classical solutions for the generalized SQG equation. Journal of Functional Analysis, 283(2):109503, 2022.
- [7] D. Cao and G. Wang. Existence of steady multiple vortex patches to the vortex-wave system. Pac. Jour. Math., 308(2):257–279, 2020.
- [8] Chae, D. and Constantin, P. and Córdoba, D. and Gancedo, F. and Wu, J. Generalized surface quasigeostrophic equations with singular velocities. Comm. Pure App. Math., 65:1037–1066, 2012.
- [9] P. Constantin, A.J. Majda, and E. Tabak. Formation of strong fronts in the 2-D quasi-geostrophic thermal active scalar. Nonlinearity, 7(6):1495–1533, 1994.
- [10] G. Crippa, M. Lopes Filho, E. Miot, and H. Nussenzveig Lopes. Flows of vector fields with point singularities and the vortex-wave system. Discrete Cont. Dyn. Syst., 36(5):2405–2417, 2016.
- [11] D. Córdoba and L. Martínez-Zoroa. Non-existence and strong ill-posedness in $C^{k,\beta}$ for the generalized Surface Quasi-geostrophic equation. Advances in Mathematics, 407:108570, 2022.
- [12] M. Donati. Two-dimensional point vortex dynamics in bounded domains: Global existence for almost every initial data. SIAM Journal on Mathematical Analysis, 54(1):79–113, 2022.
- [13] C. García. Vortex Patches Choreography for Active Scalar Equations. Jour. Nonlinear Sci., 31(75), 2021.
- [14] C. García and Gómez-Serrano J. Self-similar spirals for the generalized surface quasi-geostrophic equations. preprint, 2022. arXiv:2207.12363.
- [15] C. Geldhauser and M. Romito. Point vortices for inviscid generalized surface quasi-geostrophic models. Am. Ins. Math. Sci., 25(7):2583–2606, 2020.
- [16] O. Glass, C. Lacave, and F. Sueur. On the motion of a small body immersed in a two-dimensional incompressible perfect fluid. Bull. Soc. Math., 142(3):489–536, 2014.
- [17] O. Glass, C. Lacave, and F. Sueur. On the motion of a small light body immersed in a two dimensional incompressible perfect fluid with vorticity. Comm. Math. Phys., 341(3):1015–1065, 2016.
- [18] L. Godard-Cadillac. Vortex collapses for the euler and quasi-geostrophic models. Discrete Cont. Dyn. Syst., 42(7):3143–3168, 2022.
- [19] L. Godard-Cadillac, P. Gravejat, and D. Smets. Co-rotating vortices with n fold symmetry for the inviscid surface quasi-geostrophic equation. Indiana Univ. Math. Journ., 2020. arXiv:2010.08194.
- [20] P. Gravejat and D. Smets. Smooth travelling-wave solutions to the inviscid surface quasi-geostrophic equation. Int. Math. Res. Not., IMRN 2019(6):1744–1757, 2019.
- [21] Z. Hassainia and M.H. Wheeler. Multipole vortex patch equilibria for active scalar equations. SIAM Journal on Mathematical Analysis, 54(6):6054–6095, 2022.
- [22] T. Hmidi and J. Mateu. Existence of corotating and counter-rotating vortex pairs for active scalar equations. Comm. Math. Phys., 350:699–747, 2017.
- [23] W. Hu, I. Kukavika, and M. Ziane. Sur l'existence locale pour une équation de scalaires actifs. Comptes Rendus Math., 353(3):241–245, 2015.
- [24] C. Lacave and E. Miot. Uniqueness for the vortex-wave system when the vorticity is constant near the point vortex. SIAM J. Math. Anal., 41(3):1138–1163, 2009.
- [25] C. Lacave and E. Miot. The vortex-wave system with gyroscopic effects. Ann. Sc. Norm. Super. Pisa Cl. Sci., XXII(5):1–30, 2021.
- [26] A. Kumar M. S. Jolly and V. R. Martinez. On local well-posedness of logarithmic inviscid regularizations of generalized SQG equations in borderline sobolev spaces. Commun. Pure Appl. Anal., 21(1):101–120, 2022.
- [27] F. Marchand. Existence and regularity of weak solutions to the quasi-geostrophic equations in the spaces L^p or $\dot{H}^{-1/2}$. Comm. Math. Phys., 277(1):45–67, 2008.
- [28] C. Marchioro and M. Pulvirenti. On the vortex-wave system. Mechanics, analysis and geometry: 200 years after Lagrange. Elsevier Science, 1991.
- [29] C. Marchioro and M. Pulvirenti. Mathematical theory of incompressible nonviscous fluids, volume 96 of Applied mathematical sciences. Springer-Verlag, 1993.
- [30] Toan T. Nguyen and Trinh T. Nguyen. The inviscid limit of Navier–Stokes equations for vortex-wave data on R 2 . SIAM Jour. Math. Anal., 51(3):2575–2598, 2019.
- [31] J. Pedlosky. Geophysical Fluid Dynamics. Springer-Verlag, New-York, 1987.
- [32] R.K. Scott and D.G. Dritschel. Numerical simulation of a self-similar cascade of filament instabilities in the Surface Quasi-Geostrophic System. Phys. Rev. Lett., 112:144505, 2014.
- [33] V. N. Starovoitov. Uniqueness of a solution to the problem of evolution of a point vortex. Siberian Mathematical Journal, 35(3), 1994.
- [34] G.K. Vallis. Atmospheric and Oceanic Fluid Dynamics. Cambridge University Press, 2006.
- [35] K. Varholm, E. Wahlén, and S. Walsh. On the stability of solitary water waves with a point vortex. Comm. Pure App. Math., 73(12), 2020.
- [36] V.I. Youdovich. Non-stationary flow of an ideal incompressible liquid. Zh. Vychisl. Mat. Fiz., 3(6):1032–1066, 1963.
- [37] V.I. Youdovich. Uniqueness theorem for the basic nonstationary problem in the dynamics of an ideal incompressible fluid. Math. Res. Lett., 2(1):27–38, 1995.