



HAL
open science

Anticoagulation in COVID-19: not strong for too long?

Charles Tacquard, Alexandre Mansour, Alexandre Godon, Yves Gruel, Sophie Susen, Anne Godier, Pierre Albaladejo

► **To cite this version:**

Charles Tacquard, Alexandre Mansour, Alexandre Godon, Yves Gruel, Sophie Susen, et al.. Anticoagulation in COVID-19: not strong for too long?. *Anaesthesia Critical Care & Pain Medicine*, 2021, 40 (2), pp.100857. 10.1016/j.accpm.2021.100857 . hal-04370740

HAL Id: hal-04370740

<https://hal.science/hal-04370740>

Submitted on 3 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Anticoagulation in COVID-19: not strong for too long?

Charles Tacquard,^{a,*} Alexandre Mansour,^b Alexandre Godon,^c Yves Gruel,^d Sophie Susen,^e Anne Godier,^f and Pierre Albaladejo^c

^aDepartment of Anaesthesia and Intensive Care, Hôpitaux Universitaires de Strasbourg, 1, Place de l'Hôpital, 67091 Strasbourg Cedex, France

^bDepartment of Anaesthesiology Critical Care Medicine and Perioperative Medicine, CHU de Rennes, Rennes, France

^cDepartment of Anaesthesiology and Critical Care, Grenoble Alpes University Hospital, Grenoble, France

^dDepartment of Haematology-Haemostasis, Tours University Hospital, France

^eHeart and Lung Institute, Haemostasis Department, CHU Lille, 59037 Lille Cedex, France

^fDepartment of Anaesthesiology and Critical Care, European Georges Pompidou Hospital, Assistance Publique-Hôpitaux de Paris, Paris University, Paris, France

□ Corresponding author.

Several randomised controlled trials are currently underway to evaluate the value of increasing the dose of prophylactic anticoagulation in critically ill COVID-19 patients, either at an intermediate or therapeutic dose, in order to decrease thrombotic complications, organ failure and possibly reduce mortality in these patients.

INSPIRATION investigators recently published the results of the first large-scale randomised controlled trial evaluating the benefit of intermediate dose prophylactic anticoagulation in critically ill COVID-19 patients [1]. Their results showed that intermediate-dose prophylactic anticoagulation did not significantly reduce their composite endpoints (30-day mortality, need for extracorporeal membrane oxygenation (ECMO), thrombotic events) and therefore do not support the routine use of an increased dose of anticoagulation in unselected critically ill COVID-19 patients.

However, there are several critical points that need to be highlighted in this trial as they could limit the potential impact of these results on our daily clinical practice.

First, composite end points must be taken with caution because they combine different elements that are not necessarily all related to the same end goal. The inclusion of the need for ECMO in the primary endpoint of this study is questionable, because the link between anticoagulation and the need for ECMO is not clearly established. A similar comment can be made about the multiplatform randomised trial (ATTAC, REMAP-CAP, and ACTIV-4) on therapeutic anticoagulation. The choice of 21-day organ support free days as the primary end point is questionable because organ failure in COVID-19 is multifactorial and the direct effect of anticoagulation on this end point may be diluted, which could lead to a false conclusion of futility.

Only about 20% of the included patients underwent invasive ventilation, and the length of stay in the ICU was short, with a median of 6 [2–11] days, which is very different from the data observed in other ICU cohorts [2]. In addition, the mortality rate was strikingly high (42% of the study population), raising the question of a possible mismatch between needs and availability of ICU resources.

The rate of thrombotic events in the standard-dose prophylactic anticoagulation group was very low, at 3.5%, which differed greatly from previously published studies on critically ill patients [3], [4], [5] (*e.g.*, 27.9% (95% CI 22.1–34.1) in the recent meta-analysis by Jimenez et al. [6]), one may wonder if all thrombotic events in the INSPIRATION trial were diagnosed. This low rate of thrombotic events could have led to an underestimation of the benefit of intermediate-dose prophylactic anticoagulation.

The final concern regarding the results of the INSPIRATION trial results is the relatively high rate of bleeding in the intermediate dose group (6.2% of bleeding events) where patients were exposed to the assigned treatment for 20 [7–30] days. Several studies have reported a significant risk of bleeding in patients with COVID-19, particularly in the most severe patients and in those on enhanced-dose anticoagulation [6], [7], [8], [9]. The longitudinal evaluation of haemostatic biomarkers showed that the procoagulant status of COVID-19 patients seems to be particularly marked during the first few days of hospitalisation and then gradually decreases to normal levels [10]. This raises the question of the temporal relationship between thrombosis and the bleeding risk, and whether these risks are concomitant or staggered in time.

We performed a literature review of all studies on the thrombotic and bleeding risk associated with COVID-19. On the 5th of March 2021, we searched MEDLINE and looked for studies on COVID-19 written in English that reported a median (or mean) duration and interquartile range – IQR - (or standard deviation - SD) between hospital or ICU admission and thrombotic or bleeding events. Case series with the time between admission and the event available were also included.

Twenty-two studies were identified, 13 of which specifically focusing on critically ill patients [3], [7], [8], [9], [11], [12], [13], [14], [15], [16], [17], [18], [19], [20], [21], [22], [23], [24], [25], [26], [27], [28]. Thirteen studies reported only the time from admission to thrombotic events [3], [11], [12], [13], [14], [15], [16], [18], [19], [20], [21], [24], four studies reported only the time from admission to bleeding events [8], [25], [26], [27] and five studies reported both [7], [9], [22], [23], [28]. One study analysed two different waves of patients [15]. All but two studies reported this delay as the median and interquartile

range. Fig. 1 shows distribution of the median (or mean) and the IQR (or SD) of time from admission to event in each study. Pooled medians were estimated using a quantile estimation method. The results of the meta-analysis showed that thrombotic events occurred 7.0 [5.9–8.2] days after admission (713 events) while bleeding events occurred 11.4 [8.6–14.1] days after admission (163 events).




Fig. 1 (A) Distribution of medians and interquartile ranges for duration between admission (hospital/ICU) and events (thrombosis in blue, bleeding in red) based on selected studies. For each study, details on the type and location of recorded events and the number of events/number of patients included are shown in the right-hand column. (B) Results of the meta-analysis using a quantile estimation method. Data are expressed as median and interquartile range. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

*For these two studies, the data are expressed as a mean (standard deviation). These studies were excluded from further analysis in B.

Our results confirm that thrombotic and bleeding events are staggered in time, with thrombotic events occurring primarily in the first ten days after admission. Although some events were reported within the first few days of hospitalisation, bleeding events occurred most often late. These findings are consistent with previously published biological data [10]. Hardy et al. observed an increase in thrombin generation associated with a decrease in overall fibrinolytic capacity during the first week of hospitalisation, resulting in a strong procoagulant state. After this early stage of the disease, inflammatory markers progressively decreased, as did D-dimer levels in survivors, probably in relation to a decrease in the intensity of processes leading to microthrombosis.

Data on bleeding events are difficult to interpret correctly: (i) some bleeding is spontaneous while others are related to invasive techniques (central venous or arterial catheters, renal replacement therapy, ECMO) ; (ii) some bleeding is directly related to anticoagulation (*e.g.*, overdose) while others would have occurred independently of anticoagulation status; (iii) some patients require anticoagulation anyway for identified medical reasons (identified thrombosis, atrial fibrillation, extracorporeal circuits, etc.). Thus, data on bleeding events should be analysed not only by considering all bleeding events but also by taking into account the imputability between anticoagulation exposure and bleeding events.

These results raise the question of the optimal duration of anticoagulation (therapeutic or intermediate dose) in COVID-19 patients. Increased-dose prophylactic anticoagulation, if indicated, should be initiated very early in the course of the disease, when the thrombotic risk is high. Thereafter, in patients without a clear indication for curative anticoagulation, the dose of anticoagulation should probably be gradually reduced to a standard prophylactic dose to limit the risk of bleeding, and possibly monitored by inflammatory or coagulation markers.

References

1. Investigators I., Sadeghipour P., Talasaz A.H., Rashidi F., Sharif-Kashani B., Beigmohammadi M.T. Effect of intermediate-dose vs standard-dose prophylactic anticoagulation on thrombotic events, extracorporeal membrane oxygenation treatment, or mortality among patients with COVID-19 admitted to the intensive care unit: the inspiration randomized clinical trial. *JAMA*. 2021;18 doi: 10.1001/jama.2021.4152.
2. Network C-IGobotR, the C-ICUI Clinical characteristics and day-90 outcomes of 4244 critically ill adults with COVID-19: a prospective cohort study. *Intensive Care Med*. 2021;47(1):60–73.
3. Middeldorp S., Coppens M., van Haaps T.F., Foppen M., Vlaar A.P., Muller M.C.A. Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost*. 2020;18(8):1995–2002.
4. Suh Y.J., Hong H., Ohana M., Bompard F., Revel M.P., Valle C. Pulmonary embolism and deep vein thrombosis in COVID-19: a systematic review and meta-analysis. *Radiology*. 2020]
5. Tang N., Bai H., Chen X., Gong J., Li D., Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost*. 2020;18(5):1094–1099
6. Jimenez D., Garcia-Sanchez A., Rali P., Muriel A., Bikedli B., Ruiz-Artacho P. Incidence of VTE and bleeding among hospitalized patients with coronavirus disease 2019: a systematic review and meta-analysis. *Chest*. 2021;159(3):1182–1196. doi: 10.1016/j.chest.2020.11.005
7. Jonmarker S., Hollenberg J., Dahlberg M., Stackelberg O., Litorell J., Everhov A.H. Dosing of thromboprophylaxis and mortality in critically ill COVID-19 patients. *Crit Care*. 2020;24(1):653
8. Shah A., Donovan K., McHugh A., Pandey M., Aaron L., Bradbury C.A. Thrombotic and haemorrhagic complications in critically ill patients with COVID-19: a multicentre observational study. *Crit Care*. 2020;24(1):561
9. Godier A., Clausse D., Meslin M., Bazine M., Lang E., Huche F. Major bleeding complications in critically ill patients with COVID-19 pneumonia. *J Thromb Thrombolysis*. 2021:1–4. doi: 10.1007/s11239-021-02403-9.
10. Hardy M., Michaux I., Lessire S., Douxfils J., Dogne J.M., Bareille M. Prothrombotic disturbances of hemostasis of patients with severe COVID-19: a prospective longitudinal observational study. *Thromb Res*. 2021;197:20–23

11. Ameri P., Inciardi R.M., Di Pasquale M., Agostini P., Bellasi A., Camporotondo R. Pulmonary embolism in patients with COVID-19: characteristics and outcomes in the Cardio-COVID Italy multicenter study. *Clin Res Cardiol.* 2020;3:1–9.
12. Chen S., Zhang D., Zheng T., Yu Y., Jiang J. DVT incidence and risk factors in critically ill patients with COVID-19. *J Thromb Thrombolysis.* 2021;51(1):33–39.
13. Dalager-Pedersen M., Lund L.C., Mariager T., Winther R., Hellfritzsch M., Larsen T.B. Venous thromboembolism and major bleeding in patients with COVID-19: a nationwide population-based cohort study. *Clin Infect Dis.* 2021 ciab003.
14. Dubois-Silva A., Barbagelata-Lopez C., Mena A., Pineiro-Parga P., Llinares-Garcia D., Freire-Castro S. Pulmonary embolism and screening for concomitant proximal deep vein thrombosis in noncritically ill hospitalized patients with coronavirus disease 2019. *Intern Emerg Med.* 2020;15(5):865–870.
15. Dutch C., Thrombosis C., Kaptein F.H.J., Stals M.A.M., Grootenboers M., Braken S.J.E. Incidence of thrombotic complications and overall survival in hospitalized patients with COVID-19 in the second and first wave. *Thromb Res.* 2020;199:143–148.
16. Fournier M., Faille D., Dossier A., Mageau A., Nicaise Roland P., Ajzenberg N. Arterial thrombotic events in adult inpatients with COVID-19. *Mayo Clin Proc.* 2021;96(2):295–303
17. Fraisse M., Logre E., Pajot O., Mentec H., Plantefeve G., Contou D. Thrombotic and hemorrhagic events in critically ill COVID-19 patients: a French monocenter retrospective study. *Crit Care.* 2020;24(1):275.
18. Helms J., Severac F., Merdji H., Schenck M., Clere-Jehl R., Baldacini M. Higher anticoagulation targets and risk of thrombotic events in severe COVID-19 patients: bi-center cohort study. *Ann Intensive Care.* 2021;11(1):14. [
19. Helms J., Tacquard C., Severac F., Leonard-Lorant I., Ohana M., Delabranche X. High risk of thrombosis in patients in severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med.* 2020;4:1–10.
20. Moll M., Zon R.L., Sylvester K.W., Chen E.C., Cheng V., Connell N.T. VTE in ICU patients with COVID-19. *Chest.* 2020;158(5):2130–2135.
21. Poissy J., Goutay J., Caplan M., Parmentier E., Duburcq T., Lassalle F. Pulmonary embolism in patients with COVID-19: awareness of an increased prevalence. *Circulation.* 2020;142(2):184–186.
22. Salisbury R., Iotchkova V., Jaafar S., Morton J., Sangha G., Shah A. Incidence of symptomatic, image-confirmed venous thromboembolism following hospitalization for COVID-19 with 90-day follow-up. *Blood Adv.* 2020;4(24):6230–6239.

23. Tacquard C., Mansour A., Godon A., Godet J., Poissy J., Garrigue D. Impact of high dose prophylactic anticoagulation in critically ill patients with COVID-19 pneumonia. *Chest*. 2021 doi: 10.1016/j.chest.2021.01.017.
24. Creel-Bulos C., Liu M., Auld S.C., Gaddh M., Kempton C.L., Sharifpour M. Trends and diagnostic value of D-dimer levels in patients hospitalized with coronavirus disease 2019. *Medicine (Baltimore)* 2020;99(46)
25. Lang C.N., Dettinger J.S., Berchtold-Herz M., Utzolino S., Bemtgen X., Zotzmann V. Intracerebral hemorrhage in COVID-19 patients with pulmonary failure: a propensity score-matched registry study. *Neurocrit Care*. 2021
26. Vergori A., Pianura E., Lorenzini P., D'Abramo A., Di Stefano F., Grisetti S. Spontaneous ilio-psoas haematomas (IPHs): a warning for COVID-19 inpatients. *Ann Med*. 2021;53(1):295–301.
27. Voicu S., Chousterman B.G., Bonnin P., Deye N., Malissin I, Le Gall A. Increased anticoagulation reduces proximal deep vein thrombosis in mechanically ventilated COVID-19 patients: venous thrombosis prevention & COVID-19. *J Infect*. 2020
28. Atallah B., Sadik Z.G., Salem N., El Nekidy W.S., Almahmeed W., Park W.M. The impact of protocol-based high-intensity pharmacological thromboprophylaxis on thrombotic events in critically ill COVID-19 patients. *Anaesthesia*. 2021;76(3):327–335.