



HAL
open science

Description of a new species of *Nyssodrysis* Gilmour, 1962 (Cerambycidae: Lamiinae: Acanthocinini) from Brazil and new record, host plant, and parasitoid of *Heterachthes sablensis* Blatchley, 1920 (Cerambycinae: Tropidini: Compsina) in Panama.

Alfredo Lanuza-Garay, Marleny Rivera, Alonso Santos-Murgas

► **To cite this version:**

Alfredo Lanuza-Garay, Marleny Rivera, Alonso Santos-Murgas. Description of a new species of *Nyssodrysis* Gilmour, 1962 (Cerambycidae: Lamiinae: Acanthocinini) from Brazil and new record, host plant, and parasitoid of *Heterachthes sablensis* Blatchley, 1920 (Cerambycinae: Tropidini: Compsina) in Panama.. *Faunitaxys*, 2024, 12 (1), pp.1-4. hal-04371344

HAL Id: hal-04371344

<https://hal.science/hal-04371344>

Submitted on 3 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*



Volume 12
Numéro 01

Janvier 2024

ISSN (Print) : 2269 - 6016
ISSN (Online) : 2970 - 4960
Dépôt légal : Janvier 2024

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*

ZooBank : <https://zoobank.org/79A36B2E-F645-4F9A-AE2B-ED32CE6771CC>

Directeur de la publication, rédacteur, conception graphique et PAO:

Lionel Delaunay

Cette revue ne peut pas être vendue

Elle est distribuée par échange aux institutions (version papier)

et sur simple demande aux particuliers (format PDF)

à l'adresse suivante:

AFCFF (Association française de Cartographie de la Faune et de la Flore)

28, rue Voltaire, F- 42100 Saint Etienne

E-mail: faunitaxys.journal@gmail.com

Elle est disponible librement au téléchargement à partir du site:

<https://faunitaxys.fr/>

La parution de *Faunitaxys* est apériodique

Faunitaxys est indexé dans / *Faunitaxys* is indexed in:

- **Zoological Record**

Articles and nomenclatural novelties are referenced by:

- **ZooBank** (<https://zoobank.org>)

Online Archives:

- **HAL** (<https://hal.archives-ouvertes.fr>)

- **Internet Archive** (<https://archive.org>)

Imprimée sur les presses de SPEED COPIE, 6, rue Tréfilerie, F- 42100 Saint-Etienne

Imprimé le 02 01 2024

Description of a new species of *Nyssodrysis* Gilmour, 1962 (Cerambycidae: Lamiinae: Acanthocinini) from Brazil and new record, host plant, and parasitoid of *Heterachthes sablensis* Blatchley, 1920 (Cerambycinae: Tropicini: Compsina) in Panama

ALFREDO LANUZA-GARAY (1, 3, *), MARLENY RIVERA (4) & ALONSO SANTOS MURGAS (2, 3)

(1) Universidad de Panamá, Centro Regional Universitario de Colón, Escuela de Biología, Departamento de Zoología, Cristobal County, Randolph Avenue, Rainbow City, Colon City, PANAMA.

- E-mail: alfredo.lanusa@up.ac.pa - Orcid: <https://orcid.org/0000-0003-0480-5490> - ZooBank: <http://zoobank.org/10339FE9-0467-4C73-A9A1-6C86260811BE>

(2) Universidad de Panamá, Facultad de Ciencias Naturales, Exactas y Tecnología. Panamá, PANAMA

- E-mail: santosturgasa@gmail.com - Orcid: <https://orcid.org/0000-0001-9339-486X> - ZooBank: <https://zoobank.org/F5934143-484A-4E0C-BCA4-5ECDB53698EE>

(3) Museo de Invertebrados Graham Bell Fairchild, Universidad de Panamá, Manuel Espinosa Batista Avenue, Panama City, PANAMA.

(4) Smithsonian Tropical Research Institute, Ancon, Panama City, PANAMA.

- E-mail: riverame@si.edu - Orcid: <https://orcid.org/0000-0003-1231-4579> - ZooBank: <https://zoobank.org/F8A68827-D292-435E-AF23-A9B383661A4B>

* Corresponding author

Keywords:

Cerambycidae; description;
Lamiinae; new species;
Acanthocinini; taxonomy;
Longhorned beetles; Neotropic;
Nyssodrysis; Brazil.
esperanzae;
Heterachthes;
sablensis;
Biological interactions;

Abstract. – *Nyssodrysis esperanzae* sp. nov. (Lamiinae: Acanthocinini), a new species of long-horned beetle, is described from Brazil. Additionally, an association of braconid wasp *Cenocoelius* sp. (Braconidae: Cenocoeliinae) parasiting *Heterachthes sablensis* Blatchley, 1920 and a new host plant is reported.

Lanuzza-Garay A., Rivera M. & Santos Murgas A., 2024. – Description of a new species of *Nyssodrysis* Gilmour, 1962 (Cerambycidae: Lamiinae: Acanthocinini) from Brazil and new record, host plant, and parasitoid of *Heterachthes sablensis* Blatchley, 1920 (Cerambycinae: Tropicini: Compsina) in Panama. *Faunitaxys*, 12(01): 1 – 4.

DOI: [https://doi.org/10.57800/faunitaxys-12\(01\)](https://doi.org/10.57800/faunitaxys-12(01))

ZooBank: <https://zoobank.org/1870763D-4021-45AF-812C-F510203C58B6>

Received: 20/12/2023 – Revised: 26/12/2023 – Accepted: 27/12/2023

Introduction

Nyssodrysis Gilmour, 1962 is a small genus of American Acanthocinini including four species: *N. humeralis* Bezark & Santos Silva, 2023; *N. irrorata* (Melzer, 1927); *N. lineata* Gilmour, 1962, and *N. vittata* (Melzer, 1934). These species are distributed from northern to southern South America (Bezark & Santos Silva, 2023; Monné, 2023). Herein, we described a new species from Brazil, stored in the Smithsonian Tropical Research Institute Insect Collection in Panama. Additionally, the study of some specimens identified as *Heterachthes sablensis* Blatchley, 1920 (Cerambycinae, Tropicini) collected by the third author's research team in a mangrove forest around the community of El Libano, Chame, Panama, allowed us to report a new host plant and a parasitoid for first time.

Material and methods

Photographs of *Nyssodrysis esperanzae* sp. nov. were taken at the Insect Collection of Smithsonian Tropical Research Institute by the second author, with a Visionary camera system, equipped with a Canon

EOS6D camera, equipped with Canon 65mm f/4 Max aperture: 3 horizontal/vertical resolutions: 300 dpi and edited by Adobe Photoshop photo editor software. New species confirmation was made through the revision of original descriptions and identification keys (Gilmour, 1962, Bezark & Santos-Silva, 2023), photographs (Bezark, 2023), and comparison with specimens deposited in collections.

Identification of *H. sablensis* was made through original description (Blatchley 1920), photographs (Bezark, 2023) and identification key from Santos-Silva & Botero (2017). Its parasitoid was identified by key from Achterberg (1993, 1997). Classification and distributional data are based on Monné (2023). Both, beetle and parasitoid are deposited at the Museo de Invertebrados G. B. Fairchild de la Universidad de Panamá (MIUP). Photographs were taken in the MIUP with a Sony a6000 camera, equipped with AMSCOP 4x close-up lens and Nikkor Z MC 105mm f/2.8 VR S aperture for angular frames and edited by the “tracking” technique using Photoshop software.

Measurements (mm) were taken using Leica M165 C stereomicroscope with measuring ocular, also used in the study of the specimens. The new species are described by the first and second author, while identification of *H. sablensis* and its parasitoid are made only by the first and third author.

The collection acronyms used in the text are as follows:

STRI: Smithsonian Tropical Research Institute Insect Collection, PANAMA.

Reviewer :

Antonio Santos-Silva (Museu de Zoologia, Universidade de São Paulo, Brazil) - Orcid: <https://orcid.org/0000-0001-7128-1418>

- ZooBank: <http://zoobank.org/8E7804F2-4315-4D23-9FA5-2E5D13841364>



This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Copyright 2024. The Authors. *Faunitaxys* published by Lionel Delaunay on behalf of the AFCFF (Association française de Cartographie de la Faune et de la Flore).

Results

Acanthocinini Blanchard, 1845

Nyssodryssilla Gilmour, 1962

Nyssodryssilla esperanzae sp. nov.

(Fig. 1-5)

ZooBank: <https://zoobank.org/950006BE-DA48-4A46-8508-437BC8B2AF41>

Holotype, ♀, BRAZIL, Mato Grosso: Aripuanã, Humboldt, 18.iii.1977, H.P. Stockwell leg. (STRI_ENT_0063-018).

Description of the holotype

Coloration. – Integument mostly dark brown; ventral mouthparts dark brown, except yellowish apex of maxillary palpomere IV and apex of labial palpomere III; yellowish-brown clypeus; labrum brown on posterior 2/3, yellowish brown on anterior third. Scape dark brown, remaining antennomeres reddish brown. Pronotum with a longitudinal orangish-yellow band on each side. Ventral surface of body reddish brown, except reddish-brown coxae.

Head. – Frons densely, finely punctate; with abundant yellowish-white pubescence not obscuring integument. Vertex and area behind upper eye lobes with abundant yellowish-white pubescence almost obscuring integument, except glabrous area close to prothorax and small central area between antennal tubercles and upper eye lobes. Area behind lower eye lobes with abundant yellowish-white pubescence close to eye, pubescence denser from superior third, glabrous close to prothorax. Genae with somewhat abundant yellowish-white pubescence not obscuring integument. Antennal tubercles with pubescence as on frons frontally, pubescence slightly sparser posteriorly. Clypeus covered with abundant yellowish-white pubescence partially obscuring integument. Labrum with yellowish-white pubescence not obscuring integument on posterior 2/3, with long, erect yellowish setae interspersed; anterior third glabrous, except fringe of yellowish-white setae on anterior margin. Gulamentum glabrous except moderately abundant grayish-white pubescence anteriorly. Distance between upper eye lobes 0.31 times distance between outer margins of eyes; in frontal view, distance between lower eye lobes 0.69 times distance between outer margins of eyes. Antennae 2.4 times elytral length, reaching elytral apex at basal quarter of antennomere VI. Scape with sparse grayish-white pubescence not obscuring integument; with a few long, erect blackish setae near apex of ventral surface. Pedicel with grayish-white pubescence not obscuring integument; ventral surface with a few long, erect, thick blackish setae. Antennomeres III–XI with sparse grayish-white pubescence not obscuring integument; with short, erect, thick black setae interspersed ventrally on III–VI, short, thick black setae interspersed dorsally on V–IX, setae distinctly sparser on IX.

Antennal formula based on length of antennomere III:

– Scape = 0.78. – Pedicel = 0.07. – IV = 0.84. – V = 0.63. – VI = 0.68. – VII = 0.68. – VIII = 0.71. – IX = 0.68. – X = 0.66. – XI = 0.67.

Thorax. – Prothorax 0.51 times wider than long. Well-marked anterior constriction; sides gradually divergent from anterolateral angles to lateral tubercles, then convergent toward posterolateral angles; lateral tubercles located after the base of the posterior quarter, slightly acute. Pronotum densely, finely punctate; with transverse row of coarse punctures posteriorly; with dense, longitudinal orangish pubescence obscuring the integument on each side; central area with sparse yellowish pubescence not obscuring the integument. Prosternum covered with yellowish-white pubescence not obscuring integument; prosternal process narrowed centrally, 0.12 times procoxal width; mesanepisternum and mesepimeron with dense yellowish-white pubescence, partially obscuring integument; mesoventral process not laminiform, 0.51 times mesocoxal width, with sparse yellowish-white pubescence; metanepisternum with dense yellowish-white pubescence partially obscuring integument; metaventrite with sparse yellowish-white pubescence. Metathoracic discrimen complete. Scutellum covered with yellowish-white pubescence not obscuring integument.

Elytra. – Sides slightly widened centrally; apex obliquely truncate with outer angle slightly projected; somewhat abundantly, coarsely punctate on anterior quarter, punctures gradually finer and sparser toward apex on remaining surface; with abundant yellowish-white pubescence partially obscuring integument.

Legs. – Covered with yellowish-white pubescence not obscuring integument; meso- and metatibiae with short, erect, thick black setae interspersed. Metatarsomere I longer than II–III together.

Abdomen. – Ventrites with sparse pale-yellow pubescence. Ventrite 5 about as long as 3–4 together; sides strongly convergent on anterior 2/3, parallel-sided on posterior third; apex widely emarginate; with long dark-brown setae directed backward on apex. Last tergite rounded apically and slightly surpassing apex of ventrite 5.

Dimensions (mm) (Holotype)

- Total length, 4.90;
- Prothoracic length, 0.85;
- Anterior prothoracic width, 1.00;
- Posterior prothoracic width, 1.15;
- Maximum prothoracic width, 1.40;
- Humeral width, 1.55;
- Elytral length, 3.30.

Nyssodryssilla esperanzae sp. nov. can be included in the Key from Bezark & Santos Silva (2023) (modified) as follows:

1. Elytra marbled with small pale-yellow maculae, not vittate. Brazil (Bahia, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Paraguay *N. irrorata* (Melzer, 1927) 2
— Elytra not marbled, with or without longitudinal pubescent bands 2
- 2(1). Elytra lacking longitudinal pubescent bands; with subtriangular yellow pubescent macula on sides of anterior third. Paraguay *N. humeralis* Bezark and Santos-Silva, 2023 3
— Elytra with or without longitudinal pubescent bands; without subtriangular yellow pubescent macula on sides of anterior third 3
- 3(2). Pronotum with longitudinal orangish pubescent band on each side, without pubescent bands centrally. Brazil (Mato Grosso) *N. esperanzae* sp. nov. 4
— Pronotum more than two longitudinal pale-yellow pubescent bands 4
- 4(3). Elytra with abundant, longitudinal pale-yellow pubescent bands, without large, subelliptical brownish pubescent macula on sides of anterior half. Brazil (Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul) *N. vittata* (Melzer, 1934) 4
— Elytra with a few pale-yellow pubescent bands and with subelliptical brown pubescent macula on sides of anterior half. Peru, French Guiana, Bolivia (Pando) *N. lineata* Gilmour, 1962 4




Fig. 1-7.

1-4. *Nyssodryssilla esperanzae* sp. nov., ♀ holotype. 1. Dorsal habitus. 2. Ventral habitus showing prosternal and mesoventral process. 3. Lateral habitus. 4. Head, frontal view. 5. *Heterachthes sablensis* Blatchey, 1920, dorsal habitus. 6-7. *Cenocoelius* sp. 6. Dorsal habitus. 7. Lateral habitus.

Etymology. – Named in honor of the grandmother of the second author, Ruby Esperanza Centeno Muñoz.

Remarks. – *Nyssodryssilla esperanzae* sp. nov. is similar to *N. humeralis* (Bezark & Santos Silva, 2023). However, the main differences between them are the longitudinal orangish pubescent bands on sides of the pronotum (absent in *N. humeralis*), the absence of subtriangular pubescent maculae in *N. esperanzae* (present in *N. humeralis*), and abundance and distribution of integumental pubescence.

New record

Cerambycinae Latreille, 1802

Tropidini Martins & Galileo, 2007

Compsina Martins & Galileo, 2007

Heterachthes Newman, 1840

Heterachthes sablensis Blatchley, 1920

(Fig. 5)

Previous distribution. – United States of America (Florida), Bahamas (Andros Island), Aruba, Honduras, Guatemala, Belize, Colombia, Ecuador (Galapagos) (Monné *et al.*, 2023).

Examined material

– 2 ♂, PANAMA (new country record), Panama Province, Chame, Comunidad el Libano, 28.xi.2022-9. i.2023, C. Rivera & J. Abrego leg. (MIUP).
– 1 ♀, PANAMA, Panama Province, Chame, Comunidad el Libano, 28.xi.2022-9. i.2023, C. Rivera & J. Abrego leg. (MIUP).

Remarks. – Two male and one female emerged from red mangrove (*Rhizophora mangle* Linnaeus, 1753) fresh baits; three parasitoid wasps of the genus *Cenocoelius* Westwood, 1840 (Braconidae: Cenocoeliinae) emerged from the three specimens of *H. sablensis*. The wasps of this genus have been previously mentioned by Borgmeier (1931) and Kirch (1983) as parasitoids of larvae of *Oncideres dejeani* Thomson 1868 and *Oncideres impluviata* (Germar, 1842). Regarding its host plant, Monné (2023) indicates that *H. sablensis* is associated with *Avicennia germinans* Linnaeus, 1764 and *A. officinalis* Linnaeus, 1753 (Verbenaceae).

Acknowledgements

We thank Jean Carlos Abrego and Carlos Rivera (Universidad de Panamá) for collecting the specimens of the braconid wasp *Cenocoelius*. Also, we would like to thank Annette Aiello (STRI) for approving access to the specimen used to describe the new species in this study and for reviewing the English.

Résumé

Lanuza-Garay A., Rivera M. & Santos Murgas A., 2024. – Description d'une nouvelle espèce du genre *Nyssodryssilla* Gilmour, 1962 (Cerambycidae : Lamiinae : Acanthocinini) du Brésil et nouveau signalement, plante hôte et parasite d'*Heterachthes sablensis* Blatchley, 1920 (Cerambycinae : Tropidini : Compsina) au Panama. *Faunitaxys*, 12(01): 1 – 4.

Nyssodryssilla esperanzae sp. nov. (Lamiinae : Acanthocinini), une nouvelle espèce de longicorne, est décrite au Brésil. De plus, une association de guêpes braconides *Cenocoelius* sp. (Braconidae : Cenocoeliinae) parasitant *Heterachthes sablensis* Blatchley, 1920 et une nouvelle plante hôte sont signalées.

Mots-clés. – Cerambycidae, Lamiinae, Acanthocinini, longicornes, *Nyssodryssilla*, *esperanzae*, *Heterachthes*, *sablensis*, interactions biologiques, description, nouvelle espèce, taxonomie, Région néotropicale, Brésil.

References

- Achterberg C., 1993. – Illustrated key to the subfamilies of the Braconidae (Hymenoptera: Ichneumonoidea). *Zoologische Verhandlungen*, 283: 3-189.
- Achterberg C., 1997. – Subfamily Cenocoeliinae, pp. 185–192. In: Wharton, R.A., P.M., Marsh, and M.J., Sharkey (eds.), Manual of the New World genera of the Family Braconidae (Hymenoptera). Special publication n°1, International Society of Hymenopterists, Washington DC: 439 pp.
- Bezark L.G., 2023. – A photographic catalog of the Cerambycidae of the New World. Available from: <http://bezbycids.com/byciddb/wdefault.asp?w=n> (accessed October 20, 2023).
- Bezark L.G. & Santos-Silva A., 2023. – New species, synonymies and transfers in Neotropical Lamiinae (Coleoptera: Cerambycidae). *Insecta Mundi*, 0993: 1-26.
- Blatchley W.S., 1920. – Notes on the winter Coleoptera of western and southern Florida, with descriptions of new species. *The Canadian Entomologist*, 52(3): 68-72.
- Borgmeier T., 1931. – Uma nova espécie de *Conocoelius* (Hym. Braconidae), parasita de *Oncideres dejeani* Thoms. (Col. Cerambycidae). *Revista de Entomologia*, 1(4): 431-436.
- Gilmour E.F., 1962. – On the Neotropical Acanthocinini (Coleoptera, Cerambycidae, Lamiinae). Some new genera and generic revision. *Beiträge zur Neotropischen Fauna*, 2(4): 249-293.
- Kirch E., 1983. – Estudo dos inimigos naturais de *Oncideres impluviata* (Germar, 1824) em *Mimosa scabrella* Benth. Curitiba, Univ. Fed. do Paraná, 1983. 63 pp. Dissertação (Mestrado em Engenharia Florestal) – Universidade Federal de Santa Maria.
- Monné M.A., 2023. – Catalogue of the Cerambycidae (Coleoptera) of the Neotropical Region. Part I. Subfamily Cerambycinae. Available from: <http://www.cerambyxcat.com/> (accessed 28 November 2023).
- Santos-Silva A. & Botero J.P., 2023. – A new species of *Heterachthes* Newman, 1840 from Colombia (Coleoptera, Cerambycidae). *Zootaxa*, 4231(4): 581-584.

Authors contribution ¹

Publisher Correspondence. – LG

Writing the article. – LG

Description. – LG, RM

Article proofreading. – LG

Bibliographic work. – LG, SM

Material study. – LG, RM, SM

Author of the figures. – LG, RM, SM

¹ LG = Lanuza-Garay – RM = Rivera – SM = Santos Murgas

Derniers articles publiés

- Coache A. & Orbach E., 2023. – Contribution à la connaissance des brentides de l'archipel de Sao Tomé et Principe (Coleoptera, Curculionoidea, Brentidae: Cyphagoginae & Trachelizinae). *Faunitaxys*, 11(63) : 1 – 14. [https://doi.org/10.57800/faunitaxys-11\(63\)](https://doi.org/10.57800/faunitaxys-11(63))
- Qiu J. Y., 2023. – Flower beetles of the Asian genus *Dicronocephalus* Hope (Coleoptera: Scarabaeidae: Cetoniinae): The description of a new species from China and intraspecific variations of *D. adamsi*. *Faunitaxys*, 11(64): 1 – 11. [https://doi.org/10.57800/faunitaxys-11\(64\)](https://doi.org/10.57800/faunitaxys-11(64))
- Ballerio A. & Coache A., 2023. – A new flightless species of *Melanophilharmostes* from São Tomé (Coleoptera Scarabaeoidea Hybosoridae Ceratocanthinae). *Faunitaxys*, 11(65): 1 – 6. [https://doi.org/10.57800/faunitaxys-11\(65\)](https://doi.org/10.57800/faunitaxys-11(65))
- Minkina L., Jákl S. & Bezděk A., 2023. – A new *Rhyparus* Westwood, 1845 (Coleoptera: Scarabaeidae: Aphodiinae) species with modified pygidium and last abdominal ventrite in females. *Faunitaxys*, 11(66): 1 – 5. [https://doi.org/10.57800/faunitaxys-11\(66\)](https://doi.org/10.57800/faunitaxys-11(66))
- Lanuzza-Garay A. & Rivera M., 2023. – Description of a new species of *Sternacutus*, notes on *Pseudosparna iolandae* and *Trichotithonus curvatus*, new records and host plants in Acanthocinini (Coleoptera, Cerambycidae, Lamiinae) from Panama. *Faunitaxys*, 11(67): 1 – 6. [https://doi.org/10.57800/faunitaxys-11\(67\)](https://doi.org/10.57800/faunitaxys-11(67))
- Chen Z.-T., 2023. – *Borisnia* gen. nov., a new genus of Capniidae (Insecta: Plecoptera) from China. *Faunitaxys*, 11(68): 1 – 5. [https://doi.org/10.57800/faunitaxys-11\(68\)](https://doi.org/10.57800/faunitaxys-11(68))
- Ythier E., 2023. – A new species of *Buthus* Leach, 1815 from the Atlantic coast of Morocco (Scorpiones: Buthidae). *Faunitaxys*, 11(69): 1 – 7. [https://doi.org/10.57800/faunitaxys-11\(69\)](https://doi.org/10.57800/faunitaxys-11(69))
- Roux P. & Écharoux D., 2023. – Nouveaux *Pheropsophus* Solier 1833 d'Asie (Coleoptera, Caraboidea, Brachinini). *Faunitaxys*, 11(70): 1 – 7. [https://doi.org/10.57800/faunitaxys-11\(70\)](https://doi.org/10.57800/faunitaxys-11(70))
- Hennemann F. H., 2023. – A taxonomic review, including new species and new records of Philippine Obrimini stick insects (Insecta: Phasmatodea: Heteropterygidae: Obriminae). *Faunitaxys*, 11(71): 1 – 135. [https://doi.org/10.57800/faunitaxys-11\(71\)](https://doi.org/10.57800/faunitaxys-11(71))
- Lacroix M., Coache A. & Filippi G., 2023. – Deuxième contribution à la connaissance des Sericinae de Sao Tomé et Principe (Coleoptera, Scarabaeoidea). *Faunitaxys*, 11(72) : 1 – 11. [https://doi.org/10.57800/faunitaxys-11\(72\)](https://doi.org/10.57800/faunitaxys-11(72))
- Keith D., Pham P.V. & Sabatinelli G., 2023. – *Exolontha permirabilis*, a new species from Vietnam with notes on the genus (Col. Scarabaeidae Melolonthinae Melolonthini). *Faunitaxys*, 11(73): 1 – 5. [https://doi.org/10.57800/faunitaxys-11\(73\)](https://doi.org/10.57800/faunitaxys-11(73))
- Vives E., 2023. – Especies nuevas o interesantes de longicornios de Filipinas (Pars 24). (Coleoptera, Cerambycidae). *Faunitaxys*, 11(74): 1 – 5. [https://doi.org/10.57800/faunitaxys-11\(74\)](https://doi.org/10.57800/faunitaxys-11(74))
- Laguette M., 2023. – Generic revision of *Hypercompe* Hübner, [1818] and allied genera with descriptions of new genera and new species (Lepidoptera: Erebiidae: Arctiinae). *Faunitaxys*, 11(75): 1 – 34. [https://doi.org/10.57800/faunitaxys-11\(75\)](https://doi.org/10.57800/faunitaxys-11(75))
- Ythier E. & Lourenço W. R., 2023. – A new species of *Hadruiroides* Pocock, 1893 from Peru (Scorpiones: Caraboctonidae). *Faunitaxys*, 11(76): 1 – 7. [https://doi.org/10.57800/faunitaxys-11\(76\)](https://doi.org/10.57800/faunitaxys-11(76))
- Lourenço W. R. & Velten J., 2023. – New contribution to the knowledge of Burmite chaerilids and description of a new species (Scorpiones: Chaerilidae). *Faunitaxys*, 11(77): 1 – 4. [https://doi.org/10.57800/faunitaxys-11\(77\)](https://doi.org/10.57800/faunitaxys-11(77))
- Lourenço W. R. & Velten J., 2023. – A remarkable new species of *Betaburmesebuthus* Lourenço, 2015 from Burmite (Scorpiones: Palaeoburmesebuthidae). *Faunitaxys*, 11(78): 1 – 5. [https://doi.org/10.57800/faunitaxys-11\(78\)](https://doi.org/10.57800/faunitaxys-11(78))
- Dalens P.-H. & Robin F., 2023. – Un nouveau *Martinsia* Chemsak & Linsley, 1967 de Guyane (Coleoptera : Cerambycidae, Cerambycinae). *Faunitaxys*, 11(79): 1 – 5. [https://doi.org/10.57800/faunitaxys-11\(79\)](https://doi.org/10.57800/faunitaxys-11(79))
- Vitali F., 2023. – Taxonomical notes on the genus *Cerosterna* Dejean, 1835 with the description of a new species from the Australian realm (Coleoptera, Cerambycidae). *Faunitaxys*, 11(80): 1 – 4. [https://doi.org/10.57800/faunitaxys-11\(80\)](https://doi.org/10.57800/faunitaxys-11(80))

Faunitaxys est échangée avec les revues suivantes (« print versions ») :

- Annali del Museo Civico di Storia Naturale G. Doria (Italie)
- Boletín de la Asociación española de Entomología (Espagne)
- Boletín de la Sociedad Andaluza de Entomología (Espagne)
- Bollettino del Museo di Storia Naturale di Venezia (Italie)
- Bulletin de la Société linnéenne de Lyon (France)
- Bulletin of Insectology (Italie)
- Heteropterus Rev. Entomol. (Espagne)
- Israel Journal of Entomology (Israël)
- Klapalekiana (République Tchèque)
- Koleopterologische Rundschau (Allemagne)
- Memorie del Museo Civico di Storia Naturale di Verona (Italie)
- Nova Supplementa Entomologica (Allemagne)
- Proceedings of the Entomological Society of Washington (USA)
- Revue suisse de Zoologie (Suisse)
- Spixiana (Allemagne)
- Stuttgarter Beiträge zur Naturkunde A, Biologie (Allemagne)
- Zoosystematica Rossica (Russie)

Faunitaxys

Volume 12, Numéro 01, Janvier 2024

SOMMAIRE

Description d'une nouvelle espèce du genre *Nyssodryssilla* Gilmour, 1962 (Cerambycidae : Lamiinae : Acanthocinini) du Brésil et nouveau signalement, plante hôte et parasite d'*Heterachthes sablensis* Blatchley, 1920 (Cerambycinae : Tropidini : Compsina) au Panama.

Alfredo Lanuza-Garay, Marleny Rivera & Alonso Santos Murgas 1 – 4

CONTENTS

Description of a new species of *Nyssodryssilla* Gilmour, 1962 (Cerambycidae: Lamiinae: Acanthocinini) from Brazil and new record, host plant, and parasitoid of *Heterachthes sablensis* Blatchley, 1920 (Cerambycinae: Tropidini: Compsina) in Panama.

Alfredo Lanuza-Garay, Marleny Rivera & Alonso Santos Murgas 1 – 4



Illustration de la couverture :

The nerve center of the Smithsonian Tropical Research Institute stands in Panama City, Panama. Named for benefactor Earl S. Tupper, the center has modern laboratories, a tropical sciences library, a herbarium, an insect collection, a conference center, and the institute's administrative headquarters. Their insect collection stores 132,000 insects from Panama and other countries worldwide.

Crédits photos :

Smithsonian Tropical Research Institute (<https://stri.si.edu/es/files/tuppergallery1jpg>): Couverture

Marleny Rivera: Fig. 1-4

Alfredo Lanuza-Garay: Fig. 5

Alonso Santos Murgas: Fig. 6-7

Publié par l'Association Française de Cartographie de la Faune et de la Flore (AFCFF)