



HAL
open science

Storage protein activator controls grain protein accumulation in bread wheat in a nitrogen dependent manner

Anne Plessis, Catherine Ravel, Thierry Risacher, Nathalie Duchateau, Mireille Dardevet, Marielle Merlino, François Torney, Pierre Martre

► To cite this version:

Anne Plessis, Catherine Ravel, Thierry Risacher, Nathalie Duchateau, Mireille Dardevet, et al.. Storage protein activator controls grain protein accumulation in bread wheat in a nitrogen dependent manner. *Scientific Reports*, 2023, 13 (1), pp.22736. 10.1038/s41598-023-49139-5 . hal-04372143

HAL Id: hal-04372143

<https://hal.science/hal-04372143>

Submitted on 4 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN

Storage protein activator controls grain protein accumulation in bread wheat in a nitrogen dependent manner

Anne Plessis^{1,2}, Catherine Ravel^{1✉}, Thierry Risacher³, Nathalie Duchateau¹, Mireille Dardevet¹, Marielle Merlino¹, François Torney³ & Pierre Martre^{1,4}

The expression of cereal grain storage protein (GSP) genes is controlled by a complex network of transcription factors (TFs). Storage protein activator (SPA) is a major TF acting in this network but its specific function in wheat (*Triticum aestivum* L.) remains to be determined. Here we generated an RNAi line in which expression of the three SPA homoeologs was reduced. In this line and its null segregant we analyzed GSP accumulation and expression of GSP and regulatory TF genes under two regimes of nitrogen availability. We show that down regulation of SPA decreases grain protein concentration at maturity under low but not high nitrogen supply. Under low nitrogen supply, the decrease in SPA expression also caused a reduction in the total quantity of GSP per grain and in the ratio of GSP to albumin-globulins, without significantly affecting GSP composition. The slight reduction in GSP gene expression measured in the SPA RNAi line under low nitrogen supply did not entirely account for the more significant decrease in GSP accumulation, suggesting that SPA regulates additional levels of GSP synthesis. Our results demonstrate a clear role of SPA in the regulation of grain nitrogen metabolism when nitrogen is a limiting resource.

Bread wheat (*Triticum aestivum* L.) is the most important cereal crop in the world in terms of area harvested and commercial exchange. It provides on average 20% of the total protein in the human diet¹. Wheat grain proteins have unique properties making them suitable for use in a considerable number of food and non-food products². Both the total grain protein concentration (GPC) and the relative composition of the storage protein fraction govern the cohesiveness and viscoelasticity of gluten, the network formed by wheat grain storage proteins (GSPs) when mixed with water³. Breeding for high-yielding genotypes has decreased GPC, while modern uses of end-products require higher GPC than traditional products, limiting the potential usefulness of some varieties^{4,5}. Most wheat GSPs belong to the glutenin and gliadin prolamin families. Glutenins usually account for 35–45% of total grain protein and are composed of high-molecular-weight (HMW-GS) and low-molecular-weight (LMW-GS) sub-units, which together form very large macropolymers during grain desiccation⁶. Gliadins are monomeric proteins classed as ω 5-, ω 1,2-, α/β - or γ -gliadins⁷ and make up between 18 and 35% of total grain protein.

Transcriptional control of GSP genes plays an important role in the endosperm specific synthesis of GSPs during cereal grain development⁸ through a network of interacting transcription factors (TFs). At least twelve TFs involved in the regulation of GSP genes have been identified in different cereal species, along with the *cis*-elements they bind to. The wheat GCN4-like motif (GLM) is bound by SPA, a basic leucine zipper TF of the Opaque2 (O2) subfamily^{9,10}, with an ortholog in barley (*Hordeum vulgare* L.) named BLZ2 that binds to the same motif¹¹, as does another member of the family in barley, BLZ1¹²; the ortholog of BLZ1 in wheat is SPA Heterodimerizing Protein (SHP) and is a negative regulator of GSPs¹³. Two non-homologous DNA-binding with one finger (DOF) TFs: prolamin box binding factor (PBF) and scutellum and aleurone-expressed DOF (SAD), interact with the prolamin box^{14,15}; in bread wheat, PBF and SAD promote the transcription of glutenin genes by binding to the prolamin box and this activity is additive to the induction of the expression of glutenin genes by SPA¹⁶. In barley, an AACA motif is recognized by GAMYB (gibberellic acid-dependent of the MYB superfamily of transcriptional activators, Diaz et al. 2002). In barley, two R1MYB family TFs, Myb-related CAB

¹Université Clermont Auvergne, INRAE, UMR1095 GDEC, 63000 Clermont Ferrand, France. ²Present address: School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK. ³Centre de Recherche, Limagrain Europe, 63 720 Chappes, France. ⁴Present address: LEPSE, Université de Montpellier, INRAE, Institut SupAgro Montpellier, 34000 Montpellier, France. ✉email: catherine.ravel@inrae.fr

promoter-binding protein (MCB1) and MYBS3, can bind a GA response complex motif^{18,19} and FUSCA3, a B3-type TF, interacts with an RY box²⁰. The DOF proteins mediate the formation of several binary or ternary TF complexes^{15,17,18} in addition to the interactions between FUSCA3 and BLZ2²⁰ and between BLZ1 and BLZ2²¹. The interaction of FUSCA3 with the RY box of a glutenin gene and with SPA has been demonstrated in wheat²¹. New transcription factors regulating the expression of prolamin genes have been discovered more recently in wheat: three negative regulators of GSPs, SPR and ODORANT1^{22–24}, for which the exact location where they bind on prolamin promoters is still unclear, and one positive regulator, NAC109 that interacts with GAMYB²⁵.

SPA/O2/BLZ2 is one of the most studied TFs in this regulatory network of grain storage proteins. In wheat, a study of different SPA haplotypes has shown that this gene affects the amount of nitrogen allocated to the gliadin fraction²⁶. In maize, O2 is associated with grain lysine content²⁷ and an o2 mutant shows reduced expression of some GSP genes and corresponding proteins²⁸. The effect of modifying expression levels of RISBZ1, a SPA homolog in rice, has been tested^{29,30}. Transient overexpression of RISBZ1 in protoplasts induced trans-activation of several GSP gene promoters, which was synergistically enhanced by the simultaneous overexpression with the rice homolog of PBF³⁰. The knockdown of RISBZ1 *in planta* caused only slight changes in GSP accumulation, but when both RISBZ1 and RPBF were knocked down, GSP accumulation and gene expression were significantly reduced²⁹. In wheat, the overexpression of the copy of SPA located on the B genome led to lower accumulation of glutenin and ω -gliadin and lower expression of PBF³¹.

Nutrient availability has a major effect on GSP quantity and composition³². Higher nitrogen input increases the amount of GSP that accumulates in the grain³³. Differences in GSP composition related to nitrogen availability have been found to follow allometric scaling laws^{34–36}, which may be a consequence of the complex transcriptional regulation network controlling GSP gene expression²⁶. One particular *cis*-element on GSP gene promoters, the GLM, which is bound by SPA, plays an important role in the transcriptional response to nitrogen. It was found to be essential for the activation of GSP gene transcription in response to amino acids and ammonium, but this activation is only optimal in synergistic interaction with the endosperm box (EB), the motif formed by the GLM and the prolamin box in tandem³⁷, and other *cis*-elements³⁸. Interestingly, when nitrogen supply is low, the GLM may act as a negative regulatory motif for GSP gene transcription.

SPA orthologs thus have a prominent role in controlling GSP synthesis in cereals, particularly in response to changes in nitrogen availability. To confirm that SPA is a positive regulator of GSP gene expression in bread wheat and determine its role in the response of GSP accumulation and composition to nitrogen availability, plants with reduced expression of SPA were grown with high or low amounts of nitrogen. We analyzed the relative accumulation of the different GSP families and sub-groups and measured the expression of GSP and their regulatory TF genes throughout grain development. We show that a decrease in the expression of SPA caused a reduction in GPC, which was more significant under low nitrogen availability. GSP accounted for most of the decrease in GPC compared to other protein classes, with reductions in the amounts of all GSP families. We also describe the changes in regulatory TF expression that contribute to the response to nitrogen with notably PBF, MYBS3 and SHP being upregulated by high nitrogen. Our results suggest that transcriptional regulation is not the sole mechanism determining the response of GSP synthesis to nitrogen availability.

Methods

All methods were performed in accordance with the relevant guidelines and regulations.

Plant material and growth conditions

Immature seeds of the spring wheat (*Triticum aestivum* L.) line NB1 (a non-commercial Spring wheat variety obtained from Limagrain Europe, Saint-Beauzire, France) were transformed by *in planta* inoculation using *Agrobacterium tumefaciens* and transgenic lines were regenerated³⁹. The SPA-A full length coding sequence was previously amplified by PCR from a cDNA library from immature seeds of the bread wheat cultivar Récital. The binary vector pSCV was used to produce transgenic plants expressing both sense and antisense SPA-A cDNA separated by the first intron of rice tubulin to generate a hairpin RNA (Fig. 1). The transgene was under the control of the promoter of the subunit Dx5 of the *Glu-D1-1* HMW-GS gene and the *Nos* terminator. The plasmid includes a kanamycin resistance cassette for selection, *NptII*, controlled by the actin promoter and *Nos* terminator. For each transformation event, the number of T-DNA insertions was evaluated by qPCR performed on genomic DNA and the integrity of the transgene was verified by PCR. Transformants with several copies of the transgene were discarded. T0 plants were self-pollinated to generate the T1 generation composed of 25% homozygotes, 50% hemizygotes and 25% null segregant plants. The segregation ratio was established by cultivating 30 plants on a medium with kanamycin allowing the identification of homozygotes, hemizygotes and null segregants (Table S1). The zygosity of progenies from self-pollinated homozygotes and respective null segregants




Figure 1. Storage protein activator RNA interference construct used for wheat transformation. LB, left border; RB, Right border.

was verified by quantitative real-time PCR. Selfing of confirmed homozygotes and null segregant T2 plants gave rise to the T3 generation, i.e. the RNAi line and its null segregant to be used as a control.

T4 seeds were germinated for two to three days at room temperature on wet filter paper in Petri dishes. Germinated seeds were then transferred to soil in 50-mL PVC columns (inner diameter 7.5 cm, length 50 cm, 2 plants per column) and arranged in a greenhouse in a strip-plot design with the genotypes as rows and the nitrogen treatments (see below) as columns with four replicated blocks to form a homogeneous stand with a plant density of 261 plants m⁻². Temperature was controlled at 22 °C during the day and 18 °C during the night. Day length was 16 h, maintained with artificial light when needed. Plants received 68 mL column⁻¹ day⁻¹ of water or nutrient solution. Three nutrient solutions were used to feed the plants, N0, N3 and N15. N0 contained 1 mM KH₂PO₄, 0.5 mM NH₄NO₃, 2 mM MgSO₄, 7 mM CaCl₂, 5 mM KCl, 10 μM H₃BO₃, 0.7 μM ZnCl₂, 0.4 μM CuCl₂, 4.5 μM MnCl₂, 0.22 μM MoO₃, and 50 μM EDFS-Fe; N3 contained 1 mM KH₂PO₄, 1 mM Ca(NO₃)₂, 0.5 mM NH₄NO₃, 2 mM MgSO₄, 3 mM CaCl₂, 5 mM KCl, 10 μM H₃BO₃, 0.7 μM ZnCl₂, 0.4 μM CuCl₂, 4.5 μM MnCl₂, 0.22 μM MoO₃, and 50 μM EDFS-Fe; N15 contained 1 mM KH₂PO₄, 5 mM KNO₃, 4 mM Ca(NO₃)₂, 1 mM NH₄NO₃, 2 mM MgSO₄, 10 μM H₃BO₃, 0.7 μM ZnCl₂, 0.4 μM CuCl₂, 4.5 μM MnCl₂, 0.22 μM MoO₃, and 50 μM EDFS-Fe. All plants received N3 for four weeks, then N15 until anthesis. At anthesis, continuous water irrigation was used to remove any excess of the nutrient solution in the soil; afterwards irrigation was maintained to its previous level with water. Differences in treatment between N⁻ and N⁺ started at 300 °C days after anthesis, when the columns were rinsed again and lasted until grain ripeness: N⁺ plants received the N15 nutrient solution while N⁻ plants received N0. Main stems were tagged when the anthers of the central florets appeared. Degree-days were calculated as the sum of the average daily temperatures after anthesis with a base temperature of 0 °C.

Determination of grain dry mass and protein concentration

Grains from four ears (except at 200 °C days after anthesis, where five ears were used) were sampled from each replicate every 100 °C days from 200 °C days after anthesis to maturity (grain ripeness, 900 °C days after anthesis) and again at 1050 °C days after anthesis. Four grains per ear were sampled between 200 and 700 °C days after anthesis for RNA analysis and were immediately frozen in liquid nitrogen and stored at - 80 °C. The remaining grains of the ear were also frozen and stored at - 80 °C until they were freeze-dried. We measured the dry mass and nitrogen concentration of a sub-sample of grains (ca. 65%). The remaining grains were oven-dried at 80 °C for 48 h to calculate the percentage of remaining water. At maturity, grain yield per ear was calculated using all grains of each ear harvested.

Grains were milled for 2 min using a custom ball mill. Flour (5 mg) was weighed in tin capsules and the total N concentration was determined with the Dumas combustion method (Association of Analytical Communities International approved method no. 992.23) using a FlashEA 1112 N/Protein Analyzer (Thermo Electron Corp, Waltham, MA). Grain protein concentration (GPC) was calculated by multiplying grain N concentration by 5.62⁴⁰.

Sequential extraction, separation and quantification of grain protein fractions

Non-prolamin, gliadin and glutenin protein fractions were sequentially extracted from 60 mg of freeze-dried wholemeal flour as described by Triboui et al.³⁶ and modified by Plessis et al.⁴¹. Each 2 mL tube contained one stainless steel bead (5 mm diameter) and samples were stirred by placing the tubes on a rotating wheel (40 rpm) during each extraction and washing step. The non-prolamin protein fraction was extracted for 30 min at 4 °C from 100 mg wholemeal flour with 1.5 mL of 50 mM phosphate buffer (pH 7.8) containing 0.1 M NaCl. After centrifugation for 10 min (18,000 g) at 4 °C, the supernatant was collected and the pellet was washed twice for 10 min each time with 1.5 mL of the same buffer. After centrifugation in the same conditions, all supernatants were pooled. The same steps were used to extract the gliadin protein fraction from the previous pellet with 70% (v/v) ethanol. Finally, the glutenin protein fraction was extracted in 50 mM borate buffer (pH 8.5) containing 2% SDS (w/v) and 1% dithiothreitol (w/v). The supernatants (80 μL) of each protein fraction were oven dried overnight at 60 °C in tin capsules and their total nitrogen concentration was determined with the Dumas combustion method as described above. Protein fractions from samples of the same flour from cultivar Récital were extracted, analysed as a control in each of the 21 sets of extractions and used to determine the coefficient of variation for each of the protein fractions, which were 3.48, 5.10, 2.19, 2.61, and 1.96% for the non-prolamin, gliadin, and glutenin protein fractions, storage proteins, and total proteins, respectively.

Gliadin classes (ω 1,2-, α/β -, and γ -gliadins) and glutenin sub-units (HMW-GS and LMW-GS) were separated and quantified by HPLC (Figs. S1 and S2) using an Agilent 1290 Infinity LC system (Agilent Technologies, Santa Clara, CA, <http://www.agilent.com>) as described in Triboui et al.³⁴. The total nitrogen concentration of each protein fraction was determined by the Dumas combustion method, as described above. The gliadin extracts used were those obtained by sequential extraction, but glutenins were extracted from flour independently with a protocol adapted from Fu and Kovacs⁴². The gliadin and glutenin extracts were filtered through regenerated cellulose syringe filters (0.45-μm pore diameter, UptiDisc; Interchim, <http://www.interchim.com>), and 4 μL (gliadin) or 2 μL (glutenin) of protein extract was injected into a C8 reversed-phase Zorbax 300 StableBound column (2.1 × 9 × 100 mm, 3.5 μm, 300 Å; Agilent Technologies) maintained at 50 °C. The eluents used were ultra-pure water (solvent A) and acetonitrile (solvent B), each containing 0.1% trifluoroacetic acid. The flow rate was 1 mL min⁻¹. Proteins were separated by using a linear gradient, from 24 to 50% solvent B over 13 min for gliadin, and from 23 to 42% solvent B over 25 min for glutenin. Proteins were detected by UV absorbance at 214 nm. After the gradient, the column was washed with 80% solvent B for 2 min and then equilibrated at 24% (for gliadins) or 23% (for glutenins) solvent B for 2 min at the same flow rate. Chromatograms were processed with CHEMSTATION 10.1 software (Agilent Technologies). The signal obtained from a blank injection was subtracted from the

chromatograms before integrating the data. The HPLC peaks corresponding to each of the four gliadin classes were identified following the observations of Wieser et al.⁴³. The quantity of each gliadin class or glutenin subunit as a percentage of total gliadin or total glutenin, respectively, was calculated by dividing the areas under each HPLC peak by the total area under the chromatogram trace. The quantity of each gliadin class (or glutenin subunit) per grain was calculated by multiplying the proportion of each gliadin class (or glutenin subunit) in total gliadin (or total glutenin) by the total quantity of gliadin (or glutenin) per grain, as quantified by Dumas analysis. By subtracting the quantity of all GSPs from N_{tot} , we calculated the amount of the remaining protein fraction, mainly constituted of albumin-globulins.

RNA extraction and measurement of gene expression

Four grains per ear (same ears as for protein analysis) were sampled at 200, 300, 400, 500, 600 and 700 °C days, the embryos were cut out and the rest of the grain immediately frozen in liquid nitrogen and kept at - 80 °C. RNA was extracted from 75 mg of grain powder in 750 µL of extraction buffer (200 mM Tris-HCl pH 9, 400 mM KCl, 200 mM sucrose, 35 mM MgCl₂, 25 mM EDTA) and 600 µL phenol/chloroform (pH 8). The suspension was homogenized by vortexing for 30 s and then centrifuged for 10 min at 15,000 × g. The supernatant was collected. The pellet was resuspended in 600 µL of phenol/chloroform, centrifuged using the same conditions and the supernatant collected, and the whole step repeated. Supernatants were pooled. RNA was precipitated by adding 1 M acetic acid (1/10 volume) and ethanol (2.5 volumes). The RNA pellet was washed with 3 M Na acetate (pH 6) and resuspended in water. A second acetic acid/ethanol precipitation was performed before resuspending the pellet in 50 µL RNase free water. RNA was treated with RNase-free DNase according to the instructions of the supplier (AMBION). The RNA in solution was quantified by measuring the absorbance at 260 nm in a spectrophotometer. Approximately 2 µg of total RNA were reverse transcribed using oligo(dT)₂₀ and reverse transcriptase (Bio-rad iScript™ Select cDNA Synthesis kit) in a final volume of 40 µL. Transcript levels of four housekeeping genes and the storage protein and transcription factor (TFs) genes were quantified by real-time q-PCR using Lightcycler 480 SYBR Green I Master (Roche) in 15 µL with 5 µL of cDNA diluted 10 times. Relative expression (RE) was calculated as: $RE = \varepsilon^{\Delta C_p}$, where ε is the efficiency of the primers for the measured gene and ΔC_p is the normalized crossing point (Cp); $\Delta C_p = (C_{p1} \times C_{p2} \times C_{p3} \times C_{p4})^{0.25} - C_{pg}$, where C_{pg} is the Cp for the measured gene and C_{p1} , C_{p2} , C_{p3} and C_{p4} are the Cp values of the four housekeeping genes⁴⁴. The primer sequences are given in Table S2.

Soluble protein extraction and western blot analysis

Wheat flour (50 mg) from grains collected at 500 °C days after anthesis was dissolved in extraction buffer (10 mM sodium phosphate, 10 mM NaCl, pH 7.8 at 4 °C) supplemented with a protease inhibitor cocktail (P9599, Sigma, St. Louis, MO, USA). Proteins were precipitated from the extract supernatant with ice-cold acetone overnight at - 20 °C. The dried protein pellet was dissolved in SDS-PAGE buffer containing 80 mM Tris-HCl pH 8, 2% (w/w) SDS, 40% glycerol (v/v), 0.002% bromophenol blue (w/w), supplemented with 2% (v/v) DTT and 2.5% iodoacetamide (w/w). The protein concentration was determined using the Bradford protein assay (B6916, Sigma, St. Louis, MO, USA). In order to quantify the SPA protein, different quantities of total soluble protein extracts (10, 20, 30, and 40 µg) were separated on SDS-polyacrylamide gel (T = 10.3%, C = 1.3%).

After electrophoresis, proteins were transferred onto a nitrocellulose membrane (Hybond, ECL, GE Healthcare) using a Criterion blotter (Biorad). The membrane was incubated for 1 h at room temperature in a blocking buffer containing 10 mM Tris-HCl pH 7.6, 150 mM NaCl, 0.01% (v/v) Tween 20, and 5% (w/w) skimmed milk. The membrane was then incubated for 1 h at room temperature and overnight at 4 °C with a 1:1000 dilution of anti-SPA antibody (Eurogentec S.A., Belgium). Antirabbit IgG conjugated with horseradish peroxidase (GE Healthcare) was used as the secondary antibody (diluted 1:50,000). The signal was detected using an Immobilon™ Western Chemiluminescent HRP Substrate (ECL Millipore) following the manufacturer's protocol. Anti-SPA signals were quantified by image analysis using Image J software (<http://imagej.nih.gov/ij>). The fold-change in SPA protein abundance was calculated as the ratio of the slope of the relationship between protein amount and anti-SPA signal for the SPA RNAi and NS lines⁴⁵.

Data analysis

All statistical analyses were done in R-4.2.3 for Windows⁴⁶ (code provided in the Supplementary Information). An ANOVA model with two factors (genotype and block) was used to analyze the results. Genotype and block were regarded as fixed effects. Variance homogeneity was tested using the Bartlett test and the normality of the residuals with the Shapiro-Wilk test. The block effect was never statistically significant. Differences between NS line and SPA RNAi line were tested using the post-hoc Dunnett test, with the NS line used as control. Statistical differences were judged at the 5% level. Differences in SPA protein abundance between NS line and SPA RNAi line were tested by comparing the slopes of the standard major axis regression between the normalized volume of anti-SPA signal and the total protein mass using the 'smatr' package⁴⁷.

Results

SPA RNA and protein quantities are reduced in the SPA RNAi line

To investigate the role of SPA in regulating storage protein accumulation in bread wheat grain we generated RNAi transgenic lines to down regulate this gene. The transgene was under the control of the promoter of a HMW-GS gene. More precisely, the promoter of the allele encoding the Dx5 subunit of *Glu-D1-1* was used. This promoter is grain specific and highly induced during grain development⁴⁸ (Fig. 1). We grew SPA RNAi plants and their null-segregant siblings (NS) in the greenhouse. Grain developmental stage was measured in degree-days (°C days) after anthesis to take into account the effect of temperature on development. At 300 °C days after anthesis




Figure 2. Down-regulation of SPA in the RNAi line. SPA null segregant (NS, circles) and RNAi (RNAi, triangles) lines of bread wheat were grown in the greenhouse with low (N-, open symbols) and high (N+, closed symbols) nitrogen supply. (A–F) Relative expression of SPA homoeologs. (G) and (H) Relative expression of the sum of the three SPA homoeologs. (I) and (J) Quantification of the anti-SPA signal from western blots with different quantities of total protein extract at 500 °C days after anthesis. (A–H) Data are means for $n = 4$ independent replicates.




Figure 3. Changes in grain dry mass, total quantity of nitrogen (N_{tot}) in the grain and grain protein concentration during grain development. *SPA* null segregant (NS, circles) and RNAi (RNAi, triangles) lines of bread wheat were grown in the greenhouse with low (N $^{-}$, open symbols) and high (N $^{+}$, closed symbols) nitrogen supply. DM, dry mass. Data are means ± 1 s.e. for $n=4$ independent replicates.

Variable	Treatment			
	N–		N+	
	NS	RNAi	NS	RNAi
Single grain dry mass (mg DM grain ⁻¹)	40.0 ± 0.5	41.2 ± 0.8	40.9 ± 0.6	42.6 ± 0.7
Grain yield per ear (g DM ear ⁻¹)	0.96 ± 0.03	1.05 ± 0.05	1.04 ± 0.02	1.14 ± 0.04*
Grain N (mg N grain ⁻¹)	0.92 ± 0.02	0.86 ± 0.01**	1.24 ± 0.02	1.27 ± 0.03
Grain N yield per ear (mg N ear ⁻¹)	22.4 ± 0.5	22.3 ± 0.8	32.0 ± 0.8	35 ± 1.3
Grain protein concentration (% of DM)	13.2 ± 0.2	11.9 ± 0.2***	17.3 ± 0.2	17.0 ± 0.3

Table 1. Single grain dry mass, grain yield per ear, total quantity of N per grain (N_{tot}), grain N yield per ear, and grain protein concentration at maturity (900 and 1050 °C days) for the null segregant (NS) and SPA RNAi (RNAi) lines under low (N–) and high (N+) nitrogen availability. Data are means ± 1 s.e. for $n = 8$ independent replicates. Within a single N treatment, * ($P < 0.05$), ** ($P < 0.01$) and *** ($P < 0.001$) indicate significant differences between values for NS and RNAi lines from an ANOVA followed by the Dunnett post-hoc test.

the plants were either supplied with 15 mM nitrogen (N+ treatment) or no nitrogen (N– treatment) until the end of grain filling about 800 °C days after anthesis. We sampled grains every 100 °C days from 200 to 800 °C days after anthesis and then at maturity, 900 °C days and 1050 °C days after anthesis.

We demonstrated RNA interference in one of the transgenic lines generated by measuring the expression of the three SPA homoeologs during grain development using q-PCR (Fig. 2). In the RNAi line fewer SPA-A transcripts were detected throughout development under both nitrogen treatments compared to NS. As SPA-A is the most highly expressed of the three homoeologs, the decrease in its expression was largely responsible for the overall decrease in the relative abundance of SPA transcripts (Fig. 2G and H). For example, relative expression of total SPA transcripts was > 20% less in the RNAi line than in NS 500 °C days after anthesis for the N– treatment and 400 °C days after anthesis for the N+ treatment. Interestingly, SPA-B showed a different time-course of expression from SPA-A and SPA-D in both the NS and RNAi line. This difference has already been observed in different genetic backgrounds²⁶ and suggests distinct regulation of the different homoeologs of SPA.

We also checked the effect of RNAi on SPA protein levels. We carried out a western blot analysis on grains collected 500 °C days after anthesis (Fig. 2I and J). The quantity of SPA protein was reduced by 34% ($P = 0.071$) for the N– treatment (Fig. 2I) and 61% ($P = 0.057$) for the N+ treatment (Figs. 2J and S3). Therefore less SPA protein accumulated when expression of SPA was down regulated by RNAi.

Nitrogen accumulation is modified in the SPA RNAi line under low nitrogen availability

For plants subjected to the N– treatment, the total quantity of nitrogen per grain (N_{tot}) and grain protein content (GPC) at maturity were reduced by 6% ($P = 0.043$) and 9% ($P < 0.001$) respectively in the SPA RNAi line compared with the NS. In the case of GPC, the decrease in the RNAi line compared to the NS was already significant ($P < 0.05$) at 400 and 700 °C days after anthesis (Fig. 3C). Under these conditions of low nitrogen availability, the greatest difference in GPC between SPA RNAi line and the NS was at maturity, which can be attributed to a late increase in single grain dry mass and a decrease in N_{tot} (Fig. 3A–C).

In contrast, for plants subjected to the N+ treatment, N_{tot} was not significantly different ($P = 0.30$) between the SPA RNAi and NS lines (Fig. 3B) and GPC was only significantly reduced (– 5%) in the RNAi line at 400 and 500 °C days after anthesis (Fig. 3C). Grain yield per ear was increased by 9% while it was not significantly altered in the N– treatment.

Single grain dry mass at maturity was not significantly different between the SPA RNAi and NS lines for either of the nitrogen availability treatments (Table 1).

Storage protein accumulation but not composition is affected in the SPA RNAi line under low nitrogen availability

We determined grain protein composition throughout grain development in the NS and SPA RNAi lines. Under N– treatment, decreases in all GSP fractions were measured from around 400 °C days and onwards in the SPA RNAi line compared to NS, while a slight decrease in the albumin-globulin (AG) fraction was only detected at maturity (Fig. 4). Decreases in GSPs must have contributed more to the overall decrease in GPC under low nitrogen availability as the GSP to AG ratio was lower in the SPA RNAi line than in NS (Table 2). In the N+ treatment, the glutenin subunits, gliadin classes and AG mostly accumulated at similar rates in the SPA RNAi and NS lines (Fig. 4) and at maturity the GSP to AG ratio was the same (Table 2). At maturity, for both N treatments, the gliadin to glutenin ratio tended to be lower (– 6%) in the SPA RNAi line than NS but the difference was not statistically significant ($P = 0.5$).

For N– conditions, the relative proportions of GSP fractions in total GSP were the same in the NS and SPA RNAi lines (Table 2), showing that all GSP fractions contributed to the same relative extent to the decrease in GSP quantity when SPA was downregulated. An exception was $\omega 1,2$ -gliadin, which was reduced by 18% ($P < 0.001$) in the SPA RNAi line compared with NS. However $\omega 1,2$ -gliadin makes up less than 4% of the total amount of GSP.




Figure 4. Changes in the quantity per grain of protein fractions during grain development. SPA null segregant (NS, circles) and RNAi (RNAi, triangles) lines of bread wheat were grown in the greenhouse with low (N-, open symbols) and high (N+, closed symbols) nitrogen supply. AG, albumin-globulin; HMW-GS, high molecular weight glutenin subunits; LMW-GS, low molecular weight glutenin subunits. Data are means \pm 1 s.e. for $n = 4$ independent replicates.

Variable	Treatment			
	N–		N+	
	NS	RNAi	NS	RNAi
GSP to AG ratio	1.59 ± 0.04	1.59 ± 0.04	1.59 ± 0.04	1.59 ± 0.04
HMW-GS (% GSP)	20.4 ± 0.24	21.0 ± 0.34	19.0 ± 0.52	18.5 ± 0.18
LMW-GS (% GSP)	42.4 ± 0.50	42.6 ± 0.52	40.6 ± 0.62	40.0 ± 0.28
ω1,2-gliadins (% GSP)	2.8 ± 0.08	2.4 ± 0.12**	5.3 ± 0.28	5.1 ± 0.10
α/β-gliadins (% GSP)	17.4 ± 0.31	17.0 ± 0.37	18.9 ± 0.54	19.7 ± 0.19
γ-gliadins (% GSP)	15.8 ± 0.28	15.7 ± 0.37	15.1 ± 0.35	15.6 ± 0.19

Table 2. Grain storage protein (GSP) to albumin-globulin (AG) ratio under low (N–) and high (N+) nitrogen availability and GSP composition at maturity (900 and 1050 °C days) under low nitrogen availability, shown as the percentage of high (HMW-GS) and low (LMW-GS) molecular weight glutenin subunit, ω1,2-gliadin, α/β-gliadins and γ-gliadin fractions in total GSP. NS, null segregant line. RNAi, SPA RNAi line. Data are means ± 1 s.e. for $n = 8$ independent replicates. Within a single N treatment, ** ($P < 0.01$) indicate significant differences between NS and RNAi lines from an ANOVA followed by the Dunnett post-hoc test.

SPA under-expression decreases gliadin gene expression more than glutenin gene expression

As in bread wheat SPA is a transcriptional regulator of GSP genes¹³, we measured the expression of genes belonging to the different GSP families in the wheat SPA RNAi line (Fig. 5). For most families we used generic q-PCR primers to amplify transcripts of all the genes of a given family. For HMW-GS we amplified transcripts of the four HMW-GS genes expressed in the line used for transformation separately then summed the result. Our results show that all gene families were down regulated in the SPA RNAi line compared to NS for at least one time point of either of the treatments. In N+ conditions, the expression of some GSP genes was affected in the SPA RNAi line, mainly at the later time points (500 °C days after anthesis and onwards). The time points coincide with the largest differences in SPA expression in the RNAi line (Fig. 2). In the N+ treatment, glutenin genes were the least affected in the SPA RNAi line with no change in expression detected for LMW-GS, while gliadin genes showed the most striking decrease in expression (Fig. 5). Similar results were observed for the N– treatment but the differences between the SPA RNAi line and the NS line were smaller than for the N+ treatment.

Expression of TFs known to be part of cereal GSP regulatory network was also quantified during the linear grain filling period (Fig. 6). The expression of *PBF*, *MYBS3*, *GAMYB* and *MCB1* was lower in N+ than in N– conditions in both SPA RNAi and NS lines. The expression of *SAD* and *SHP* was upregulated in the N+ treatment compared with the N– treatment in NS for at least two time points, but in the SPA RNAi line *SHP* did not respond to nitrogen supply. *PBF* and *GAMYB* expression was downregulated for one or more time points in the SPA RNAi line compared with NS under N– conditions, while *SHP* and *GAMYB* were downregulated in the SPA RNAi line compared with NS under N+ conditions for at least one time point.

Discussion

In this study, we show that the under-expression of SPA in bread wheat can result in reduced expression of GSP genes and when nitrogen is limiting a decrease in GPC and the GSP to AG ratio of grain. These results confirm previous indications that SPA has a role in the transcriptional regulation of GSP genes¹³ like its orthologs in other cereals^{11,28,30}. However, down-regulating SPA had a fairly limited effect on GSP gene expression. This may have been because the decrease in SPA gene expression was not large enough to induce a stronger effect or because of functional redundancy like in rice where the under-expression of both *RISBZ1*, the SPA ortholog, and *RPBF* resulted in a much more significant decrease in GSP gene expression than in the *RISBZ1* knock-down line²⁹.

Our results show that SPA under-expression has a stronger effect on the expression of gliadin than glutenin genes. This is consistent with a study of natural genetic variation in wheat where two haplotypes of SPA-A were identified. Different levels of SPA-A expression from each haplotype led to the allocation of different quantities of total grain nitrogen to the gliadin fraction, but equal amounts to the glutenin fraction²⁶.

SPA activates the expression of LMW-GS and HMW-GS by binding the GLM^{9,10,49}, so we expected to observe a lower level of glutenin expression in the SPA RNAi line. However, HMW-GS and LMW-GS gene expression was barely affected by SPA under-expression (Fig. 5A and B). For both HMW-GS and LMW-GS, the level of gene expression we measured here was the combined expression of several members of the gene family. Nucleotide diversity in promoters of LMW-GS gene family members⁵⁰ could however result in differences in expression patterns. As we were not able to monitor the expression of each individual LMW-GS gene, particularly the one used in previous studies^{9,49}, the known effect of SPA regulation might have been masked by different responses from the rest of the gene family. The GLM motifs identified in the promoters of HMW-GS genes were shown to be functional as they are activated after SPA binding¹⁰; the same applies to the G-like box with bZIP proteins⁴⁸. As these boxes belong to a common regulatory framework shared by all the HMW-GS promoters¹⁰, it is expected that the entirety of HMW-GS genes respond in the same way.

Changes in GSP gene expression alone did not account for all of the modifications in GSP accumulation observed in the SPA RNAi line under low nitrogen supply. In addition differences in gene expression did not always lead to differences in protein accumulation. For example, gene expression of gliadins was lower in the SPA RNAi line than in the WT under high nitrogen availability at some time points, but this did not result in




Figure 5. Changes in gene expression of grain storage proteins during grain development. *SPA* null segregant (NS, circles) and RNAi (RNAi, triangles) lines of bread wheat were grown in the greenhouse with low (N-, open symbols) and high (N+, closed symbols) nitrogen supply. HMW-GS, high molecular weight glutenin subunits; LMW-GS, low molecular weight glutenin subunits. Data are means \pm 1 s.e. for $n=4$ independent replicates.

any detectable change in grain protein accumulation or composition. This indicates that mechanisms other than transcription regulate GSP synthesis compensating for the deregulation of GSP expression. Discrepancies between gene expression and protein accumulation in wheat have already been observed for γ -gliadins in a study of the effect of nitrogen and sulfur availability on the regulation of wheat GSP synthesis³⁴ and for gliadins as part of the characterization of *nac019* triple mutants²⁵. Moreover, an association study of wheat grain protein composition found that several nitrogen assimilation and metabolism genes were associated with GPC and N_{tot} , further evidence of levels of non-transcriptional regulation of grain protein accumulation⁴¹. Field studies have shown that N_{tot} is mainly determined by the supply of N to grains and its accumulation is therefore mainly source driven^{35,51,52}.

The effect of under-expressing *SPA* was dependent on nitrogen availability as we only observed changes in grain protein content and composition under the low nitrogen treatment. This does not seem to be related to *SPA* being more highly expressed in the RNAi line when nitrogen was available, as on the contrary, there was still an obvious decrease in *SPA* protein concentration compared to the control. GLM binding by *SPA* has already been shown to have a role in integrating the effects of nitrogen availability at the transcriptional level on GSP synthesis in barley grain³⁸. Here we saw in wheat that this occurs mostly at the protein synthesis level. In *o2*, a maize mutant for the *SPA* ortholog *O2*, genes involved in amino acid metabolism are differentially expressed²⁸. Under varying levels of nitrogen and sulfur availability, amino acid transport and metabolism are modulated to adjust wheat




Figure 6. Changes in gene expression of transcription factors of the grain storage protein transcriptional regulation network during grain development. *SPA* null segregant (NS, circles) and RNAi (RNAi, triangles) lines of bread wheat were grown in the greenhouse with low (N⁻, open symbols) and high (N⁺, closed symbols) nitrogen supply. Full details of gene names are provided in the text. Data are means \pm 1 s.e. for $n=4$ independent replicates.

GSP synthesis and composition³⁴. Thus *SPA* probably modulates the nitrogen response at different levels, both directly by regulating GSP gene expression and indirectly by controlling the expression of other genes involved in regulating GSPs either transcriptionally, as suggested by our results, or translationally⁵³.

We attempted to generate plants over-expressing SPA but none of the transformed lines showed the expected increase in SPA gene expression. This could mean that over-expression of SPA is detrimental to the development of the embryo or to germination. Our use of an HMW-GS gene promoter for the SPA transgene may have resulted in a lethal dose of SPA as it contains an activation domain for SPA itself that might have generated a feed-forward regulatory loop. Another study was more successful in producing SPA over-expressors in bread wheat, achieving ten to 20-fold increases in the expression of *TaSPA-B*³¹; while they used an HMW-GS promoter like us, it came from a different allele and it is unclear which part of the promoter they used, therefore it is possible the absence of some boxes led to lower, and thus non-lethal, levels of SPA expression than in our transformants. Unexpectedly in regards to our results and previous studies¹³, Guo et al.³¹ found that the over-expression of SPA did not lead to higher accumulation of GSP, and on the contrary diminished the quantity of glutenin and ω -gliadin. This could be at least partly due to the indirect effect of SPA over-expression reducing the expression of the glutenin activator *PBF*¹⁶ and increasing the expression of the glutenin repressor *SHP*¹³. The study by Guo et al.³¹ was done at a single level of nitrogen supply and our results, along with previous work³⁸, show that SPA regulation of GSP accumulation and the expression of other transcription factors in the regulatory network is dependent on nitrogen availability. It is thus possible that different consequences of increasing the expression of SPA would have been obtained varying nitrogen supply and that using plants with modified expression of SPA might improve GPC under certain conditions of fertilization. More generally, any attempt at improving GPC in cereals should involve testing in a wide range of nutritional conditions.

Data availability

The datasets generated during the current study are available from the corresponding author on reasonable request.

Received: 12 September 2023; Accepted: 5 December 2023

Published online: 20 December 2023

References

1. FAOSTAT. <https://www.fao.org/faostat/en/#data/SCL> (2020).
2. Shewry, P. R. Wheat. *J. Exp. Bot.* **60**, 1537–1553 (2009).
3. MacRitchie, F. Wheat proteins: Characterization and role in flour functionality. *Cereal Foods World* **44**, 188–193 (1999).
4. Oury, F.-X. & Godin, C. Yield and grain protein concentration in bread wheat: How to use the negative relationship between the two characters to identify favourable genotypes?. *Euphytica* **157**, 45–57 (2007).
5. Aguirrezabal, L., Martre, P., Pereyra-Irujo, G., Izquierdo, N. & Allard, V. *Management and Breeding Strategies for the Improvement of Grain and Oil Quality* (Elsevier Academic Press Inc, 2009).
6. Don, C., Mann, G., Bekes, F. & Hamer, R. J. HMW-GS affect the properties of glutenin particles in GMP and thus flour quality. *J. Cereal Sci.* **44**, 127–136 (2006).
7. Wieser, H. Chemistry of gluten proteins. *Food Microbiol.* **24**, 115–119 (2007).
8. Bartels, D. & Thompson, R. Synthesis of messenger-Rnas coding for abundant endosperm proteins during wheat-grain development. *Plant Sci.* **46**, 117–125 (1986).
9. Albani, D. et al. The wheat transcriptional activator SPA: A seed-specific bZIP protein that recognizes the GCN4-like motif in the bifactorial endosperm box of prolamin genes. *Plant Cell* **9**, 171–184 (1997).
10. Ravel, C. et al. Conserved cis-regulatory modules in promoters of genes encoding wheat high-molecular-weight glutenin subunits. *Front. Plant Sci.* **5**, 621 (2014).
11. Onate, L., Vicente-Carbajosa, J., Lara, P., Diaz, I. & Carbonero, P. Barley BLZ2, a seed-specific bZIP protein that interacts with BLZ1 in vivo and activates transcription from the GCN4-like motif of B-hordein promoters in barley endosperm. *J. Biol. Chem.* **274**, 9175–9182 (1999).
12. Vicente-Carbajosa, J., Onate, L., Lara, P., Diaz, I. & Carbonero, P. Barley BLZ1: A bZIP transcriptional activator that interacts with endosperm-specific gene promoters. *Plant J.* **13**, 629–640 (1998).
13. Boudet, J. et al. The bZIP transcription factor SPA Heterodimerizing Protein represses glutenin synthesis in *Triticum aestivum*. *Plant J.* <https://doi.org/10.1111/tpj.14163> (2019).
14. Diaz, I., Martinez, M., Isabel-LaMoneda, I., Rubio-Somoza, I. & Carbonero, P. The DOF protein, SAD, interacts with GAMYB in plant nuclei and activates transcription of endosperm-specific genes during barley seed development. *Plant J.* **42**, 652–662 (2005).
15. Vicente-Carbajosa, J., Moose, S. P., Parsons, R. L. & Schmidt, R. J. A maize zinc-finger protein binds the prolamin box in zein gene promoters and interacts with the basic leucine zipper transcriptional activator Opaque2. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 7685–7690 (1997).
16. Merlino, M. et al. Wheat DOF transcription factor TaSAD regulates glutenin gene expression with WPBF and in cooperation with SPA. *PLoS ONE* **18**, 6 (2023).
17. Diaz, I. et al. The GAMYB protein from barley interacts with the DOF transcription factor BPBF and activates endosperm-specific genes during seed development. *Plant J.* **29**, 453–464 (2002).
18. Rubio-Somoza, I., Martinez, M., Abraham, Z., Diaz, I. & Carbonero, P. Ternary complex formation between HvMYB3 and other factors involved in transcriptional control in barley seeds. *Plant J.* **47**, 269–281 (2006).
19. Rubio-Somoza, I., Martinez, M., Diaz, I. & Carbonero, P. HvMCB1, a R1MYB transcription factor from barley with antagonistic regulatory functions during seed development and germination. *Plant J.* **45**, 17–30 (2006).
20. Moreno-Risueno, M. A. et al. FUSCA3 from barley unveils a common transcriptional regulation of seed-specific genes between cereals and Arabidopsis. *Plant J.* **53**, 882–894 (2008).
21. Sun, F. et al. Functional characterization of TaFUSCA3, a B3-superfamily transcription factor gene in the wheat. *Front. Plant Sci.* **8**, 141 (2017).
22. Li, J. et al. TaNAC100 acts as an integrator of seed protein and starch synthesis exerting pleiotropic effects on agronomic traits in wheat. *Plant J.* **108**, 829–840 (2021).
23. Luo, G. et al. The MYB family transcription factor TuODORANT1 from *Triticum urartu* and the homolog TaODORANT1 from *Triticum aestivum* inhibit seed storage protein synthesis in wheat. *Plant Biotechnol. J.* **19**, 1863–1877 (2021).
24. Shen, L. et al. A novel NAC family transcription factor SPR suppresses seed storage protein synthesis in wheat. *Plant Biotechnol. J.* **19**, 992–1007 (2021).
25. Gao, Y. et al. The endosperm-specific transcription factor TaNAC019 regulates glutenin and starch accumulation and its elite allele improves wheat grain quality. *Plant Cell* **33**, 603–622 (2021).

26. Ravel, C. *et al.* Nucleotide polymorphism in the wheat transcriptional activator *spa* influences its pattern of expression and has pleiotropic effects on grain protein composition, dough viscoelasticity, and grain hardness. *Plant Physiol.* **151**, 2133–2144 (2009).
27. Manicacci, D. *et al.* Epistatic Interactions between Opaque2 transcriptional activator and its target gene *CyPPDK1* control kernel trait variation in maize. *Plant Physiol.* **150**, 506–520 (2009).
28. Hunter, B. G. *et al.* Maize opaque endosperm mutations create extensive changes in patterns of gene expression. *Plant Cell* **14**, 2591–2612 (2002).
29. Kawakatsu, T., Yamamoto, M. P., Touno, S. M., Yasuda, H. & Takaiwa, F. Compensation and interaction between *RISBZ1* and *RPBF* during grain filling in rice. *Plant J. Cell Mol. Biol.* **59**, 908–920 (2009).
30. Yamamoto, M. P., Onodera, Y., Touno, S. M. & Takaiwa, F. Synergism between *RPBF* *Dof* and *RISBZ1* *bZIP* activators in the regulation of rice seed expression genes. *Plant Physiol.* **141**, 1694–1707 (2006).
31. Guo, D. *et al.* Over-expressing *TaSPA-B* reduces prolamin and starch accumulation in wheat (*Triticum aestivum* L.) grains. *Int. J. Mol. Sci.* **21**, 3257 (2020).
32. Tabe, L., Hagan, N. & Higgins, T. J. V. Plasticity of seed protein composition in response to nitrogen and sulfur availability. *Curr. Opin. Plant Biol.* **5**, 212–217 (2002).
33. Shewry, P. R., Tatham, A. S. & Halford, N. G. Nutritional control of storage protein synthesis in developing grain of wheat and barley. *Plant Growth Regul.* **34**, 105–111 (2001).
34. Dai, Z. *et al.* Transcriptional and metabolic alternations rebalance wheat grain storage protein accumulation under variable nitrogen and sulfur supply. *Plant J.* **83**, 326–343 (2015).
35. Martre, P., Porter, J. R., Jamieson, P. D. & Triboui, E. Modeling grain nitrogen accumulation and protein composition to understand the Sink/Source regulations of nitrogen remobilization for wheat. *Plant Physiol.* **133**, 1959–1967 (2003).
36. Triboui, E., Martre, P. & Triboui-Blondel, A. M. Environmentally-induced changes in protein composition in developing grains of wheat are related to changes in total protein content. *J. Exp. Bot.* **54**, 1731–1742 (2003).
37. Colot, V., Robert, L., Kavanagh, T., Bevan, M. & Thompson, R. Localization of sequences in wheat endosperm protein genes which confer tissue-specific expression in tobacco. *Embo J.* **6**, 3559–3564 (1987).
38. Muller, M. & Knudsen, S. The nitrogen response of a barley *C-Hordein* promoter is controlled by positive and negative regulation of the *Gcn4* and endosperm box. *Plant J.* **4**, 343–355 (1993).
39. Risacher, T., Craze, M., Bowden, S., Paul, W. & Barsby, T. Highly efficient agrobacterium-mediated transformation of wheat via in planta inoculation. In *Methods in Molecular Biology*, vol. 478 (eds. Jones, H. D. & Shewry, P. R.), 115–124 (Springer, 2009).
40. Mosse, J., Huet, J. & Baudet, J. The amino-acid composition of wheat-grain as a function of nitrogen-content. *J. Cereal Sci.* **3**, 115–130 (1985).
41. Plessis, A., Ravel, C., Bordes, J., Balfourier, F. & Martre, P. Association study of wheat grain protein composition reveals that gliadin and glutenin composition are trans-regulated by different chromosome regions. *J. Exp. Bot.* **64**, 3627–3644 (2013).
42. Fu, B. X. & Kovacs, M. I. P. Rapid single-step procedure for isolating total glutenin proteins of wheat flour. *J. Cereal Sci.* **29**, 113–116 (1999).
43. Wieser, H., Antes, S. & Seilmeier, W. Quantitative determination of gluten protein types in wheat flour by reversed-phase high-performance liquid chromatography. *Cereal Chem.* **75**, 644–650 (1998).
44. Pfaffl, M. W., Tichopad, A., Prgomet, C. & Neuvians, T. P. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper–Excel-based tool using pair-wise correlations. *Biotechnol. Lett.* **26**, 509–515 (2004).
45. Charette, S. J., Lambert, H., Nadeau, P. J. & Landry, J. Protein quantification by chemiluminescent Western blotting: Elimination of the antibody factor by dilution series and calibration curve. *J. Immunol. Methods* **353**, 148–150 (2010).
46. R Core Team. R: A language and environment for statistical computing. <http://www.R-project.org/>, <http://www.gbif.org/resources/2585> (2015).
47. Warton, D. I., Duursma, R. A., Falster, D. S. & Taskinen, S. *smatr* 3-an R package for estimation and inference about allometric lines. *Methods Ecol. Evol.* **3**, 257–259 (2012).
48. Norre, F. *et al.* Powerful effect of an atypical bifactorial endosperm box from wheat *HMWG-Dx5* promoter in maize endosperm. *Plant Mol. Biol.* **50**, 699–712 (2002).
49. Conlan, R. S., Hammond-Kosack, M. & Bevan, M. Transcription activation mediated by the *bZIP* factor *SPA* on the endosperm box is modulated by *ESBF-1* in vitro. *Plant J.* **19**, 173–181 (1999).
50. Juhasz, A., Makai, S., Sebestyen, E., Tamas, L. & Balazs, E. Role of conserved non-coding regulatory elements in LMW glutenin gene expression. *Plos One* **6**, e29501 (2011).
51. Triboui, E. & Triboui-Blondel, A. M. Productivity and grain or seed composition: A new approach to an old problem—invited paper. *Eur. J. Agron.* **16**, 163–186 (2002).
52. Gaju, O. *et al.* Nitrogen partitioning and remobilization in relation to leaf senescence, grain yield and grain nitrogen concentration in wheat cultivars. *Field Crops Res.* **155**, 213–223 (2014).
53. Shewry, P. R. *et al.* Improving the end use properties of wheat by manipulating the grain protein composition. *Euphytica* **119**, 45–48 (2001).

Acknowledgements

The authors are grateful to David Cormier, David Alvarez, Virginie Jonval and Sibille Perrochon for technical support.

Author contributions

T.R. and F.T. created and selected the transgenic lines. P.M., C.R. and A.P. designed the experiment. A.P. supervised the cultivation of the plants and harvested grain samples. A.P. and N.D. analysed grain composition. A.P. and M.D. carried the RNA extraction and measured gene expression. M.M. extracted grain proteins, performed and analysed the western blot. A.P. and P.M. analysed the data. A.P., P.M. and C.R. wrote the manuscript. All authors commented on the manuscript.

Funding

Funding provided by the Fonds Unique Interministériel (grant no. 08 2 90 6227-EJ:0001547).

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-49139-5>.

Correspondence and requests for materials should be addressed to C.R.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023