



Optimization of phosphorus availability in soil: impact of Phosphate-Solubilizing Microorganisms on rapeseed growth and nutrient use efficiency

Charlotte Amy, Mélanie Bressan, Karine Laval, Jean-Christophe Avice

► To cite this version:

Charlotte Amy, Mélanie Bressan, Karine Laval, Jean-Christophe Avice. Optimization of phosphorus availability in soil: impact of Phosphate-Solubilizing Microorganisms on rapeseed growth and nutrient use efficiency. 6th International Symposium Phosphorus in Soils and Plants, Sep 2018, Leuven, Belgium. hal-04374569

HAL Id: hal-04374569

<https://hal.science/hal-04374569>

Submitted on 5 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimization of phosphorus availability in soil: impact of Phosphate-Solubilizing Microorganisms on rapeseed growth and nutrient use efficiency

Charlotte AMY^{1,2}, Mélanie BRESSAN², Karine LAVAL², Jean-Christophe AVICE¹

¹ UMR INRA/UCN 950 Ecophysiologie Végétale et Agronomie (EVA), Université de Caen Normandie, F-14032 Caen, France
² UniLaSalle, unité de recherche AGHYLE, UP 2018.C101, 4277, 3 Rue du Tronquet - CS 40118, F-76134 Mont-Saint-Aignan, France

Context and Hypotheses :

Available Phosphorus (P) is a scarce element in soils for plants and micro-organisms growth due to its quick complexation in soils to organomineral particles such as clay, Iron or Aluminium^[1]. P-Solubilizing Microorganisms (PSM) are able to solubilize complexed P. To face nutrient limitations, plants and micro-organisms interact through rhizodeposition for instance to enhance availability of limiting elements^[2]. Rapeseed (*Brassica napus* L.) is a major oleoproteaginous crop which requires high Nitrogen (N) ^[3] and P inputs and presents a low N and P use efficiencies. Furthermore, little is known concerning rapeseed interactions with soil micro-organisms in relation with P and N availabilities in soils.

Our hypotheses are : i) Optimisation of P nutrition in rapeseed can enhance P and N use efficiency, ii) Rapeseed can recruit PSM through rhizodeposition iii) Changes in P availability in soils can modify N cycle dynamics.


Nutrients required for a yield of 35 q/ha^[4] :

	Exported	Restituted	Total
K ₂ O	35	315	350
N	110	134	244
CaO	17	157	174
MgO	14	73	87
SO ₃	72	143	215
P ₂ O ₅	47	41	88

Unit : kg / ha

Objectives, experimental design and preliminary results

Objective 1: Interactions of N x P nutrition in rapeseed



Preliminary results :





Figure 1 : P content in organs of rapeseed at vegetative stage in response to various P treatments. Plants were grown on Hoagland (3,75mM NO₃) solution. P was measured by X-Ray Fluorescence (XRF)

Objective 2: Crop effect on potential beneficial soil microorganisms



Preliminary results :


- 299 strains were isolated from soils under winter pea (*Pisum sativum*), rapeseed (*Brassica napus*) and **faba bean** (*Vicia faba*)
- 166 strains have shown a positive solubilization index
- Positive strains have been tested on **(for?)** auxin (IAA) production

Table 1 : Isolated strains presenting positive solubilization index and potential **other** growth promoting properties **as** IAA **or** **siderophore** production.

PVK = Pikovskaya medium, NBRIP = NBRIP medium. Solubilization index = solubilization halo size / colony size

Crop	Strain	PVK solubilization index	NBRIP solubilization index	IAA production (ng/μL)	siderophore production
<i>Pisum sativum</i>	P21	1,176	1,297	208,400	+
	Pc12	1,000	1,698	97,200	+
<i>Vicia faba</i>	F46	3,674	1,600	76,300	+
	Fc30	1,368	1,314	99,700	-
	Fc48	1,210	1,000	147,600	-
	C17	1,000	1,401	297,126	NA
<i>Brassica napus</i>	C32	1,824	2,540	211,600	NA
	Cc15	1,468	1,885	122,900	NA
	Cc21	1,358	2,711	113,700	NA

Objective 3: Effect of PSM inoculum on rapeseed growth and N/P cycles



Perspectives :

- Quantitative P solubilization and organic acids production assays will be carried out in order to better discriminate PSM under each crop
- Other analyses will be done to test **other growth promotion properties of the isolated strains** : **HCN production**, **ACC deaminase activity**, **germination test**
- Interesting microorganisms will be **then** identified by **DNA sequencing**