

Predicting tiger mosquito breeding sites through earth observation and statistical analysis

Thibault Catry, Annelise Tran, Ophélie Hoarau, Benjamin Pillot, Nausicaa Habchi-Hanriot, Claire Teillet

► To cite this version:

Thibault Catry, Annelise Tran, Ophélie Hoarau, Benjamin Pillot, Nausicaa Habchi-Hanriot, et al.. Predicting tiger mosquito breeding sites through earth observation and statistical analysis. GEOMED 2022, Oct 2022, Irvine (CA), United States. hal-04380501

HAL Id: hal-04380501 https://hal.science/hal-04380501

Submitted on 13 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Predicting Tiger mosquito breeding sites through Earth Observation and statistical analysis

Thibault Catry¹, Annelise Tran², Ophélie Hoarau², Benjamin Pillot¹, Nausicaa Habchi-Hanriot³ and Claire Teillet¹

¹ ESPACE-DEV, Univ Montpellier, IRD, Univ Antilles, Univ Guyane, Univ Réunion, Montpellier, France ² UMR TETIS, Univ Montpellier, AgroParisTech, CIRAD, CNRS, INRAE, Montpellier, France

3 Regional Health Agency, Reunion Island, France



Espace DEV

DBSERVATION SPATIALE, MODÈL & SCIENCE IMPLIQUÉE

Introduction

Aedes albopictus is the main vector of arboviroses like dengue and chikungunya in Reunion Island. Recently, vector control services have been collecting in-situ data on potentiel breeding sites covering about 10% of the districts of the island.

This study investigates the capability of environmental features extracted from remote sensing data to be used in the prediction of the distribution of potential breeding sites of Ae. albopictus at **Reunion Island.**

A 3 steps approach:

Extract relevant landscape indicators of the urban environment from very high spatial resolution imagery



Identify correlations between landscape indicators, climate variables and breeding site data




Predict spatially the number of breeding sites over the entire territory of Reunion Island







In situ Data



Environmental, climatic and remotely-sensed data





July 2020 Mosaique of Pleaides images

Very High Spatial Resolution image 0.5 m resolution


In situ data on types and number of breeding sites were collected between January 2020 and March 2022 over 10 % of the urban territory of Reunion Island by the Regional Health Agency (ARS).

Spectral indices were calculated from a very-high resolution Pléaides mosaic from 2020 over the island, while data on the annual cumulative rainfall and the annual average temperature were obtained from the Météo-France / CIRAD network (aware.cirad.fr). Elevation and built up areas come from the BDALTI and BDTOPO

Methods


For each district sampled in 2020-2022, we analyzed the correlations between :

Textural information over urban areas was extracted from the Pléaides mosaic, using the FOTOTEX Method (Teillet et al., 2021).





the number of potential observed breeding sites

- ii. the mean values of textural and spectral indices, as well as the mean precipitation and annual temperature, through both univariate and multivariate analysis
- \rightarrow The most significant variables were eventually combined using a generalized linear model.









Positive correlation with NDVI, temperature, % built up and vegetation, and texture **Negative correlation** NDWI, Brightness Index and rainfall Using the best model identified by the GLM and AIC, we predicted the number of breeding over the urban areas of **Reunion Island**

Conclusions

- Textural indices highlight different district typologies based on spatial arrangements between buildings and vegetation.
- Using a GLM for statistical analysis, we found that the correlation between combined environmental variables and in situ data on breeding sites of Ae. albopictus is significant.
- This statistical relationship makes it possible to use earth observation environmental variables as predictors of the number of potential breeding sites for all areas in the island.

Acknowledgments

project ANISETTE This funded work was by the (anisette.cirad.fr), a project supported by the French Space Agency, CNES (DAR 4800001029), under the platform in partnership One Health Indian Ocean (<u>www.onehealth-oi.org</u>). The authors thank Thierry Baldet (medical entomologist, Cirad Astre) for fruitful discussions