GeoOneHealth Symposium

Remote sensing analysis of the links between urban landscapes and the risk of exposure to *Aedes* mosquitoes, vectors of arboviruses. Claire Teillet

Directors : Rodolphe Devillers and Emmanuel Roux

claire.teillet@ird.fr

Research Institute for development, UMR ESPACE-DEV, Montpellier Funded by CNES & Région Occitanie













PhD Context

Regional Cooperation Project for the Observation of the Guyana Shield by SATellite (PROGYSAT)





Health: Malaria and mosquito-borne arboviruses - Emmanuel Roux (IRD), Margarete Gomes (SVS-AP)



Urban Axis: Analysis of Urban Space Dynamics by Satellite - Nadine Dessay (IRD), Gutemberg Silva (UNIFAP), Paulo Peiter (Fiocruz)





PhD labelled by RIVOC in MUSE program which deals with the sustainable management of vectors risk as an issue for global health in France



Project financed by the French Spatial Agency to work on several methods combining remote sensing and spatial modelling to predict the dynamics of mosquito vectors and associated diseases.

Annelise Tran, CIRAD, UMR Tetis

Context

ARBOVIRUSES:

viral diseases caused by a virus transmitted by an arthropod vector (focus on *Aedes* mosquitoes)

- Increasing risk to global health
- Distribution of vectors spatial boundaries of transmission of this diseases
- Great diversity of factors affecting the distribution of *Aedes* vectors at different scales



Context

- Spatial modelling of vector-borne diseases
 - $\circ~$ based on a number of heterogeneous data (entomological, epidemiological, environmental, socio-economic) and at different scales
- Potential of remote sensing (proxy extraction, diversity of data, genericity of methods)



Risk map identifying habitats suitable for *Aedes albopictus* provinces of Iran. Nejati et al., 2017

Risk map identifying habitats suitable for *Aedes aegypti*, in Cordoba City, Agentina. Estallo et al., 2018

Scientific question and objective



Limited access to epidemiological and entomological data



Land cover / spectral indices vs. Landscape structure



Need for simple tools for spatialization of risks



Unsuitable global products for city scale applications, especially for health issues

Objective

To develop an approach to spatialize the risk of exposure to Aedes mosquitoes that makes the best use of satellite data and available data (entomological, epidemiological, etc.) in order to make it reproducible, generic and adapted to the needs of health actors.

Two contrasting study sites



Equatorial Climate Regular outbreaks Aedes aegypti

No outbreak, imported and autochthonous cases detected and controlled Crisis risk area Aedes albopictus

Public health partners



Collaboration with public health partners to identify needs in terms of cartography and tools

- Better identify risks between outbreaks and at a finer scale
- Target priority areas so that their efforts and resources can be directed according to the need.



Differents scale of analysis (Global, Regional, City, Neighborhoods...)

 \rightarrow Focus on the city and neighborhoods scales



Project axis

Axis 1 : Urban variables based on satellite imagery Axis 2 : Breeding sites and mosquito density modeling Axis 3 : Degree of populations exposure to mosquitoes Axis 4 : Spatialized index of population exposure to Aedes vectors



Axis 1: Urban variables from satellite imagery to characterise urban landscapes in relation to *Aedes* mosquitoes

Objectives:

- Identify factors that influence the presence of breeding sites and *Aedes* mosquitoes
- Identify which urban variables are related to this factors and how to extract them from thanks to several satellite imagery





Life cycle of mosquitoes (illustration for *Aedes albopictus*) Marti, Teillet et al., 2022



Axis 1: Urban variables from satellite imagery to characterise urban landscapes in relation to Aedes mosquitoes

• Review focus on articles that use remote sensing and geomatics, which variables are used for modeling *Aedes* distribution ?





Axis 1: Urban variables from satellite imagery to characterise urban landscapes in relation to Aedes mosquitoes



- Imagery allows to study urban landscapes at different scale
- Majority of sensors allows to work at the city scale but much less at neighborhoods scale which can be interesting for health studies
- Climatic variables is at low spatial resolution no matter the scale of analysis, other variables depends on the scale and resolution

Spectral indices Land cover Temperature Precipitations Digital elevation model Land use Landscapes metrics Texture index



- Different types of spatial modelling of mosquito densities:
 - "Mechanistic" differential equation model based on the bio-ecology of the *Aedes* vector (Tran et al., 2013) ARBOCARTO - tool CIRAD & French Health Ministry
- Input of this model :



→ Objective: To refine and improve the estimation of the number of potential breeding sites based on remote sensing information



Focus on characterization on urban landscape :

- Texture analysis with **FOTOTEX** algorithm (<u>Teillet et al., 2021</u>)
- Allows to improve knowledge of landscape by analysing the texture of differents images at differents scales





(b)

What is a texture in image processing ?

(a) Texture can be defined as a function of spatial variation of the brightness intensity of the pixels.

In other terms : arrangement, disposition of elements in relation to each other create a pattern that you can see with your eyes



(c)



• Principle of the FOTOTEX method



A- Partitioning the image into analysis windows block or sliding window

B- Fourier transform analysis

Decomposition of the signal linked to the repeating patterns as a sum of sinusoidal functions (texture-to-frequency conversion)

C- PCA on the spatial frequency matrix each window is characterised by a large number of frequency variables (repetition of patterns) which are reduced

D- RGB colour composition

Spatial representation of the distribution of the frequencies that make up the initial image R=PC1, G=PC2, B=PC3



- Modelling breeding sites with the help of remote sensing
 - Statistical model developed in Reunion (*Aedes albopictus*, specific spatial delimitation)





- Modelling breeding sites with the help of remote sensing
 - \circ Statistical model developed in Reunion (Aedes albopictus, specific spatial delimitation)



Correlation between predicted values and observed values

Estimation du nombre de gîtes prédits dans chaque zone ALIZES



SCR : RGR 92 / UTM zone 40S (EPSG : 2975) Données : ARS et BD topo de l'IGN



- Application of the statistical model in Cayenne:
 - \circ value of breeding sites very high
 - \circ heterogeneous areas especially close to dense urban areas



N





• Exploratory approach to estimate the number of potential breeding sites over study sites



Axis 3: Spatialize the areas where the population is most exposed to mosquitoes using urban variables from remote sensing and spatialized information

stribution and population density cors in relationship to population exposure have been found in literature



Atlas des tissus urbains, AUDEG, 2019

 \rightarrow Objective : Integrate more information on population exposure into modelling risk

Axis 3: Spatialize the areas where the population is most exposed to mosquitoes using urban variables from remote sensing and spatialized information

Internship of Nicholas de Kock (3 months)

• Creating socio-economic classes from remote sensing information combined with OSM data and global data (GHSL)







- Spatialized risk indicator
 - Combine predictions on mosquitoes and the degree of exposure of populations
 - Apply all the approach and validate on Cayenne and Montpellier
 - Apply to other study sites to test the reproducibility, automatic and generality of the entire approach





TÉLÉDÉTECTION ET MODÉLISATION SPATIALE

Applications à la surveillance et au contrôle des maladies liées aux moustiques







https://www.quae.com/produit/1784/97827592 36299/teledetection-et-modelisation-spatiale