



HAL
open science

How does trait variance partitioning help us to understand plant community assembly? The example of pond communities in the Kerguelen Islands

Pauline Douce, D Renault, Laurent Simon, Florian Mermillod-Blondin, Felix Vallier, Anne-kristel Bittebière

► To cite this version:

Pauline Douce, D Renault, Laurent Simon, Florian Mermillod-Blondin, Felix Vallier, et al.. How does trait variance partitioning help us to understand plant community assembly? The example of pond communities in the Kerguelen Islands. *Journal of Vegetation Science*, 2023, 34 (6), pp.e13217. 10.1111/jvs.13217 . hal-04382947

HAL Id: hal-04382947

<https://hal.science/hal-04382947>

Submitted on 7 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **How does trait variance partitioning help us to understand plant community**
2 **assembly? The example of pond communities in the Kerguelen Islands**

3
4 We focused on the macrophyte communities living in the ponds of the Iles Kerguelen
5 (sub-Antarctic region). Partitioning trait variance allows to detect at which scale operate
6 the decisive processes in plant community assembly. In addition, we simultaneously
7 considered biotic and abiotic variables to better understand the effects of
8 environmental variations on traits, subsequently impacting plant performance and
9 species fate.

10

11 Pauline Douce^{1*}, David Renault^{2,3}, Laurent Simon¹, Florian Mermillod-Blondin¹, Felix Vallier¹,
12 Anne-Kristel Bittebiere¹

13

14 *Corresponding author: pauline.douce@univ-lyon1.fr

15 L. Simon: laurent.simon@univ-lyon1.fr

16 F. Mermillod-Blondin: [florian.mermillod-blondin@univ-lyon1](mailto:florian.mermillod-blondin@univ-lyon1.fr)

17 D. Renault: david.renault@univ-rennes1.fr

18 F. Vallier: felix.vallier@univ-lyon1.fr

19 A.-K. Bittebiere: anne-kristel.bittebiere@univ-lyon1.fr

20

21 ¹Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR5023 LEHNA, F-69622,
22 Villeurbanne, France

23 ²Univ Rennes, CNRS, ECOBIO [(Ecosystèmes, biodiversité, évolution)], - UMR 6553, F 35000
24 Rennes, France

25 ³Institut Universitaire de France, 1 Rue Descartes, 75231 Paris cedex 05, France

26

27 **Funding information**

28 This study was funded by the French National Research Agency, 'PONDS' project (ANR-21-
29 CE02-0003-01, JCJC call 2020–2021), the BiodivERsA 'ASICS' project (ANR-20-EBI5-0004,
30 BiodivClim call 2019–2020), the Paul-Emile Victor French Polar Institute (IPEV 136
31 SUBANTECO project), and the long-term research network on biodiversity in Antarctic and
32 sub-Antarctic ecosystems (Zone Atelier InEE-CNRS Antarctique et Terres Australes).

33 **Abstract**

34 Question: in the current context of biodiversity erosion, functional approaches to the
35 examination of community assembly mechanisms and better prediction of plant species fates
36 have emerged. The assessment of trait variation patterns should be a powerful means of
37 identifying community assembly mechanisms. However, most studies of trait variations and
38 their consequences for individual performance (*i.e.*, vegetative biomass) are usually
39 incomplete as they focused on single ecological scales or filters, with no consideration of
40 relationships between traits. Such research has provided a fragmented view of plant
41 community assembly.

42 Location: We examined macrophyte communities living in ponds of the sub-Antarctic Iles
43 Kerguelen.

44 Methods: We measured traits related to resource acquisition and conservation on all
45 occurring species, and examined their variation across temporal (years), spatial (sites), and
46 taxonomic (between- and within-species) scales and in response to multiple abiotic and biotic
47 habitat variables. The consequences of these trait variations and the effects of their correlation
48 for plant individual performance were also explored.

49 Results: Trait distributions were fairly conserved among sites, whereas we observed a
50 large amount of intraspecific trait variation enabling individuals to resist filters. Responses to
51 biotic and/or abiotic variables were trait dependent, and simultaneous trait responses should
52 enable individual plants to face several simultaneous constraints. Almost all traits had direct or
53 indirect effects on individual performance, indicating the need to consider trait relationships.

54 Conclusion: The partitioning of trait variance is a relevant approach to the identification of
55 the scale at which the most decisive processes for plant community assembly operate without
56 the interference of scale dependency issues, and should orient further research. In addition,
57 several biotic and abiotic variables should be considered in future studies to better understand
58 the effects of environmental changes on plant communities.

59 **Keywords**

60 Functional trait, plant performance, variance partitioning, intraspecific trait variability,
61 macrophyte, plant community assembly, filter, pond ecosystem

62 **Introduction**

63 In the current context of biodiversity erosion, it is crucial to improve our understanding of the
64 mechanisms underlying plant community dynamics for the prediction of plant species fates,
65 particularly in aquatic ecosystems, which are especially sensitive to ongoing changes ([Pan et
66 al., 2023](#)). Two decades ago, functional approaches emerged that allowed us to rebuild
67 community ecology and establish general principles regarding assembly mechanisms ([Calow,
68 1987](#); [McGill et al., 2006](#)). Since then, studies based on functional traits (morphological,
69 physiological, and phenological characteristics that directly or indirectly impact individual
70 performance through growth, reproduction, or survival) have been conducted in terrestrial
71 (e.g., [Garnier et al., 2001](#); [Albert, Thuiller, Yoccoz, Douzet, et al., 2010](#); [Violle et al., 2007](#)),
72 and aquatic plant communities ([Chmara et al., 2021](#); [Fu et al., 2013, 2014](#); [Luo et al., 2016](#);
73 [Ma et al., 2022](#); [Su et al., 2019](#); [Yang et al., 2004](#); [Zhang et al., 2019](#)). Functional approaches
74 have revealed that marked trait variation occurs in the field at inter- and intraspecific levels in
75 response to a range of biological processes operating at different spatial and temporal scales,
76 ultimately leading to species coexistence ([Albert, Thuiller, Yoccoz, Douzet, et al., 2010](#);
77 [Messier et al., 2010](#); [Taudiere & Violle, 2016](#)).

78 Individual trait variation is usually observed independently through the lens of space,
79 time, or taxonomy, resulting in a fragmented view of community assembly, with the exception
80 of a few studies in which two of these scales have been considered ([Fu et al., 2013](#); [Fyllas et
81 al., 2020](#); [Johnson & Toprak, 2021](#); [Kim & Nishihiro, 2020](#)). Temporal trait variation is reported
82 as a part of a seasonal strategy allowing plants to permanently adjust trait performance to deal
83 with external stresses; examples are the adjustment of physiological (e.g., photosynthetic
84 capacity; [Grassi et al., 2005](#); [Muraoka et al., 2010](#)) and morphological (e.g., leaf thickness;
85 [Fullana-Pericas et al., 2017](#)) traits. Spatial scales also shape overall trait variations, which can
86 arise as species-specific responses to various environmental filtering processes. In
87 macrophytes species for example, shoot height and branch length have been found to vary
88 primarily among lakes (in connection with trophic status and water depth), whereas the
89 root/shoot ratio and leaf number are linked to within-lake variations ([Fu et al., 2013](#)).The

90 decomposition of trait variations across taxonomic scales (McGill, 2008; Swenson et al., 2006)
91 allows for the consideration of inter- vs. intraspecific trait variations to determine whether
92 species identity is a critical filtering factor, or whether individuals are filtered out according to
93 their trait values. Overall, studies providing information on the scale that carries the most
94 important trait variations are of particular interest, as the results should aid the understanding
95 of many community assembly patterns and processes (McGill, 2008; Taudiere & Violle, 2016;
96 Violle et al., 2007, 2012) and the assessment of the scale dependency issue in community
97 ecology (e.g., Cavender-Bares et al., 2009; Swenson et al., 2006).

98 In the last 10 years, studies reported the major role of intraspecific trait variability (ITV)
99 in ecological processes, and showed that ITV can account for up to 40% of trait variance in
100 plant communities (Albert, Thuiller, Yoccoz, Soudant, et al., 2010; Jung et al., 2010). ITV would
101 allow a better resistance to filters (Jung et al., 2010; Violle et al., 2012) through adaptation
102 (genetic variability) and/or phenotypic plasticity (Albert et al., 2011; Byars et al., 2007;
103 Ghalambor et al., 2007). In aquatic plant communities, the high investment of macrophyte
104 species in clonal growth suggests that resistance to filters mainly relies on plastic responses
105 (De Wilde et al., 2014; Gao et al., 2021; Han et al., 2021; Li et al., 2019; Pilon & Santamaría,
106 2002; Wolfer & Straile, 2004). Most plant morphological traits are key for coping with
107 environmental filters (see, e.g., Lavorel & Garnier, 2002). Responses to both biotic and abiotic
108 filters have been evidenced extensively in macrophyte traits involved in growth or resource
109 acquisition [*i.e.*, the specific leaf area (SLA), height, and root and internode lengths; Fu et al.,
110 2013; Gao et al., 2021; Pilon & Santamaría, 2002; Wolfer & Straile, 2004] and in resource
111 conservation [*i.e.*, the leaf dry matter content (LDMC) and specific connection mass; Elger &
112 Willby, 2003; Gao et al., 2021; Koleszár et al., 2022; Pilon & Santamaría, 2002]. How traits
113 respond to simultaneously varying biotic and abiotic conditions, however, remains unclear
114 (Chalmandrier et al., 2022). This may lead to the partial interpretation or even misinterpretation
115 of trait variation patterns, especially in *in-situ* studies with such simultaneous variation (Cadotte
116 & Tucker, 2017; Kraft et al., 2015). Indeed, similar patterns of trait variations (convergence or
117 divergence in trait values) can be related to abiotic and/or biotic conditions (Cadotte & Tucker,

118 2017). We thus argue that the partitioning of intraspecific trait variations should aid the
119 determination of the roles of traits in species' biotic and abiotic resistance to environmental
120 filters (Le Bagousse-Pinguet et al., 2017; Liu et al., 2018). We expect traits to respond
121 simultaneously to biotic and abiotic conditions, but with varying strength depending on the trait
122 category (resource conservation vs. acquisition). In particular, we expect traits related to
123 resource acquisition to respond to water, sediment nutrients, and biotic conditions, determining
124 competitive abilities (Garnier et al., 2004; Reich, 2014; Violle et al., 2007; Wright et al., 2004),
125 and those related to resource conservation to respond mostly to harsh abiotic conditions,
126 conferring individual plant resistance to environmental constraints (Funk et al., 2021; Levitt,
127 1980; Reich, 2014).

128 Variation in functional traits should ultimately influence individual performance (*i.e.*,
129 vegetative biomass production; Geber & Griffen, 2003), and thus species fate in the plant
130 community (Ghalambor et al., 2007), but with varying intensity depending on the trait.
131 Moreover, several traits co-vary and/or are subjected to tradeoffs in individual plants to
132 coordinate different biological functions (Maire et al., 2013; Valladares et al., 2007; Violle et
133 al., 2007). For instance, resource conservation– and acquisition-related traits have been found
134 to negatively co-vary (Fort et al., 2013; Garnier et al., 2001; Kleyer et al., 2019; Reich et al.,
135 1997, 1999); low SLAs are generally associated with high LDMCs, reflecting a functional
136 relationship with the leaf life span (Roche et al., 2004; Wright et al., 2004). However, such trait
137 relationships, which can indirectly influence individual performance, are usually neglected in
138 relevant studies. The identification of the direct and indirect influences of traits on performance
139 likely imposes hierarchies among plant traits, which can be revealed with structural equation
140 modeling [SEM; Ackerly et al., 2000; Dwyer & Laughlin, 2017; Saiz et al., 2018; Vile et al.,
141 2006]. We expect resource conservation–related traits to have more direct impacts on
142 individual biomass than do resource acquisition–related traits, as they are related directly to
143 individual size and thus affect individual performance.

144 A growing number of theories and concepts in ecology has been tested recently in the
145 sub-Antarctic Iles Kerguelen region, considered to contain sentinel ecosystems and open-air

146 laboratories (Bergstrom & Chown, 1999). Climate change, particularly warming, is especially
147 rapid in this region (Walther et al., 2002) and has strong impacts on aquatic ecosystems,
148 including ponds that harbor remarkably species-poor macrophyte communities (Douce et al.,
149 2023a). In this study, we used macrophyte communities to construct a simple model that is
150 more realistic than the use of mesocosms (which include all variations in abiotic and biotic
151 conditions). We measured traits related to resource acquisition and conservation and
152 performance in all macrophyte species. Trait variations were studied across spatial, temporal,
153 and taxonomic scales, and covered multiple abiotic and biotic habitat conditions.
154 Consequences for individual plant performance were also explored. We tested the following
155 hypotheses (Fig. 1):

- 156 i. that the partitioning of trait variance among spatial, temporal, and taxonomic scales
157 would enable the detection of mechanisms involved in plant community assembly,
158 revealing a large amount of ITV related to coexistence mechanisms and within-site
159 heterogeneity;
- 160 ii. that all traits would respond to both biotic and abiotic conditions, but with trait-
161 dependent intensity, with traits related to resource acquisition (SLA, height, and root
162 and internode length) responding to water, sediment nutrients, and biotic conditions
163 and those related to resource conservation (LDMC and specific root and internode
164 masses) responding mostly to harsh abiotic conditions; and
- 165 iii. that traits related to resource conservation would affect individual vegetative biomass
166 more directly than those related to resource acquisition, with the potential modulation
167 of relationships between traits, *i.e.*, negative co-variation of resource acquisition– and
168 conservation-related traits, modifying their role in individual performance.

169 **Materials and methods**

170 **Study model**

171 The study was carried out in the field at the French sub-Antarctic Iles Kerguelen (South Indian
172 Ocean; 48°30–50°S, 68°27–70°35E). The climate on this archipelago is characterized by cold

173 temperatures (annual mean of 4.6°C with little variation; [Frenot et al., 2006](#); [Lebouvier et al.](#)
174 [2011](#)) and abundant precipitation (500–3200 mm along an east–west gradient; [Frenot et al.,](#)
175 [1998](#); [Meteo France](#)). Plant communities in polar regions are paucispecific (and thus have
176 simplified plant–plant interaction) and subjected to highly reduced anthropogenically induced
177 environmental pressure. As such, they constitute simplified models relative to temperate plant
178 communities, which is especially relevant for the inference of mechanisms of community
179 assembly in constrained abiotic environments ([Bergstrom & Chown, 1999](#)).

180 The study was conducted in ponds at three sites located along the shore of the
181 archipelago’s main island (Appendix S1): Cap Molloy, Isthme Bas, and Cap Ratmanoff. These
182 three sites are less than 40 km apart and have similar climates, and their overall abiotic
183 conditions are more variable locally than among sites (Appendix S2; [Douce et al., 2023a](#)). At
184 each site, 15 ponds of variable areas were selected to overcome bias related to system size,
185 which influences the water temperature ([Bornette & Puijalon, 2011](#)). All but one of the ponds
186 investigated were small (average area = 181.57 ± 382.39 m²; exception, 180,800 m²; [Douce](#)
187 [et al., 2023a](#)). They are shallow freshwater systems enriched by nutrient inputs from marine
188 fauna, such as seabirds and seals, due to their littoral proximity ([Smith, 2008](#)). The ponds’
189 water depth was similar (average of the 45 ponds = 17.29 ± 8.28 cm).

190 The macrophyte communities at the study sites were composed mainly of six native
191 species that can all locally co-occur: *Limosella australis* R.Br. (Scrophulariaceae), *Callitriche*
192 *antarctica* Engelm (Plantaginaceae), *Juncus scheuchzerioides* Gaudich. (Juncaceae),
193 *Ranunculus biternatus* Smith, *Ranunculus pseudotrullifolius* Skottsb., and *Ranunculus*
194 *moseleyi* Hook.f (all Ranunculaceae). All of these species are perennials (*i.e.*, they remain
195 present during winter) with clonal network structures, *i.e.*, the plants are composed of related
196 ramets (shoots with leaves and roots) that are connected by plagiotropic stems. These clonal
197 structures display slow growth rate, and ramet lifespan generally exceed two years with low
198 investment in sexual reproduction. *C. antarctica*, *L. australis*, *R. pseudotrullifolius*, and *R.*
199 *moseleyi* are hydrophytes, and *R. biternatus* and *J. scheuchzerioides* are helophytes. Only *C.*

200 *antarctica* is caulescent and harbors a floating canopy; the other five species have rosette
201 architectures and are rooted in pond sediments (Douce et al. 2023a). As the water level in
202 ponds is highly variable (Douce et al 2023a), individual leaves can be alternately submerged
203 or emerged, without any variations in their morphology. Individuals then display leaves of
204 homogeneous morphology (Appendix S3). These macrophyte communities may experience
205 different episodes of disturbances through the year, as for instance, the freezing of pond
206 surface in winter (although this phenomenon remains limited to the first 5-10 cm of water
207 column), the temporary drying in summer (for less than a month), and the seal trampling, that
208 may altogether lead to the low macrophyte cover in ponds (low community resilience).
209 Nevertheless, the studied communities remain very stable through years and seasons (Douce
210 et al. 2023a).

211

212 Field sampling

213 The study was conducted in 2017, 2018, 2019, and 2020, with plant ramet sampling at the
214 three sites conducted each year in November. In each pond, we randomly positioned three to
215 five 1 × 1-m quadrats, depending on the pond area, and usually including pond edges. In each
216 quadrat, we collected one mature ramet of each of the six studied species, when present, for
217 the measurement of plant traits. One ramet was composed of leaves, roots, and one
218 connection internode for the five rosette species, and a fully developed caulescent structure
219 (with attached leaves and roots and no ramification arising from axillary buds), and one
220 connection internode for the caulescent species. All collected ramets, including those of the
221 helophyte species, were submerged completely in the ponds (mean ramet height = 7.14 ± 2.99
222 cm, mean water depth = 17.29 ± 8.28 cm). Given the low variability in water depth between
223 ponds, the ramets were collected at similar depths. In addition, we recorded species
224 abundances in each quadrat.

225 Pond abiotic conditions were monitored every 3 months from November 2019 to
226 November 2020. For each sampling date and pond, water and sediment samples were hand
227 collected at similar depths for nutrient and phytoplankton concentration analyses. These

228 samples were collected at three points per pond and then mixed together to account for
229 possible heterogeneity in nutrient spatial distribution. One composite water sample (30 mL)
230 and one composite sediment sample (50 mL) were obtained per pond. The sediment was a
231 mixture of sand and organic matter, with similar grain and composition among ponds. As
232 abiotic variables are likely to vary within ponds, in each quadrat at the ramet collection points,
233 we measured the mean water depth (based on three measurements) and mean dissolved
234 oxygen (DO) concentration (based on measurements taken at three water depths). We also
235 measured the pH and electrical conductivity (EC), since they are not related to the
236 phytoplankton concentration in the studied systems (Douce et al., 2023b). These
237 measurements were performed using a multiparameter HQ40D HACH sensor (EC ± 0.1
238 $\text{mS}\cdot\text{cm}^{-1}$, DO concentration $\pm 0.01 \text{ mg}\cdot\text{L}^{-1}$). From September 2020 onward, whole-pond water
239 temperatures were recorded every 30 minutes using loggers (Hobo MX2202) positioned in the
240 middle of the water column. Water temperatures in November 2019–August 2020 were
241 simulated using their relationship to air temperatures recorded by Meteo France (2020 records,
242 Port-aux-Français; see Douce et al., 2023b). Each pond area (length \times width) was measured
243 in November 2019. All of these measured parameters have been found to structure the studied
244 communities (Douce et al., 2023a).

245

246 Trait measurements

247 Seven functional traits of the mature ramet samples were measured, following Cornelissen et
248 al. (2003). The selected traits characterize ramet growth strategies and are known to be
249 responsive to biotic and abiotic conditions (Table 1), especially in macrophyte species (Dalla
250 Vecchia et al., 2020). They are related to individual resource acquisition (height, SLA, and
251 internode and root lengths) and conservation (LDMC and internode and root specific masses;
252 see Table 1 for details).

253 For each sampled ramet, the SLA and LDMC of one randomly selected healthy
254 mature leaf were first measured. Second, the length and specific mass of one randomly
255 selected attached internode were measured. The specific internode mass was calculated as

256 the ratio between the internode dry mass and length ($\text{g}\cdot\text{cm}^{-1}$). Then, the longest primary root
257 attached to the ramet was used to measure the maximum root length and specific root mass
258 [calculated as the ratio between the dry mass and root length ($\text{g}\cdot\text{cm}^{-1}$)]. We also estimated
259 ramet performance resulting from its growth strategy by measuring its total biomass (*i.e.*, dry
260 masses of leaves, stems, roots, and one connection internode; see, *e.g.*, [Adomako et al., 2021](#);
261 [Keser et al., 2014](#); [Younginger et al., 2017](#); [Zhao et al., 2020](#)). Dry mass measurements were
262 performed after the plant parts had been dried for 48 hours in an oven at 65°C . All plants were
263 cleaned before measurement.

264

265 [Measurement of nutrient and phytoplankton concentrations](#)

266 For the measurement of water nutrient and phytoplankton concentrations (as reflected by the
267 amount of chlorophyll a per milliliter water), water samples were passed through GF/F filters
268 ($0.7\ \mu\text{m}$; Whatman) to remove coarse and fine particles and phytoplanktonic organisms within
269 24 hours of sampling. All samples (filtered water and sediments, and filters) were stored at –
270 20°C for several weeks before chemical analysis in metropolitan France.

271 Chlorophyll a pigments were quantified from the GF/F Whatman filters using the
272 Unesco method ([Vohra, 1966](#)) and a spectrophotometer. N-NH_4^+ , N-NO_3^- , and P-PO_4^{3-}
273 concentrations in the water samples were determined using colorimetric methods and a
274 sequential analyzer (SmartChem 200; AMS Alliance) ([Grasshoff et al., 1999](#)). The organic C:N
275 ratio in sediment samples was measured using the capsule method ([Brodie et al., 2011](#)). An
276 aliquot of $5\ \text{mg} \pm 10\%$ homogeneous sediment sample was acidified into $100\ \mu\text{L}$ $2\ \text{M}$ HCl in
277 silver capsules to eliminate carbonates. The capsule liquids were evaporated on a 65°C hot
278 plate for 12 hours, and the capsules were then oven dried at 80°C for 2 days. The total organic
279 C and N contents were then measured with an elementary analyzer (FlashEA 1112 NC
280 Analyzer[®]; Thermo Fisher Scientific, Waltham, MA, USA). The bioavailable P concentration in
281 sediments was determined following the protocol of [Ni et al. \(2016\)](#). Bioavailable P was
282 extracted from $50\ \text{mg}$ dry sediment samples with $5\ \text{mL}$ NaOH ($1\ \text{M}$). Then, the supernatant
283 was collected, its pH was stabilized with HCl ($3.5\ \text{M}$), and the extracted P that had been

284 converted to orthophosphate was quantified using the molybdate/ascorbic acid blue method
285 (Murphy & Riley, 1962); results are reported as milligrams of available P per gram dry
286 sediment.

287

288 Data analysis

289 We tested for correlations between all pairs of traits using Spearman rank correlation analysis.

290 We detected no strong correlation ($r < 0.7$) between traits (Dormann et al., 2013). Thus, each
291 trait was indicative of a part of a global growth strategy.

292

293 *Abiotic and biotic characterization of ponds*

294

295 To characterize the abiotic conditions within ponds, the mean of daily (day and night) water
296 temperatures from November 2019 to November 2020 was calculated for each pond (Douce
297 et al., 2023b). Abiotic variables measured at the quadrat scale were averaged at the pond
298 scale. Depending on the analysis, we performed several reductions of abiotic variables. Water
299 nutrient dimensions (N-NH₄⁺, N-NO₃⁻, and P-PO₄³⁻) were reduced to the first axis of a principal
300 component analysis (PCA), in which data were centered and standardized by standard
301 deviation (Appendix S4), hereafter referred to as “PCAwater.” PCAwater axis 1 (42.06%) was
302 related negatively to the N-NH₄⁺ and N-NO₃⁻ concentrations. Similar dimension reduction was
303 performed for sediment nutrients (C:N ratio and bioavailable P; Appendix S5), hereafter
304 referred to as “PCAsediments.” PCAsediments axis 1 (58.59%) was related negatively to the
305 C:N ratio and positively to the bioavailable P concentration.

306 To characterize the biotic conditions within ponds, we calculated two non-correlated
307 indices: macrophyte species richness and Pielou evenness. Biotic dimensions describing plant
308 species abundances in each pond were reduced to the first three axes of a factorial
309 correspondence analysis (“FCAplants”; Appendix S6). The first FCAplants axis (33.81%) was
310 driven positively by the abundance of *R. moseleyi* and negatively by that of *C. antarctica*. The

311 second axis (24.99%) related positively with the abundance of *R. pseudotrullifolius* and
312 negatively with that of *J. scheuchzerioides*. The third axis (16.99%) was related positively to
313 the abundance of *L. australis* and negatively to that of *R. biternatus*. To assess the functional
314 characteristics of pond communities, we calculated the overall functional dispersion (FDis)
315 using seven traits (height, SLA, LDMC, internode length, specific internode mass, maximum
316 root length, and specific root mass) and the *fdisp* function of the FD package (Laliberté &
317 Legendre, 2010).

318

319 *Variance partitioning across nested scales*

320 We used trait measurements from 2017–2020 to partition trait variance across four nested
321 scales. The trait data were normalized (log or square-root transformed) when needed, and
322 standardized to equalize the weight of each trait dimension. A variance component analysis
323 was performed for each of the seven studied traits using the *Parvart* procedure (Cati package)
324 across temporal (four years), spatial (three sites), and taxonomic (inter- and intraspecific
325 variance) scales introduced into the model (including measurement error) in that order. As
326 individuals do not display any heterophyllia, the intraspecific trait variation only accounts for
327 trait variations between individuals. The significance of the variance components was
328 assessed by determining 95% confidence intervals (CIs) through bootstrapping. We randomly
329 selected 700 of the 1430 individuals in our dataset with replacement (Messier et al., 2010),
330 and partitioned the trait variance among the four nested scales. We repeated this procedure
331 500 times to ultimately calculate a 95% CI for each variance component. To get rid of the
332 imperfect nesting of species within sites (Appendix S7), we ran alternative models of variance
333 partitioning without the species scale, and obtained similar results (Appendix S8). Similarly, as
334 some species were less abundant (*R. moseleyi*, *R. pseudotrullifolius*, and *L. australis*;
335 Appendix S7), we ran alternative tests of variance partitioning without the individual data of
336 these scarce species, to test the robustness of our results. We obtained similar results with
337 and without the inclusion of scarce species data (Appendix S9).

338

339 *Analysis of trait responses to biotic and abiotic filters*

340 We aimed to determine which traits and categories of traits were involved in plant biotic and/or
341 abiotic resistance. We considered univariate responses and tested for the effects of abiotic
342 and biotic variables measured or calculated at the pond scale on trait values measured in
343 November 2020 and averaged at the pond scale for all ramets of all species. In the full models,
344 we included as explanatory variables the species identity (a categorical variable indicating the
345 ramet species), macrophyte species richness, Pielou evenness, community FDis, first three
346 FCAplants axes (Appendix S6), and phytoplankton concentration for biotic variables, and the
347 water depth, pH, EC, DO concentration, mean temperature, pond area, and first PCAwater
348 and PCAsediments axes (Appendices S3 and S4) for abiotic variables. The trait data were
349 normalized (log or square-root transformed) as needed and standardized. We applied stepwise
350 model reduction based on the Akaike information criterion (AIC) using the MuMIn package
351 (Bartoń, 2013) to select the most parsimonious full model, and performed a type II analysis of
352 variance.

353 A similar procedure was applied to test for the effects of the above-listed abiotic and
354 biotic variables on traits grouped into multivariate-response resource acquisition (height, SLA,
355 internode length, and maximum root length), and resource conservation (LDMC, specific root
356 mass, and specific internode mass) categories. The trait data were normalized (log or square-
357 root transformed) as needed and standardized. We selected variables by backward elimination
358 from the full model, removing those with non-significant effects, and then performed a type II
359 multivariate analysis of variance (yielding the Pillai statistic) of the reduced model.

360

361 *Analysis of trait and abiotic variable effects on species performance*

362 The ramet total biomass (reflecting performance) and trait values recorded in November 2020
363 were averaged for each species at the pond level ($n = 97$) and standardized to eliminate

364 variance due to species identity. We tested for direct and indirect effects of the seven traits on
365 individual performance using SEM (Grace et al., 2010). All possible relationships between
366 traits, based on the literature (Appendix S10), were included. We performed a path analysis
367 with the lavaan package (Rosseel, 2012), and reduced the full model by variable selection
368 based on AIC values.

369 All statistical analyses were performed with R 4.0.3.

370

371 **Results**

372 *Traits varied primarily within macrophyte species*

373 All studied traits exhibited larger variances within sites than between sites, since space (site)
374 had no influence at all. Time (year) had little influence (explaining less than 8% of variance in
375 all traits) when significant. Conversely, taxonomy explained most of the variance within sites
376 (Fig. 2). Most traits (with the exception of internode mass and LDMC) exhibited higher trait
377 variance at the intra- as compared with the interspecific scale. The intraspecific variance of the
378 trait was even maximal in the internode length (79.47%) and the maximum root length
379 (65.52%).

380 *The variation of most traits was affected by abiotic and biotic variables*

381 Within ponds, growth strategies relied on resource acquisition and conservation traits.
382 Resource acquisition traits as a whole responded to the abiotic variables of water nutrients
383 (PCAwater) and depth. Resource conservation traits were impacted by biotic and abiotic
384 variables, namely the phytoplankton concentration, species abundance, and sediment
385 nutrients (PCAsediments; Table 2).

386 Four traits responded to only abiotic (height, SLA, and internode length, respectively
387 explaining 3.15%, 14.31%, and 5.03% of the variance) or biotic (specific internode mass,
388 explaining 10.28% of the variance) pond variables (Table 2). Variation in the LDMC, maximum
389 root length, and specific root mass was influenced by both biotic and abiotic variables (Table

390 2). The responses of these three traits to the variables examined were trait dependent, with a
391 greater cumulative influence of biotic variables (Table 2). The LDMC and maximum root length
392 were predicted largely by species abundances (explaining 5.90% and 4.37%, respectively, of
393 the variance), and the specific root mass was predicted mostly by the phytoplankton
394 concentration (explaining 10.49% of the variance). To lesser extents, the water physico-
395 chemistry (*i.e.*, temperature, pH, and nutrients) influenced both root traits and sediment
396 nutrients (PCA_{sediments}) influenced the LDMC.

397

398 [Plant traits directly and indirectly affected individual performance](#)

399 We identified multiple direct or indirect effects of all traits, except that the LDMC did not affect
400 the individual total biomass (Fig. 3). Traits directly influencing the individual total biomass
401 (height and all clonal and root traits) had positive effects, with the height (0.38, $p < 0.001$) and
402 specific internode mass (0.43, $p < 0.001$) having the strongest influence. These direct effects
403 on individual performance were modulated (*i.e.*, amplified or reduced) by their correlations (Fig.
404 3). For example, the internode length and specific root mass correlated with height, indirectly
405 supporting their total positive effect on individual performance. The SLA had no direct effect
406 on individual performance, but had a negative influence through its relationship with the
407 specific internode mass (-0.20 , $p < 0.05$). When total trait effects on individual performance
408 were considered (Fig. 3b), the SLA had a negative impact and all other traits had overall
409 positive effects.

410

411 **Discussion**

412 [Taxonomy explains most trait variance](#)

413 The linking of patterns to processes in community ecology is often challenging, especially in
414 field-based studies. Several authors have suggested that trait variance be used to infer

415 community assembly mechanisms (McGill, 2008; Taudiere & Violle, 2016; Violle et al., 2007,
416 2012). To tease apart the multiple factors that drive community assembly in aquatic systems,
417 we partitioned the variance of several traits among three distinct, yet interacting, scales (space,
418 time, and taxonomy). Studies in which different scales are taken into account simultaneously
419 remain rare (but see, e.g., Henn et al., 2018; Liu et al., 2018; Luo et al., 2016), and to our
420 knowledge, none of them was performed over several years.

421 In the present study, variance partitioning among the scales examined was similar
422 among traits, indicating that similar processes determined the different trait values. Community
423 assembly mechanisms operating over time (year) and space (region) were responsible for a
424 small percentage of the total trait variance, in line with the observations of Fu et al. (2013).
425 Those authors partitioned trait variance across spatial scales alone, and found that a large
426 spatial scale (*i.e.*, a regional scale including different lakes) did not account for much variance
427 in the root/shoot ratio and leaf number for the macrophyte *Potamogeton maackianus* (Fu et
428 al., 2013). In our work, trait distributions were fairly conserved among sites, which suggests
429 exchanges, therefore dispersion, between studied sites. The small size of the study area
430 probably explains the occurrence of most variance within and not among sites; a larger study
431 area should be considered in future research. Intrasite variance (at inter- and intraspecific
432 scales) overwhelmingly dominated intersite variance, for all traits. Thus, a mechanism
433 participated in the spreading of trait values around the optimum within sites. This pattern may
434 arise from several non-exclusive processes. First, adaptation to habitat heterogeneity within
435 sites can lead to differences in optimal trait values at the micro-site scale. Second, biotic
436 interactions can disperse trait values to limit similarity among individuals, thereby reducing
437 competition. As highlighted by Messier et al. (2010), niche and neutral theories can contribute
438 significantly to the explanation of community assembly.

439 Interestingly, taxonomy accounted for most variance in all traits, with in particular
440 intraspecific trait variance being greater (56.29% on average) than interspecific variance for
441 most traits. However, the relative importance of interspecific *versus* intraspecific trait variability

442 was trait dependent, consistent with previous findings (Albert, Thuiller, Yoccoz, Douzet, et al.,
443 2010; Fajardo & McIntire, 2010; Roche et al., 2004). We found LDMC and specific internode
444 mass to vary more between species than within species This indicates that traits related to
445 resources conservation are highly conserved among species, allowing different macrophyte
446 species to specialize and co-occur along habitat gradients (different conditions among ponds).
447 However, resource acquisition traits (height, SLA, internode length, maximum root length)
448 varied more within species than between species, in line with Siefert et al. (2015) who found
449 that “whole-plant traits” (*i.e.* plant height, and architecture) were more variable within
450 population than between species in a global-scale meta-analysis. This high amount of
451 intraspecific trait variability would mostly rely on phenotypic plasticity, already known to allow
452 macrophyte individuals to resist to filters, and to environmental heterogeneity at very fine scale
453 (within-pond habitat variances) (Fu et al., 2012, 2013; L. Ni, 2001; Yang et al., 2004). Another
454 hypothesis is that, in macrophytes, architectural traits can display multiple trait values since
455 individuals are much less constrained by gravitational force than terrestrial species (Fu et al.,
456 2013).

457 We confirmed that partitioning trait variance is a relevant approach to detect at which
458 scale operate the most decisive processes in plant community assembly without scale
459 dependency issues (McGill, 2008). Moreover, we underlined the strong importance of ITV in
460 shaping community assembly (Jiang et al., 2016; Violle et al., 2012), and we then argue that
461 not only species mean and community-aggregated trait values (Schamp et al., 2007; Stubbs
462 & Bastow Wilson, 2004), but also trait distributions have to be analyzed when studying
463 community dynamics and structure (de la Riva et al., 2016; Henn et al., 2018). Including ITV
464 into community ecology can improve the detection of processes underlying species
465 coexistence at local scale (Doudová & Douda, 2020; Fu et al., 2014; Violle et al., 2009).

466

467 [Intraspecific trait variation occurred in response to biotic and abiotic variables](#)

468 As studies on aquatic systems overall focus on macrophyte species responses to a specific,
469 and usually limited, number of environmental variables ([Barker et al., 2008](#); [Fu, 2020](#); [Fu et](#)
470 [al., 2014](#); [Li et al., 2018](#); [Ma et al., 2022a](#); [Martin & Coetzee, 2014](#); [Shields & Moore, 2016](#)),
471 our vision of ITV in the context of multiple constraints remains partial. With this study, we
472 sought to fill this gap by simultaneously examining trait responses to multiple abiotic and biotic
473 variables.

474 Contrary to our expectations, not all traits of the pond-dwelling individuals responded
475 to both abiotic and biotic variables. Three resource acquisition traits (height, SLA, and
476 internode length) and one resource conservation trait (specific internode mass) were
477 influenced by abiotic or biotic variables alone, although they are known to usually respond to
478 both variable types ([Fu et al., 2013](#); [Gao et al., 2021](#); [Pilon & Santamaría, 2002](#)). However, all
479 other trait variations (*i.e.*, the LDMC, maximum root length, and specific root mass) responded
480 to biotic and abiotic variables as predicted. This difference in trait response patterns indicates
481 that individual plants do not respond as a whole, but instead that each trait has a singular role,
482 in the context of multiple constraints. This finding challenges the validity of conclusions that
483 assume responses at the whole-plant level ([De Kroon et al., 2005](#)), and suggests that survival
484 within multi-constrained habitats relies on low phenotypic integration ([Murren, 2002](#)), enabling
485 the expression of multivariate phenotypes. Overall, our findings underline the relevance of
486 multi-trait approaches integrating biotic and abiotic variables, as also stated by [Chalmandrier](#)
487 [et al. \(2022\)](#), to efficiently capture plant response strategies and ultimately identify mechanisms
488 of species coexistence at local scales. In addition, our observations suggest that patterns of
489 trait variation measured *in situ* should be interpreted carefully, as they may not reflect
490 responses to only biotic or abiotic filters ([Duarte et al., 1986](#); [Körs et al., s. d.](#); [Su et al., 2019](#)).

491 In the ponds of the Iles Kerguelen, each trait category was involved in the response to
492 specific habitat variables. First, traits related to resource acquisition, as a whole, responded to
493 water nutrients. The SLA decreased when water nutrients increased, suggesting that

494 macrophyte species are plastic, maximizing either light capture when higher nutrient
495 concentrations stimulate phytoplankton growth, reducing light penetration into water (Rabalais,
496 2002), or nutrient capture (Zervas et al., 2019). In addition, the root length, also involved in
497 nutrient acquisition, was impacted positively by species abundances. Competition for nutrient
498 acquisition favors root trait divergence and forces plants to optimize resource uptake
499 (Ferguson et al., 2016; Manschadi et al., 2006). This result indicates that the consideration of
500 root trait variation can aid the detection of responses to biotic competition, although it is not
501 incorporated well into field studies due to the difficulty of sampling (Colom & Baucom, 2020).
502 Lastly, internodes elongated with water depth, probably as part of a strategy to avoid harsh
503 conditions (Bittebiere & Mony, 2015; Callaghan et al., 1992; Louâpre et al., 2012). Second,
504 traits related to resource conservation (the LDMC and specific internode and root masses)
505 were impacted negatively by competition. Resource storage in connection internodes and
506 roots decreased in response to species abundance and/or phytoplankton biomass, likely
507 reflecting a lack of available resources. To a lesser extent, the LDMC responded to sediment
508 nutrients, consistent with previous findings (Fu, 2020; Garnier et al., 2001). Previous studies
509 have suggested that an increase in nutrient availability induces leaf trait variability toward low
510 nutrient conservation, for instance by reducing the LDMC (Poorter & Garnier, 2007).

511 Based on our results, we recommend that several biotic and abiotic variables should
512 be considered in future studies to better understand the effects of environmental changes on
513 plant communities (Cavieres et al., 2014; Crain et al., 2004; Kim & Nishihiro, 2020; Klanderud
514 et al., 2015; Mitchell et al., 2017; Morris et al., 2020; Wood et al., 2012).

515

516 Effects of trait values and their relationships on macrophyte performance

517 The SEM approach allowed us to demonstrate that all traits considered except the LDMC had
518 direct or indirect effects on individual performance. This observation is especially important, as
519 the relationships between traits can influence individual performance (Engbersen et al., 2022;

520 [Pywell et al., 2003](#)), in particular in macrophyte species ([Fu et al., 2014](#); [Ma et al., 2022b](#)). Our
521 findings support the current view in functional ecology of the relevance of multi-trait approaches
522 in which trait relationships are taken into account ([Albert, Thuiller, Yoccoz, Douzet, et al., 2010](#);
523 [Bittebiere et al., 2019](#); [Blonder et al., 2014](#); [Gustafsson & Norkko, 2019](#)).

524 As traits are interrelated, adjustments in trait values must be coordinated to produce
525 properly functioning organisms; in addition, patterns of trait correlation can vary greatly across
526 environments ([Murren, 2002](#)). To maintain optimal fitness, functionally related traits of
527 individual plants should correlate more strongly (*i.e.*, high level of integration within the
528 functional unit) than do functionally different traits (*i.e.*, among functional units), conferring
529 flexibility as well as coordination ([Nicotra et al., 1997](#)). This pattern is consistent with our
530 results, as we detected the strongest correlations among traits related to resource
531 conservation (the specific internode and root masses) and leaf traits (the SLA and LDMC).

532 With the pooling of direct and indirect effects, we found that height and resource
533 conservation traits (the specific internode and root masses) had stronger positive effects than
534 did the other resource acquisition traits (root and internode lengths and SLA), consistent with
535 the results reported by [Wildová et al. \(2007\)](#). Individual growth and resource conservation traits
536 are related directly to individual size, thus affecting the individual biomass, whereas resource
537 acquisition traits are related more to colonization ability and allocation between organs, thus
538 affecting individual performance more indirectly ([Wildová et al., 2007](#)). Thus, individuals would
539 prioritize investment in resource conservation in constrained habitats, such as sub-Antarctic
540 ponds.

541 Finally, abiotic variables can have cascading effects on individual performance through
542 their influence on trait variation. Seasonal and annual changes in abiotic values in constrained
543 habitats such as polar ponds would greatly impact plant trait values. For example, an increase
544 in the water temperature reduces root length, indirectly and negatively affecting individual
545 performance. Thus, although increases in the water temperature can stimulate macrophyte
546 performance within the optimum thermal range ([Dar et al., 2014](#)), they may have negative

547 impacts in sub-Antarctic ponds, suggesting that the optimal thermal range can be exceeded
548 ([Frenot et al., 2006](#)). Similarly, variation in water depth affects plant performance through its
549 effect on internode length. With continued increases in the water temperature and reductions
550 in precipitation ([Walther et al., 2002](#)), one can predict that the consequences for the
551 macrophyte community structure and its productivity will be severe.

552

553 **Conclusion**

554 The partitioning of trait variance among multiple ecological scales provides a less fragmented
555 picture of variables that drive community assembly and avoids the misinterpretation of trait
556 variation in field studies. This study underlines the strong importance of ITV in the shaping of
557 community assembly ([Jiang et al., 2016](#); [Violle et al., 2012](#)), and more precisely, in species
558 coexistence and responses to multiple abiotic filters. We recommend the simultaneous
559 consideration of several biotic and abiotic variables in future studies to better understand the
560 effects of environmental changes on plant communities ([Cavieres et al., 2014](#); [Crain et al.,](#)
561 [2004](#); [Klanderud et al., 2015](#); [Mitchell et al., 2017](#); [Morris et al., 2020](#); [Wood et al., 2012](#)).
562 Additionally, we emphasize the relevance of multi-trait approaches in which trait relationships
563 are taken into account in the examination of trait effects on individual performance.

564

565 **Acknowledgments**

566 We thank all of the volunteers for their invaluable help in the field and laboratory during the 4
567 years of data collection.

568 **Author contributions**

569 A-K.B. and D.R. acquired the financial support for the study. A-K.B. and D.R. lead the
570 administration of the project (Management and coordination responsibility for the research
571 activity planning and execution, including obtention of sampling permits). P.D. and A-K.B.
572 conceived the ideas and designed methodology. P.D. and A-K.B. collected the data. P.D., L.S.,

573 F.M-B., F.V. and A-K.B. performed the nutrient analyses. P.D. and A-K.B. analyzed the data.
574 P.D., D.R., L.S., F.M-B. and A-K.B. led to the writing of the manuscript. All authors contributed
575 critically to the drafts and gave final approval for publication.

576 **Conflict of interest**

577 The authors declare that they have no conflict of interest.

578

579 **Data availability statement**

580 The data that support the findings of this study are openly available in figshare at
581 https://figshare.com/articles/dataset/Data_ponds_Kerguelen/21904257

582 **Supporting information**

583 Appendix S1. Locations of the three sampling sites in the French sub-Antarctic Iles Kerguelen.

584 Appendix S2. Means \pm standard deviations for abiotic and biotic variables at the site scale.

585 Appendix S3. Photos of the aerial parts of the six studied macrophyte species showing the
586 absence of heterophylly.

587 Appendix S4. Eigenvector scores for the first axis of a principal component analysis (PCA 1)
588 of water nutrient parameters measured at the pond level.

589 Appendix S5. Eigenvector scores for the first axis of a principal component analysis (PCA 1)
590 of abiotic parameters measured at the pond level.

591 Appendix S6. Eigenvector scores for the first three axes of a factorial correspondence analysis
592 (FCAs 1–3) of water nutrient parameters measured at the pond level.

593 Appendix S7. Numbers of collected and measured samples per species, site, and year.

594 Appendix S8. Variance partitioning of individual traits across three nested scales: year, site,
595 and within-site.

596 Appendix S9. Variance partitioning of individual traits across three nested scales: year, site,
597 and within-site when deleting the less abundant species.

598 Appendix S10. Theoretical structural equation model representing the direct and indirect
599 effects of trait values measured in ponds on individual performance (total biomass).

600 **References**

- 601 Ackerly, D. D., Dudley, S. A., Sultan, S. E., Schmitt, J., Coleman, J. S., Linder, C. R., Sandquist, D. R.,
602 Geber, M. A., Evans, A. S., Dawson, T. E., & Lechowicz, M. J. (2000). The Evolution of Plant
603 Ecophysiological Traits : Recent Advances and Future Directions: New research addresses
604 natural selection, genetic constraints, and the adaptive evolution of plant ecophysiological
605 traits. *BioScience*, *50*(11), 979-995. [https://doi.org/10.1641/0006-](https://doi.org/10.1641/0006-3568(2000)050[0979:TEOPET]2.0.CO;2)
606 [3568\(2000\)050\[0979:TEOPET\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2000)050[0979:TEOPET]2.0.CO;2)
- 607 Adomako, M. O., Xue, W., Du, D.-L., & Yu, F.-H. (2021). Soil biota and soil substrates influence
608 responses of the rhizomatous clonal grass *Leymus chinensis* to nutrient heterogeneity. *Plant*
609 *and Soil*, *465*(1), 19-29. <https://doi.org/10.1007/s11104-021-04967-0>
- 610 Albert, C. H., Grassein, F., Schurr, F. M., Vieilledent, G., & Violle, C. (2011). When and how should
611 intraspecific variability be considered in trait-based plant ecology? *Perspectives in Plant*
612 *Ecology, Evolution and Systematics*, *13*(3), 217-225.
613 <https://doi.org/10.1016/j.ppees.2011.04.003>
- 614 Albert, C. H., Thuiller, W., Yoccoz, N. G., Douzet, R., Aubert, S., & Lavorel, S. (2010). A multi-trait
615 approach reveals the structure and the relative importance of intra- vs. interspecific
616 variability in plant traits : Intra- vs. interspecific variability in plant traits. *Functional Ecology*,
617 *24*(6), 1192-1201. <https://doi.org/10.1111/j.1365-2435.2010.01727.x>
- 618 Albert, C. H., Thuiller, W., Yoccoz, N. G., Soudant, A., Boucher, F., Saccone, P., & Lavorel, S. (2010).
619 Intraspecific functional variability : Extent, structure and sources of variation. *Journal of*
620 *Ecology*, *98*(3), 604-613. <https://doi.org/10.1111/j.1365-2745.2010.01651.x>
- 621 Barker, T., Hatton, K. O., Connor, L., & Moss, B. (2008). Effects of nitrate load on submerged plant
622 biomass and species richness : Results of a mesocosm experiment. *Fundamental and Applied*
623 *Limnology*, *173*(2), 89-100. <https://doi.org/10.1127/1863-9135/2008/0173-0089>
- 624 Bartoń, K. (2013). MuMIn : Multi-model inference. In *R package version 1.10.0*. (Vol. 1).

625 Bergstrom, D. M., & Chown, S. L. (1999). Life at the front : History, ecology and change on southern
626 ocean islands. *Trends in Ecology & Evolution*, *14*(12), 472-477.
627 [https://doi.org/10.1016/S0169-5347\(99\)01688-2](https://doi.org/10.1016/S0169-5347(99)01688-2)

628 Bittebiere, A., Saiz, H., & Mony, C. (2019). New insights from multidimensional trait space responses
629 to competition in two clonal plant species. *Functional Ecology*, *33*(2), 297-307.
630 <https://doi.org/10.1111/1365-2435.13220>

631 Bittebiere, A.-K., & Mony, C. (2015). Plant traits respond to the competitive neighbourhood at
632 different spatial and temporal scales. *Annals of Botany*, *115*(1), 117-126.
633 <https://doi.org/10.1093/aob/mcu206>

634 Blonder, B., Lamanna, C., Violle, C., & Enquist, B. J. (2014). The n -dimensional hypervolume : The n -
635 dimensional hypervolume. *Global Ecology and Biogeography*, *23*(5), 595-609.
636 <https://doi.org/10.1111/geb.12146>

637 Bornette, G., & Puijalon, S. (2011). Response of aquatic plants to abiotic factors : A review. *Aquatic
638 Sciences*, *73*(1), 1-14. <https://doi.org/10.1007/s00027-010-0162-7>

639 Brodie, C. R., Leng, M. J., Casford, J. S. L., Kendrick, C. P., Lloyd, J. M., Yongqiang, Z., & Bird, M. I.
640 (2011). Evidence for bias in C and N concentrations and $\delta^{13}\text{C}$ composition of terrestrial and
641 aquatic organic materials due to pre-analysis acid preparation methods. *Chemical Geology*,
642 *282*(3-4), 67-83. <https://doi.org/10.1016/j.chemgeo.2011.01.007>

643 Butterfield, B. J., & Callaway, R. M. (2013). A functional comparative approach to facilitation and its
644 context dependence. *Functional Ecology*, *27*(4), 907-917. [https://doi.org/10.1111/1365-
645 2435.12019](https://doi.org/10.1111/1365-2435.12019)

646 Byars, S. G., Papst, W., & Hoffmann, A. A. (2007). LOCAL ADAPTATION AND COGRADIENT SELECTION
647 IN THE ALPINE PLANT, POA HIEMATA, ALONG A NARROW ALTITUDINAL GRADIENT. *Evolution*,
648 *61*(12), 2925-2941. <https://doi.org/10.1111/j.1558-5646.2007.00248.x>

649 Cadotte, M. W., & Tucker, C. M. (2017). Should Environmental Filtering be Abandoned? *Trends in
650 Ecology & Evolution*, *32*(6), 429-437. <https://doi.org/10.1016/j.tree.2017.03.004>

651 Callaghan, T. V., Carlsson, B. Å., Jónsdóttir, I. S., Svensson, B. M., Jonasson, S., Carlsson, B. A., &
652 Jonsdottir, I. S. (1992). Clonal Plants and Environmental Change : Introduction to the
653 Proceedings and Summary. *Oikos*, 63(3), 341. <https://doi.org/10.2307/3544959>
654 Calow, P. (1987). Towards a Definition of Functional Ecology. *Functional Ecology*, 1(1), 57-61.
655 <https://doi.org/10.2307/2389358>
656 Cavender-Bares, J., Kozak, K. H., Fine, P. V. A., & Kembel, S. W. (2009). The merging of community
657 ecology and phylogenetic biology. *Ecology Letters*, 12(7), 693-715.
658 <https://doi.org/10.1111/j.1461-0248.2009.01314.x>
659 Cavieres, L. A., Brooker, R. W., Butterfield, B. J., Cook, B. J., Kikvidze, Z., Lortie, C. J., Michalet, R.,
660 Pugnaire, F. I., Schöb, C., Xiao, S., Anthelme, F., Björk, R. G., Dickinson, K. J. M., Cranston, B.
661 H., Gavilán, R., Gutiérrez-Girón, A., Kanka, R., Maalouf, J.-P., Mark, A. F., ... Callaway, R. M.
662 (2014). Facilitative plant interactions and climate simultaneously drive alpine plant diversity.
663 *Ecology Letters*, 17(2), 193-202. <https://doi.org/10.1111/ele.12217>
664 Chalmardrier, L., Stouffer, D. B., Purcell, A. S. T., Lee, W. G., Tanentzap, A. J., & Laughlin, D. C. (2022).
665 Predictions of biodiversity are improved by integrating trait-based competition with abiotic
666 filtering. *Ecology Letters*, ele.13980. <https://doi.org/10.1111/ele.13980>
667 Chmara, R., Pronin, E., & Szmeja, J. (2021). Functional macrophyte trait variation as a response to the
668 source of inorganic carbon acquisition. *PeerJ*, 9, e12584.
669 <https://doi.org/10.7717/peerj.12584>
670 Colom, S. M., & Baucom, R. S. (2020). Belowground Competition Can Influence the Evolution of Root
671 Traits. *The American Naturalist*, 195(4), 577-590. <https://doi.org/10.1086/707597>
672 Cornelissen, J. H. C., Lavorel, S., Garnier, E., Díaz, S., Buchmann, N., Gurvich, D. E., Reich, P. B., Steege,
673 H. ter, Morgan, H. D., Heijden, M. G. A. van der, Pausas, J. G., & Poorter, H. (2003). A
674 handbook of protocols for standardised and easy measurement of plant functional traits
675 worldwide. *Australian Journal of Botany*, 51(4), 335. <https://doi.org/10.1071/BT02124>

676 Crain, C. M., Silliman, B. R., Bertness, S. L., & Bertness, M. D. (2004). PHYSICAL AND BIOTIC DRIVERS
677 OF PLANT DISTRIBUTION ACROSS ESTUARINE SALINITY GRADIENTS. *Ecology*, *85*(9),
678 2539-2549. <https://doi.org/10.1890/03-0745>

679 Dalla Vecchia, A., Villa, P., & Bolpagni, R. (2020). Functional traits in macrophyte studies : Current
680 trends and future research agenda. *Aquatic Botany*, *167*, 103290.
681 <https://doi.org/10.1016/j.aquabot.2020.103290>

682 Dar, N. A., Pandit, A. K., & Ganai, B. A. (2014). Factors affecting the distribution patterns of aquatic
683 macrophytes. *Limnological Review*, *14*(2), 75-81. <https://doi.org/10.2478/limre-2014-0008>

684 De Kroon, H., Huber, H., Stuefer, J. F., & Van Groenendael, J. M. (2005). A modular concept of
685 phenotypic plasticity in plants. *New Phytologist*, *166*(1), 73-82.
686 <https://doi.org/10.1111/j.1469-8137.2004.01310.x>

687 de la Riva, E. G., Pérez-Ramos, I. M., Tosto, A., Navarro-Fernández, C. M., Olmo, M., Marañón, T., &
688 Villar, R. (2016). Disentangling the relative importance of species occurrence, abundance and
689 intraspecific variability in community assembly : A trait-based approach at the whole-plant
690 level in Mediterranean forests. *Oikos*, *125*(3), 354-363. <https://doi.org/10.1111/oik.01875>

691 De Wilde, M., Sebei, N., Puijalon, S., & Bornette, G. (2014). Responses of macrophytes to
692 dewatering : Effects of phylogeny and phenotypic plasticity on species performance.
693 *Evolutionary Ecology*, *28*(6), 1155-1167. <https://doi.org/10.1007/s10682-014-9725-8>

694 Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B.,
695 Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E., Reineking, B.,
696 Schröder, B., Skidmore, A. K., Zurell, D., & Lautenbach, S. (2013). Collinearity : A review of
697 methods to deal with it and a simulation study evaluating their performance. *Ecography*,
698 *36*(1), 27-46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>

699 Dostálek, T., Rokaya, M. B., & Münzbergová, Z. (2020). Plant palatability and trait responses to
700 experimental warming. *Scientific Reports*, *10*(1), 10526. [https://doi.org/10.1038/s41598-020-](https://doi.org/10.1038/s41598-020-67437-0)
701 [67437-0](https://doi.org/10.1038/s41598-020-67437-0)

702 Douce, P., Mermillod-Blondin, F., Simon, L., Dolédec, S., Eymar-Dauphin, P., Renault, D., Sulmon, C.,
703 Vallier, F., & Bittebiere, A.-K. (2023a). Biotic and abiotic drivers of aquatic plant communities
704 in shallow pools and wallows on the sub-Antarctic Iles Kerguelen. *Polar Biology*, *46*(4),
705 303-318. <https://doi.org/10.1007/s00300-023-03122-y>

706 Douce, P., Saiz, H., Benot, M.-L., Mermillod-Blondin, F., Simon, L., Renault, D., Vallier, F., Oury, Y.,
707 Fontaine, M., & Bittebiere, A.-K. (2023b). Functional characteristics rather than co-
708 occurrences determine the outcome of interactions between neighbouring plants in sub-
709 Antarctic ponds : Consequences for macrophyte community biomass. *Freshwater Biology*,
710 *68*(4), 561-576. <https://doi.org/10.1111/fwb.14047>

711 Doudová, J., & Douda, J. (2020). Along with intraspecific functional trait variation, individual
712 performance is key to resolving community assembly processes. *Functional Ecology*, *34*(11),
713 2362-2374. <https://doi.org/10.1111/1365-2435.13646>

714 Duarte, C. M., Kalff, J., & Peters, R. H. (1986). Patterns in Biomass and Cover of Aquatic Macrophytes
715 in Lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, *43*(10), 1900-1908.
716 <https://doi.org/10.1139/f86-235>

717 Dwyer, J. M., & Laughlin, D. C. (2017). Constraints on trait combinations explain climatic drivers of
718 biodiversity : The importance of trait covariance in community assembly. *Ecology Letters*,
719 *20*(7), 872-882. <https://doi.org/10.1111/ele.12781>

720 Elger, A., & Willby, N. J. (2003). Leaf Dry Matter Content as an Integrative Expression of Plant
721 Palatability : The Case of Freshwater Macrophytes. *Functional Ecology*, *17*(1), 58-65.

722 Engbersen, N., Stefan, L., Brooker, R. W., & Schöb, C. (2022). Using plant traits to understand the
723 contribution of biodiversity effects to annual crop community productivity. *Ecological*
724 *Applications*, *32*(1). <https://doi.org/10.1002/eap.2479>

725 Fajardo, A., & McIntire, E. J. B. (2010). Under strong niche overlap conspecifics do not compete but
726 help each other to survive : Facilitation at the intraspecific level: Intraspecific facilitation in

727 Nothofagus pumilio. *Journal of Ecology*, no-no. <https://doi.org/10.1111/j.1365->
728 2745.2010.01771.x

729 Ferguson, L., Sancho, G., Rutter, M. T., & Murren, C. J. (2016). Root architecture, plant size and soil
730 nutrient variation in natural populations of *Arabidopsis thaliana*. *Evolutionary Ecology*, 30(1),
731 155-171. <https://doi.org/10.1007/s10682-015-9808-1>

732 Fort, F., Jouany, C., & Cruz, P. (2013). Root and leaf functional trait relations in Poaceae species :
733 Implications of differing resource-acquisition strategies. *Journal of Plant Ecology*, 6(3),
734 211-219. <https://doi.org/10.1093/jpe/rts034>

735 Frenot, Y., Gloaguen, J. C., Cannavacciuolo, M., & Bellido, A. (1998). Primary succession on glacier
736 forelands in the subantarctic Kerguelen Islands. *Journal of Vegetation Science*, 9(1), 75-84.
737 <https://doi.org/10.2307/3237225>

738 Frenot, Y., Lebouvier, M., Gloaguen, J.-C., Hennion, F., Vernon, P., & Chapuis, J.-L. (2006). Impact des
739 changements climatiques et de la fréquentation humaine sur la biodiversité des îles
740 subantarctiques françaises. *Belgeo. Revue belge de géographie*, 3, 363-372.
741 <https://doi.org/10.4000/belgeo.12097>

742 Freschet, G. T., Valverde-Barrantes, O. J., Tucker, C. M., Craine, J. M., McCormack, M. L., Violle, C.,
743 Fort, F., Blackwood, C. B., Urban-Mead, K. R., Iversen, C. M., Bonis, A., Comas, L. H.,
744 Cornelissen, J. H. C., Dong, M., Guo, D., Hobbie, S. E., Holdaway, R. J., Kembel, S. W., Makita,
745 N., ... Roumet, C. (2017). Climate, soil and plant functional types as drivers of global fine-root
746 trait variation. *Journal of Ecology*, 105(5), 1182-1196. <https://doi.org/10.1111/1365->
747 2745.12769

748 Fu, H. (2020). Trait-based community assembly of submersed macrophytes subjected to nutrient
749 enrichment in freshwater lakes_ Do traits at the individual level matter? *Ecological*
750 *Indicators*, 9.

751 Fu, H., Yuan, G., Cao, T., Ni, L., Zhang, M., & Wang, S. (2012). An alternative mechanism for shade
752 adaptation : Implication of allometric responses of three submersed macrophytes to water
753 depth. *Ecological Research*, 27(6), 1087-1094. <https://doi.org/10.1007/s11284-012-0991-z>

754 Fu, H., Yuan, G., Lou, Q., Dai, T., Xu, J., Cao, T., Ni, L., Zhong, J., & Fang, S. (2018). Functional traits
755 mediated cascading effects of water depth and light availability on temporal stability of a
756 macrophyte species. *Ecological Indicators*, 89, 168-174.
757 <https://doi.org/10.1016/j.ecolind.2018.02.010>

758 Fu, H., Yuan, G., Zhong, J., Cao, T., Ni, L., & Xie, P. (2013). Environmental and Ontogenetic Effects on
759 Intraspecific Trait Variation of a Macrophyte Species across Five Ecological Scales. *PLOS ONE*,
760 8(4), e62794. <https://doi.org/10.1371/journal.pone.0062794>

761 Fu, H., Zhong, J., Yuan, G., Xie, P., Guo, L., Zhang, X., Xu, J., Li, Z., Li, W., Zhang, M., Cao, T., & Ni, L.
762 (2014). Trait-based community assembly of aquatic macrophytes along a water depth
763 gradient in a freshwater lake. *Freshwater Biology*, 59(12), 2462-2471.
764 <https://doi.org/10.1111/fwb.12443>

765 Fullana-Pericas, M., Conesa, M. A., Soler, S., Ribas-Carbo, M., Granell, A., & Galmes, J. (2017).
766 Variations of leaf morphology, photosynthetic traits and water-use efficiency in Western-
767 Mediterranean tomato landraces. *Photosynthetica*, 55(1), 121-133.
768 <https://doi.org/10.1007/s11099-016-0653-4>

769 Funk, J. L., Larson, J. E., & Ricks-Oddie, J. (2021). Plant traits are differentially linked to performance
770 in a semiarid ecosystem. *Ecology*, 102(5), e03318. <https://doi.org/10.1002/ecy.3318>

771 Fyllas, N. M., Michelaki, C., Galanidis, A., Evangelou, E., Zaragoza-Castells, J., Dimitrakopoulos, P. G.,
772 Tsadilas, C., Arianoutsou, M., & Lloyd, J. (2020). Functional Trait Variation Among and Within
773 Species and Plant Functional Types in Mountainous Mediterranean Forests. *Frontiers in Plant*
774 *Science*, 11, 212. <https://doi.org/10.3389/fpls.2020.00212>

775 Gao, Y., Wang, L., Hu, X., Zhang, Z., Liu, B., Zhang, X., & Wang, G. (2021). Rapid adaptive responses of
776 rosette-type macrophyte *Vallisneria natans* juveniles to varying water depths : The role of

777 leaf trait plasticity. *Ecology and Evolution*, 11(20), 14268-14281.
778 <https://doi.org/10.1002/ece3.8142>

779 Garnier, E., Cortez, J., Billès, G., Navas, M.-L., Roumet, C., Debussche, M., Laurent, G., Blanchard, A.,
780 Aubry, D., Bellmann, A., Neill, C., & Toussaint, J.-P. (2004). PLANT FUNCTIONAL MARKERS
781 CAPTURE ECOSYSTEM PROPERTIES DURING SECONDARY SUCCESSION. *Ecology*, 85(9),
782 2630-2637. <https://doi.org/10.1890/03-0799>

783 Garnier, E., Shipley, B., Roumet, C., & Laurent, G. (2001). A Standardized Protocol for the
784 Determination of Specific Leaf Area and Leaf Dry Matter Content. *Functional Ecology*, 15(5),
785 688-695.

786 Geber, M. A., & Griffen, L. R. (2003). Inheritance and Natural Selection on Functional Traits.
787 *International Journal of Plant Sciences*, 164(S3), S21-S42. <https://doi.org/10.1086/368233>

788 Ghalambor, C. K., McKAY, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-adaptive
789 phenotypic plasticity and the potential for contemporary adaptation in new environments.
790 *Functional Ecology*, 21(3), 394-407. <https://doi.org/10.1111/j.1365-2435.2007.01283.x>

791 Gong, H., & Gao, J. (2019). Soil and climatic drivers of plant SLA (specific leaf area). *Global Ecology*
792 *and Conservation*, 20, e00696. <https://doi.org/10.1016/j.gecco.2019.e00696>

793 Grace, J. B., Anderson, T. M., Olff, H., & Scheiner, S. M. (2010). On the specification of structural
794 equation models for ecological systems. *Ecological Monographs*, 80(1), 67-87.
795 <https://doi.org/10.1890/09-0464.1>

796 Grasshoff, K., Ehrhardt, M., Kremling, K., & Anderson, L. G. (Éds.). (1999). *Methods of seawater*
797 *analysis* (3rd, completely rev. and extended ed éd.). Wiley-VCH.

798 Grassi, G., Vicinelli, E., Ponti, F., Cantoni, L., & Magnani, F. (2005). Seasonal and interannual variability
799 of photosynthetic capacity in relation to leaf nitrogen in a deciduous forest plantation in
800 northern Italy. *Tree Physiology*, 25(3), 349-360. <https://doi.org/10.1093/treephys/25.3.349>

801 Gray, S. B., & Brady, S. M. (2016). Plant developmental responses to climate change. *Developmental*
802 *Biology*, 419(1), 64-77. <https://doi.org/10.1016/j.ydbio.2016.07.023>

803 Gustafsson, C., & Norkko, A. (2019). Quantifying the importance of functional traits for primary
804 production in aquatic plant communities. *Journal of Ecology*, *107*(1), 154-166.
805 <https://doi.org/10.1111/1365-2745.13011>

806 Han, S., Xing, Z., Jiang, H., Li, W., & Huang, W. (2021). Biological adaptive mechanisms displayed by a
807 freshwater plant to live in aquatic and terrestrial environments. *Environmental and*
808 *Experimental Botany*, *191*, 104623. <https://doi.org/10.1016/j.envexpbot.2021.104623>

809 Henn, J. J., Buzzard, V., Enquist, B. J., Halbritter, A. H., Klanderud, K., Maitner, B. S., Michaletz, S. T.,
810 Pötsch, C., Seltzer, L., Telford, R. J., Yang, Y., Zhang, L., & Vandvik, V. (2018). Intraspecific Trait
811 Variation and Phenotypic Plasticity Mediate Alpine Plant Species Response to Climate
812 Change. *Frontiers in Plant Science*, *9*.
813 <https://www.frontiersin.org/articles/10.3389/fpls.2018.01548>

814 Hutchings, M. J., Turkington, R., Klein, E., & Carey, P. (1997). Morphological plasticity in *Trifolium*
815 *repens* L. : The effects of clone genotype, soil nutrient level, and the genotype of conspecific
816 neighbours. *Canadian Journal of Botany*, *75*(8), 1382-1393. <https://doi.org/10.1139/b97-852>

817 Jiang, Y., Chen, X., Ma, J., Liang, S., Huang, J., Liu, R., & Pan, Y. (2016). Interspecific and Intraspecific
818 Variation in Functional Traits of Subtropical Evergreen and Deciduous Broadleaved Mixed
819 Forests in Karst Topography, Guilin, Southwest China. *Tropical Conservation Science*, *9*(4),
820 194008291668021. <https://doi.org/10.1177/1940082916680211>

821 Johnson, R. K., & Toprak, V. (2021). Local habitat is a strong determinant of spatial and temporal
822 patterns of macrophyte diversity and composition in boreal lakes. *Freshwater Biology*, *66*(8),
823 1490-1501. <https://doi.org/10.1111/fwb.13733>

824 Jung, V., Violle, C., Mondy, C., Hoffmann, L., & Muller, S. (2010). Intraspecific variability and trait-
825 based community assembly : Intraspecific variability and community assembly. *Journal of*
826 *Ecology*, *98*(5), 1134-1140. <https://doi.org/10.1111/j.1365-2745.2010.01687.x>

827 Keser, L. H., Dawson, W., Song, Y.-B., Yu, F.-H., Fischer, M., Dong, M., & van Kleunen, M. (2014).
828 Invasive clonal plant species have a greater root-foraging plasticity than non-invasive ones.
829 *Oecologia*, 174(3), 1055-1064. <https://doi.org/10.1007/s00442-013-2829-y>

830 Kim, J. Y., & Nishihiro, J. (2020). Responses of lake macrophyte species and functional traits to
831 climate and land use changes. *Science of The Total Environment*, 736, 139628.
832 <https://doi.org/10.1016/j.scitotenv.2020.139628>

833 Klanderud, K., Vandvik, V., & Goldberg, D. (2015). The Importance of Biotic vs. Abiotic Drivers of Local
834 Plant Community Composition Along Regional Bioclimatic Gradients. *PLOS ONE*, 10(6),
835 e0130205. <https://doi.org/10.1371/journal.pone.0130205>

836 Kleyer, M., Trinogga, J., Cebrián-Piqueras, M. A., Trenkamp, A., Fløjgaard, C., Ejrnæs, R., Bouma, T. J.,
837 Minden, V., Maier, M., Mantilla-Contreras, J., Albach, D. C., & Blasius, B. (2019). Trait
838 correlation network analysis identifies biomass allocation traits and stem specific length as
839 hub traits in herbaceous perennial plants. *Journal of Ecology*, 107(2), 829-842.
840 <https://doi.org/10.1111/1365-2745.13066>

841 Koleszár, G., Lukács, B. A., Nagy, P. T., & Szabó, S. (2022). Shade tolerance as a key trait in invasion
842 success of submerged macrophyte *Cabomba caroliniana* over *Myriophyllum spicatum*.
843 *Ecology and Evolution*, 12(9), e9306. <https://doi.org/10.1002/ece3.9306>

844 Kõrs, A., Vilbaste, S., Käiro, K., Pall, P., Piirsoo, K., Truu, J., & Viik, M. (s. d.). *Temporal changes in the*
845 *composition of macrophyte communities and environmental factors governing the*
846 *distribution of aquatic plants in an unregulated lowland river (Emajõgi, Estonia)*. 17, 14.

847 Kraft, N. J. B., Adler, P. B., Godoy, O., James, E. C., Fuller, S., & Levine, J. M. (2015). Community
848 assembly, coexistence and the environmental filtering metaphor. *Functional Ecology*, 29(5),
849 592-599. <https://doi.org/10.1111/1365-2435.12345>

850 Laliberté, E., & Legendre, P. (2010). A distance-based framework for measuring functional diversity
851 from multiple traits. *Ecology*, 91(1), 299-305. <https://doi.org/10.1890/08-2244.1>

852 Lavorel, S., & Garnier, E. (2002). Predicting changes in community composition and ecosystem
853 functioning from plant traits : Revisiting the Holy Grail. *Functional Ecology*, 16(5), 545-556.
854 <https://doi.org/10.1046/j.1365-2435.2002.00664.x>

855 Le Bagousse-Pinguet, Y., Gross, N., Maestre, F. T., Maire, V., de Bello, F., Fonseca, C. R., Kattge, J.,
856 Valencia, E., Leps, J., & Liancourt, P. (2017). Testing the environmental filtering concept in
857 global drylands. *Journal of Ecology*, 105(4), 1058-1069. <https://doi.org/10.1111/1365-2745.12735>

859 Lebouvier, M. (2007). Conservation and management in the French sub-Antarctic islands and
860 surrounding seas. *Papers and Proceedings of the Royal Society of Tasmania*, 23-28.
861 <https://doi.org/10.26749/rstpp.141.1.23>

862 Levitt, J. (1980). *Water, Radiation, Salt, and Other Stresses*. Elsevier.

863 Li, F., Yang, N., Zhu, L., Xie, Y., Yang, G., Hu, C., Chen, X., & Deng, Z. (2018). Competition and
864 facilitation of two wetland macrophytes under different water levels and nutrient-
865 heterogeneous conditions. *Freshwater Science*. <https://doi.org/10.1086/697964>

866 Li, L., Barrett, S. C. H., Song, Z., & Chen, J. (2019). Sex-specific plasticity of reproductive allocation in
867 response to water depth in a clonal, dioecious macrophyte. *American Journal of Botany*,
868 106(1), 42-50. <https://doi.org/10.1002/ajb2.1218>

869 Liu, R., Liang, S., Long, W., & Jiang, Y. (2018). Variations in Leaf Functional Traits Across Ecological
870 Scales in Riparian Plant Communities of the Lijiang River, Guilin, Southwest China. *Tropical
871 Conservation Science*, 11, 194008291880468. <https://doi.org/10.1177/1940082918804680>

872 Louâpre, P., Bittebière, A.-K., Clément, B., Pierre, J.-S., & Mony, C. (2012). How Past and Present
873 Influence the Foraging of Clonal Plants? *PLoS ONE*, 7(6), e38288.
874 <https://doi.org/10.1371/journal.pone.0038288>

875 Loughnan, D., & Gilbert, B. (2017). Trait-mediated community assembly : Distinguishing the
876 signatures of biotic and abiotic filters. *Oikos*, 126(8), 1112-1122.
877 <https://doi.org/10.1111/oik.03945>

878 Luo, Y., Liu, J., Tan, S., Cadotte, M. W., Xu, K., Gao, L., & Li, D. (2016). Trait variation and functional
879 diversity maintenance of understory herbaceous species coexisting along an elevational
880 gradient in Yulong Mountain, Southwest China. *Plant Diversity*, 38(6), 303-311.
881 <https://doi.org/10.1016/j.pld.2016.11.002>

882 Ma, F., Yang, L., Zhang, C., Tao, M., Yu, H., Liu, C., & Yu, D. (2022). Functional evenness and
883 community-weighted mean traits have strong correlation with macrophyte community
884 productivity. *Aquatic Sciences*, 84(1), 2. <https://doi.org/10.1007/s00027-021-00833-4>

885 Ma, F., Zuo, Z., Yang, L., Li, D., Wang, H., Li, F., Fan, S., Liu, C., & Yu, D. (2022). The effect of trait-based
886 diversity on productivity results mainly from intraspecific trait variability in the macrophyte
887 community. *Freshwater Biology*, 67(7), 1137-1149. <https://doi.org/10.1111/fwb.13906>

888 Maire, V., Gross, N., Hill, D., Martin, R., Wirth, C., Wright, I. J., & Soussana, J.-F. (2013). Disentangling
889 Coordination among Functional Traits Using an Individual-Centred Model : Impact on Plant
890 Performance at Intra- and Inter-Specific Levels. *PLoS ONE*, 8(10), e77372.
891 <https://doi.org/10.1371/journal.pone.0077372>

892 Manschadi, A. M., Christopher, J., deVoil, P., & Hammer, G. L. (2006). The role of root architectural
893 traits in adaptation of wheat to water-limited environments. *Functional Plant Biology*, 33(9),
894 823. <https://doi.org/10.1071/FP06055>

895 Martin, G. D., & Coetsee, J. A. (2014). Competition between two aquatic macrophytes, *Lagarosiphon*
896 major (Ridley) Moss (Hydrocharitaceae) and *Myriophyllum spicatum* Linnaeus (Haloragaceae)
897 as influenced by substrate sediment and nutrients. *Aquatic Botany*, 114, 1-11.
898 <https://doi.org/10.1016/j.aquabot.2013.11.001>

899 Martinez-Almoyna, C., Piton, G., Abdulhak, S., Boulangeat, L., Choler, P., Delahaye, T., Dentant, C.,
900 Foulquier, A., Poulenard, J., Noble, V., Renaud, J., Rome, M., Saillard, A., The ORCHAMP
901 Consortium, Thuiller, W., & Münkemüller, T. (2020). Climate, soil resources and microbial
902 activity shape the distributions of mountain plants based on their functional traits.
903 *Ecography*, 43(10), 1550-1559. <https://doi.org/10.1111/ecog.05269>

904 McGill, B., Enquist, B., Weiher, E., & Westoby, M. (2006). Rebuilding community ecology from
905 functional traits. *Trends in Ecology & Evolution*, *21*(4), 178-185.
906 <https://doi.org/10.1016/j.tree.2006.02.002>

907 McGill, B. J. (2008). Exploring Predictions of Abundance from Body Mass Using Hierarchical
908 Comparative Approaches. *The American Naturalist*, *172*(1), 88-101.
909 <https://doi.org/10.1086/588044>

910 Meng, T.-T., Wang, H., Harrison, S. P., Prentice, I. C., Ni, J., & Wang, G. (2015). Responses of leaf traits
911 to climatic gradients : Adaptive variation versus compositional shifts. *Biogeosciences*, *12*(18),
912 5339-5352. <https://doi.org/10.5194/bg-12-5339-2015>

913 Messier, J., McGill, B. J., & Lechowicz, M. J. (2010). How do traits vary across ecological scales? A case
914 for trait-based ecology: How do traits vary across ecological scales? *Ecology Letters*, *13*(7),
915 838-848. <https://doi.org/10.1111/j.1461-0248.2010.01476.x>

916 Mitchell, R. M., Bakker, J. D., Vincent, J. B., & Davies, G. M. (2017). Relative importance of abiotic,
917 biotic, and disturbance drivers of plant community structure in the sagebrush steppe.
918 *Ecological Applications*, *27*(3), 756-768. <https://doi.org/10.1002/eap.1479>

919 Moles, A. T., Warton, D. I., Warman, L., Swenson, N. G., Laffan, S. W., Zanne, A. E., Pitman, A.,
920 Hemmings, F. A., & Leishman, M. R. (2009). Global patterns in plant height. *Journal of*
921 *Ecology*, *97*(5), 923-932. <https://doi.org/10.1111/j.1365-2745.2009.01526.x>

922 Morris, W. F., Ehrlén, J., Dahlgren, J. P., Loomis, A. K., & Louthan, A. M. (2020). Biotic and
923 anthropogenic forces rival climatic/abiotic factors in determining global plant population
924 growth and fitness. *Proceedings of the National Academy of Sciences*, *117*(2), 1107-1112.
925 <https://doi.org/10.1073/pnas.1918363117>

926 Muraoka, H., Saigusa, N., Nasahara, K. N., Noda, H., Yoshino, J., Saitoh, T. M., Nagai, S., Murayama, S.,
927 & Koizumi, H. (2010). Effects of seasonal and interannual variations in leaf photosynthesis
928 and canopy leaf area index on gross primary production of a cool-temperate deciduous

929 broadleaf forest in Takayama, Japan. *Journal of Plant Research*, 123(4), 563-576.
930 <https://doi.org/10.1007/s10265-009-0270-4>

931 Murphy, J., & Riley, J. P. (1962). A modified single solution method for the determination of
932 phosphate in natural waters. *Analytica Chimica Acta*, 27, 31-36.

933 Murren, C. J. (2002). Phenotypic integration in plants. *Plant Species Biology*, 17(2-3), 89-99.
934 <https://doi.org/10.1046/j.1442-1984.2002.00079.x>

935 Ni, L. (2001). Growth of *Potamogeton maackianus* under Low-Light Stress in Eutrophic Water. *Journal*
936 *of Freshwater Ecology*, 16(2), 249-256. <https://doi.org/10.1080/02705060.2001.9663809>

937 Ni, Z., Wang, S., & Wang, Y. (2016). Characteristics of bioavailable organic phosphorus in sediment
938 and its contribution to lake eutrophication in China. *Environmental Pollution*, 219, 537-544.
939 <https://doi.org/10.1016/j.envpol.2016.05.087>

940 Nicotra, A. B., Chazdon, R. L., & Schlichting, C. D. (1997). Patterns of genotypic variation and
941 phenotypic plasticity of light response in two tropical *Piper* (Piperaceae) species. *American*
942 *Journal of Botany*, 84(11), 1542-1552. <https://doi.org/10.2307/2446616>

943 Pan, Y., García-Girón, J., & Iversen, L. L. (2023). Global change and plant-ecosystem functioning in
944 freshwaters. *Trends in Plant Science*. <https://doi.org/10.1016/j.tplants.2022.12.013>

945 Pilon, J., & Santamaría, L. (2002). Clonal variation in the thermal response of the submerged aquatic
946 macrophyte *Potamogeton pectinatus*. *Journal of Ecology*, 90(1), 141-152.
947 <https://doi.org/10.1046/j.0022-0477.2001.00645.x>

948 Poorter, H., & Garnier, E. (2007). Ecological Significance of Inherent Variation in Relative Growth Rate
949 and Its Components. In *Functional Plant Ecology* (2^e éd.). CRC Press.

950 Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J., & Villar, R. (2009). Causes and consequences of
951 variation in leaf mass per area (LMA) : A meta-analysis. *New Phytologist*, 182(3), 565-588.
952 <https://doi.org/10.1111/j.1469-8137.2009.02830.x>

953 Puijalon, S., Bouma, T. J., Van Groenendael, J., & Bornette, G. (2008). Clonal Plasticity of Aquatic Plant
954 Species Submitted to Mechanical Stress : Escape versus Resistance Strategy. *Annals of*
955 *Botany*, 102(6), 989-996. <https://doi.org/10.1093/aob/mcn190>

956 Pywell, R. F., Bullock, J. M., Roy, D. B., Warman, L., Walker, K. J., & Rothery, P. (2003). Plant traits as
957 predictors of performance in ecological restoration : *Plant traits as predictors of*
958 *performance. Journal of Applied Ecology*, 40(1), 65-77. [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2664.2003.00762.x)
959 [2664.2003.00762.x](https://doi.org/10.1046/j.1365-2664.2003.00762.x)

960 Rabalais, N. N. (2002). *Nitrogen in Aquatic Ecosystems*. 31(2), 11.

961 Reich, P. B. (2014). The world-wide 'fast-slow' plant economics spectrum : A traits manifesto. *Journal*
962 *of Ecology*, 102(2), 275-301. <https://doi.org/10.1111/1365-2745.12211>

963 Reich, P. B., Ellsworth, D. S., Walters, M. B., Vose, J. M., Gresham, C., Volin, J. C., & Bowman, W. D.
964 (1999). Generality of Leaf Trait Relationships : A Test Across Six Biomes. *Ecology*, 80(6),
965 1955-1969. [https://doi.org/10.1890/0012-9658\(1999\)080\[1955:GOLTRA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1955:GOLTRA]2.0.CO;2)

966 Reich, P. B., Walters, M. B., & Ellsworth, D. S. (1997). From tropics to tundra : Global convergence in
967 plant functioning. *Proceedings of the National Academy of Sciences*, 94(25), 13730-13734.
968 <https://doi.org/10.1073/pnas.94.25.13730>

969 Roche, P., Díaz-Burlinson, N., & Gachet, S. (2004). Congruency analysis of species ranking based on
970 leaf traits : Which traits are the more reliable? *Plant Ecology Formerly 'Vegetatio'*, 174(1),
971 37-48. <https://doi.org/10.1023/B:VEGE.0000046056.94523.57>

972 Rooney, N., & Kalff, J. (2000). Inter-annual variation in submerged macrophyte community biomass
973 and distribution : The influence of temperature and lake morphometry. *Aquatic Botany*,
974 68(4), 321-335. [https://doi.org/10.1016/S0304-3770\(00\)00126-1](https://doi.org/10.1016/S0304-3770(00)00126-1)

975 Rosseel, Y. (2012). **lavaan** : An R Package for Structural Equation Modeling. *Journal of Statistical*
976 *Software*, 48(2). <https://doi.org/10.18637/jss.v048.i02>

977 Rusch, G. M., Wilmann, B., Klimešová, J., & Evju, M. (2011). Do Clonal and Bud Bank Traits Vary in
978 Correspondence with Soil Properties and Resource Acquisition Strategies? Patterns in Alpine

979 Communities in the Scandian Mountains. *Folia Geobotanica*, 46(2-3), 237-254.
980 <https://doi.org/10.1007/s12224-010-9072-7>

981 Saiz, H., Gómez-Gardeñes, J., Borda, J. P., & Maestre, F. T. (2018). The structure of plant spatial
982 association networks is linked to plant diversity in global drylands. *Journal of Ecology*, 106(4),
983 1443-1453. <https://doi.org/10.1111/1365-2745.12935>

984 Santamaría, L. (2002). Why are most aquatic plants widely distributed? Dispersal, clonal growth and
985 small-scale heterogeneity in a stressful environment. *Acta Oecologica*, 23(3), 137-154.
986 [https://doi.org/10.1016/S1146-609X\(02\)01146-3](https://doi.org/10.1016/S1146-609X(02)01146-3)

987 Schamp, B. S., Chau, J., & Aarssen, L. W. (2007). Dispersion of traits related to competitive ability in
988 an old-field plant community. *Journal of Ecology*, 0(0), 071119203335008-???
989 <https://doi.org/10.1111/j.1365-2745.2007.01328.x>

990 Shields, E. C., & Moore, K. A. (2016). Effects of sediment and salinity on the growth and competitive
991 abilities of three submersed macrophytes. *Aquatic Botany*, 132, 24-29.
992 <https://doi.org/10.1016/j.aquabot.2016.03.005>

993 Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., Aarssen, L. W., Baraloto,
994 C., Carlucci, M. B., Cianciaruso, M. V., de L. Dantas, V., de Bello, F., Duarte, L. D. S., Fonseca,
995 C. R., Freschet, G. T., Gaucherand, S., Gross, N., Hikosaka, K., Jackson, B., ... Wardle, D. A.
996 (2015). A global meta-analysis of the relative extent of intraspecific trait variation in plant
997 communities. *Ecology Letters*, 18(12), 1406-1419. <https://doi.org/10.1111/ele.12508>

998 Slade, A. J., & Hutchings, M. J. (1987a). The Effects of Light Intensity on Foraging in the Clonal Herb
999 *Glechoma Hederacea*. *Journal of Ecology*, 75(3), 639-650. <https://doi.org/10.2307/2260196>

1000 Slade, A. J., & Hutchings, M. J. (1987b). The Effects of Nutrient Availability on Foraging in the Clonal
1001 Herb *Glechoma Hederacea*. *The Journal of Ecology*, 75(1), 95.
1002 <https://doi.org/10.2307/2260538>

1003 Smith, V. R. (2008). Energy flow and nutrient cycling in the Marion Island terrestrial ecosystem : 30
1004 years on. *Polar Record*, 44(3), 211-226. <https://doi.org/10.1017/S0032247407007218>

- 1005 Stubbs, W. J., & Bastow Wilson, J. (2004). Evidence for limiting similarity in a sand dune community :
1006 *Limiting similarity within a dune community. Journal of Ecology*, 92(4), 557-567.
1007 <https://doi.org/10.1111/j.0022-0477.2004.00898.x>
- 1008 Su, H., Chen, J., Wu, Y., Chen, J., Guo, X., Yan, Z., Tian, D., Fang, J., & Xie, P. (2019). Morphological
1009 traits of submerged macrophytes reveal specific positive feedbacks to water clarity in
1010 freshwater ecosystems. *Science of The Total Environment*, 684, 578-586.
1011 <https://doi.org/10.1016/j.scitotenv.2019.05.267>
- 1012 Swenson, N. G., Enquist, B. J., Pither, J., Thompson, J., & Zimmerman, J. K. (2006). *THE PROBLEM AND*
1013 *PROMISE OF SCALE DEPENDENCY IN COMMUNITY PHYLOGENETICS*. 87(10), 7.
- 1014 Tao, S., Guo, Q., Li, C., Wang, Z., & Fang, J. (2016). Global patterns and determinants of forest canopy
1015 height. *Ecology*, 97(12), 3265-3270. <https://doi.org/10.1002/ecy.1580>
- 1016 Taudiere, A., & Violle, C. (2016). cati : An R package using functional traits to detect and quantify
1017 multi-level community assembly processes. *Ecography*, 39(7), 699-708.
1018 <https://doi.org/10.1111/ecog.01433>
- 1019 Valladares, F., Gianoli, E., & Gómez, J. M. (2007). Ecological limits to plant phenotypic plasticity. *The*
1020 *New Phytologist*, 176(4), 749-763. <https://doi.org/10.1111/j.1469-8137.2007.02275.x>
- 1021 Vile, D., Shipley, B., & Garnier, E. (2006). A STRUCTURAL EQUATION MODEL TO INTEGRATE CHANGES
1022 IN FUNCTIONAL STRATEGIES DURING OLD-FIELD SUCCESSION. *Ecology*, 87(2), 504-517.
1023 <https://doi.org/10.1890/05-0822>
- 1024 Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., Jung, V., & Messier, J. (2012).
1025 The return of the variance : Intraspecific variability in community ecology. *Trends in Ecology*
1026 *& Evolution*, 27(4), 244-252. <https://doi.org/10.1016/j.tree.2011.11.014>
- 1027 Violle, C., Garnier, E., Leconteur, J., Roumet, C., Pouteur, C., Blanchard, A., & Navas, M.-L. (2009).
1028 Competition, traits and resource depletion in plant communities. *Oecologia*, 160(4), 747-755.
1029 <https://doi.org/10.1007/s00442-009-1333-x>

1030 Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the
1031 concept of trait be functional! *Oikos*, *116*(5), 882-892. <https://doi.org/10.1111/j.0030->
1032 [1299.2007.15559.x](https://doi.org/10.1111/j.0030-1299.2007.15559.x)

1033 Vohra, F. (1966). Determination of photosynthetic pigments in sea-water. *Monographs*
1034 *Onocéanographie Methodology*, *66*.

1035 Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.-M.,
1036 Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological responses to recent climate change.
1037 *Nature*, *416*(6879), 389-395. <https://doi.org/10.1038/416389a>

1038 Wang, P., Zhang, Q., Xu, Y.-S., & Yu, F.-H. (2016). Effects of water level fluctuation on the growth of
1039 submerged macrophyte communities. *Flora*, *223*, 83-89.
1040 <https://doi.org/10.1016/j.flora.2016.05.005>


1041 Wehn, S., Taugourdeau, S., Johansen, L., & Hovstad, K. A. (2017). Effects of abandonment on plant
1042 diversity in semi-natural grasslands along soil and climate gradients. *Journal of Vegetation*
1043 *Science*, *28*(4), 838-847. <https://doi.org/10.1111/jvs.12543>

1044 Wildová, R., Gough, L., Herben, T., Hershock, C., & Goldberg, D. E. (2007). Architectural and growth
1045 traits differ in effects on performance of clonal plants : An analysis using a field-
1046 parameterized simulation model. *Oikos*, *116*(5), 836-852. <https://doi.org/10.1111/j.0030->
1047 [1299.2007.15430.x](https://doi.org/10.1111/j.0030-1299.2007.15430.x)

1048 Wolfer, S. R., & Straile, D. (2004). Spatio-temporal dynamics and plasticity of clonal architecture in
1049 *Potamogeton perfoliatus*. *Aquatic Botany*, *78*(4), 307-318.
1050 <https://doi.org/10.1016/j.aquabot.2003.11.005>

1051 Wood, K. A., Stillman, R. A., Clarke, R. T., Daunt, F., & O'Hare, M. T. (2012). Understanding Plant
1052 Community Responses to Combinations of Biotic and Abiotic Factors in Different Phases of
1053 the Plant Growth Cycle. *PLoS ONE*, *7*(11), e49824.
1054 <https://doi.org/10.1371/journal.pone.0049824>

1055 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J.,
1056 Chapin, T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J.,
1057 Hikosaka, K., Lamont, B. B., Lee, T., Lee, W., Lusk, C., ... Villar, R. (2004). The worldwide leaf
1058 economics spectrum. *Nature*, 428(6985), 821-827. <https://doi.org/10.1038/nature02403>
1059 Xiao, K., Yu, D., & Wang, J. (2006). Habitat selection in spatially heterogeneous environments : A test
1060 of foraging behaviour in the clonal submerged macrophyte *Vallisneria spiralis*. *Freshwater*
1061 *Biology*, 51(8), 1552-1559. <https://doi.org/10.1111/j.1365-2427.2006.01590.x>
1062 Yang, Y., Yu, D., Li, Y., Xie, Y., & Geng, X. (2004). Phenotypic Plasticity of Two Submersed Plants in
1063 Response to Flooding. *Journal of Freshwater Ecology*, 19(1), 69-76.
1064 <https://doi.org/10.1080/02705060.2004.9664514>
1065 Younginger, B. S., Sirová, D., Cruzan, M. B., & Ballhorn, D. J. (2017). Is biomass a reliable estimate of
1066 plant fitness? *Applications in Plant Sciences*, 5(2), 1600094.
1067 <https://doi.org/10.3732/apps.1600094>
1068 Zervas, D., Tsiaoussi, V., Kallimanis, A. S., Dimopoulos, P., & Tsiripidis, I. (2019). Exploring the
1069 relationships between aquatic macrophyte functional traits and anthropogenic pressures in
1070 freshwater lakes. *Acta Oecologica*, 99, 103443. <https://doi.org/10.1016/j.actao.2019.103443>
1071 Zhang, P., Grutters, B. M. C., van Leeuwen, C. H. A., Xu, J., Petruzzella, A., van den Berg, R. F., &
1072 Bakker, E. S. (2019). Effects of Rising Temperature on the Growth, Stoichiometry, and
1073 Palatability of Aquatic Plants. *Frontiers in Plant Science*, 9, 1947.
1074 <https://doi.org/10.3389/fpls.2018.01947>
1075 Zhao, C.-Y., Liu, Y.-Y., Shi, X.-P., & Wang, Y.-J. (2020). Effects of soil nutrient variability and competitor
1076 identify on growth and co-existence among invasive alien and native clonal plants.
1077 *Environmental Pollution*, 261, 113894. <https://doi.org/10.1016/j.envpol.2019.113894>
1078
1079



1081

1082 **Figure 1.** Conceptual graphic setting hypotheses.


1083

Table 1. Measured plant functional traits and associated responses to abiotic and biotic conditions.

Traits	Description	Calculation	Abiotic variables with evidenced trait shifts	Biotic interactions with evidenced trait shifts
<i>Resource acquisition traits</i>				
Specific Leaf Area (SLA)	Proxy of plant photosynthetic capacity and resource acquisition	Ratio of area and dry mass of leaf ($\text{mm}^2 \cdot \text{mg}^{-1}$)	Temperature [a, ab, b, c, d], nutrients [a, e, ab], moisture [f, g], light [ac]	Competition for light [h, i], above-ground competition [j, g], facilitation [i]
Height	Allow optimal light exposure	Shortest distance between the upper limit of the main photosynthetic tissues and the ground level (cm)	Temperature [a, n], water availability [o, p, g], water depth [q], soil nutrients [a, r]	Aerial competition [m, i], belowground competition [i, g], facilitation [i]
Internode length	Plant ability to exploit local environment and to disperse at short distance	Distance between two successive ramets (cm)	Soil nutrients [t, u, v, s], light [ac, ae, af, w]	Spatial competition [m, x, s]
Maximum root length	Acquisition of nutrients	Length of the longest root (cm)	Temperature [a, ad, n], soil nutrients [a, e]	Competition [y]
<i>Resource conservation traits</i>				
Leaf Dry Matter Content (LDMC)	Proxy of physical resistance of leaves and resources storage	Ratio of dry and fresh masses of leaf ($\text{mg} \cdot \text{g}^{-1}$)	Temperature [b, c], moisture [k, g], soil nutrients [e, k], water nutrients [l]	Competition [m, g]
Specific internode mass	Proxy of resources allocation for clonal multiplication and storage	Ratio of dry mass and length of connection ($\text{mg} \cdot \text{cm}^{-1}$)	Soil nutrients [s, t]	Competition [m]
Specific root mass	Proxy of resources storage	Ratio of dry mass and length of root ($\text{mg} \cdot \text{cm}^{-1}$)	Temperature [a, q], nutrients [a, e], Moisture [a, n]	

1085 a = (Cornelissen et al., 2003); ab = (Gong & Gao, 2019); ac = (Santamaría, 2002); ad = (Freschet et al., 2017); ae = (Xiao et al.,
1086 2006); af = (Puijalón et al., 2008); b = (Meng et al., 2015); c = (Dostálek et al., 2020); d = (Poorter et al., 2009); e = (Lavorel &
1087 Garnier, 2002); f = (Jung et al., 2010); g = (Loughnan & Gilbert, 2017); h = (Violle et al., 2009); i = (Butterfield & Callaway, 2013);
1088 j = (Rooney & Kalff, 2000); k = (Wehn et al., 2017); l = (Fu et al., 2018); m = (Bittebiere & Mony, 2015); n = (Gray & Brady,
1089 2016); o = (Moles et al., 2009); p = (Tao et al., 2016); q = (Wang et al., 2016); r = (Martinez-Almoyna et al., 2020); s = (Hutchings
1090 et al., 1997); t = (Rusch et al., 2011); u = (Louâpre et al., 2012); v = (Slade & Hutchings, 1987b); w = (Slade & Hutchings, 1987a);
1091 x = (Bittebiere et al., 2019); y = (Colom & Baucom, 2020).

1092
1093



1094

1095

1096

1097

1098

1099

1100

1101

Figure 2. Partitioning of individual trait variance across four nested scales, obtained using the Partvar procedure on the data collected from 2017 to 2020. All data were log or square-root transformed when needed, and standardized to equalize the weight of each trait dimension. * $p < 0.05$ (bootstrapping, $n = 1430$). SLA, specific leaf area; LDMC, leaf dry matter content. Time: four years of sampling; space: three sampled sites *i.e.* Cap Molloy, Isthme Bas, and Cap Ratmanoff; Between species: the six different macrophyte species of pond communities *i.e.* *Callitriche antarctica*, *Limosella australis*, *Juncus scheuchzerioides*, *Ranunculus biternatus*, *R. pseudotrullifolius*, and *R. moseleyi*; Within species: intraspecific trait variation.

1102

1103

Trait	Explanatory variables of selected model					
	<i>n</i>	Predictor	F	d.f.	Model predictors coefficient	Part of explained variance (%)
<i>Resource acquisition</i>						
All	99	Species identity	8.80***	5		
		PCAwater	8.67***	1		
		Water depth	2.64*	1		
Height	101	Functional diversity	3.49.	1	0.12	1.55
		pH	8.77**	1	-0.20	3.15
		Species richness	3.30.	5	-0.13	1.46
		Species identity	23.62***	1		52.24
		Water depth	3.06.	1	0.12	1.35
		Residuals			91	
SLA	102	FCAplants 2	3.50.	1	-0.14	1.97
		Electric conductivity	8.66**	1	0.27	4.86
		Water temperature	4.56*	1	0.18	2.57
		Phytoplankton concentration	2.38 ^{n.s.}	1	-0.13	1.34
		PCAwater	12.24***	1	-0.32	6.88
		Species identity	11.54***	5		32.40
		Residuals			89	
Internode length	100	pH	2.33 ^{n.s.}	1	-0.11	1.78
		Species richness	3.71.	1	-0.15	2.83
		Species identity	5.79***	5		22.20
		Water depth	6.22*	1	0.19	5.03
		Residuals			89	
Maximum root length		FCAplants 1	9.21**	1	0.24	4.37
		FCAplants 3	3.61.	1	-0.14	1.71
		Water temperature	7.70**	1	-0.21	3.66
		Species identity	19.10***	5		45.39
		Water depth	2.42	1	0.11	1.18
		Residuals			92	
<i>Resource conservation</i>						
All	97	Species identity	8.34***	5		
		Phytoplankton concentration	3.33*	1		
		FCAplants 3	7.13***	1		
		PCAsediments	5.37**	1		


LDMC		FCApplants 3	9.87**	1	-0.23	5.90
		Functional diversity	2.65 ^{n.s.}	1	0.11	1.58
		PCAsediments	7.62**	1	-0.22	4.55
		Species identity	11.22***	5		33.56
		Residuals		91		54.41
Specific internode mass	98	FCApplants 1	5.60*	1	-0.17	2.71
		Water conductivity	2.89.	1	-0.12	1.39
		Phytoplankton concentration	7.33**	1	-0.18	7.57
		Species identity	18.37***	5		44.37
		Residuals		91		43.96
Specific root mass	102	pH	4.11*	1	-0.15	1.92
		Phytoplankton concentration	22.42***	1	-0.41	10.49
		Pielou evenness	3.77.	1	-0.13	1.76
		PCAwater	7.26**	1	-0.24	3.40
		Species identity	17.03***	5		39.85
		Residuals		91		42.58

1105 Water and sediment nutrient dimensions were reduced to first principal component axes (PCAwater and PCAsediments;
1106 Appendices S3 and S4). To account for the influence of within-pond species abundance, FCAplants represents the first three
1107 axes from a factorial correspondence analysis (Appendix S5). Variables in bold had significant effects. * $p < 0.05$, ** $p < 0.01$,
1108 *** $p < 0.001$, "." $0.05 < p < 0.1$, n.s. not significant ($p > 0.1$), multivariate (all) and individual analyses of variance of the most
1109 probable models (2020 data). n , number of points used to calculate the relationship. Height and maximum root length were
1110 square-root transformed and all other data were log transformed. FCAplants 1 is related positively to the abundance of *R.*
1111 *moseleyi* and negatively to the abundance of *C. antarctica*, FCAplants 2 is related positively to the abundance of *R.*
1112 *pseudotrullifolius* and negatively to the abundance of *L. australis*, and FCAplants 3 is related positively to the abundance of *L.*
1113 *australis* and negatively to the abundance of *R. biternatus*.

1114

1115

1116 **Figure 3.** a) Structural equation model representing the direct and indirect pond-level effects of trait values on individual
 1117 performance of all species [November 2020 data, reduced model (lavaan package, variable selection based on Akaike
 1118 information criteria); full model presented in Appendix S8]. Trait values were standardized for each species to eliminate
 1119 variance due to species identity. Standardized path coefficients are given with p values: *** $p < 0.001$, ** $p < 0.01$, " $.$ " $0.05 < p$
 1120 < 0.1 , no asterisk $p \geq 0.1$. Line widths represent relationship strength. R^2 values reflect the proportion of response variable
 1121 variance explained. b) Absolute direct and indirect effects of trait values on individual performance.



1122