

### Herd factors associated with levels of parasitism in alternative pig farms

Maxime Delsart, Nicolas Rose, Barbara Dufour, Jean-Michel Répérant, Radu Blaga, Françoise Pol, Christelle Fablet

#### ▶ To cite this version:

Maxime Delsart, Nicolas Rose, Barbara Dufour, Jean-Michel Répérant, Radu Blaga, et al.. Herd factors associated with levels of parasitism in alternative pig farms. Annual meeting of the European Veterinary Parasitology, Jun 2023, Paris, France. 2023. hal-04383298

#### HAL Id: hal-04383298 https://hal.science/hal-04383298

Submitted on 9 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# HERD FACTORS ASSOCIATED WITH LEVELS OF ANSES PARASITISM IN ALTERNATIVE PIG FARMS



DELSART Maxime<sup>1</sup>, ROSE Nicolas<sup>2</sup>, DUFOUR Barbara<sup>1</sup>, REPERANT Jean-Michel<sup>3</sup>, BLAGA Radu<sup>4</sup>, POL Françoise<sup>5</sup>, FABLET Christelle<sup>2</sup>

1- Anses, École Nationale Vétérinaire d'Alfort, Laboratoire de Santé Animale USC EPIMAI, 94700 Maisons-Alfort, France

2- Anses Ploufragan-Plouzané-Niort, Unité Épidémiologie, Santé et Bien-Être, 22440 Ploufragan, France

3- Anses Ploufragan-Plouzané-Niort, Unité Virologie, Immunologie, Parasitologie Aviaires et Cunicoles, 22440 Ploufragan, France

4- Anses, INRAE, École Nationale Vétérinaire d'Alfort, Laboratoire de Santé Animale, BIPAR, 94700 Maisons-Alfort, France

5- ONIRIS, 101 Rte de Gachet, 44300 Nantes, France

### **BACKGROUND & AIM**

Pig farms on litter or with outdoor access (runs, free-range) are widely popular with consumers and citizens, particularly for reasons of animal welfare<sup>1</sup>. However, parasitism may constitute a critical point in these kinds of farms.





This study is part of a large research project (PIGAL) jointly studying animal health, biosecurity, animal welfare and veterinary public health in alternative pig farms. It aims to identify the herd factors associated with the level of parasite infestation.

## RESULTS

Farm description:

- 80 farrow-to-finish farms, 20
   fatteners and 12 farrow-to
   wean herds
   Number of farms
   0
   1-2
- 69% of the herds were

   69% of the herds were
   7-10
   7-10
   11-14

   the French quality program

   Label Rouge'.

Figure 1: Distribution of farms

Coccidia oocysts were observed in pig faeces in the largest proportion of farms (84%) followed by eggs of strongyles (55%), *Trichuris suis* (32%) and *A. suum* (16%). The rates of *A. suum* and *T. gondii* seropositive farms were 89% and 71% respectively. Hygiene and especially decontamination of facilities were factors associated with low levels of parasitism in farms (Figure 2). Conversely, keeping animals outdoors or on litter, poor building maintenance, farms with small numbers of animals and summer were parameters associated with high levels of parasitism. The use of multiple anthelmintic treatments on growing pigs was associated with low levels of *T. suis* egg infestation and high levels of *A. suum* seroprevalence.

## CONCLUSIONS

This study highlighted the internal parasite burden in alternative farms. Even if some factors are not under farmers' control (season, outdoor rearing), there is still some room for improvement via hygiene and appropriate use of treatments.

### **MATERIAL & METHODS**



Figure 2: Results of the multiple correspondence analysis between the descriptive variables of the farm, the season during which the visit was made and the classification of the farms according to (A) their level of parasitism, (B) their level of parasitism with regard to strongyles, (C) to *Trichuris suis*, (D) to coccidia, (E) to their level of seropositivity to *Ascaris suum* or to (F) *Toxoplasma gondii* on animals.

Abbreviations: Cluster TP: 1, low level of parasitism; 2, high level of parasitism; Cluster Strg: 1, low level of strongyles; 2, high level of strongyles; Cluster Ts: 1, low level of T. suis; 2, high level of T. suis; Cluster Co: 1, low level of coccidia; 2, high level of coccidia; Cluster Ser2: 0, low level of seropositivity to A. suum; 1, high level of seropositivity to A. suum; Cluster Tg: 1, low level of seropositivity to T. gondii; 2, high level of seropositivity to T. gondii; BuildMaint: correct maintenance of farm building (0, No; 1, Yes); **Dw\_Q**; deworming in quarantine (0, No; 1, Yes); Farm\_size: farm size (1, ≤50 sows; 2, between 50 and 100 sows; 3, >100 sows); Farm\_Type: Type of farm (1, Farrow-to-finish; 2, Partial farrow-to-finish); Farrow\_Lit: presence of litter in farrowing pens (0, No; 1, Yes); Farrow\_Out: farrowing outdoors (0, No; 1, Yes); Farrow\_outyard: farrowing indoors with an outdoor courtyard (0, No; 1, Yes); Fat\_Out: fatteners units outdoors (0, No; 1, Yes); **Fat\_Outyard**: fatteners units indoors with an outdoor courtyard (0, No; 1, Yes); **Int\_Bt**: interval in weeks between batches of animals (1, <5 weeks; 2, 5 or 6 weeks, 3, >6 weeks); Int\_DW\_Copro22: Interval in weeks between faecal samples taken from finishers and the last deworming treatment administered to these animals (1, <6 weeks; 2, ≥6 weeks); **M&P\_Out**: mating and pregnancy outdoors (0, No; 1, Yes); **M&P\_Outyard**: mating and pregnancy indoors with an outdoor courtyard (0, No; 1, Yes); **M&P\_Lit**: mating and pregnancy on litter (0, No; 1, Yes); **NbDw\_Gwp2**: number of deworming treatments on grower/finisher pigs  $(1, \leq 1; 2, \geq 2)$ .; **NbDw\_Gwp3**: number of deworming treatments on grower/finisher pigs  $(1, \leq 1; 2, 2 \text{ and } 3, \geq 3);$ **NbDw\_Sow**: number of deworming treatments on grower/finisher pigs  $(1, \le 1; 2, \ge 2)$ .; **Q\_In**: indoor quarantine (0, No; 1, Yes); Q\_Out: quarantine outdoors (0, No; 1, Yes); Season: the season in which the visit and the samples were taken; Wsyst\_Nur: nursery rooms washed systematically between each batch (0, No; 1, Yes); Wsyst\_Fat: fatteners rooms washed systematically between each batch (0, No; 1, Yes); Wsyst\_Farrow: farrowing rooms washed systematically between each batch (0, No; 1, Yes).

The study was conducted in 112 alternative pig farms (size > 30 sows and/or 100 pigs), in continental France (Figure 1). **Samples**:

- Faeces from 10 sows, 10 growers (10-12 weeks old) and/or 10 finishers (≥22 weeks old)
- Blood samples from 10 sows and/or 10 finishers (≥22 weeks old)

#### Laboratory analyses:

• Coprology (McMaster technique using saturated NaCl as a flotation fluid)



Data on the facilities and management of the farm were collected during an on-farm visit and were analyzed using multiple correspondence analysis to determine farm profiles regarding infestation and farm characteristics.

DELSART M., POL F., DUFOUR B., ROSE N., FABLET C. (2020) Pig Farming in Alternative Systems: Strengths and Challenges in Terms of Animal Welfare, Biosecurity, Animal Health and Pork Safety. Agriculture 10(7), 261
 DUBEY J.P., DESMONTS G. (1987) Serological responses of equids fed Toxoplasma gondii oocysts. Equine Vet. J. 19(4), 337-339

### **Contact :** Maxime DELSART

+331.43.96.72.98 - maxime.delsart@vet-alfort.fr

Financial partners: Avril Nutrition Animale SAS, Cooperl, Elanco, FranceAgriMer, Herta, Inaporc, Sofral le Gouessant, and Zoetis France

With technical help: Chantal Benoit, Amandine Blaizot, Édouard Boudin, Jean-François Da-Costa, Virginie Dorenlor, Florent Eono, Éric Eveno, Stéphane Kerphérique, Gilles Poulain, Marie Souquière, Martine Thomas-Hénaff and Sandra Thoumire

