



**HAL**  
open science

***Pseudomonas aeruginosa* antimicrobial susceptibility profiles, resistance mechanisms and international clonal lineages: update from ESGARS-ESCMID/ISARPAE Group**

Antonio Oliver, Estrella Rojo-Molinero, Jorge Arca-Suarez, Yeşim Beşli, Pierre Bogaerts, Rafael Cantón, Cansu Cimen, Peter Croughs, Olivier Denis, Christian Giske, et al.

► **To cite this version:**

Antonio Oliver, Estrella Rojo-Molinero, Jorge Arca-Suarez, Yeşim Beşli, Pierre Bogaerts, et al.. *Pseudomonas aeruginosa* antimicrobial susceptibility profiles, resistance mechanisms and international clonal lineages: update from ESGARS-ESCMID/ISARPAE Group. *Clinical Microbiology and Infection*, In press, 10.1016/j.cmi.2023.12.026 . hal-04384072

**HAL Id: hal-04384072**

**<https://hal.science/hal-04384072>**

Submitted on 10 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 ***Pseudomonas aeruginosa* antimicrobial susceptibility profiles, resistance**  
2 **mechanisms and international clonal lineages: update from ESGARS-**  
3 **ESCMID/ISARPAE Group**

4 **Antonio Oliver<sup>1</sup>, Estrella Rojo-Molinero<sup>1</sup>, Jorge Arca-Suarez<sup>2</sup>, Yeşim Beşli<sup>3</sup>**  
5 **Pierre Bogaerts<sup>4</sup>, Rafael Cantón<sup>5</sup>, Cansu Cimen<sup>6</sup>, Peter D Croughs<sup>7</sup>, Olivier**  
6 **Denis<sup>8</sup>, Christian G. Giske<sup>9</sup>, Tíscar Graells<sup>10</sup>, Te-Din Daniel Huang<sup>4</sup>, Bogdan**  
7 **I. Iorga<sup>11</sup>, Onur Karatuna<sup>12</sup>, Béla Kocsis<sup>13</sup>, Andreas Kronenberg<sup>14</sup>, Carla**  
8 **López-Causapé<sup>1</sup>, Surbhi Malhotra-Kumar<sup>15</sup>, Luis Martínez Martínez<sup>16</sup>,**  
9 **Annarita Mazzariol<sup>17</sup>, Sylvain Meyer<sup>18</sup>, Thierry Naas<sup>19</sup>, Daan W Notermans<sup>20</sup>,**  
10 **Jesús Oteo-Iglesias<sup>21</sup>, Torunn Pedersen<sup>22</sup>, Mateja Pirš<sup>23</sup>, Patricia Poeta<sup>24</sup>,**  
11 **Laurent Poirel<sup>25</sup>, Spyros Pournaras<sup>26</sup>, Arnfinn Sundsfjord<sup>22, 27</sup>, Dora Szabó<sup>13,</sup>**  
12 **<sup>28</sup>, Arjana Tambić-Andrašević<sup>29</sup>, Rossitza Vatcheva-Dobrevska<sup>30</sup>, Astra**  
13 **Vitkauskiene<sup>31</sup> and Katy Jeannot<sup>32</sup> on behalf of ESGARS-ISARPAE**  
14 **members.**

15 <sup>1</sup>Servicio de Microbiología, Hospital Universitario Son Espases, Instituto de  
16 Investigación Sanitaria Illes Balears (IdISBa), Palma de Mallorca, Spain; CIBER de  
17 Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos III, Madrid, Spain.

18 <sup>2</sup>Servicio de Microbiología and Instituto de Investigación Biomédica A Coruña (INIBIC),  
19 Complejo Hospitalario Universitario A Coruña, A Coruña, Spain; CIBER de  
20 Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos III, Madrid, Spain.

21 <sup>3</sup>Department of Medical Microbiology, Amerikan Hastanesi, Istanbul, Turkey.

22 <sup>4</sup>National Center for Antimicrobial Resistance in Gram-, CHU UCL Namur, Yvoir,  
23 Belgium.

24 <sup>5</sup>Servicio de Microbiología, Hospital Universitario Ramón y Cajal-IRYCIS, Madrid,  
25 Spain; CIBER de Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos  
26 III, Madrid, Spain.

27 <sup>6</sup>Institute for Medical Microbiology and Virology, University of Oldenburg, Oldenburg,  
28 Germany. Department of Medical Microbiology and Infection Prevention, University of  
29 Groningen, University Medical Center Groningen, Groningen, The Netherlands

30 <sup>7</sup>Department of Medical Microbiology and Infectious Diseases, Erasmus Medical  
31 Center, Rotterdam, the Netherlands.

32 <sup>8</sup>Department of Microbiology, CHU Namur site-Godinne, Université Catholique de  
33 Louvain, Yvoir, Belgium; Ecole de Santé Publique, Université Libre de Bruxelles,  
34 Brussels, Belgium.

35 <sup>9</sup>Department of Clinical Microbiology, Karolinska University Hospital; Department of  
36 Laboratory Medicine, Division of Clinical Microbiology, Karolinska Institutet, Solna,  
37 Stockholm, Sweden.

38 <sup>10</sup>Department of Neurobiology, Care Sciences and Society (NVS), Division of Family  
39 Medicine and Primary Care, Karolinska Institutet, Huddinge, Stockholm, Sweden.

40  
41 <sup>11</sup>Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, Gif-  
42 sur-Yvette, France.

43 <sup>12</sup>EUCAST Development Laboratory, Clinical Microbiology, Central Hospital, Växjö,  
44 Sweden.

45 <sup>13</sup>Institute of Medical Microbiology, Semmelweis University, Budapest, Hungary.

46 <sup>14</sup>Institute for Infectious Diseases, University of Bern, Bern, Switzerland.

47 <sup>15</sup>Laboratory of Medical Microbiology, Vaccine & Infectious Disease Institute, University  
48 of Antwerp, Antwerpen, Belgium.

49 <sup>16</sup>Unidad de Microbiología, Hospital Universitario Reina Sofía, Departamento de  
50 Química Agrícola, Edafología y Microbiología, Universidad de Córdoba, e Instituto  
51 Maimonides de Investigación Biomédica de Córdoba (IMIBIC), Spain; CIBER de  
52 Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos III, Madrid, Spain.

53 <sup>17</sup>Microbiology and Virology Section, Department of Diagnostic and Public Health,  
54 University of Verona, Verona, Italy.

55 <sup>18</sup>Inserm UMR 1092 and Université of Limoges, Limoges, France.

56 <sup>19</sup>Laboratoire Associé du Centre National de Référence de la Résistance aux  
57 Antibiotiques: Entérobactéries Résistantes aux Carbapénèmes, Le Kremlin-Bicêtre,  
58 France; Université Paris-Saclay, Équipe INSERM ReSIST, Faculté de Médecine.

59 <sup>20</sup>Centre for Infectious Disease Control. National Institute for Public Health and the  
60 Environment (RIVM), Bilthoven, The Netherlands.

61 <sup>21</sup>Reference and Research Laboratory in Resistance to Antibiotics and Infections  
62 Related to Healthcare, National Centre for Microbiology, Instituto de Salud Carlos III,  
63 Madrid, Spain; CIBER de Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud  
64 Carlos III, Madrid, Spain.

65 <sup>22</sup>Norwegian National Advisory Unit on Detection of Antimicrobial Resistance,  
66 Department of Microbiology and Infection Control, University Hospital of North Norway,  
67 Tromsø, Norway.

68 <sup>23</sup>Institute of Microbiology and Immunology, Faculty of Medicine, University of  
69 Ljubljana, Ljubljana, Slovenia.

70 <sup>24</sup>MicroART-Microbiology and Antibiotic Resistance Team, Department of Veterinary  
71 Sciences, University of Trás-os-Montes and Alto Douro (UTAD), Vila Real, Portugal.;  
72 Associated Laboratory for Green Chemistry (LAQV-REQUIMTE), University NOVA of  
73 Lisboa, Lisboa, Portugal; Veterinary and Animal Research Centre (CECAV), University

74 of Trás-os-Montes and Alto Douro (UTAD), Vila Real, Portugal; Associate Laboratory  
75 for Animal and Veterinary Sciences (AL4AnimalS), Vila Real, Portugal.

76 <sup>25</sup>Emerging Antibiotic Resistance Unit, Medical and Molecular Microbiology,  
77 Department of Medicine, University of Fribourg, Fribourg, Switzerland; Swiss National  
78 Reference Center for Emerging Antibiotic Resistance, Fribourg, Switzerland.

79 <sup>26</sup>Laboratory of Clinical Microbiology, Attikon University Hospital, Medical School,  
80 National and Kapodistrian University of Athens, Athens, Greece.

81 <sup>27</sup>Research Group on Host-Microbe Interactions, Department of Medical Biology, UiT  
82 The Arctic University of Norway, Tromsø, Norway.

83 <sup>28</sup>Human Microbiota Study Group, Semmelweis University-Eötvös Lóránd Research  
84 Network, Budapest, Hungary.

85 <sup>29</sup>Department of Clinical Microbiology, University Hospital for Infectious Diseases,  
86 Zagreb, Croatia; School of Dental Medicine, University of Zagreb, Zagreb, Croatia.

87 <sup>30</sup>Department of Microbiology and Virology, University Hospital 'Tsaritsa Yoanna- ISUL'  
88 Sofia, Bulgaria.

89 <sup>31</sup>Department of Laboratory Medicine, Faculty of Medicine, Medical Academy,  
90 Lithuanian University of Health Science, Kaunas, Lithuania.

91 <sup>32</sup>Laboratoire de Bactériologie, Centre Hospitalier Universitaire de Besançon,  
92 Besançon, France; Laboratoire associé du Centre National de Référence de la  
93 Résistance aux Antibiotiques, Centre Hospitalier Universitaire de Besançon, France;  
94 Chrono-environnement UMR 6249, CNRS, Université Franche-Comté, Besançon,  
95 France.

96

97 **Abstract**

98

99 **Scope:** *Pseudomonas aeruginosa*, a ubiquitous opportunistic pathogen  
100 considered one of the paradigms of antimicrobial resistance, is among the main  
101 causes of hospital-acquired and chronic infections associated with significant  
102 morbidity and mortality. This growing threat results from the extraordinary  
103 capacity of *P. aeruginosa* to develop antimicrobial resistance through  
104 chromosomal mutations, the increasing prevalence of transferable resistance  
105 determinants (such as the carbapenemases and the extended spectrum  $\beta$ -  
106 lactamases), and the global expansion of epidemic lineages. The general  
107 objective of this initiative is to provide a comprehensive update of *P. aeruginosa*  
108 resistance mechanisms, especially for the extensively drug-resistant (XDR)/  
109 difficult to treat resistance (DTR) international high-risk epidemic lineages, and  
110 how the recently approved  $\beta$ -lactams and  $\beta$ -lactam/  $\beta$ -lactamase inhibitor  
111 combinations may affect resistance mechanisms and the definition of  
112 susceptibility profiles. **Methods:** To address this challenge, the European Study  
113 Group for Antimicrobial Resistance Surveillance (ESGARS) from the European  
114 Society of Clinical Microbiology and Infectious Diseases (ESCMID) launched  
115 the “Improving Surveillance of Antibiotic-Resistant *Pseudomonas aeruginosa* in  
116 Europe” (ISARPAE) initiative in 2022, supported by the Joint programming  
117 initiative on antimicrobial resistance (JPIAMR) network call and included a panel  
118 of over 40 researchers from 18 European Countries. Thus, an ESGARS-  
119 ISARPAE position paper was designed and the final version agreed after four  
120 rounds of revision and discussion by all panel members. **Questions addressed**  
121 **in the position paper:** To provide an update on (i) the emerging resistance  
122 mechanisms to classical and novel antipseudomonal agents, with a particular  
123 focus on  $\beta$ -lactams, (ii) the susceptibility profiles associated with the most  
124 relevant  $\beta$ -lactam resistance mechanisms, (iii) the impact of the novel agents  
125 and resistance mechanisms on the definitions of resistance profiles and (iv)  
126 the globally expanding XDR/DTR high-risk lineages and their association with  
127 transferable resistance mechanisms. **Implication:** The evidence presented  
128 herein can be used for coordinated epidemiological surveillance and decision-  
129 making at the European and global level.

130

131 **Scope and context**

132 *Pseudomonas aeruginosa*, a ubiquitous opportunistic pathogen  
133 considered one of the paradigms of antimicrobial resistance, is among the main  
134 causes of hospital-acquired and chronic infections associated with significant  
135 morbidity and mortality (1). Accordingly, *P. aeruginosa* infections are estimated  
136 to be associated with over 300,000 annual deaths and are at the top of the  
137 WHO priority list for the need for research and development of new antibiotics  
138 (2,3). This growing threat results from the extraordinary capacity of this  
139 pathogen to develop antimicrobial resistance through chromosomal mutations  
140 and from the increasing prevalence of transferable resistance determinants,  
141 particularly those encoding carbapenemases or extended-spectrum  $\beta$ -  
142 lactamases (ESBLs) (4,5). Combinations of such mechanisms lead to  
143 concerning and complex resistance profiles, defined by the European Centre for  
144 Disease Prevention and Control (ECDC) and the Centers for Disease Control  
145 and Prevention (CDC) as multidrug-resistant (MDR), extensively drug-resistant  
146 (XDR), and pandrug-resistant (PDR), while the Infectious Diseases Society of  
147 America/National Institutes of Health (IDSA/NIH) defines them as difficult-to-  
148 treat resistance (DTR) (6,7). *P. aeruginosa* possesses a non-clonal epidemic  
149 population structure, comprising a limited number of widespread lineages,  
150 selected from a background of numerous rare and unrelated genotypes  
151 recombined at high frequency (8). In fact, several surveys have provided  
152 evidence for the existence of XDR/DTR international high-risk clonal lineages,  
153 which have disseminated in hospitals worldwide (9–11). Beyond classical  
154 molecular epidemiology analysis and phenotypic assessment of resistance  
155 mechanisms, whole genome sequencing (WGS) studies are providing pertinent  
156 information to elucidate the complex and evolving resistome of MDR/XDR/DTR  
157 *P. aeruginosa* high-risk lineages (12–15).

158 The recent introduction of novel  $\beta$ -lactam/ $\beta$ -lactamase inhibitor  
159 combinations (BLBLIs) such as ceftolozane/tazobactam, ceftazidime/avibactam,  
160 meropenem/vaborbactam or imipenem/relebactam and the siderophore-  
161 cephalosporin cefiderocol, has contributed to mitigate, to some extent, the  
162 problem of XDR/DTR *P. aeruginosa* (16–19). These agents exhibit enhanced  
163 stability against intrinsically- and chromosomally-encoded  $\beta$ -lactam resistance

164 mechanisms in *P. aeruginosa*, such as overexpression of the AmpC  $\beta$ -  
165 lactamase encoding gene, overproduction of efflux pumps, or inactivation of the  
166 OprD porin. However, they are not exempt from resistance development  
167 through emerging mutational mechanisms (20–24). These include modification  
168 (quantitative or qualitative) of AmpC hydrolytic activity or efflux pumps  
169 substrate specificity, which were observed shortly after their introduction into  
170 clinical practice. Moreover, BLBLIs are not currently effective against the most  
171 potent transferable carbapenemases, particularly class B or metallo- $\beta$ -  
172 lactamases [MBLs] such as VIM, IMP or NDM enzymes (25). Consequently,  
173 use of BLBLIs could lead to the selection of these concerning resistance  
174 mechanisms (26). Besides the approved options, several novel BLBLIs are  
175 undergoing clinical trials (25). These agents, such as aztreonam/avibactam,  
176 cefepime/zidebactam or cefepime/taniborbactam, promise additional  
177 therapeutic choices and the ability to counteract already established resistance  
178 mechanisms (17).

179 The introduction of novel BLBLIs is therefore significantly broadening the  
180 range of treatment options for XDR/DTR *P. aeruginosa* infections(17,25).  
181 However, this expansion will also have a major impact on antimicrobial  
182 resistance epidemiology, including both novel and existing mutation-driven  
183 resistance mechanisms, transferable resistance determinants and epidemic  
184 high-risk clonal lineages. A comprehensive understanding of *P. aeruginosa*  
185 resistance mechanisms and susceptibility profiles, especially of the XDR/DTR  
186 high-risk lineages, and how these promising novel agents may affect resistance  
187 mechanisms and, in turn, the definition of resistance profiles, is needed to have  
188 a common ground and may help to anticipate and coordinate epidemiological  
189 information in the future.

#### 190 **Questions addressed in the position paper**

191 To address this challenge, the European Study Group for Antimicrobial  
192 Resistance Surveillance (ESGARS) from the European Society of Clinical  
193 Microbiology and Infectious Diseases (ESCMID) launched the “Improving  
194 Surveillance of Antibiotic-Resistant *Pseudomonas aeruginosa* in Europe”  
195 (ISARPAE) initiative in 2022, supported by the Joint programming initiative on

196 antimicrobial resistance (JPIAMR) network. Thus, this position document from  
197 the ESGARS-ISARPAE Group aimed to provide an update on (i) the emerging  
198 resistance mechanisms to classical and novel anti-pseudomonal agents, with a  
199 particular focus on  $\beta$ -lactams, (ii) the susceptibility profiles associated with the  
200 most relevant  $\beta$ -lactam resistance mechanisms, (iii) the impact of the novel  
201 agents and resistance mechanisms on the definitions of resistance profiles, and  
202 (iv) the globally expanding XDR/DTR high-risk lineages and their association  
203 with transferable  $\beta$ -lactamases.

## 204 **Methods**

205 All ESGARS-ESCMID members were contacted and invited to participate in the  
206 ISARPAE initiative, according to their interest and experience in the topic. This  
207 resulted in the generation of a panel of over 40 researchers from 18 European  
208 countries in June 2022. The panel agreed the above objectives to be addressed  
209 in the position paper and AO and ERM prepared a first draft of the documented  
210 after extensive literature review helped by other panel members. In July 2023  
211 the first draft of the document was sent to all ISARPAE members for revision  
212 and specific contributions, leading to a second draft version that was  
213 extensively revised and discussed during an ISARPAE hybrid (onsite/online)  
214 meeting that took place at Hospital Son Espases-IdISBa (Mallorca, Spain) on  
215 September 8<sup>th</sup> 2023. The third resulting draft was then sent for review by panel  
216 members and final version of the document was approved in October 6<sup>th</sup>2023.

## 217 **Emerging resistance mechanisms to classical and novel anti-** 218 **pseudomonal agents and associated susceptibility profiles**

219 **Table 1** shows the main categories and agents showing  
220 antipseudomonal activity, including those recently introduced and those that will  
221 be clinically available in the next few years, and presents the respective  
222 mutation-driven and horizontally-acquired resistance mechanisms. On the other  
223 hand, **Figure 1** shows the susceptibility profiles associated with the most  
224 relevant  $\beta$ -lactam resistance mechanisms in *P. aeruginosa*.

225

## 226 ***Pseudomonas aeruginosa* $\beta$ -lactam resistome**



227

228 *Pseudomonas aeruginosa* is intrinsically resistant to aminopenicillins,  
229 alone and combined with clavulanic acid, as well as to most of the older  
230 cephalosporins, notably including the third generation cephalosporin  
231 cefotaxime, due to the production of an inducible AmpC  $\beta$ -lactamase (27).  
232 Moreover, AmpC plays a major role in the basal resistance level (MIC) of *P.*  
233 *aeruginosa* to the potent AmpC inducer imipenem. On the other hand, the  
234 constitutive expression of the efflux pump MexAB-OprM plays a major role in  
235 the basal resistance level to most other  $\beta$ -lactams except imipenem.

236

237 The most frequent mutation-driven resistance mechanism to classical  
238 antipseudomonal penicillins (such as piperacillin) and cephalosporins (such as  
239 ceftazidime or cefepime) is the overproduction of the chromosomal  
240 cephalosporinase AmpC, involving a large number of genes belonging to cell-  
241 wall recycling regulatory pathways (28). Notably, among these genes, the  
242 mutational inactivation of *dacB*, encoding the nonessential penicillin-binding  
243 protein (PBP] PBP4 and *ampD*, encoding a N-acetyl-muramyl-L-alanine  
244 amidase, have been found to be the most frequent cause of derepressed *ampC*  
245 gene expression, and subsequent broad-spectrum  $\beta$ -lactam resistance (29,30).  
246 Additionally, specific point mutations causing a conformational change in the  
247 transcriptional regulator AmpR, leading to *ampC* upregulation and resistance to  
248 broad-spectrum  $\beta$ -lactams, have been noted among clinical strains. These  
249 mutations include the D135N amino acid replacement, described in several  
250 species (28) and the G154H mutation linked to the disseminated MDR/XDR  
251 ST175 high-risk lineage (14). Mutation of several other genes, including those  
252 encoding amidases (AmpDh2 and AmpDh3), PBPs, such as PBP5 or PBP7,  
253 lytic transglycosylases, MPL, or NuoN have also been shown to enhance *ampC*  
254 expression, either alone or in combination with other mutations. Nevertheless,  
255 their impact on  $\beta$ -lactam resistance among clinical strains still needs to be  
256 further analysed (28).

257 In addition to *ampC* overexpression, recent studies have revealed that  
258 increased levels of  $\beta$ -lactam resistance, involving the novel BLBLIs  
259 ceftolozane/tazobactam and ceftazidime/avibactam, may result from mutations  
260 leading to the modification of the catalytic center of AmpC, currently mainly

261 occurring in (up to 10-15%) patients treated with these agents (20,31–33).  
262 Additional studies identified diverse AmpC variants associated with high-level  
263 resistance to BLBLIs, including the above mentioned ceftolozane/tazobactam  
264 and ceftazidime/avibactam, in a small proportion (around 1%) of clinical *P.*  
265 *aeruginosa* isolates (34). Over 500 variants of those AmpC enzymes, also  
266 called *Pseudomonas* Derived Cephalosporinases (PDC), have been described  
267 so far, including those associated with increased ceftolozane/tazobactam and  
268 ceftazidime/avibactam resistance. Moreover, some of these variants, such as  
269 those showing the L320P substitution, have a significant impact on cefiderocol  
270 MICs, but only a marginal effect on susceptibility to ceftolozane/tazobactam and  
271 ceftazidime/avibactam (35). An updated database of PDC variants is  
272 maintained at IdISBa and is freely available at  
273 ([https://arpbigidisba.com/pseudomonas-aeruginosa-derived-cephalosporinase-](https://arpbigidisba.com/pseudomonas-aeruginosa-derived-cephalosporinase-pdc-database/)  
274 [pdc-database/](https://arpbigidisba.com/pseudomonas-aeruginosa-derived-cephalosporinase-pdc-database/)) and at the Beta-Lactamase Data Base  
275 (<http://www.blbd.eu/BLDB.php?prot=C#PDC>) (36). Typically, the strains  
276 producing these AmpC variants show collateral susceptibility to imipenem  
277 (decreased MICs) and also to antipseudomonal penicillins such as piperacillin.  
278 Additionally, resistance development to ceftolozane/tazobactam and/or  
279 ceftazidime/avibactam may involve mutations leading to the structural  
280 modification of narrow spectrum OXA-2 and OXA-10 acquired oxacillinases  
281 (20,37,38). Interestingly, these mutations may lead to collateral susceptibility to  
282 meropenem. Thus, imipenem/relebactam, and to a lesser extent, cefiderocol,  
283 meropenem/vaborbactam and the novel combinations under development  
284 cefepime/zidebactam and cefepime/taniborbactam might be interesting options  
285 to treat infections by strains that have developed ceftolozane/tazobactam and/or  
286 ceftazidime/avibactam resistance through mutations in AmpC or OXA-2/10 (39).

287       Horizontally-acquired  $\beta$ -lactamase genes are obviously a major source of  
288 resistance, including to the novel  $\beta$ -lactams and BLBLI (**Figure 1**). An extensive  
289 revision of the nature and prevalence of the different horizontally-acquired  $\beta$ -  
290 lactamases detected in *P. aeruginosa* is beyond the scope of this document.  
291 However, globally, MBLs are arguably the most frequent carbapenemases in *P.*  
292 *aeruginosa*, but very large geographical differences in prevalence and nature  
293 have been documented (40,41). At European level, VIM, and particularly VIM-2,

294 are likely the most frequently reported enzymes, but with major differences  
295 across different countries, and with an increasing prevalence of NDM enzymes  
296 (42,43). Moreover, GES class A carbapenemases variants such as GES-5 are  
297 also increasingly reported in European countries (43,44). Classical  
298 antipseudomonal penicillins, cephalosporins and carbapenems lack significant  
299 activity and should be avoided against strains producing class A or MBL  
300 carbapenemases, even if MICs close to the clinical breakpoints are obtained for  
301 piperacillin/tazobactam, cefepime or even carbapenems for some VIM-2-  
302 producing isolates (12). Moreover, the production of MBLs is a frequent  
303 mechanism of resistance to ceftolozane/tazobactam, ceftazidime/avibactam,  
304 meropenem/vaborbactam and imipenem/relebactam (26). However, with a few  
305 exceptions such as some NDM variants, cefiderocol retains activity due to its  
306 higher stability against hydrolysis and efficient uptake through the iron transport  
307 systems (45). The combination of aztreonam with avibactam may also be a  
308 useful future alternative for MBL producing strains, particularly when additionally  
309 hyperproducing AmpC and/or coproducing acquired class A enzymes (46,47).  
310 Likewise, the novel combinations under development cefepime/zidebactam and  
311 cefepime/taniborbactam also remain active. The underlying mechanism for  
312 cefepime/zidebactam activity against MBL producing strains is based on the  
313 fact that zidebactam has direct antipseudomonal activity by targeting PBP2, and  
314 therefore provides synergy with  $\beta$ -lactams targeting PBP3 such as the  
315 cephalosporins (48). On the other hand, the activity of cefepime/taniborbactam  
316 relies on the fact that taniborbactam inhibits MBL hydrolytic activity, except for  
317 IMPs (49). In addition to these three antimicrobials (cefiderocol,  
318 cefepime/zidebactam and cefepime/taniborbactam), ceftazidime/avibactam, and  
319 to a lower extent imipenem/relebactam and meropenem/vaborbactam show  
320 activity against producers of Ambler class A carbapenemases (such as GES-5  
321 and KPCs) (50–52). However, the frequent concomitant OprD deficiency and/or  
322 MexAB-OprM overexpression limits the activity of imipenem/relebactam and  
323 meropenem/vaborbactam against clinical *P. aeruginosa* strains producing class  
324 A carbapenemases (52,53). On the other hand, resistance development to  
325 ceftazidime/avibactam caused by the selection of mutations within the catalytic  
326 site of KPC and GES enzymes has been described (54–56). Interestingly, these  
327 mutations restore carbapenem susceptibility (if the strain is not oprD deficient)

328 leading to an ESBL phenotype (54). In addition to those of classes A and B, a  
329 few cases of class D carbapenemase production have been reported in *P.*  
330 *aeruginosa*, including the epidemic dissemination OXA-198 in a hospital from  
331 Belgium (57).

332 In addition to  $\beta$ -lactamases, there is growing evidence on the role of  
333 target modification in *P. aeruginosa*  $\beta$ -lactam resistance. Of particular relevance  
334 are the mutations in *ftsI*, encoding PBP3, an essential class B PBP with  
335 transpeptidase activity (58). Indeed, data from cystic fibrosis (CF) patients  
336 (59,60), epidemic high-risk clonal lineages (12,14) as well as from *in vitro*  
337 studies (61) have shown that PBP3 is under strong mutational pressure, with  
338 specific mutations in this PBP contributing to  $\beta$ -lactam resistance development.  
339 R504C/H and F533L mutations are those being most commonly reported and  
340 located within the protein domains implicated in the formation and stabilization  
341 of the inactivating complex  $\beta$ -lactam-PBP3 (62). Moreover, these specific  
342 mutations have been documented to emerge *in vivo* during chronic respiratory  
343 infection in CF patients (59,60) and upon exposure to meropenem (61),  
344 aztreonam (63) and ceftazidime (64) *in vitro*. However, the detailed effect of  
345 PBP3 mutations on  $\beta$ -lactam resistance phenotypes needs to be further  
346 investigated using isogenic strains. Likewise, despite unique polymorphisms  
347 having been detected in some clinical strains for other PBPs, their potential role  
348 in  $\beta$ -lactam resistance still needs to be experimentally determined. Also  
349 noteworthy are the specific PBP2 mutations involved in resistance to  
350 zidebactam (65), that obviate the  $\beta$ -lactam enhancer activity of this BLI.

351 Other relevant components of the  $\beta$ -lactam mutational resistome are the  
352 genes encoding OprD and efflux pumps. The inactivation of OprD is known to  
353 be the most frequent imipenem resistance mechanisms in *P. aeruginosa*  
354 (66,67). OprD inactivation typically results from indels or nonsense mutations,  
355 including the Q142X mutation, characteristic of the widespread ST175 high-risk  
356 clonal lineage (14). Additionally, some amino-acid replacements have been  
357 associated with OprD-driven resistance, particularly in the CF setting (68).  
358 However, it should be noted that the presence of OprD inactivating mutations  
359 has also been identified in some carbapenem-susceptible isolates (69). On the  
360 other hand, imipenem resistance may also result from repression of *oprD*

361 caused by mutations in the MexEF-OprN efflux pump regulators (*mexS/T*) or  
362 the ParRS two-component system (70). Overexpression of MexAB-OprM,  
363 caused by mutation of several genes involved in its regulation (*mexR*, *nalC* or  
364 *nalD*) increases MICs of most  $\beta$ -lactams including meropenem but not  
365 imipenem, whereas overexpression of genes encoding MexXY (*mexZ*, *parRS*,  
366 *amgS* mutations) is involved in cefepime resistance (70)

367 Efflux pumps may also play a major role in resistance to the novel  
368 BLBLIs, not only because of their capacity to extrude the  $\beta$ -lactam components  
369 but, particularly, for their capacity to accommodate their partner  $\beta$ -lactamase  
370 inhibitor. Indeed MexAB-OprM overexpression plays a role in resistance to  
371 ceftazidime/avibactam, aztreonam/avibactam, cefepime/zidebactam,  
372 imipenem/relebactam, and meropenem/vaborbactam (65,71–73). Likewise,  
373 MexXY overexpression should also impact cefepime combinations with  
374 zidebactam or taniborbactam (65). Moreover, mutations leading to the  
375 modification of the substrate recognition domain of the efflux pump MexCD-  
376 OprJ have been shown to drive ceftolozane/tazobactam resistance  
377 development *in vivo* (23)

378 Additionally, another potentially relevant mutational  $\beta$ -lactam resistance  
379 mechanism is the selection of large [up to 600 kb] deletions affecting specific  
380 parts of the chromosome (61,64). Although the basis of the conferred resistance  
381 phenotype still needs to be further clarified, these mutants can be recognized by  
382 the characteristic brown pigment (pyomelanine) caused by the deletion of one  
383 of the included genes, *hmgA*, coding for a homogentisate-1,2-dioxygenase.  
384 These deletions has been documented in both *in vitro* evolved  $\beta$ -lactam-  
385 resistant mutants and CF isolates (61,74). However, the deletion of *hmgA* is not  
386 responsible for the resistance phenotype, which could be linked to the deletion  
387 of another of the affected genes, *galU*. This gene codes for a UDP-glucose  
388 pyrophosphorylase involved in the synthesis of the lipopolysaccharide (LPS)  
389 core. Indeed, analysis of transposon mutant libraries has revealed that  
390 inactivation of *galU* increases the MICs of ceftazidime and meropenem (75,76).

391 Lastly, specific cefiderocol resistance development mechanisms involve  
392 the selection of mutations in iron uptake systems, particularly in TonB-

393 dependent receptors such as *piuA/piuC*, *pirA/pirR* or *fptA* [pyochelin receptor]  
394 (35). Among these, mutations seem to be particularly frequent in *piuC*, an iron-  
395 dependent oxygenase involved in the expression of the adjacent *piuA* [or its  
396 homolog *piuD* depending on the strain] iron receptor. On the other hand,  
397 mutations in the *fptA* gene, despite being frequent, do not seem to have a direct  
398 significant impact on cefiderocol MICs, and thus selection might reflect adaptive  
399 mutations for growing in the presence of cefiderocol.

400

#### 401 ***Pseudomonas aeruginosa* aminoglycoside resistome**

402 Primary aminoglycoside resistance is typically linked to the production of  
403 horizontally-acquired aminoglycoside modifying enzymes, including  
404 acetyltransferases, adenylyltransferases and phosphoryltransferases, frequently  
405 co-transferred with ESBLs or carbapenemases (77). The specific pattern of  
406 aminoglycoside resistance depends on the specific enzymes involved, with  
407 amikacin showing an overall higher activity than tobramycin (78). However, the  
408 more recently described transferable 16S rRNA methylases, which modify the  
409 cellular target of aminoglycosides, are further concerning since they confer  
410 resistance to all clinically available members of this antibiotic family and are  
411 also cotransferred with ESBLs or carbapenemases (79–81).

412 On the other hand, the development of resistance to aminoglycosides  
413 has been particularly linked to the overexpression of genes encoding the  
414 MexXY-OprM system upon some mutations in the regulatory machinery.  
415 Indeed, mutational overexpression of this pump, mainly caused by *mexZ*,  
416 *amgS*, or *parRS* mutations, is very frequent among clinical isolates, from both  
417 CF patients and nosocomial infections (82,83). Moreover, recent studies show  
418 that the epidemic high-risk clone ST175 hyperproduces MexXY due to a  
419 specific mutation in *mexZ* (G195E) (14). However, recent data suggests that the  
420 aminoglycoside mutational resistome extends far beyond MexXY  
421 hyperproduction, and high-level resistance may result from the accumulation of  
422 multiple mutations. The involvement of several novel resistance determinants  
423 has been documented (84–86). Among them, is noteworthy *fusA1*, coding for  
424 the elongation factor G. Indeed, specific *fusA1* mutations have been linked to

425 aminoglycoside resistance *in vitro* (4,86) and among clinical, strains, particularly  
426 from CF patients (4,60,87–89). Moreover, the implication of *fusA1* mutations in  
427 aminoglycoside resistance has been demonstrated through site-directed  
428 mutagenesis (90).

429

### 430 ***Pseudomonas P. aeruginosa* fluoroquinolone resistome**

431

432 Fluoroquinolone resistance in *P. aeruginosa* is primarily driven by  
433 mutational mechanisms. The fluoroquinolone mutational resistome generally  
434 includes specific missense mutations in DNA gyrase (*gyrA* and/or *gyrB*) and  
435 topoisomerase IV (*parC* and/or *parE*) Quinolone Resistance-Determining  
436 Regions (QRDRs) (13,91). High-level fluoroquinolone resistance in *P.*  
437 *aeruginosa* high-risk lineages is nearly universal, and typically involves  
438 combinations of mutations in GyrA T83 and ParC S87 (12)QRDR mutations  
439 involved in fluoroquinolone resistance in CF might be more variable (60). It is  
440 also well-known that the mutational overexpression of efflux pumps modulates  
441 fluoroquinolone resistance [Table 1]. While the overexpression of MexAB-OprM  
442 and MexXY-OprM is globally frequent among clinical strains, its contribution to  
443 clinical fluoroquinolone resistance is likely to be modest (91)On the other hand,  
444 the mutational overproduction of MexEF-OprN or MexCD-OprJ is associated  
445 with clinical fluoroquinolone resistance. Although their prevalence has been  
446 considered low, except in the settings of CF chronic infections, recent data  
447 show that it might be higher than expected (68). Lastly, the transferable  
448 quinolone resistance determinant QnrVC has also been reported, linked to  
449 some epidemic strains producing acquired carbapenemases such as ST175  
450 and ST244 (92,93).

451

### 452 ***Pseudomonas aeruginosa* polymyxin resistome**

453

454 Due to its limited efficacy, toxicity and high ECOFF values (4 mg/L),  
455 colistin is not considered an optimal treatment for wild-type *P. aeruginosa*, at  
456 least in monotherapy [[www.eucast.org](http://www.eucast.org)]. Moreover, whereas the prevalence of  
457 polymyxin [colistin and polymyxin B] resistance is still globally low (<5%), it has

458 increased in the last years because of the frequent use of these last-resource  
459 antibiotics for the treatment of MDR/XDR/DTR nosocomial and CF isolates,  
460 particularly in countries with no access to novel BLBLIs (94). Polymyxin  
461 resistance results most frequently from the modification of the LPS caused by  
462 the addition of a 4-amino-4-deoxy-L-arabinose moiety in the lipid A structure  
463 (95,96). The involved mutations are frequently located in the PmrAB or PhoPQ  
464 two-component regulators, which lead to the activation of the *arnBCADTEF*  
465 operon (97). More recent studies have revealed that mutations in the ParRS  
466 two-component regulator, not only produce polymyxin resistance due to the  
467 activation of the *arnBCADTEF* operon, but also lead to a MDR phenotype  
468 determined by the hyperproduction of MexXY and the repression of *oprD* (98).  
469 Moreover, two additional two-component regulators, ColRS and CprRS, have  
470 also been determined to be involved in colistin resistance (99). The analysis of  
471 colistin resistance mechanisms among clinical strains is not always  
472 straightforward, since the presence of mutations in these two-component  
473 regulators is not always associated with clinical colistin resistance, probably  
474 denoting partial complementation between the different regulators (60,99,100).  
475 Moreover, recent *in vitro* evolution assays have revealed the implication of  
476 additional mutations in high level colistin resistance, facilitated by the  
477 emergence of *mutS* deficient mutator (phenotypes such as those occurring in  
478 LptD, LpxC or MigA (101). On the other hand, the role of phosphoethanolamine  
479 modification of LPS in *P. aeruginosa* seems marginal, including both, that are  
480 driven by intrinsic *eptA* gene expression (102) as well as that are driven by  
481 transferable determinants (103).

482

### 483 ***Pseudomonas aeruginosa* fosfomicin resistome**

484

485 Although not classified as an antipseudomonal agent (ECOFF of 256  
486 mg/L), fosfomicin has been considered in the last decade as a potentially  
487 useful antibiotic in urinary tract infections and as combined therapy for  
488 MDR/XDR/DTR *P. aeruginosa* in other infection sites (104). However,  
489 spontaneous mutation rates for fosfomicin resistance are high and the  
490 mechanism involved is typically the mutational inactivation of *glpT*, coding for a  
491 glycerol-3-phosphate permease required for fosfomicin uptake (105,106).



492 Mutations in *gfpT* are also frequently found among MDR/XDR/DTR strains  
493 (107). Certain specific mutations, like T211P, have become fixed in some  
494 widespread lineages as described for ST175 (14)

495

#### 496 **Definitions of resistance profiles in *Pseudomonas aeruginosa***

497 According to established recommendations by ECDC (6) the MDR profile  
498 is defined as resistance to at least one agent in at least three of eight antibiotic  
499 categories. These categories include antipseudomonal penicillins +  $\beta$ -lactamase  
500 inhibitor combinations (ticarcillin/clavulanate, piperacillin/tazobactam),  
501 antipseudomonal cephalosporins (ceftazidime and cefepime), monobactams  
502 (aztreonam), antipseudomonal carbapenems (imipenem, meropenem,  
503 doripenem), fluoroquinolones (ciprofloxacin, levofloxacin), aminoglycosides  
504 (gentamicin, tobramycin, amikacin, netilmicin), polymyxins (colistin, polymyxin  
505 B) and fosfonic acids (fosfomicin). The XDR profile is defined as resistance to  
506 at least one agent in all antibiotic classes except one or two. Likewise, PDR  
507 profile is defined as resistance to all agents in the eight antibiotic categories.  
508 The eighth category (fosfonic acids, fosfomicin) included in the ECDC  
509 recommendations should be likely not considered, given the lack of current  
510 EUCAST clinical breakpoints. Likewise, the inclusion of gentamicin as  
511 antipseudomonal agents is questionable according to current EUCAST  
512 breakpoints, and the activity of ticarcillin/clavulanate likely not comparable to  
513 that of piperacillin/tazobactam in *P. aeruginosa*. On the other hand, the DTR  
514 (difficult to treat resistance) profile is defined according to IDSA/NIH  
515 recommendations as resistance to all first line (classical) agents:  
516 antipseudomonal penicillins +  $\beta$ -lactamase inhibitor combinations,  
517 cephalosporins, monobactams, carbapenems and fluoroquinolones (7). Thus, if  
518 fosfomicin is not considered, all DTR isolates would meet the XDR criteria,  
519 since they are resistant to at least five of seven categories, but not the other  
520 way around.

521 However, neither the ECDC or IDSA/NIH definitions take into  
522 consideration the novel  $\beta$ -lactams and BLBLIs. The inclusion of these novel  
523 agents is challenging, starting by grouping them into meaningful “categories”

524 since their properties, spectrum and mechanisms of resistance show similarities  
525 but also marked differences. As shown in **Table 1**, at least 5 novel categories  
526 could be considered to include the novel  $\beta$ -lactams already approved: fifth  
527 generation antipseudomonal cephalosporins + classical  $\beta$ -lactamase inhibitors  
528 (ceftolozane/tazobactam), antipseudomonal cephalosporins +  
529 diazabicycloctanes  $\beta$ -lactamase inhibitors (ceftazidime/avibactam),  
530 antipseudomonal carbapenems + diazabicycloctanes  $\beta$ -lactamase inhibitors  
531 (imipenem/relebactam), antipseudomonal carbapenems + boronic acid  $\beta$ -  
532 lactamase inhibitors (meropenem/vaborbactam) and siderophore  
533 antipseudomonal cephalosporins (cefiderocol). Additionally, there are at least  
534 three further classes to be considered in the future if the corresponding  
535 antibiotics are approved: monobactams + diazabicycloctanes  $\beta$ -lactamase  
536 inhibitors (aztreonam/avibactam), antipseudomonal cephalosporins+  
537 diazabicycloctanes  $\beta$ -lactamase and PBP2 inhibitors (cefepime/zidebactam)  
538 and antipseudomonal cephalosporins + boronic acid  $\beta$ -lactamase inhibitors  
539 including MBLs (cefepime/taniborbactam).

540 Within the framework of the ECDC definitions, these novel categories  
541 could potentially align with MDR implying resistance to at least three classes (of  
542 up to 13), XDR indicating resistance to all but one or two and PDR indicating  
543 resistance to all. Regarding DTR definition, it would imply resistance to all the  
544 novel  $\beta$ -lactams approved. However, the practical application of this definition is  
545 likely to encounter challenges due to limited access to these antibiotics for  
546 treatment and to the capacity to perform antimicrobial susceptibility testing in  
547 several countries. Moreover, the classification of the resistance profiles for the  
548 novel agents under development into clinical SIR categories will need to  
549 consider PK/PD data, not yet available in some cases, in addition to existing  
550 phenotypic and genomic information.

551

552 **Update on *Pseudomonas aeruginosa* high-risk lineages and their**  
553 **association with transferable  $\beta$ -lactamases**

554 In a recent review (10), according to their prevalence, global spread and  
555 association with MDR/XDR/DTR profiles, and specially with concerning  
556 horizontally-acquired  $\beta$ -lactamases such as ESBLs and carbapenemases, the  
557 worldwide top ten *P. aeruginosa* high-risk lineages were established to be, by  
558 order of relevance, ST235, ST111, ST233, ST244, ST357, ST308, ST175,  
559 ST277, ST654 and ST298. **Figure 2** shows updated information for these top  
560 ten high-risk lineages, including their virulence profile (presence of the genes  
561 coding the type III secretion system exotoxins ExoS and/or ExoU), worldwide  
562 distribution and association with acquired carbapenemases from key  
563 publications in the last three years (40–42,93,108–112). Particularly noteworthy  
564 is the expansion of KPC enzymes in several of these lineages (ST233, ST277  
565 and ST654 in addition to the previous detection in ST235, ST111 and ST244),  
566 followed by NDM (ST244 and ST357 in addition to ST235, ST233, ST308 and  
567 ST654). Moreover, coproduction of various carbapenemases is not infrequent  
568 among those lineages (43). Besides these top ten lineages, a few others have  
569 gained relevance in the last few years, including globally expanding ST309,  
570 associated with the production of VIM-2, ST773 linked to NDM-1, or ST463  
571 associated with the production of KPC-2, particularly in China (113–118).

572

### 573 **Concluding remarks and future challenges**

574 *P. aeruginosa* infections rank among the foremost global resistance  
575 threats, associated with significant morbidity and mortality. *P. aeruginosa*  
576 resistance mechanisms and epidemiology are complex and ever-evolving, with  
577 a significant impact on novel and forthcoming  $\beta$ -lactams. The interplay between  
578 novel antibiotics and resistance is notably challenging, as certain mechanisms  
579 can lead to cross-resistance to multiple agents, while others may confer  
580 collateral susceptibility to relevant antipseudomonals such as carbapenems.  
581 The global dissemination of XDR/DTR high-risk lineages are also a major  
582 challenge, particularly when coupled with increased virulence and capacity to  
583 acquire exogenous resistance elements as documented for ST235 (11). In this  
584 sense, a recent nation-wide survey of *P. aeruginosa* susceptibility profiles and  
585 resistance genomics has revealed in one hand a significant generalized

586 decrease of resistance rates and XDR/DTR profiles in Spain in the last five  
587 years, but in the other, a significant increase in the proportion of the concerning  
588 carbapenemase-producing ST235 high-risk lineage (44).

589 Therefore, there is a major need for establishing comprehensive  
590 resistance surveillance initiatives, integrating both phenotypic and genomic  
591 data, as well as metadata. However, our current capacity to predict the  
592 susceptibility profiles and emerging high-risk clonal lineages from genomic  
593 sequences still needs to be improved, potentially through the incorporation of  
594 machine learning, knowledge-based approaches, or so-called artificial  
595 intelligence tools (43,119,120). Nevertheless, current achievable surveillance  
596 strategies at European level should at least integrate: (1) monitoring of  
597 concerning high-risk lineages (particularly ST235); (2) analyses of resistance  
598 prevalence trends to recently introduced agents (like the novel BLBLIs) in  
599 addition to classical antipseudomonals; (3) monitoring of strains producing  
600 horizontally-acquired resistance mechanisms (particularly carbapenemases and  
601 ESBLs); and (4) monitoring of noteworthy chromosomal resistance mechanisms  
602 such as the AmpC (PDC) derivatives involved in resistance to the novel BLBLIs.  
603 Likewise, in this scenario, antimicrobial stewardship and infection control are of  
604 paramount importance. Nevertheless, these aspects are equally challenging  
605 and should be guided by rapid diagnostics and antimicrobial susceptibility  
606 testing, including the detection of resistance mechanisms and specific high-risk  
607 clonal lineages (121). Thus, efforts should also be directed to the  
608 implementation and scaling of personalized precision medicine that allows us to  
609 establish early targeted treatments and specific epidemiological control  
610 measures adapted to the strain/mechanism involved.

611

612 **Conflict of interests**

613

614 AO received grants, consulting fees and honoraria for lectures from MSD,  
615 Shionogi, Pfizer and Wockhardt. ERM received honoraria for lectures from  
616 Menarini and Shionogi. JAS received honoraria for lectures from Shionogi. RC  
617 received grants and honoraria for lectures from MSD, Menarini and Shionogi.  
618 PDC received grants from MSD and Shionogi. All other authors declare no  
619 conflict of interests

620

621 **Authors contributions.**

622

623 All authors agreed the questions to be addressed in the position paper. AO and  
624 ERM drafted a first version of the document that was extensively revised by all  
625 other authors.

626

627 **Acknowledgements**

628 This work was done under the auspices of the European Study Group on  
629 Antimicrobial Resistance Surveillance (ESGARS) from the European Society of  
630 Clinical Microbiology and Infectious Diseases (ESCMID).

631

632 **Funding**

633 This work was supported by an EU JPIAMR Grant (ISARPAE). AO, CLC, SMK,  
634 AT and DS are supported by the EU Horizon 2020 research and innovation  
635 program (952491-AmReSu). AO, ERM, JAS, CLC, RC and JOI are supported  
636 by the Instituto de Salud Carlos III, Ministerio de Ciencia e Innovación and  
637 Unión Europea – NextGenerationEU through the Centers for network research I  
638 Personalized and precision medicine grant (MePRAM Project, PMP22/00092).  
639 BK was supported by János Bolyai Research Scholarship (BO/00286/22/5) of  
640 the Hungarian Academy of Sciences. PP is supported by the projects  
641 UIDP/00772/2020 and LA/P/0059/2020 funded by the Portuguese Foundation  
642 for Science and Technology (FCT).

643

644

645

646

647 **References**

- 648 1. Horcajada JP, Montero M, Oliver A, Sorlí L, Luque S, Gómez-Zorrilla S, et  
649 al. Epidemiology and Treatment of Multidrug-Resistant and Extensively  
650 Drug-Resistant *Pseudomonas aeruginosa* Infections. *Clin Microbiol Rev.*  
651 2019 ;32(4).
- 652 2. Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray  
653 A, et al. Global burden of bacterial antimicrobial resistance in 2019: a  
654 systematic analysis. *Lancet.* 2022 Feb 12 ;399(10325):629–55.
- 655 3. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL,  
656 et al. Discovery, research, and development of new antibiotics: the WHO  
657 priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect*  
658 *Dis.* 2018 Mar 1 ;18(3):318–27.
- 659 4. López-Causapé C, Cabot G, Del Barrio-Tofiño E, Oliver A. The Versatile  
660 Mutational Resistome of *Pseudomonas aeruginosa*. *Front Microbiol.*  
661 2018;9:685.
- 662 5. Tenover FC, Nicolau DP, Gill CM. Carbapenemase-producing  
663 *Pseudomonas aeruginosa* -an emerging challenge. *Emerg Microbes*  
664 *Infect.* 2022 ;11(1):811–4.
- 665 6. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske  
666 CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-  
667 resistant bacteria: an international expert proposal for interim standard  
668 definitions for acquired resistance. *Clin Microbiol Infect.* 2012 ;18(3):268–  
669 81.
- 670 7. Kadri SS, Adjemian J, Lai YL, Spaulding AB, Ricotta E, Rebecca Prevots  
671 D, et al. Difficult-to-Treat Resistance in Gram-negative Bacteremia at 173  
672 US Hospitals: Retrospective Cohort Analysis of Prevalence, Predictors,  
673 and Outcome of Resistance to All First-line Agents. *Clin Infect Dis.* 2018  
674 ;67(12):1803–14.
- 675 8. Pelegri AC, Palmieri M, Mirande C, Oliver A, Moons P, Goossens H, et  
676 al. *Pseudomonas aeruginosa*: a clinical and genomics update. *FEMS*  
677 *Microbiol Rev.* 2021 Nov 1 ;45(6).
- 678 9. Oliver A, Mulet X, López-Causapé C, Juan C. The increasing threat of  
679 *Pseudomonas aeruginosa* high-risk clones. *Drug Resist Updat.* 2015 Jul 1  
680 ;21–22:41–59.
- 681 10. del Barrio-Tofiño E, López-Causapé C, Oliver A. *Pseudomonas*  
682 *aeruginosa* epidemic high-risk clones and their association with  
683 horizontally-acquired  $\beta$ -lactamases: 2020 update. *Int J Antimicrob Agents.*  
684 2020 Dec 1 ;56(6).
- 685 11. Treepong P, Kos VN, Guyeux C, Blanc DS, Bertrand X, Valot B, et al.  
686 Global emergence of the widespread *Pseudomonas aeruginosa* ST235  
687 clone. *Clin Microbiol Infect.* 2018 Mar 1 ;24(3):258–66.
- 688 12. Del Barrio-Tofiño E, Zamorano L, Cortes-Lara S, López-Causape C,  
689 Sánchez-Diener I, Cabot G, et al. Spanish nationwide survey on

- 690 Pseudomonas aeruginosa antimicrobial resistance mechanisms and  
691 epidemiology. J Antimicrob Chemother. 2019 Jul 1 ;74(7):1825–35.
- 692 13. Kos VN, Déraspe M, McLaughlin RE, Whiteaker JD, Roy PH, Alm RA, et  
693 al. The resistome of Pseudomonas aeruginosa in relationship to  
694 phenotypic susceptibility. Antimicrob Agents Chemother. 2015 Jan 1  
695 ;59(1):427–36.
- 696 14. Cabot G, López-Causapé C, Ocampo-Sosa AA, Sommer LM, Domínguez  
697 MÁ, Zamorano L, et al. Deciphering the Resistome of the Widespread  
698 Pseudomonas aeruginosa Sequence Type 175 International High-Risk  
699 Clone through Whole-Genome Sequencing. Antimicrob Agents  
700 Chemother. 2016 Dec 1 ;60(12):7415–23.
- 701 15. Jaillard M, van Belkum A, Cady KC, Creely D, Shortridge D, Blanc B, et al.  
702 Correlation between phenotypic antibiotic susceptibility and the resistome  
703 in Pseudomonas aeruginosa. Int J Antimicrob Agents. 2017 Aug 1  
704 ;50(2):210–8.
- 705 16. Wright H, Bonomo RA, Paterson DL. New agents for the treatment of  
706 infections with Gram-negative bacteria: restoring the miracle or false  
707 dawn? Clin Microbiol Infect. 2017 Oct 1 ;23(10):704–12.
- 708 17. Yahav D, Giske CG, Gramatniece A, Abodakpi H, Tam VH, Leibovici L.  
709 New  $\beta$ -Lactam- $\beta$ -Lactamase Inhibitor Combinations. Clin Microbiol Rev.  
710 2020 Jan 1 ;34(1):1–61.
- 711 18. Bassetti M, Echols R, Matsunaga Y, Ariyasu M, Doi Y, Ferrer R, et al.  
712 Efficacy and safety of ceftiderocol or best available therapy for the  
713 treatment of serious infections caused by carbapenem-resistant Gram-  
714 negative bacteria (CREDIBLE-CR): a randomised, open-label, multicentre,  
715 pathogen-focused, descriptive, phase 3 trial. Lancet Infect Dis. 2021 Feb 1  
716 ;21(2):226–40.
- 717 19. Viale P, Sandrock CE, Ramirez P, Rossolini GM, Lodise TP. Treatment of  
718 critically ill patients with ceftiderocol for infections caused by multidrug-  
719 resistant pathogens: review of the evidence. Ann Intensive Care. 2023  
720 Dec 1 ;13(1).
- 721 20. Fraile-Ribot PA, Cabot G, Mulet X, Periañez L, Luisa Martín-Pena M, Juan  
722 C, et al. Mechanisms leading to in vivo ceftolozane/tazobactam resistance  
723 development during the treatment of infections caused by MDR  
724 Pseudomonas aeruginosa. J Antimicrob Chemother. 2018 Mar 1  
725 ;73(3):658–63.
- 726 21. Mojica MF, De La Cadena E, García-Betancur JC, Porras J, Novoa-  
727 Caicedo I, Páez-Zamora L, et al. Molecular Mechanisms of Resistance to  
728 Ceftazidime/Avibactam in Clinical Isolates of Enterobacterales and  
729 Pseudomonas aeruginosa in Latin American Hospitals. mSphere. 2023  
730 Apr 20 ;8(2).
- 731 22. Alonso-García I, Vázquez-Ucha JC, Lasarte-Monterrubio C, González-  
732 Mayo E, Lada-Salvador P, Vela-Fernández R, et al. Simultaneous and  
733 divergent evolution of resistance to cephalosporin/ $\beta$ -lactamase inhibitor

- 734 combinations and imipenem/relebactam following ceftazidime/avibactam  
735 treatment of MDR *Pseudomonas aeruginosa* infections. *J Antimicrob*  
736 *Chemother.* 2023 May 1 ;78(5):1195–200.
- 737 23. Gomis-Font MA, Pitart C, del Barrio-Tofiño E, Zboromyrska Y, Cortes-Lara  
738 S, Mulet X, et al. Emergence of Resistance to Novel Cephalosporin- $\beta$ -  
739 Lactamase Inhibitor Combinations through the Modification of the  
740 *Pseudomonas aeruginosa* MexCD-OprJ Efflux Pump. *Antimicrob Agents*  
741 *Chemother.* 2021 Aug 1 ;65(8).
- 742 24. Shields RK, Stellfox ME, Kline EG, Samanta P, Van Tyne D. Evolution of  
743 Imipenem-Relebactam Resistance Following Treatment of Multidrug-  
744 Resistant *Pseudomonas aeruginosa* Pneumonia. *Clin Infect Dis.* 2022 Aug  
745 15 ;75(4):710–4.
- 746 25. Bahr G, González LJ, Vila AJ. Metallo- $\beta$ -lactamases in the Age of  
747 Multidrug Resistance: From Structure and Mechanism to Evolution,  
748 Dissemination, and Inhibitor Design. *Chem Rev.* 2021 Jul 14  
749 ;121(13):7957–8094.
- 750 26. Ruedas-López A, Alonso-García I, Lasarte-Monterrubbio C, Guijarro-  
751 Sánchez P, Gato E, Vázquez-Ucha JC, et al. Selection of AmpC  $\beta$ -  
752 Lactamase Variants and Metallo- $\beta$ -Lactamases Leading to  
753 Ceftolozane/Tazobactam and Ceftazidime/Avibactam Resistance during  
754 Treatment of MDR/XDR *Pseudomonas aeruginosa* Infections. *Antimicrob*  
755 *Agents Chemother.* 2022 Feb 1 ;66(2).
- 756 27. Lister PD, Wolter DJ, Hanson ND. Antibacterial-resistant *Pseudomonas*  
757 *aeruginosa*: Clinical impact and complex regulation of chromosomally  
758 encoded resistance mechanisms. Vol. 22, *Clinical Microbiology Reviews.*  
759 American Society for Microbiology (ASM); 2009. p. 582–610.
- 760 28. Juan C, Torrens G, González-Nicolau M, Oliver A. Diversity and regulation  
761 of intrinsic  $\beta$ -lactamases from non-fermenting and other Gram-negative  
762 opportunistic pathogens. *FEMS Microbiol Rev.* 2017 Nov 1 ;41(6):781–  
763 815.
- 764 29. Juan C, Maciá MD, Gutiérrez O, Vidal C, Pérez JL, Oliver A. Molecular  
765 mechanisms of beta-lactam resistance mediated by AmpC  
766 hyperproduction in *Pseudomonas aeruginosa* clinical strains. *Antimicrob*  
767 *Agents Chemother.* 2005 Nov ;49(11):4733–8.
- 768 30. Moya B, Dötsch A, Juan C, Blázquez J, Zamorano L, Haussler S, et al.  
769 Beta-lactam resistance response triggered by inactivation of a  
770 nonessential penicillin-binding protein. *PLoS Pathog.* 2009 ;5(3).
- 771 31. Cabot G, Bruchmann S, Mulet X, Zamorano L, Moyá B, Juan C, et al.  
772 *Pseudomonas aeruginosa* ceftolozane-tazobactam resistance  
773 development requires multiple mutations leading to overexpression and  
774 structural modification of AmpC. *Antimicrob Agents Chemother.* 2014  
775 ;58(6):3091–9.
- 776 32. Lahiri SD, Johnstone MR, Ross PL, McLaughlin RE, Olivier NB, Alm RA.  
777 Avibactam and class C  $\beta$ -lactamases: mechanism of inhibition,



- 778 conservation of the binding pocket, and implications for resistance.  
779 Antimicrob Agents Chemother. 2014 Oct 1 ;58(10):5704–13.
- 780 33. Haidar G, Philips NJ, Shields RK, Snyder D, Cheng S, Potoski BA, et al.  
781 Ceftolozane-Tazobactam for the Treatment of Multidrug-Resistant  
782 *Pseudomonas aeruginosa* Infections: Clinical Effectiveness and Evolution  
783 of Resistance. Clin Infect Dis. 2017 Jul 1 ;65(1):110–20.
- 784 34. Berrazeg M, Jeannot K, Ntsogo Enguéné VY, Broutin I, Loeffert S,  
785 Fournier D, et al. Mutations in  $\beta$ -Lactamase AmpC Increase Resistance of  
786 *Pseudomonas aeruginosa* Isolates to Antipseudomonal Cephalosporins.  
787 Antimicrob Agents Chemother. 2015 Oct 1 ;59(10):6248–55.
- 788 35. Gomis-Font MA, Sastre-Femenia MÀ, Taltavull B, Cabot G, Oliver A. In  
789 vitro dynamics and mechanisms of cefiderocol resistance development in  
790 wild-type, mutator and XDR *Pseudomonas aeruginosa*. J Antimicrob  
791 Chemother. 2023 Jul 5 ;78(7):1785–94.
- 792 36. Naas T, Oueslati S, Bonnin RA, Dabos ML, Zavala A, Dortet L, et al. Beta-  
793 lactamase database (BLDB) - structure and function. J Enzyme Inhib Med  
794 Chem. 2017 Jan 1;32(1):917–9.
- 795 37. Fraile-Ribot PA, Mulet X, Cabot G, Del Barrio-Tofiño E, Juan C, Pérez JL,  
796 et al. In Vivo Emergence of Resistance to Novel Cephalosporin- $\beta$ -  
797 Lactamase Inhibitor Combinations through the Duplication of Amino Acid  
798 D149 from OXA-2  $\beta$ -Lactamase (OXA-539) in Sequence Type 235  
799 *Pseudomonas aeruginosa*. Antimicrob Agents Chemother. 2017 Sep 1  
800 ;61(9).
- 801 38. Arca-Suárez J, Lasarte-Monterrubbio C, Rodiño-Janeiro BK, Cabot G,  
802 Vázquez-Ucha JC, Rodríguez-Iglesias M, et al. Molecular mechanisms  
803 driving the in vivo development of OXA-10-mediated resistance to  
804 ceftolozane/tazobactam and ceftazidime/avibactam during treatment of  
805 XDR *Pseudomonas aeruginosa* infections. J Antimicrob Chemother. 2021  
806 ;76(1):91–100.
- 807 39. Lasarte-Monterrubbio C, Fraile-Ribot PA, Vázquez-Ucha JC, Cabot G,  
808 Guijarro-Sánchez P, Alonso-García I, et al. Activity of cefiderocol,  
809 imipenem/relebactam, cefepime/taniborbactam and cefepime/zidebactam  
810 against ceftolozane/tazobactam- and ceftazidime/avibactam-resistant  
811 *Pseudomonas aeruginosa*. J Antimicrob Chemother. 2022 Oct 1  
812 ;77(10):2809–15.
- 813 40. Reyes J, Komarow L, Chen L, Ge L, Hanson BM, Cober E, et al. Global  
814 epidemiology and clinical outcomes of carbapenem-resistant  
815 *Pseudomonas aeruginosa* and associated carbapenemases (POP): a  
816 prospective cohort study. Lancet Microbe. 2023 Mar 1 ;4(3):e159–70.
- 817 41. Wang MG, Liu ZY, Liao XP, Sun RY, Li RB, Liu Y, et al. Retrospective  
818 Data Insight into the Global Distribution of Carbapenemase-Producing  
819 *Pseudomonas aeruginosa*. Antibiotics (Basel). 2021 May 1 ;10(5).
- 820 42. Fortunato G, Vaz-Moreira I, Gajic I, Manaia CM. Insight into phylogenomic  
821 bias of blaVIM-2 or blaNDM-1 dissemination amongst carbapenem-

- 822 resistant *Pseudomonas aeruginosa*. *Int J Antimicrob Agents*. 2023 May 1  
823 ;61(5).
- 824 43. Torrens G, Van Der Schalk TE, Cortes-Lara S, Timbermont L, Del Barrio-  
825 Tofiño E, Xavier BB, et al. Susceptibility profiles and resistance genomics  
826 of *Pseudomonas aeruginosa* isolates from European ICUs participating in  
827 the ASPIRE-ICU trial. *J Antimicrob Chemother*. 2022 Jul 1 ;77(7):1862–  
828 72.
- 829 44. Àngel Sastre-Femenia M, Fernández-Muñoz A, Gomis-Font MA, Taltavull  
830 B, López-Causapé C, Arca-Suárez J, et al. *Pseudomonas aeruginosa*  
831 antibiotic susceptibility profiles, genomic epidemiology and resistance  
832 mechanisms: a nation-wide five-year time lapse analysis. *The Lancet*  
833 *Regional Health - Europe*. 2023;34:100736.
- 834 45. Takemura M, Wise MG, Hackel MA, Sahn DF, Yamano Y. In vitro activity  
835 of cefiderocol against MBL-producing Gram-negative bacteria collected in  
836 North America and Europe in five consecutive annual multinational  
837 SIDERO-WT surveillance studies (2014-2019). *J Antimicrob Chemother*.  
838 2023 Aug 2 ;78(8).
- 839 46. Le Terrier C, Nordmann P, Poirel L. In vitro activity of aztreonam in  
840 combination with newly developed  $\beta$ -lactamase inhibitors against MDR  
841 Enterobacterales and *Pseudomonas aeruginosa* producing metallo- $\beta$ -  
842 lactamases. *J Antimicrob Chemother* 2022 Jan 1;78(1):101–7.
- 843 47. Le Terrier C, Nordmann P, Freret C, Seigneur M, Poirel L. Impact of  
844 Acquired Broad Spectrum  $\beta$ -Lactamases on Susceptibility to Novel  
845 Combinations Made of  $\beta$ -Lactams (Aztreonam, Cefepime, Meropenem,  
846 and Imipenem) and Novel  $\beta$ -Lactamase Inhibitors in *Escherichia coli* and  
847 *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2023 Jul  
848 1;67(7).
- 849 48. Moya B, Barcelo IM, Bhagwat S, Patel M, Bou G, Papp-Wallace  
850 KM, et al. WCK 5107 (Zidebactam) and WCK 5153 Are Novel Inhibitors of  
851 PBP2 Showing Potent “ $\beta$ -Lactam Enhancer” Activity against  
852 *Pseudomonas aeruginosa*, Including Multidrug-Resistant Metallo- $\beta$ -  
853 Lactamase-Producing High-Risk Clones. *Antimicrob Agents Chemother*.  
854 2017 Jun 1 ;61(6).
- 855 49. Meletiadis J, Paranos P, Georgiou PC, Vourli S, Antonopoulou S,  
856 Michelaki A, et al. In vitro comparative activity of the new beta-lactamase  
857 inhibitor taniborbactam with cefepime or meropenem against *Klebsiella*  
858 *pneumoniae* and cefepime against *Pseudomonas aeruginosa* metallo-  
859 beta-lactamase-producing clinical isolates. *Int J Antimicrob Agents*. 2021  
860 Nov 1 ;58(5).
- 861 50. Poirel L, De la Rosa JMO, Sadek M, Nordmann P. Impact of Acquired  
862 Broad-Spectrum  $\beta$ -Lactamases on Susceptibility to Cefiderocol and Newly  
863 Developed  $\beta$ -Lactam/ $\beta$ -Lactamase Inhibitor Combinations in *Escherichia*  
864 *coli* and *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2022  
865 Apr 1 ;66(4).

- 866 51. Shortridge D, Carvalhaes C, Deshpande L, Castanheira M. Activity of  
867 meropenem/vaborbactam and comparators against Gram-negative  
868 isolates from Eastern and Western European patients hospitalized with  
869 pneumonia including ventilator-associated pneumonia (2014-19). *J*  
870 *Antimicrob Chemother.* 2021 Oct 1 ;76(10):2600–5.
- 871 52. Mushtaq S, Meunier D, Vickers A, Woodford N, Livermore DM. Activity of  
872 imipenem/relebactam against *Pseudomonas aeruginosa* producing ESBLs  
873 and carbapenemases. *J Antimicrob Chemother.* 2021 Feb 1 ;76(2):434–  
874 42.
- 875 53. Fraile-Ribot PA, Zamorano L, Orellana R, Del Barrio-Tofiño E, Sánchez-  
876 Diener I, Cortes-Lara S, et al. Activity of Imipenem-Relebactam against a  
877 Large Collection of *Pseudomonas aeruginosa* Clinical Isolates and  
878 Isogenic  $\beta$ -Lactam-Resistant Mutants. *Antimicrob Agents Chemother.*  
879 2020 Jan 27 ;64(2).
- 880 54. Fraile-Ribot PA, Fernández J, Gomis-Font MA, Forcelledo L, Mulet X,  
881 López-Causapé C, et al. In Vivo Evolution of GES  $\beta$ -Lactamases Driven  
882 by Ceftazidime/Avibactam Treatment of *Pseudomonas aeruginosa*  
883 Infections. *Antimicrob Agents Chemother.* 2021 Sep 1 ;65(9).
- 884 55. Faccone D, de Mendieta JM, Albornoz E, Chavez M, Genero F, Echegorry  
885 M, et al. Emergence of KPC-31, a KPC-3 Variant Associated with  
886 Ceftazidime-Avibactam Resistance, in an Extensively Drug-Resistant  
887 ST235 *Pseudomonas aeruginosa* Clinical Isolate. *Antimicrob Agents*  
888 *Chemother.* 2022 Nov 1 ;66(11).
- 889 56. Recio R, Villa J, González-Bodí S, Brañas P, Orellana MÁ, Mancheño-  
890 Losa M, et al. Genomic Analysis of Ceftazidime/Avibactam-Resistant  
891 GES-Producing Sequence Type 235 *Pseudomonas aeruginosa* Isolates.  
892 *Antibiotics (Basel).* 2022 Jul 1 ;11(7).
- 893 57. Bonnin RA, Bogaerts P, Girlich D, Huang TD, Dortet L, Glupczynski Y, et  
894 al. Molecular Characterization of OXA-198 Carbapenemase-Producing  
895 *Pseudomonas aeruginosa* Clinical Isolates. *Antimicrob Agents Chemother.*  
896 2018 Jun 1 ;62(6).
- 897 58. Chen W, Zhang YM, Davies C. Penicillin-Binding Protein 3 Is Essential for  
898 Growth of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother.* 2016  
899 Jan 1 ;61(1).
- 900 59. Caballero JD, Clark ST, Coburn B, Zhang Y, Wang PW, Donaldson SL, et  
901 al. Selective Sweeps and Parallel Pathoadaptation Drive *Pseudomonas*  
902 *aeruginosa* Evolution in the Cystic Fibrosis Lung. *mBio.* 2015 ;6(5).
- 903 60. López-Causapé C, Sommer LM, Cabot G, Rubio R, Ocampo-Sosa AA,  
904 Johansen HK, et al. Evolution of the *Pseudomonas aeruginosa* mutational  
905 resistome in an international Cystic Fibrosis clone. *Sci Rep.* 2017 Dec 1  
906 ;7(1).
- 907 61. Cabot G, Zamorano L, Moyà B, Juan C, Navas A, Blázquez J, et al.  
908 Evolution of *Pseudomonas aeruginosa* Antimicrobial Resistance and

- 909 Fitness under Low and High Mutation Rates. *Antimicrob Agents*  
910 *Chemother.* 2016 Mar 1 ;60(3):1767–78.
- 911 62. Han S, Zaniewski RP, Marr ES, Lacey BM, Tomaras AP, Evdokimov A, et  
912 al. Structural basis for effectiveness of siderophore-conjugated  
913 monocarbams against clinically relevant strains of *Pseudomonas*  
914 *aeruginosa*. *Proc Natl Acad Sci U S A.* 2010 Dec 21 ;107(51):22002–7.
- 915 63. Jorth P, McLean K, Ratjen A, Secor PR, Bautista GE, Ravishankar S, et  
916 al. Evolved Aztreonam Resistance Is Multifactorial and Can Produce  
917 Hypervirulence in *Pseudomonas aeruginosa*. *mBio.* 2017 Sep 1 ;8(5).
- 918 64. Cabot G, Florit-Mendoza L, Sánchez-Diener I, Zamorano L, Oliver A.  
919 Deciphering  $\beta$ -lactamase-independent  $\beta$ -lactam resistance evolution  
920 trajectories in *Pseudomonas aeruginosa*. *J Antimicrob Chemother.* 2018  
921 Dec 1 ;73(12):3322–31.
- 922 65. Barceló I, Cabot G, Palwe S, Joshi P, Takalkar S, Periasamy H, et al. In  
923 vitro evolution of cefepime/zidebactam (WCK 5222) resistance in  
924 *Pseudomonas aeruginosa*: dynamics, mechanisms, fitness trade-off and  
925 impact on in vivo efficacy. *J Antimicrob Chemother.* 2021 Oct 1  
926 ;76(10):2546–57.
- 927 66. Lister PD, Wolter DJ, Hanson ND. Antibacterial-resistant *Pseudomonas*  
928 *aeruginosa*: clinical impact and complex regulation of chromosomally  
929 encoded resistance mechanisms. *Clin Microbiol Rev.* 2009 Oct  
930 ;22(4):582–610
- 931 67. Castanheira M, Deshpande LM, Costello A, Davies TA, Jones RN.  
932 Epidemiology and carbapenem resistance mechanisms of carbapenem-  
933 non-susceptible *Pseudomonas aeruginosa* collected during 2009-11 in 14  
934 European and Mediterranean countries. *J Antimicrob Chemother.* 2014  
935 ;69(7):1804–14.
- 936 68. Richardot C, Plésiat P, Fournier D, Monlezun L, Broutin I, Llanes C.  
937 Carbapenem resistance in cystic fibrosis strains of *Pseudomonas*  
938 *aeruginosa* as a result of amino acid substitutions in porin OprD. *Int J*  
939 *Antimicrob Agents.* 2015 May 1 ;45(5):529–32.
- 940 69. Ocampo-Sosa AA, Cabot G, Rodríguez C, Roman E, Tubau F, Macia MD,  
941 et al. Alterations of OprD in carbapenem-intermediate and -susceptible  
942 strains of *Pseudomonas aeruginosa* isolated from patients with bacteremia  
943 in a Spanish multicenter study. *Antimicrob Agents Chemother.* 2012 Apr  
944 ;56(4):1703–13.
- 945 70. Li XZ, Plésiat P, Nikaido H. The challenge of efflux-mediated antibiotic  
946 resistance in Gram-negative bacteria. *Clin Microbiol Rev.* 2015  
947 ;28(2):337–418
- 948 71. Gomis-Font MA, Cabot G, Sánchez-Diener I, Fraile-Ribot PA, Juan C,  
949 Moya B, et al. In vitro dynamics and mechanisms of resistance  
950 development to imipenem and imipenem/relebactam in *Pseudomonas*  
951 *aeruginosa*. *J Antimicrob Chemother.* 2020 Sep 1 ;75(9):2508–15.

- 952 72. Gomis-Font MA, Cabot G, López-Argüello S, Zamorano L, Juan C, Moyá  
953 B, et al. Comparative analysis of in vitro dynamics and mechanisms of  
954 ceftolozane/tazobactam and imipenem/relebactam resistance  
955 development in *Pseudomonas aeruginosa* XDR high-risk clones. *J*  
956 *Antimicrob Chemother.* 2022 Apr 1 ;77(4):957–68.
- 957 73. Sanz-García F, Hernando-Amado S, Martínez JL. Mutation-Driven  
958 Evolution of *Pseudomonas aeruginosa* in the Presence of either  
959 Ceftazidime or Ceftazidime-Avibactam. *Antimicrob Agents Chemother.*  
960 2018 Oct 1 ;62(10).
- 961 74. Hocquet D, Petitjean M, Rohmer L, Valot B, Kulasekara HD, Bedel E, et  
962 al. Pyomelanin-producing *Pseudomonas aeruginosa* selected during  
963 chronic infections have a large chromosomal deletion which confers  
964 resistance to pyocins. *Environ Microbiol.* 2016 Oct 1 ;18(10):3482–93.
- 965 75. Dötsch A, Becker T, Pommerenke C, Magnowska Z, Jänsch L, Häussler  
966 S. Genomewide identification of genetic determinants of antimicrobial drug  
967 resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother.*  
968 2009 Jun ;53(6):2522–31.
- 969 76. Alvarez-Ortega C, Wiegand I, Olivares J, Hancock REW, Martínez JL.  
970 Genetic determinants involved in the susceptibility of *Pseudomonas*  
971 *aeruginosa* to beta-lactam antibiotics. *Antimicrob Agents Chemother.* 2010  
972 Oct ;54(10):4159–67.
- 973 77. Potron A, Poirel L, Nordmann P. Emerging broad-spectrum resistance in  
974 *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: Mechanisms and  
975 epidemiology. *Int J Antimicrob Agents.* 2015 May 16 ;45(6):568–85.
- 976 78. Costello SE, Deshpande LM, Davis AP, Mendes RE, Castanheira M.  
977 Aminoglycoside-modifying enzyme and 16S ribosomal RNA  
978 methyltransferase genes among a global collection of Gram-negative  
979 isolates. *J Glob Antimicrob Resist.* 2019 Mar 1 ;16:278–85.
- 980 79. Yokoyama K, Doi Y, Yamane K, Kurokawa H, Shibata N, Shibayama K, et  
981 al. Acquisition of 16S rRNA methylase gene in *Pseudomonas aeruginosa*.  
982 *Lancet.* 2003 Dec 6 ;362(9399):1888–93.
- 983 80. Choi YJ, Kim YA, Junglim K, Jeong SH, Shin JH, Shin KS, et al.  
984 Emergence of NDM-1-producing *Pseudomonas aeruginosa* Sequence  
985 Type 773 Clone: Shift of Carbapenemase Molecular Epidemiology and  
986 Spread of 16S rRNA Methylase Genes in Korea. *Ann Lab Med.* 2023 Mar  
987 1;43(2):196–9.
- 988 81. Mc Gann PT, Lebreton F, Jones BT, Dao HD, Martin MJ, Nelson MJ, et al.  
989 Six Extensively Drug-Resistant Bacteria in an Injured Soldier, Ukraine.  
990 *Emerg Infect Dis.* 2023 Aug 1;29(8):1692–5.
- 991 82. Guénard S, Muller C, Monlezun L, Benas P, Broutin I, Jeannot K, et al.  
992 Multiple mutations lead to MexXY-OprM-dependent aminoglycoside  
993 resistance in clinical strains of *Pseudomonas aeruginosa*. *Antimicrob*  
994 *Agents Chemother.* 2014 Jan ;58(1):221–8.

- 995 83. Prickett MH, Hauser AR, McColley SA, Cullina J, Potter E, Powers C, et al.  
996 Aminoglycoside resistance of *Pseudomonas aeruginosa* in cystic fibrosis  
997 results from convergent evolution in the *mexZ* gene. *Thorax*. 2017 Jan 1  
998 ;72(1):40–7.
- 999 84. El’Garch F, Jeannot K, Hocquet D, Llanes-Barakat C, Plésiat P.  
1000 Cumulative effects of several nonenzymatic mechanisms on the resistance  
1001 of *Pseudomonas aeruginosa* to aminoglycosides. *Antimicrob Agents*  
1002 *Chemother*. 2007 Mar ;51(3):1016–21.
- 1003 85. Schurek KN, Marr AK, Taylor PK, Wiegand I, Semene L, Khaira BK, et al.  
1004 Novel genetic determinants of low-level aminoglycoside resistance in  
1005 *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2008 Dec  
1006 ;52(12):4213–9.
- 1007 86. Feng Y, Jonker MJ, Moustakas I, Brul S, Ter Kuile BH. Dynamics of  
1008 Mutations during Development of Resistance by *Pseudomonas*  
1009 *aeruginosa* against Five Antibiotics. *Antimicrob Agents Chemother*. 2016  
1010 Jul 1 ;60(7):4229–36.
- 1011 87. Chung JCS, Becq J, Fraser L, Schulz-Trieglaff O, Bond NJ, Foweraker J,  
1012 et al. Genomic variation among contemporary *Pseudomonas aeruginosa*  
1013 isolates from chronically infected cystic fibrosis patients. *J Bacteriol*. 2012  
1014 Sep ;194(18):4857–66.
- 1015 88. Markussen T, Marvig RL, Gómez-Lozano M, Aanæs K, Burleigh AE, Høiby  
1016 N, et al. Environmental heterogeneity drives within-host diversification and  
1017 evolution of *Pseudomonas aeruginosa*. *mBio*. 2014 Sep 16 ;5(5).
- 1018 89. Greipel L, Fischer S, Klockgether J, Dorda M, Mielke S, Wiehlmann L, et  
1019 al. Molecular Epidemiology of Mutations in Antimicrobial Resistance Loci  
1020 of *Pseudomonas aeruginosa* Isolates from Airways of Cystic Fibrosis  
1021 Patients. *Antimicrob Agents Chemother*. 2016 Nov 1 ;60(11):6726–34.
- 1022 90. Bolard A, Plésiat P, Jeannot K. Mutations in Gene *fusA1* as a Novel  
1023 Mechanism of Aminoglycoside Resistance in Clinical Strains of  
1024 *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2018 Feb  
1025 ;62(2).
- 1026 91. Bruchmann S, Dötsch A, Nouri B, Chaberny IF, Häussler S. Quantitative  
1027 contributions of target alteration and decreased drug accumulation to  
1028 *Pseudomonas aeruginosa* fluoroquinolone resistance. *Antimicrob Agents*  
1029 *Chemother*. 2013 Mar ;57(3):1361–8.
- 1030 92. Belotti PT, Thabet L, Laffargue A, André C, Coulange-Mayonnove L, Arpin  
1031 C, et al. Description of an original integron encompassing *blaVIM-2*,  
1032 *qnrVC1* and genes encoding bacterial group II intron proteins in  
1033 *Pseudomonas aeruginosa*. *J Antimicrob Chemother*. 2015 Aug 1  
1034 ;70(8):2237–40.
- 1035 93. Pérez-Vázquez M, Sola-Campoy PJ, Zurita ÁM, Ávila A, Gómez-Bertomeu  
1036 F, Solís S, et al. Carbapenemase-producing *Pseudomonas aeruginosa* in  
1037 Spain: interregional dissemination of the high-risk clones ST175 and

- 1038 ST244 carrying blaVIM-2, blaVIM-1, blaIMP-8, blaVIM-20 and blaKPC-2.  
1039 Int J Antimicrob Agents. 2020 Jul 1 ;56(1).
- 1040 94. Shortridge D, Gales AC, Streit JM, Huband MD, Tsakris A, Jones RN.  
1041 Geographic and Temporal Patterns of Antimicrobial Resistance in  
1042 *Pseudomonas aeruginosa* Over 20 Years From the SENTRY Antimicrobial  
1043 Surveillance Program, 1997-2016. Open Forum Infect Dis. 2019 Mar 15  
1044 ;6(Suppl 1):S63–8.
- 1045 95. Olaitan AO, Morand S, Rolain JM. Mechanisms of polymyxin resistance:  
1046 acquired and intrinsic resistance in bacteria. Front Microbiol. 2014  
1047 ;5(NOV). Available from: <https://pubmed.ncbi.nlm.nih.gov/25505462/>
- 1048 96. Jeannot K, Bolard A, Plésiat P. Resistance to polymyxins in Gram-  
1049 negative organisms. Int J Antimicrob Agents. 2017 May 1 ;49(5):526–35.
- 1050 97. Barrow K, Kwon DH. Alterations in two-component regulatory systems of  
1051 phoPQ and pmrAB are associated with polymyxin B resistance in clinical  
1052 isolates of *Pseudomonas aeruginosa*. Antimicrob Agents Chemother.  
1053 2009 Dec ;53(12):5150–4.
- 1054 98. Muller C, Plésiat P, Jeannot K. A two-component regulatory system  
1055 interconnects resistance to polymyxins, aminoglycosides,  
1056 fluoroquinolones, and  $\beta$ -lactams in *Pseudomonas aeruginosa*. Antimicrob  
1057 Agents Chemother. 2011 Mar ;55(3):1211–21.
- 1058 99. Gutu AD, Sgambati N, Strasbourger P, Brannon MK, Jacobs MA, Haugen  
1059 E, et al. Polymyxin resistance of *Pseudomonas aeruginosa* phoQ mutants  
1060 is dependent on additional two-component regulatory systems. Antimicrob  
1061 Agents Chemother. 2013 May ;57(5):2204–15.
- 1062 100. Moskowitz SM, Brannon MK, Dasgupta N, Pier M, Sgambati N, Miller AK,  
1063 et al. PmrB mutations promote polymyxin resistance of *Pseudomonas*  
1064 *aeruginosa* isolated from colistin-treated cystic fibrosis patients. Antimicrob  
1065 Agents Chemother. 2012 Feb ;56(2):1019–30.
- 1066 101. Döbelmann B, Willmann M, Steglich M, Bunk B, Nübel U, Peter S, et al.  
1067 Rapid and Consistent Evolution of Colistin Resistance in Extensively Drug-  
1068 Resistant *Pseudomonas aeruginosa* during Morbidostat Culture.  
1069 Antimicrob Agents Chemother. 2017 Sep 1 ;61(9).
- 1070 102. Cervoni M, Sposato D, Lo Sciuto A, Imperi F. Regulatory Landscape of  
1071 the *Pseudomonas aeruginosa* Phosphoethanolamine Transferase Gene  
1072 *eptA* in the Context of Colistin Resistance. Antibiotics (Basel). 2023 Feb 1  
1073 ;12(2).
- 1074 103. Snesrud E, Maybank R, Kwak YI, Jones AR, Hinkle MK, McGann P.  
1075 Chromosomally Encoded *mcr-5* in Colistin-Nonsusceptible *Pseudomonas*  
1076 *aeruginosa*. Antimicrob Agents Chemother. 2018 Aug 1 ;62(8).
- 1077 104. Michalopoulos AS, Livaditis IG, Gougoutas V. The revival of fosfomycin.  
1078 Int J Infect Dis. 2011 Nov ;15(11).
- 1079 105. Castañeda-García A, Rodríguez-Rojas A, Guelfo JR, Blázquez J. The  
1080 glycerol-3-phosphate permease GlpT is the only fosfomycin transporter in  
1081 *Pseudomonas aeruginosa*. J Bacteriol. 2009 Nov ;191(22):6968–74.

- 1082 106. Rodríguez-Rojas A, Maciá MD, Couce A, Gómez C, Castañeda-García  
1083 A, Oliver A, et al. Assessing the emergence of resistance: the absence of  
1084 biological cost in vivo may compromise fosfomicin treatments for *P.*  
1085 *aeruginosa* infections. *PLoS One*. 2010 ;5(4).
- 1086 107. Del Barrio-Tofiño E, López-Causapé C, Cabot G, Rivera A, Benito N,  
1087 Segura C, et al. Genomics and Susceptibility Profiles of Extensively Drug-  
1088 Resistant *Pseudomonas aeruginosa* Isolates from Spain. *Antimicrob*  
1089 *Agents Chemother*. 2017 Nov 1 ;61(11).
- 1090 108. Silveira MC, Rocha-de-Souza CM, de Oliveira Santos IC, Pontes L da S,  
1091 Oliveira TRT e., Tavares-Teixeira CB, et al. Genetic Basis of Antimicrobial  
1092 Resistant Gram-Negative Bacteria Isolated From Bloodstream in Brazil.  
1093 *Front Med (Lausanne)*. 2021 Mar 15 ;8.
- 1094 109. Takahashi T, Tada T, Shrestha S, Hishinuma T, Sherchan JB, Tohya M,  
1095 et al. Molecular characterisation of carbapenem-resistant *Pseudomonas*  
1096 *aeruginosa* clinical isolates in Nepal. *J Glob Antimicrob Resist*. 2021 Sep  
1097 1;26:279–84.
- 1098 110. Lebreton F, Snesrud E, Hall L, Mills E, Galac M, Stam J, et al. A panel of  
1099 diverse *Pseudomonas aeruginosa* clinical isolates for research and  
1100 development. *JAC Antimicrob Resist*. 2021 Dec 1 ;3(4).
- 1101 111. Cejas D, Elena A, González-Espinosa FE, Pallecchi L, Vay C, Rossolini  
1102 GM, et al. Characterisation of blaKPC-2-harbouring plasmids recovered  
1103 from *Pseudomonas aeruginosa* ST654 and ST235 high-risk clones. *J Glob*  
1104 *Antimicrob Resist*. 2022 Jun 1;29:310–2.
- 1105 112. Pincus NB, Bachta KER, Ozer EA, Allen JP, Pura ON, Qi C, et al. Long-  
1106 term Persistence of an Extensively Drug-Resistant Subclade of Globally  
1107 Distributed *Pseudomonas aeruginosa* Clonal Complex 446 in an  
1108 Academic Medical Center. *Clin Infect Dis*. 2020 Sep 15 ;71(6):1524–31.
- 1109 113. Hu Y, Peng W, Wu Y, Li H, Wang Q, Yi H, et al. A Potential High-Risk  
1110 Clone of *Pseudomonas aeruginosa* ST463. *Front Microbiol*. 2021 May 28  
1111 ;12.
- 1112 114. Martak D, Gbaguidi-Haore H, Meunier A, Valot B, Conzelmann N, Eib M,  
1113 et al. High prevalence of *Pseudomonas aeruginosa* carriage in residents  
1114 of French and German long-term care facilities. *Clin Microbiol Infect*. 2022  
1115 Oct 1 ;28(10):1353–8.
- 1116 115. Fonseca ÉL, Morgado SM, Caldart R V., Freitas F, Vicente ACP.  
1117 Emergence of a VIM-2-producing extensively drug-resistant (XDR)  
1118 *Pseudomonas aeruginosa* ST309 in South America: a comparative  
1119 genomic analysis. *Int J Antimicrob Agents*. 2022 Feb 1 ;59(2).
- 1120 116. Chilam J, Argimón S, Limas MT, Masim ML, Gayeta JM, Lagrada ML, et  
1121 al. Genomic surveillance of *Pseudomonas aeruginosa* in the Philippines,  
1122 2013-2014. *Western Pac Surveill Response J*. 2021 Apr 1 ;12(2).
- 1123 117. Morales-Espinosa R, Delgado G, Espinosa LF, Isselo D, Méndez JL,  
1124 Rodríguez C, et al. Fingerprint Analysis and Identification of Strains ST309  
1125 as a Potential High Risk Clone in a *Pseudomonas aeruginosa* Population



1126 Isolated from Children with Bacteremia in Mexico City. *Front Microbiol.*  
1127 2017 Mar 1 ;8(MAR).

1128 118. Khan A, Tran TT, Rios R, Hanson B, Shropshire WC, Sun Z, et al.  
1129 Extensively Drug-Resistant *Pseudomonas aeruginosa* ST309 Harboring  
1130 Tandem Guiana Extended Spectrum  $\beta$ -Lactamase Enzymes: A Newly  
1131 Emerging Threat in the United States. *Open Forum Infect Dis.* 2019 Jul 1  
1132 ;6(7).

1133 119. Cortes-Lara S, Barrio-Tofiño E del, López-Causapé C, Oliver A,  
1134 Martínez-Martínez L, Bou G, et al. Predicting *Pseudomonas aeruginosa*  
1135 susceptibility phenotypes from whole genome sequence resistome  
1136 analysis. *Clin Microbiol Infect.* 2021 Nov 1 ;27(11):1631–7.

1137 120. Khaledi A, Weimann A, Schniederjans M, Asgari E, Kuo T, Oliver A, et al.  
1138 Predicting antimicrobial resistance in *Pseudomonas aeruginosa* with  
1139 machine learning-enabled molecular diagnostics. *EMBO Mol Med.* 2020  
1140 Mar 6 ;12(3).

1141 121. Cabot G, Lara-Esbrí P, Mulet X, Oliver A. Whole-genome sequence-  
1142 guided PCR for the rapid identification of the *Pseudomonas aeruginosa*  
1143 ST175 high-risk clone directly from clinical samples. *J Antimicrob*  
1144 *Chemother.* 2021 Apr 1 ;76(4):945–9.

1145

1146

**Table 1. Main resistance mechanisms to classical and novel antibiotics in *Pseudomonas aeruginosa***

Antipseudomonal categories	Antipseudomonal agents	Main mutational resistance mechanisms	Alternative mutational resistance mechanisms	Mutational resistance on horizontally acquired determinants	Horizontally-acquired resistance mechanisms
Penicillins + $\beta$ -lactamase inhibitors	Piperacillin/tazobactam	AmpC $\uparrow$	PBP3, GalU		ESBLs, class A and B carbapenemases
Cephalosporins	Ceftazidime	AmpC $\uparrow$	PBP3, GalU	OXA-2/10	ESBLs, class A and B carbapenemases
	Cefepime	MexXY $\uparrow$ , AmpC $\uparrow$	PBP3, GalU	OXA-2/10	ESBLs, class A and B carbapenemases
Monobactams	Aztreonam	MexAB $\uparrow$ , AmpC $\uparrow$	PBP3, GalU	OXA-2/10	ESBLs and class A carbapenemases
Carbapenems	Imipenem	OprD-	MexST, PBP2, PBP1a		Class A and B carbapenemases
	Meropenem	OprD-, MexAB $\uparrow$	PBP3, GalU		Class A and B carbapenemases
Fifth generation cephalosporins+ classical $\beta$ -lactamase inhibitors	Ceftolozane/tazobactam	AmpC $\Omega$ -loop	PBP3, GalU Efflux pumps	OXA-2/10	ESBLs, class A and B carbapenemases
Cephalosporins + diazabicycloctanes $\beta$ -lactamase inhibitors	Ceftazidime/avibactam	AmpC $\Omega$ -loop, MexAB $\uparrow$	PBP3, GalU	OXA-2/10, GES, KPC	Class B carbapenemases
carbapenems + diazabicycloctanes $\beta$ -lactamase inhibitors	Imipenem/relebactam	OprD-, MexAB $\uparrow$ **	MexST, ParRS PBP2, PBP1a		Class A and B carbapenemases
carbapenems + boronic acid $\beta$ -lactamase inhibitors	Meropenem/vaborbactam	OprD-, MexAB $\uparrow$	PBP3, GalU		Class A and B carbapenemases
Siderophore cephalosporins	Cefiderocol	Iron transporters, AmpC $\Omega$ -loop	PBP3, GalU	OXA-2/10**	ESBLs, class A and B carbapenemases**
Monobactams + diazabicycloctanes $\beta$ -lactamase inhibitors	Aztreonam/avibactam*	MexAB $\uparrow$	PBP3, GalU		ESBLs and class A carbapenemases**
Cephalosporins+	Cefepime/zidebactam*	MexXY $\uparrow$ , MexAB $\uparrow$	PBP3, GalU		ESBLs, class A and B

diazabicyclooctanes $\beta$ -lactamase and PBP2 inhibitors			PBP2		carbapenemases**
cephalosporins+ boronic acid $\beta$ -lactamase inhibitors including MBLs	Cefepime/taniborbactam*	MexXY $\uparrow$ , MexAB $\uparrow$	PBP3, GalU		IMPs
Fluoroquinolones	Ciprofloxacin, levofloxacin	QRDR	MexAB/XY/CD/EF $\uparrow$		Qnr
Aminoglycosides	Tobramycin, amikacin	MexXY $\uparrow$ **	FusA1		Aminoglycoside modifying enzymes, 16S rRNA methylases
Polymyxins	Colistin, polymyxin B	PmrAB/PhoPQ/ParRS			MCR (Very uncommon)
Fosfonic acids	Fosfomicin	GlpT			FosA

\* Not yet approved

\*\*Low level resistance. Clinical resistance requires additional mechanisms

$\uparrow$  Hyperproduction

Antibiotic	AmpC ↑	MexAB↑	OprD-	AmpC Ω-loop*	OXA ESBL	ESBL	CarbA	CarbA Mut**	CarbB	Iron transp.
Piperacillin/tazobactam	R	r	S	S/r	R	R	R	R	R	S
Ceftazidime	R	r	S	R	R	R	R	R	R	S
Cefepime	r/R	r/R	S	R	R	R	R	R	R	S
Aztreonam	r/R	R	S	R	r/R	R	R	R	S	S
Imipenem	S	S	r/R	S	S	S	R	S	R	S
Meropenem	S	r	r	S	S	S	R	S	R	S
Ceftolozane/tazobactam	S	S	S	R	R	r/R	R	R	R	S
Ceftazidime/avibactam	S/r	r	S	r/R	r/R	S/r	S	R	R	S
Meropenem/vaborbactam	S	r	r	S	S	S	r/R	S	R	S
Imipenem/relebactam	S	r	r	S	S	S	r/R	S	R	S
Cefiderocol	S	S	S	S/r	S/r	S/r	S/r	S/r	S/r	r
Aztreonam/avibactam	S	R	S	r/R	r/R	S/r	S/r	r/R	S	S
Cefepime/zidebactam	S	r/R	S	S/r	S/r	S/r	S/r	S/r	r/R	S
Cefepime/taniborbactam	S	r/R	S	S/r	S/r	S/r	S/r	S/r	r/R	S

**Figure 1.** Antimicrobial spectrum expected for classical and novel  $\beta$ -lactams and  $\beta$ -lactam- $\beta$ -lactamase inhibitor combinations against most relevant *P. aeruginosa* resistance mechanisms when present alone in clinical strains. To reduce complexity, combinations of resistance mechanisms are not considered, but acknowledged to be frequent among clinical strains. S (green), fully susceptible; r (orange), reduced susceptibility; R (red) clinical resistance. For some antibiotics-mechanisms combinations a range of effect S/r or r/R is considered depending on the specific mechanism or mutation; in such cases, the specific color chosen was that of the most likely phenotype. It is noted however, that variation in the quantitative effect on resistance does occur according to the specific nature of the mechanisms or their expression. \* AmpC (PDC) variants associated with ceftolozane/tazobactam and/or ceftazidime/avibactam resistance. \*\*KPC or GES mutations associated with ceftazidime/avibactam resistance and collateral carbapenem susceptibility.

ST	Clonal Complex	O-antigen Serotype	T3SS	CONTINENT						CARBAPENEMASE							
				N. America	S. America	Europe	Africa	Asia	Oceania	<i>bla</i> GES	<i>bla</i> KPC	<i>bla</i> FIM	<i>bla</i> GIM	<i>bla</i> IMP	<i>bla</i> NDM	<i>bla</i> SPM	<i>bla</i> VIM
ST235	CC235	O11	ExoU+														
ST111	CC111	O12 (O4)	ExoS+														
ST233	CC233	O6	ExoS+														
ST244	CC244	O2	ExoS+														
ST357	CC357	O11	ExoU+														
ST308	CC308	O11	ExoU+														
ST175	CC175	O4	ExoS+														
ST277	CC277	O2	ExoS+														
ST654	CC654	O11	ExoS+														
ST298	CC446	O11	ExoU+														

**Figure 2.** Summary of the main characteristics of the top 10 *P. aeruginosa* high-risk clones. Updated in July 2023 from Del Barrio-Tofiño 2020 (10). Novel descriptions since 2020 are shown in red.