



HAL
open science

ACTION OF THE AUTOMORPHISM GROUP ON THE JACOBIAN OF KLEIN'S QUARTIC CURVE II: INVARIANT THETA-FUNCTIONS

Dimitri Markushevich, Anne Moreau

► **To cite this version:**

Dimitri Markushevich, Anne Moreau. ACTION OF THE AUTOMORPHISM GROUP ON THE JACOBIAN OF KLEIN'S QUARTIC CURVE II: INVARIANT THETA-FUNCTIONS. *Épjournal de Géométrie Algébrique*, In press. hal-04391974

HAL Id: hal-04391974

<https://hal.science/hal-04391974v1>

Submitted on 12 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ACTION OF THE AUTOMORPHISM GROUP ON THE JACOBIAN OF KLEIN'S QUARTIC CURVE II: INVARIANT THETA-FUNCTIONS

DIMITRI MARKUSHEVICH AND ANNE MOREAU

ABSTRACT. Bernstein–Schwarzman conjectured that the quotient of a complex affine space by an irreducible complex crystallographic group generated by reflections is a weighted projective space. The conjecture was proved by Schwarzman and Tokunaga–Yoshida in dimension 2 for almost all such groups, and for all crystallographic reflection groups of Coxeter type by Looijenga, Bernstein–Schwarzman and Kac–Peterson in any dimension. We prove that the conjecture is true for the crystallographic reflection group in dimension 3 for which the associated collineation group is Klein's simple group of order 168. In this case the quotient is the 3-dimensional weighted projective space with weights 1, 2, 4, 7. The main ingredient in the proof is the computation of the algebra of invariant theta-functions. Unlike the Coxeter case, the invariant algebra is not free polynomial, and this was the major stumbling block.

CONTENTS

Introduction	1
1. The group, the lattice and the associated theta-functions	5
2. Theta transformation formula	7
3. Unitary action of G on theta-functions and its character	10
4. Hypersurface in a weighted projective spaces of dimension 4	13
5. Degree-8 hypersurfaces in $\mathbb{P}(1, 1, 2, 4, 7)$ with correct singularities	15
6. Relation to quotients by the Weyl group for the root systems B_3, C_3	19
References	23

INTRODUCTION

A general conjecture of Bernstein and Schwarzman [BS] claims that the quotient of \mathbb{C}^n by the action of an irreducible complex crystallographic group generated by reflections (complex crystallographic reflection group for short) is a weighted projective space. The conjecture is proved for almost all complex crystallographic reflection group for $n = 2$, as well as for irreducible complex crystallographic reflection groups of Coxeter type of any rank $n \geq 2$. A complex crystallographic reflection group is said to be *of Coxeter type* if it is obtained by complexification of a real crystallographic group or, in other words, if the group of its linear parts is conjugate to a finite subgroup of the orthogonal group $\mathbf{O}(n)$. See [MM] for more historical comments and further references.

Since the 80's, the conjecture remained widely open for any irreducible complex crystallographic reflection group of rank $n \geq 3$ which is genuinely complex, that is not of Coxeter type. In the present paper we prove the conjecture for the rank-3 complex crystallographic reflection group Γ of type $[K_{24}]$ in the classification of Popov [Po]. We show that the quotient

$X = \mathbb{C}^3/\Gamma$ is isomorphic to a weighted projective space.¹ More explicitly, the main result of the article is the following:

Main Theorem (Theorem 5.3). The quotient variety \mathcal{J}/G , where $\mathcal{J} = \mathcal{J}(C)$ is the 3-dimensional Jacobian of the plane Klein quartic curve C , by the full automorphism group G of order 336 is isomorphic to the weighted projective space $\mathbb{P}(1, 2, 4, 7)$.

The group $[K_{24}]$ is in several regards the most intriguing one among the rank-3 complex crystallographic reflection groups. Firstly, it is the only one whose projectivized group of linear parts is simple, namely, is equal to Klein's group H of order 168. The groups of linear parts in all the other cases are solvable. Secondly, the quotient \mathbb{C}^3/Γ is isomorphic to the quotient of the Jacobian $\mathcal{J}(C)$ of Klein's quartic curve

$$C := \{x^3y + y^3z + z^3x = 0\} \subset \mathbb{P}^2$$

by the full group

$$G = \{\pm 1\} \times H$$

of its automorphisms as a principally polarized abelian variety. As follows from Hurwitz' bound, C has a maximal number of automorphisms for a curve of genus 3. Algebraic varieties acted on by Klein's group, not only curves, are a recurrent subject of interest in algebraic geometry. Thirdly, $[K_{24}]$ encloses a very rich number-theoretic content, for Klein's curve C is nothing else but the modular curve $X(7)$, or else it can be viewed as a Shimura's curve, while its Jacobian is isomorphic, as an abstract abelian variety, to the cube of the elliptic modular curve $X_0(49)$. The representation of the translation lattice of Γ that we use was given in [Ma].

An important ingredient of our proof for $[K_{24}]$ is the computation of the Hilbert function of the algebra of Γ -invariant theta-functions. The proofs of the conjecture in the Coxeter case obtained in 80's passed through showing that the invariant algebra is free. As the quotient variety is the spectrum of the invariant algebra, the freeness of the latter implies that the quotient is a weighted projective space. But this idea does not work for genuinely complex complex crystallographic reflection groups, since the invariant algebra is not anymore free. In the case of $[K_{24}]$, we succeed to understand the structure of relations between generators of this algebra. It turns out that the ideal of relations is principal and defines a hypersurface of degree 8 in the 4-dimensional weighted projective space $\mathbb{P}(1, 1, 2, 4, 7)$.

On the other hand, in [MM], we have determined the singularities of X : they are images of the orbits whose stabilizers are not generated by reflections. We observed that the singularities of X are analytically equivalent to those of $\mathbb{P}(1, 2, 4, 7)$, which prompted us to conjecture that X is isomorphic to this weighted projective space. The latter also embeds in $\mathbb{P}(1, 1, 2, 4, 7)$ as an octic hypersurface! It can be given by the equation $y_0y_4 = y_3^2$. The last step of our proof is to show that all degree-8 hypersurfaces in $\mathbb{P}(1, 1, 2, 4, 7)$ whose singularities are those of $\mathbb{P}(1, 2, 4, 7)$ are equivalent to $y_0y_4 = y_3^2$ by automorphisms of $\mathbb{P}(1, 1, 2, 4, 7)$.

As a byproduct, we see that $\mathbb{P}(1, 2, 4, 7)$ possesses a nontrivial deformation, obtained by varying the coefficients of the octic in the 4-dimensional weighted projective space. This deformation turns out to be versal, and is a partial smoothing, so that the general member of the

¹The proof of the Bernstein–Schwarzman conjecture for $[K_{24}]$ appeared in the second version of the present article, posted on arxiv in November 2022, and at that moment it was the only non-Coxeter complex crystallographic reflection group of rank > 2 for which the conjecture was established. In March 2023, Eric Rains posted the preprint [Rains], in which he proves the conjecture in full generality.

deformation family is a 2-Gorenstein Fano 3-fold with Picard group \mathbb{Z} and only rigid isolated singularities. Deformations of 1-Gorenstein weighted projective spaces in dimension 3 have been studied in [DS].

Another noteworthy feature of our quotient X is the existence of a natural double cover $\bar{Y} \rightarrow X$, a Calabi–Yau orbifold that can be used as a target space for the superstring compactification; in particular, an interesting question is what its mirror family is. This double cover is obtained as the quotient $\bar{Y} = \mathbb{C}^3/\Gamma_0$, where Γ_0 is the subgroup of index 2 in Γ , the unimodular part of Γ . It can also be realized as an anticanonical hypersurface in $\mathbb{P}(1, 2, 4, 7, 14)$ of dimension 4. However, the known procedure of constructing mirrors of generic hypersurfaces in a weighted projective space does not apply to this case, for \bar{Y} is by no means generic, it is a very special member of the anticanonical system on this weighted projective space having non-isolated singularities. We hope to return to the study of this Calabi–Yau orbifold in future.

In another direction, we expect that the other complex crystallographic reflection groups of rank 3 can be treated in much a similar way; the other classes should be easier to handle, since their groups of linear parts are all solvable. However, it remains unclear how one can predict the weights in general, except for the case of Coxeter type, where they are the exponents of the extended Dynkin diagram of the associated affine root system.

Let us now detail our strategy to prove the main theorem. In general, a *complex crystallographic group* Γ of rank n is a group of affine transformations of \mathbb{C}^n which fits in an exact triple

$$0 \longrightarrow L \longrightarrow \Gamma \longrightarrow d\Gamma \longrightarrow 1,$$

where $L \simeq \mathbb{Z}^{2n}$ is a lattice of maximal rank $2n$ in \mathbb{C}^n , acting by translations, and $d\Gamma$ is a finite subgroup of the unitary group $\mathbf{U}(n)$. A complex crystallographic group is a *complex crystallographic reflection group* if it is generated by complex affine reflections, where an affine transformation of \mathbb{C}^n is called a *reflection* if it is of finite order and its fixed locus is an affine hyperplane. In our case $d\Gamma = G$; there is a unique G -invariant rank-6 lattice L in \mathbb{C}^3 , which turns out to be the period lattice Λ of Klein's quartic C , and the above extension is necessarily split, so that $\Gamma = \Lambda \rtimes G$.

The quotient \mathbb{C}^n/Γ can be thought of as the quotient of the abelian variety $\mathbb{C}^n/L = A$ by the induced action of the finite group of linear parts: $\mathbb{C}^n/\Gamma \simeq A/d\Gamma$. The idea applied in the case of irreducible complex crystallographic reflection groups of Coxeter type in [BS, Lo] is to represent A as the Proj of the graded algebra $S(\mathcal{L})$ of sections of the powers of an ample line bundle \mathcal{L} on A , linearizable by the action of $d\Gamma$, and then \mathbb{C}^n/Γ is the Proj of the $d\Gamma$ -invariant part:

$$S(\mathcal{L}) = \bigoplus_{k=0}^{\infty} H^0(A, \mathcal{L}^k), \quad A = \text{Proj } S(\mathcal{L}), \quad \mathbb{C}^n/\Gamma \simeq A/d\Gamma = \text{Proj} \left(S(\mathcal{L})^{d\Gamma} \right).$$

The sections of the powers of \mathcal{L} are given by theta-functions for the lattice L , and it is proven that the invariant part is a polynomial algebra in $n + 1$ free generators.

Mimicking this approach, we introduce the theta-functions $\theta_{m,k} : \mathbb{C}^3 \rightarrow \mathbb{C}$ of degree $k \geq 0$ for the lattice Λ in such a way that $\{\theta_{m,k}\}_{m \in P_k}$ is a basis of the space $H^0(\mathcal{J}, \mathcal{L}^k)$ of global sections of the k -th power of an appropriate line bundle \mathcal{L} on $\mathcal{J} = \mathbb{C}^3/\Lambda$ defining a principal polarization, where m runs over a set P_k of representatives of $\frac{1}{k}\mathbb{Z}^3/\mathbb{Z}^3$. Unlike the Coxeter case, the thus defined line bundle \mathcal{L} is not $d\Gamma$ -invariant, only even powers of \mathcal{L} can be $d\Gamma$ -linearized, and thus we have to work with the second Veronese subalgebra $S(\mathcal{L}^2)$ of $S(\mathcal{L})$.

We have

$$X = A/d\Gamma \simeq \text{Proj } S(\mathcal{L}^2)^{d\Gamma}, \quad S(\mathcal{L}^2)^{d\Gamma} = \bigoplus_{p=0}^{\infty} H^0(A, \mathcal{L}^{2p})^{d\Gamma}.$$

Thus one cannot expect $S(\mathcal{L}^2)^{d\Gamma}$ to be polynomial; what we prove is the isomorphism between $S(\mathcal{L}^2)^{d\Gamma}$ and the second Veronese algebra of $\mathbb{P}(1, 2, 4, 7)$, the latter being non-polynomial.

Now we describe the content of the paper by sections. Section 1 gathers definitions and preliminary results on the complex crystallographic reflection group Γ , the lattice Λ and associated theta functions.

In Section 2, we compute the transformation formula for the action of the elements of the modular group on our theta functions (Theorem 2.4).

In Section 3, we determine the Hilbert function of the algebra $S(\mathcal{L}^2)^G$, proving that it coincides with the Hilbert function of the second Veronese algebra of $\mathbb{P}(1, 2, 4, 7)$, see Theorem 3.3. This is the second step of the proof of the isomorphism $X \simeq \mathbb{P}(1, 2, 4, 7)$, in which the first one was the study of the singularities of X , done in [MM, Theorem 4.3].

Next, we prove that X admits an embedding in the $\mathbb{P}(1, 1, 2, 4, 7)$ as a degree-8 hypersurface. This is done in Section 4. We first show that there exist 4 homogeneous elements $\varphi_0, \dots, \varphi_3$ of $S(\mathcal{L}^2)^G$ of degrees 1, 1, 2, 4 which are algebraically independent. The four invariant theta-functions φ_i are chosen in an ad hoc way in Lemma 4.2, and their algebraic independence is proved by evaluating their Jacobian. Using the known Hilbert function, we deduce from this that there is a fifth element φ_4 of $S(\mathcal{L}^2)^G$ of degree 7, such that the five functions φ_i generate $S(\mathcal{L}^2)^G$, and this makes X into a degree-8 hypersurface in $\mathbb{P}(1, 1, 2, 4, 7)$ (Theorem 4.1).

Section 5 contains the last step of the proof of the main result: in Proposition 5.2, we show that the degree-8 hypersurfaces in $\mathbb{P}(1, 1, 2, 4, 7)$ whose singularities are those of $\mathbb{P}(1, 2, 4, 7)$ form just one orbit under the action of the group of coordinate changes in $\mathbb{P}(1, 1, 2, 4, 7)$. Thus X is equivalent, modulo a coordinate change, to $\mathbb{P}(1, 2, 4, 7)$ embedded in $\mathbb{P}(1, 1, 2, 4, 7)$ as a degree-8 hypersurface. This implies the main result of the paper (Theorem 5.3). In Remark 5.4, we discuss the nontrivial deformations of $\mathbb{P}(1, 2, 4, 7)$ smoothing out the non-isolated singularity, provided by the family of weighted octics in $\mathbb{P}(1, 1, 2, 4, 7)$.

In Section 6 we look at the quotient $\mathbb{C}^3/\Lambda \rtimes W$, where W is a maximal real reflection subgroup of G , of order 48. The group $\Lambda \rtimes W$ is complex crystallographic, but is not generated by reflections. It is plausible that this quotient is a weak weighted projective space (see Proposition 6.3), and we formulate a conjecture, generalizing this example, which says that if Γ, Γ_1 are commensurable complex crystallographic groups with the same linear parts, such that Γ is complex crystallographic reflection, then \mathbb{C}^n/Γ_1 is a weak weighted projective space (Conjecture 6.4). We also provide in Section 6 some heuristic explanation of our ad hoc choice for generators of $S(\mathcal{L}^2)^G$ (see Remark 6.1).

Acknowledgements. The authors thank Thomas Dedieu et Xavier Roulleau for discussions. D.M. was partially supported by the PRCI SMAGP (ANR-20-CE40-0026-01) and the Labex CEMPI (ANR-11-LABX-0007-01). A.M. is partially supported by the European Research Council (ERC) under the European Union's Horizon 2020 research innovation programme (ERC-2020-SyG-854361-HyperK).

1. THE GROUP, THE LATTICE AND THE ASSOCIATED THETA-FUNCTIONS

Let the complex vector space $V = \mathbb{C}^3$, with coordinates (z_1, z_2, z_3) , be endowed with its standard Hermitian product $\langle | \rangle$ and its standard symmetric bilinear form $(|)$, and consider the complex root system Φ in it, consisting of the 42 vectors obtained from $(2, 0, 0)$, $(0, \alpha, \alpha)$ and $(1, 1, \bar{\alpha})$, where $\alpha = \frac{1+i\sqrt{7}}{2}$, by sign changes and permutations of coordinates. For each $\phi \in \Phi$, the reflection of order 2

$$r_\phi: V \longrightarrow V, \quad z \longmapsto z - 2 \frac{\langle \phi | z \rangle}{\langle \phi | \phi \rangle} \phi$$

sends Φ onto Φ . As $r_\phi = r_{-\phi}$, we obtain 21 reflections in this way. We choose

$$\phi_1 = (0, \alpha, \alpha), \quad \phi_2 = (0, 0, 2), \quad \phi_3 = (1, 1, \bar{\alpha})$$

as the basic roots and denote $r_i = r_{\phi_i}$, $i = 1, 2, 3$. We define the lattice Λ as the root lattice $Q(\Phi)$ of Φ , that is $\Lambda = \sum_{\phi \in \Phi} \mathbb{Z}\phi$. We have

$$\Lambda = \mathcal{O}\phi_1 + \mathcal{O}\phi_2 + \mathcal{O}\phi_3, \quad \mathcal{O} = \mathbb{Z}[\alpha] = \mathbb{Z} + \alpha\mathbb{Z}.$$

By [Ma, 235–236], Λ is the period lattice of Klein's quartic; it can also be represented in the form

$$\Lambda = \{(z_1, z_2, z_3) \in \mathcal{O}^3 : z_1 \equiv z_2 \equiv z_3 \pmod{\alpha}, z_1 + z_2 + z_3 \equiv 0 \pmod{\bar{\alpha}}\}.$$

The group G is the full group of complex-linear automorphisms of Λ . It contains 21 reflections, all of order two, and is generated by the three basic reflections: $G = \langle r_1, r_2, r_3 \rangle$. According to [ST, (10.1)], the following relations are defining for G :

$$(1) \quad r_1^2 = r_2^2 = r_3^2 = (r_1 r_2)^4 = (r_2 r_3)^4 = (r_3 r_1)^3 = (r_1 r_2 r_1 r_3)^3 = 1.$$

Klein's simple group of order 168 is the unimodular part of G :

$$H = \{h \in G : \det(h) = 1\}.$$

It is generated by the antireflections $\rho_\phi := -r_\phi$; of course, the antireflections $\rho_i = -r_i$ associated to the three basic roots ϕ_i ($i = 1, 2, 3$) suffice to generate H . According to [Po], there is a unique extension of Λ by G , the split one, or the semi-direct product $\Gamma = \Lambda \rtimes G$, and it is a complex crystallographic reflection group.

Remark that Λ contains the sublattice αQ , homothetic to the root lattice $Q = Q(C_3)$ of the real root system C_3 , and thus G contains the Weil group $W = W(C_3)$ of order 48. The latter consists of all the monomial matrices of size 3 whose only nonzero elements are ± 1 . We have $W = G \cap \mathbf{O}(3)$, and G is the union of 7 cosets $g_7^i W$ ($i = 0, \dots, 6$) for some element g_7 of order 7. We can choose $g_7 = \rho_1 \rho_2 \rho_3 = -r_1 r_2 r_3 = \frac{1}{2} \begin{pmatrix} -1 & 1 & \alpha \\ -\bar{\alpha} & -\bar{\alpha} & 0 \\ 1 & -1 & \alpha \end{pmatrix}$. This is a handy way to enumerate all the elements of G , say, for computer checks of some properties.

Let M denote the weight lattice of C_3 , $M = Q^* = \mathbb{Z}^3 + \frac{1}{2}(1, 1, 1)$. Then $\Lambda = 2\bar{\alpha}M + \alpha Q$ is generated by the columns of the 6-by-3 period matrix $\Omega = (\omega_1 | \omega_2)$, where ω_1, ω_2 are square blocks of size 3, the columns of ω_2 being α times the elements of a basis of C_3 , and we set:

$$(2) \quad \omega_2 = \alpha C, \quad \omega_1 = -2\bar{\alpha}(C^\top)^{-1}, \quad C = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 2 \end{pmatrix}.$$

The normalized period matrix of \mathcal{J} is obtained by multiplying Ω on the left by ω_2^{-1} :

$$(3) \quad \omega_2^{-1}\Omega = (Z | I), \quad Z = \omega_2^{-1}\omega_1 = \tau B,$$

where $\tau = -\bar{\alpha}^2$ and

$$(4) \quad B = (C^\top C)^{-1} = \frac{1}{4} \begin{pmatrix} 4 & 4 & 2 \\ 4 & 8 & 4 \\ 2 & 4 & 3 \end{pmatrix}.$$

is real symmetric and positive definite, so that $Z = \tau B \in \mathfrak{H}_3$, where \mathfrak{H}_r denotes the Siegel half-space of complex symmetric matrices of size r with positive definite imaginary part.

Let $c_1, c_2 \in \mathbb{R}^r$ be two vectors, $k \in \mathbb{Z}$, $k \geq 0$, $Z \in \mathfrak{H}_r$. The classical theta-function with period Z , characteristic $c_1 Z + c_2$ and of degree k is the complex-valued function

$$\mathbb{C}^r \ni v \mapsto \theta_k^{[c_1]}(v, Z) = \sum_{u \in \mathbb{Z}^r} e^{2\pi i k [(v+c_2)^\top (u+c_1) + \frac{1}{2}(u+c_1)^\top Z (u+c_1)]}.$$

When $k = 1$, the subscript k is usually omitted. For $k = 1$ and $c_1 = c_2 = 0$, the function $\theta_{[0]}^{[0]}(\bullet, Z)$ represents a section of a uniquely determined line bundle \mathcal{L} on the principally polarized abelian variety $A = \mathbb{C}^r / (ZZ^r + \mathbb{Z}^r)$, and then for any k, c_1, c_2 , the function $\theta_k^{[c_1]}(\bullet, Z)$ represents a section of the line bundle $T_{c_1 Z + c_2}^*(\mathcal{L}^k)$, the pullback of the tensor power \mathcal{L}^k of \mathcal{L} by the translation by the point $c_1 Z + c_2 \bmod (ZZ^r + \mathbb{Z}^r)$ of A . If we choose a set P_k of representatives of $\frac{1}{k}\mathbb{Z}^r / \mathbb{Z}^r$, then the k^3 theta-functions from $\{\theta_k^{[c_1]}(\bullet, Z)\}_{m \in P_k}$ represent a basis of $H^0(A, T_{c_1 Z + c_2}^*(\mathcal{L}^k))$. See e.g. [BL].

Definition 1.1. Let Λ be as above, $Z \in \mathfrak{H}_3$ as in (3), $r = 3$, $k \in \mathbb{Z}$, $k \geq 1$, and $m \in \frac{1}{k}\mathbb{Z}^3$. We define the theta-function for Λ of degree k with characteristic m by the formula

$$\theta_{m,k}(z) = \theta_k^{[m]}(\omega_2^{-1}z, Z) \text{ for any } z \in V.$$

There is a unique line bundle \mathcal{L} on $\mathcal{J} = V/\Lambda \simeq V/(ZZ^3 + \mathbb{Z}^3)$ defining a principal polarization such that the functions $\{\theta_{m,k}\}_{m \in P_k}$ form a basis of $H^0(\mathcal{J}, \mathcal{L}^k)$, where P_k is a set of representatives of $\frac{1}{k}\mathbb{Z}^3 / \mathbb{Z}^3$, for all $k \geq 1$.

We denote by $\mathbf{Sp}(2r, \mathbb{Z})$ the symplectic group of automorphisms of \mathbb{Z}^{2r} preserving the skew-symmetric bilinear form given by the matrix E :

$$\mathbf{Sp}(2r, \mathbb{Z}) = \{A \in M_{2r}(\mathbb{Z}) : A^\top E A = E\}, \quad E = E_r = \begin{pmatrix} 0 & -I_r \\ I_r & 0 \end{pmatrix},$$

where I (or I_r) denotes the identity matrix (of size r). We represent E and the matrices $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{Sp}(2r, \mathbb{Z})$ by their blocks of size r . A crucial ingredient of our computation of the action of G on the theta-functions $\theta_{m,k}$ is a transformation formula under modular transformations. The action of the modular transformation γ on a classical theta-function is defined by

$$(5) \quad (\theta_k^{[c_1]})^\gamma(v, Z) = \theta_k^{[c_1]}(((cZ + d)^\top)^{-1}v, (aZ + b)(cZ + d)^{-1}).$$

The following is a particular case of Igusa's Theorem for a polarization of type (k, k, \dots, k) :

Theorem 1.2 (Igusa, [Ig, Theorem II.5.6]). For every $(v, Z) \in \mathbb{C}^r \times \mathfrak{H}_r$, $c_1, c_2 \in \mathbb{R}^g$, $m \in P_k$ and $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{Sp}(2r, \mathbb{Z})$, the matrix $cZ + d$ is invertible and we have:

$$\left(\theta_k \begin{bmatrix} m+c_1 \\ c_2' \end{bmatrix} \right)^\gamma (v, Z) = e^{\pi i k v^\top (cZ+d)^{-1} c v} \det(cZ + d)^{\frac{1}{2}} \cdot \sum_{m' \in P_k} u_{m, m'} \theta_k \begin{bmatrix} m'+c_1 \\ c_2' \end{bmatrix} (v, Z),$$

where

$$\begin{bmatrix} c_1' \\ c_2' \end{bmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} (cd^\top)_0 \\ (ab^\top)_0 \end{bmatrix},$$

$(u_{m, m'}) \in \mathbf{U}(k^r)$, and $(h)_0$ denotes the column vector of the diagonal elements of h for any square matrix h .

In the next section, we will represent the automorphisms from G by modular transformations and compute explicitly the matrices $(u_{m, m'})$ for even k . As we will see, the presence of the half-integer inhomogeneous term in the transformation formula for the characteristics c_1, c_2 implies the non-invariance of \mathcal{L} under the action of G ; however, the even powers of \mathcal{L} are G -invariant.

2. THETA TRANSFORMATION FORMULA

We keep the notation of the previous section. The elements of G are complex 3-by-3 matrices leaving invariant the lattice Λ . As Λ is generated by the columns of the 6-by-3 matrix $\Omega = (\omega_1 | \omega_2)$, we can associate to each $g \in G$ a matrix $\gamma = \gamma_g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{Sp}(6, \mathbb{Z})$ in such a way that

$$(\omega_1 | \omega_2) \begin{pmatrix} a^\top & c^\top \\ b^\top & d^\top \end{pmatrix} = g(\omega_1 | \omega_2).$$

Obviously, the map $g \mapsto \gamma_g$ is a group homomorphism.

Lemma 2.1. Let $g \in G$, and let $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be as above. Then the following properties are verified:

- (i) $(aZ + b)(cZ + d)^{-1} = Z$;
- (ii) $\det(cZ + d) = \det g = \pm 1$;
- (iii) $\det d = \pm 1$.

Proof. (i) We have:

$$\begin{aligned} Z^\top &= Z = \omega_2^{-1} \omega_1 = (g\omega_2)^{-1} (g\omega_1) = (\omega_1 c^\top + \omega_2 d^\top)^{-1} \cdot (\omega_1 a^\top + \omega_2 b^\top) = \\ &= (Zc^\top + d^\top)^{-1} (Za^\top + b^\top) = ((aZ + b)(cZ + d)^{-1})^\top. \end{aligned}$$

(ii) By definition of γ , $Zc^\top + d^\top = (cZ + d)^\top$ is the matrix of g in the basis of \mathbb{C}^3 given by the columns of ω_2 , so $\det g = \det (cZ + d)^\top$.

(iii) This is verified by a direct computation of the matrices γ_g for all elements $g \in G$. \square

Corollary 2.2. For $z \in V$, we have

$$g \cdot \theta_{0,1}(z) = (\theta_{[0]}^0)^{\gamma_g} (v, Z) = \chi_g \theta_{[\nu']}^{\nu'} (v, Z),$$

where

$$v = \omega_2^{-1} z, \quad \chi_g = e^{\pi i v^\top (cZ+d)^{-1} c v} (\det g)^{1/2}, \quad \begin{bmatrix} \nu' \\ \nu'' \end{bmatrix} = -\frac{1}{2} \begin{pmatrix} d^\top & -b^\top \\ -c^\top & a^\top \end{pmatrix} \begin{bmatrix} (cd^\top)_0 \\ (ab^\top)_0 \end{bmatrix}.$$

Proof. This immediately follows from the previous lemma, Igusa's Theorem with $k = 1$ and the inversion formula

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \begin{pmatrix} d^\top & -b^\top \\ -c^\top & a^\top \end{pmatrix}$$

for matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{Sp}(2r, \mathbb{Z})$. \square

We see that the theta-function $\theta_{0,1}$, representing a section of \mathcal{L} , acquires a half-integer characteristic upon the action by an element $g \in G$ whenever the diagonal elements of the integer matrices cd^\top , ab^\top are not all even. In this case $g^*\mathcal{L}$ is not \mathcal{L} , but the translation of \mathcal{L} by a point of order 2.

Corollary 2.3. (i) For $g \in G$ we have the equivalence $g^*\mathcal{L} \simeq \mathcal{L} \iff g \in W$, where $W = W(C_3) = G \cap \mathbf{O}(3)$ is the subgroup of real matrices in G .

(ii) $g^*\mathcal{L}^k \simeq \mathcal{L}^k$ for all $g \in G \iff k$ is even.

Proof. By a direct computation we verify that for $g \in G$, the diagonal elements of the integer matrices cd^\top , ab^\top are all even if and only if $g \in W$. This implies both assertions. \square

Thus the problem of calculating the action of G on $H^0(\mathcal{J}, \mathcal{L}^k)$ has sense only for even k . We are now going to calculate, in a particular case and only for even k , the matrix $(u_{m,m'})$ from Igusa's Theorem up to proportionality; we denote the matrix we find by $(\tilde{u}_{m,m'})$, as it is a multiple of Igusa's matrix $(u_{m,m'})$ which is not necessarily unitary.

For $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{Sp}(2r, \mathbb{Z})$ with $\det d = \pm 1$ we define:

$$\tilde{a} = a - bd^{-1}c = (d^\top)^{-1}, \quad \tilde{b} = \tilde{b}^\top = b\tilde{a}^\top, \quad \tilde{c} = \tilde{c}^\top = -cd^\top.$$

Theorem 2.4. Let k be a positive even integer; P_k a set of representatives of $\frac{1}{k}\mathbb{Z}^r/\mathbb{Z}^r$, $\gamma \in \mathbf{Sp}(2r, \mathbb{Z})$ such that $\det d = \pm 1$, and $Z = \tau B$ for a real symmetric positive definite matrix B of size r , where $\tau \in \mathbb{C}$, $\text{Im } \tau > 0$. Then $\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix}^\gamma = \chi \sum_{m' \in P_k} \tilde{u}_{m,m'} \theta_k \begin{bmatrix} m' \\ 0 \end{bmatrix}$, where $\chi = \chi_\gamma(v, Z)$ is a nowhere vanishing analytic function on $\mathbb{C}^r \times \mathfrak{H}_r$, depending on γ , and

$$\tilde{u}_{m,m'} = e^{\pi i k \tilde{b}[m]} \sum_{\hat{m} \in P_k} e^{2\pi i k (m - dm' + \frac{1}{2} \tilde{c} \hat{m})^\top \hat{m}}.$$

Proof. We decompose γ in a product of elementary transformations as follows:

$$\gamma = \begin{pmatrix} 1 & \tilde{b} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & \tilde{c} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \tilde{a} & 0 \\ 0 & d \end{pmatrix},$$

$\parallel \qquad \parallel \qquad \parallel \qquad \parallel \qquad \parallel$
 $\sigma_1 \qquad \sigma_2 \qquad \sigma_3 \qquad \sigma_4 \qquad \sigma_5$

and apply the factors of this decomposition successively.

Step 1. $(\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix})^{\sigma_1}(v, Z) = \theta_k \begin{bmatrix} m \\ 0 \end{bmatrix}(v, Z + \tilde{b}) =$

$$\sum_{u \in \mathbb{Z}^r} e^{2\pi i k (v^\top(u+m) + \frac{1}{2}(u+m)^\top Z(u+m))} \cdot e^{\pi i k \tilde{b}[u]} \cdot e^{2\pi i k m^\top \tilde{b}u} \cdot e^{\pi i k \tilde{b}[m]} = e^{\pi i k \tilde{b}[m]} \theta_k \begin{bmatrix} m \\ 0 \end{bmatrix}(v, Z),$$

where we use the notation $A[u] = u^\top A u$ for any symmetric matrix A of size r and any vector $u \in \mathbb{C}^r$, and $e^{\pi i k \tilde{b}[u]} = e^{2\pi i k m^\top \tilde{b}u} = 1$ for any $u \in \mathbb{Z}^r$, k even.

Step 2. We have

$$(\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix})^{\sigma_1 \sigma_2}(v, Z) = e^{\pi i k \tilde{b}[m]} \theta_k \begin{bmatrix} m \\ 0 \end{bmatrix}(-Z^{-1}v, -Z^{-1}) = e^{\pi i k \tilde{b}[m]} \theta_k \begin{bmatrix} -m \\ 0 \end{bmatrix}(Z^{-1}v, -Z^{-1});$$

$$\theta_k \begin{bmatrix} -m \\ 0 \end{bmatrix} (Z^{-1}v, -Z^{-1}) = \sum_{u \in \mathbb{Z}^r} e^{\frac{2\pi ik}{\tau} (v^\top B^{-1}(u-m) - \frac{1}{2}(u-m)^\top B^{-1}(u-m))} = e^{\frac{\pi ik}{\tau} A[v]} \sum_{u \in \mathbb{Z}^r} e^{-\frac{\pi ik}{\tau} A[u-m-v]},$$

where $A = B^{-1}$ and we used the hypothesis that $Z = \tau B$. To transform the latter expression, we apply the Jacobi inversion formula (see for example [Gu, Ch. VI]):

$$\sum_{u \in \mathbb{Z}^r} e^{\pi i t A[x+u]} = \frac{1}{\sqrt{(-it)^r \det A}} \sum_{u \in \mathbb{Z}^r} e^{-\frac{\pi i}{t} A^{-1}[u] + 2\pi i x^\top u},$$

for $x \in \mathbb{R}^r$, $t \in \mathbb{C}$, $\text{Im } t > 0$, $A \in M_r(\mathbb{R})$, $A^\top = A$, $A > 0$. We set $t = -k/\tau$ and $x = -m - v$, and we obtain:

$$(\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix})^{\sigma_2} (v, Z) = \chi_0 \sum_{u \in \mathbb{Z}^r} e^{\pi i k Z \begin{bmatrix} u \\ k \end{bmatrix} - 2\pi i k (m+v)^\top \frac{u}{k}} = \chi_0 \sum_{u' \in \frac{1}{k}\mathbb{Z}^r} e^{\pi i k Z[u'] - 2\pi i k (m+v)^\top u'},$$

where $\chi_0(v, Z) = \frac{e^{\pi i k Z^{-1}[v]}}{\sqrt{(-ik)^r \det Z^{-1}}}$. Now each $u' \in \frac{1}{k}\mathbb{Z}^r$ has a unique representation $u' = u + m'$ with $u \in \mathbb{Z}^r$ and $m' \in P_k$, and we obtain

$$(\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix})^{\sigma_1 \sigma_2} (v, Z) = \chi_0 e^{\pi i k \tilde{b}[m]} \sum_{m' \in P_k} e^{2\pi i k m^\top m'} \theta_k \begin{bmatrix} m' \\ 0 \end{bmatrix} (v, Z),$$

where $\chi_0 = \chi_0(v, Z)$ is the nowhere vanishing analytic function in v, Z defined above.

Step 3. We have

$$(\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix})^{\sigma_1 \sigma_2 \sigma_3} (v, Z) = \chi_1 e^{\pi i k \tilde{b}[m]} \sum_{m' \in P_k} e^{2\pi i k (m^\top m' + \frac{1}{2} \tilde{c}[m'])} \theta_k \begin{bmatrix} m' \\ 0 \end{bmatrix} (v, Z),$$

where $\chi_1(v, Z) = \chi_0(v, Z + \tilde{c})$.

Step 4. The calculation is similar to that in Step 2:

$$\begin{aligned} & (\theta_k \begin{bmatrix} m \\ 0 \end{bmatrix})^{\sigma_1 \sigma_2 \sigma_3 \sigma_4} (v, Z) = \\ & \chi_1 (Z^{-1}v, -Z^{-1}) e^{\pi i k \tilde{b}[m]} \sum_{m' \in P_k} e^{2\pi i k (m + \frac{1}{2} \tilde{c} m')^\top m'} \theta_k \begin{bmatrix} m' \\ 0 \end{bmatrix} (Z^{-1}v, -Z^{-1}) = \\ & \chi_2 \sum_{m' \in P_k} \sum_{m'' \in P_k} \zeta_{m', m''} \theta_k \begin{bmatrix} m'' \\ 0 \end{bmatrix} (v, Z), \text{ where } \zeta_{m', m''} = e^{\pi i k \tilde{b}[m]} e^{2\pi i k (m + \frac{1}{2} \tilde{c} m' - m'')^\top m'} \end{aligned}$$

and $\chi_2(v, Z) = \chi_1(Z^{-1}v, -Z^{-1})\chi_0(v, Z)$.

$$\text{Step 5. } \theta_k \begin{bmatrix} m \\ 0 \end{bmatrix}^\gamma (v, Z) = \chi \sum_{m', m''} \zeta_{m', m''} \theta_k \begin{bmatrix} m'' \\ 0 \end{bmatrix} \begin{pmatrix} \tilde{a} & 0 \\ 0 & d \end{pmatrix} (v, Z) = \chi \sum_{m', m''} \zeta_{m', m''} \theta_k \begin{bmatrix} \tilde{a}^\top m'' \\ 0 \end{bmatrix} (v, Z) =$$

$$\chi \sum_{m', m''} \zeta_{m', (\tilde{a}^\top)^{-1} m''} \theta_k \begin{bmatrix} m'' \\ 0 \end{bmatrix} (v, Z) = \chi \sum_{m''} \tilde{u}_{m, m''} \theta_k \begin{bmatrix} m'' \\ 0 \end{bmatrix} (v, Z),$$

where $\tilde{u}_{m, m''}$ is as in the statement of the theorem and $\chi(v, Z) = \chi_2((d^\top)^{-1}v, (d^\top)^{-1}Zd^{-1})$. \square

Together with Igusa's Theorem and Lemma 2.1, this theorem obviously implies the following corollary.

Corollary 2.5. *For every even $k \geq 0$, the application of Theorem 2.4 to the theta-functions $\theta_{m,k}$ for the lattice Λ introduced in Definition 1.1 provides a map $g \mapsto \tilde{U}_g$, where \tilde{U}_g is the complex matrix $(\tilde{u}_{m,m'})$ of size k^3 defined in the statement of Theorem 2.4 with γ_g in place of γ , and this map provides a group homomorphism from G to the projective unitary group $\mathbf{PU}(k^3) = \mathbf{U}(k^3)/(\text{homotheties})$.*

3. UNITARY ACTION OF G ON THETA-FUNCTIONS AND ITS CHARACTER

Neither Igusa's Theorem, nor our approach applied in Theorem 2.4, allow us to conclude that the matrices \tilde{U}_g can be normalized by multiplying by some constants ϵ_g in such a way that the normalized map $g \mapsto U_g = \epsilon_g \tilde{U}_g$ is a group homomorphism $G \rightarrow \mathbf{U}(k^3)$. However we managed to find convenient constants ϵ_g by trial and error. We will write $U_g^{(k)}, \tilde{U}_g^{(k)}$ when we want to specify the degree k of theta-functions on which U_g, \tilde{U}_g act.

Proposition 3.1. *Let r_1, r_2, r_3 be the basic reflections generating G introduced in Section 1. Set $U_j = \frac{1}{k^3} \tilde{U}_{r_j}$ for $j = 1, 2$, $U_3 = \frac{1}{ik^3} \tilde{U}_{r_3}$. Then $U_1, U_2, U_3 \in \mathbf{U}(k^3)$, and we denote by $U_G^{(k)}$ the subgroup of $\mathbf{U}(k^3)$ generated by these three matrices.*

(1) *The U_j satisfy the same relations (1) as the basic reflections r_j :*

$$U_1^2 = U_2^2 = U_3^2 = (U_1 U_2)^4 = (U_2 U_3)^4 = (U_3 U_1)^3 = (U_1 U_2 U_1 U_3)^3 = 1.$$

(2) *$U_G^{(2)} \simeq H$ and $U_G^{(k)} \simeq G$ for all even $k \geq 4$.*

Proof. We verify this for $k = 2, 4, 6$ by direct computation using the computer algebra system Macaulay2 [M2]; the result for all even k follows from the fact that the algebra of even-degree theta-functions on a p.p.a.v. is generated in degrees ≤ 6 . \square

We thus have a unitary representation ρ_k of G on the space $H^0(\mathcal{J}, \mathcal{L}^k)$ of dimension k^3 for each even $k \geq 0$, defined by substituting the U_j in place of the r_j in the words in the r_j defining all the elements of G . We denote by χ_k the character of this representation. In order to determine it, we start by fixing the choice of representatives of the conjugacy classes of G . The Klein's simple group H has 6 conjugacy classes, represented by the following elements:

$$g_1 = 1, \quad g_2 = \rho_1, \quad g_3 = \rho_1 \rho_3 \rho_1 \rho_2, \quad g_4 = \rho_1 \rho_2, \quad g_7 = \rho_1 \rho_2 \rho_3, \quad g_7^{-1},$$

where $\rho_i = -r_i$ are the basic antireflections and the subscript p in g_p stands for the order of g_p .

The conjugacy classes of G are deduced from these in an obvious way: to every conjugacy class $\text{Cl}_H(g)$ in H correspond two conjugacy classes in G of the same length: $\text{Cl}_G(g) = \text{Cl}_H(g)$ and $\text{Cl}_G(-g) = -\text{Cl}_H(g)$. Also to each irreducible representation f of H correspond two irreducible representations of G , $\tilde{f} = f \circ \pi$ and $\tilde{f} \otimes \det$, where $\pi : G \rightarrow H \simeq G/\langle -1 \rangle$ is the natural surjection.

The lengths of the conjugacy classes of H are given by the next table, providing the characters of H . The characters of G are easily deduced from it.

g	g_1	g_2	g_3	g_4	g_7	g_7^{-1}
$ \text{Cl}_H(\gamma) $	1	21	56	42	24	24
χ_1	1	1	1	1	1	1
χ_3	3	-1	0	1	$-\bar{\alpha}$	$-\alpha$
$\chi_{\bar{3}}$	3	-1	0	1	$-\alpha$	$-\bar{\alpha}$
χ_6	6	2	0	0	-1	-1
χ_7	7	-1	1	-1	0	0
χ_8	8	0	-1	0	1	1

Theorem 3.2. For any even $k > 0$, the character χ_k of ρ_k takes the following values on the above representatives of the conjugacy classes:

g_1	$-g_1$	g_2	$-g_2$	g_3	$-g_3$	g_4	$-g_4$	g_7	$-g_7$	g_7^{-1}	$-g_7^{-1}$
k^3	8	$2k$	k^2	k	2	k	$3 + (-1)^{\frac{k}{2}}$	$\begin{pmatrix} k \\ 7 \end{pmatrix}$ if $7 \nmid k$, $-i\sqrt{7}$ if $7 \mid k$	1	$\begin{pmatrix} k \\ 7 \end{pmatrix}$ if $7 \nmid k$, $i\sqrt{7}$ if $7 \mid k$	1

where

$$\begin{pmatrix} k \\ 7 \end{pmatrix} = \begin{cases} 1 & \text{if } k \equiv 1, 2 \text{ or } 4 \pmod{7} \\ -1 & \text{if } k \equiv 3, 5 \text{ or } 6 \pmod{7} \\ 0 & \text{if } k \equiv 0 \pmod{7}. \end{cases}$$

is the Legendre symbol.

Proof. The result is obvious for g_1 and needs some reasoning, following the same pattern, in the other cases. We will illustrate this reasoning on the example of $g = g_7$, where the details of the calculation are most involved. We first observe that the normalization constant for $g = g_7$ is $\epsilon_g = \frac{1}{ik^3}$, so that

$$\chi_k(g) = \text{tr } U_g = \frac{1}{ik^3} \Sigma_k, \text{ where } \Sigma_k = \sum_{m \in P_k} \tilde{u}_{m,m}.$$

By Theorem 2.4,

$$\Sigma_k = \sum_{m, m' \in P_k} e^{\pi i k (2m'^T (\text{id} - d)m + \tilde{b}[m] + \tilde{c}[m'])} = \sum_{0 \leq x_i \leq k-1} e^{\frac{\pi i}{k} K[x]},$$

where we pass to the summation over the integer column vector $x = (x_1, \dots, x_6)^T = k(m', m)^T \in \mathbb{Z}^6$, $K[x] = x^T K x$, as before, and K is the following integer matrix of size 6:

$$K = \begin{pmatrix} \tilde{c} & I_3 - d \\ I_3 - d^T & \tilde{b} \end{pmatrix}.$$

Explicitly, we have:

$$\gamma_g = \left(\begin{array}{ccc|ccc} -1 & 0 & 1 & 0 & -2 & -1 \\ -1 & 1 & 0 & 0 & -4 & -2 \\ 0 & 0 & 1 & -1 & -3 & -2 \\ \hline 0 & 0 & 0 & -1 & -1 & 0 \\ -1 & 1 & -1 & 1 & 0 & 0 \\ 1 & -1 & 2 & -1 & -1 & -1 \end{array} \right), \quad K = \left(\begin{array}{ccc|ccc} 0 & 0 & 0 & 2 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & -1 & 2 & 1 & 1 & 2 \\ \hline 2 & -1 & 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 & 4 & 2 \\ 0 & 0 & 2 & 1 & 2 & 2 \end{array} \right).$$

By Gauss diagonalization over \mathbb{Z} , we reduce the quadratic form $x \mapsto K[x]$ to the diagonal representation $y = (y_1, \dots, y_6) \mapsto y_1^2 + y_2^2 + y_3^2 - y_4^2 - y_5^2 - 7y_6^2$. Thus

$$\begin{aligned} \Sigma_k &= \sum_{0 \leq x_i \leq k-1} e^{\frac{\pi i}{k} K[x]} = \sum_{0 \leq y_i \leq k-1} e^{\frac{\pi i}{k} (y_1^2 + y_2^2 + y_3^2 - y_4^2 - y_5^2 - 7y_6^2)} = \\ &= \left(\sum_{y=0}^{k-1} e^{\frac{\pi i}{k} y^2} \right)^3 \left(\sum_{y=0}^{k-1} e^{-\frac{\pi i}{k} y^2} \right)^2 \left(\sum_{y=0}^{k-1} e^{-\frac{7\pi i}{k} y^2} \right). \end{aligned}$$

The exponential sums in the last line belong to the class of Gauss sums. We use the following formula for Gauss sums from [BEW]:

$$G(q, r) = \sum_{n=0}^{r-1} e^{\frac{2\pi i q}{r} n^2} = (1+i) \kappa_q^{-1} \sqrt{r} \left(\frac{r}{q} \right),$$

if $2 \nmid q$, $4 \mid r$ and $\gcd(q, r) = 1$, where $\kappa_q = \begin{cases} 1 & \text{if } q \equiv 1 \pmod{4} \\ i & \text{if } q \equiv 3 \pmod{4}, \end{cases}$ and $\left(\frac{r}{q} \right)$ is the Jacobi

symbol (which coincides with the Legendre symbol when q is prime). Applying this formula, we obtain:

$$\begin{aligned} \sum_{y=0}^{k-1} e^{\frac{\pi i}{k} y^2} &= \frac{1}{2} G(1, 2k) = \frac{1+i}{\sqrt{2}} \sqrt{k}, \\ \sum_{y=0}^{k-1} e^{-\frac{7\pi i}{k} y^2} &= \frac{1}{2} \overline{G(7, 2k)} = \frac{1+i}{\sqrt{2}} \left(\frac{k}{7} \right) \sqrt{k} \text{ if } 7 \nmid k, \\ \sum_{y=0}^{k-1} e^{-\frac{7\pi i}{k} y^2} &= \frac{7}{2} \overline{G(1, 2k_1)} = \frac{1-i}{\sqrt{2}} \sqrt{7k} \text{ if } k = 7k_1. \end{aligned}$$

We thus finish the calculation of Σ_k and obtain the value of $\chi_k(g)$ given in the table. \square

By taking the scalar product of χ_k with the trivial character, we obtain:

Theorem 3.3. *The Hilbert function of the algebra of invariant theta-functions*

$$S(\mathcal{L}^2)^G = \bigoplus_{p=0}^{\infty} H^0(\mathcal{J}, \mathcal{L}^{2p})^G,$$

is given by:

$$h_{S(\mathcal{L}^2)^G} \left(\frac{k}{2} \right) = \frac{1}{336} \left[k^3 + 21k^2 + 140k + 294 + (-1)^{\frac{k}{2}} \times 42 + 48 \left(\frac{k}{7} \right) \right].$$

This coincides with the Hilbert function of the second Veronese algebra of $\mathbb{P}(1, 2, 4, 7)$.

Proof. The formula for the Hilbert function is an immediate consequence of Theorem 3.2. The Hilbert series for $\mathbb{P} = \mathbb{P}(1, 2, 4, 7)$ is given by

$$h_{\mathbb{P}}(t) = \frac{1}{(1-t)(1-t^2)(1-t^4)(1-t^7)},$$

and that of the second Veronese of \mathbb{P} is

$$h_{\mathbb{P}}^{(2)}(t) = \frac{1}{2} (h_{\mathbb{P}}(t^{1/2}) + h_{\mathbb{P}}(-t^{1/2})) = \frac{1+t^4}{(1-t)^2(1-t^2)(1-t^7)}.$$

Denote h_p the coefficients of the latter series, so that $h_{\mathbb{P}}^{(2)}(t) = \sum_{p \geq 0} h_p t^p$; by [St, Theorem 4.4.1], the Hilbert function $p \mapsto h_p$ is a rational quasipolynomial of degree 3 whose coefficients are periodic with period 14. The dominant coefficient being constant, it suffices to compare the initial segments of the sequences $h_{S(\mathcal{L}^2)^G}(p), h_p$ of length 42 to see that they coincide for all $p \geq 0$. In this way we conclude the proof. \square

4. HYPERSURFACE IN A WEIGHTED PROJECTIVE SPACES OF DIMENSION 4

For any theta-function θ of even degree $k > 0$, we denote by $\mathcal{R}_G^{(k)}$ the Reynolds operator

$$(6) \quad \mathcal{R}_G^{(k)}(\theta) = \frac{1}{336} \sum_{g \in G} U_g^{(k)}(\theta).$$

We will call $\mathcal{R}_G^{(k)}(\theta)$ the G -average of θ ; for given k , the G -averages of theta-functions of degree k represent sections of \mathcal{L}^k which generate $H^0(\mathcal{J}, \mathcal{L}^k)^G$ over \mathbb{C} .

We will identify sections of \mathcal{L}^k with regular functions on the total space $\mathbf{V}(\mathcal{L}^{-1})$ of the line bundle \mathcal{L}^{-1} which are fiber-homogeneous of degree k . Explicitly, $\mathbf{V}(\mathcal{L}^{-1})$ is the quotient of the trivial line bundle $\mathbb{C} \times V$ over $V = \mathbb{C}^3$ by the action of $\Lambda \simeq \{Zu' + u'' \mid u', u'' \in \mathbb{Z}^3\}$:

$$Zu' + u'' : (t, v) \mapsto \left(e^{2\pi i(v^T u' + \frac{1}{2}Z[u'])} t, v + Zu' + u'' \right),$$

and to each degree- k theta-function θ we associate the Λ -invariant function

$$(7) \quad \tilde{\theta} : \mathbb{C} \times V \rightarrow \mathbb{C}, (t, v) \mapsto t^k \theta(v),$$

which descends to a fiber-homogeneous function of degree k on $\mathbf{V}(\mathcal{L}^{-1})$ denoted by the same symbol $\tilde{\theta}$.

Theorem 4.1. *There exist five G -invariant theta-functions $\varphi_i = \varphi_{i, k_i}$ of degrees k_i , where $(k_0, \dots, k_4) = (2, 2, 4, 8, 14)$, such that the associated sections of the powers \mathcal{L}^{k_i} of \mathcal{L} generate $S(\mathcal{L}^2)^G$. The ideal of relations between these generators is generated by a single relation of weighted degree 16, so that the quotient $X = \mathcal{J}/G$ is isomorphic to a hypersurface in the weighted projective space $\mathbb{P}(1, 1, 2, 4, 7)$ of weighted degree 8.*

Proof. It suffices to show that we can find G -invariant theta-functions $\varphi_0, \varphi_1, \varphi_2, \varphi_3$ of respective degrees 2, 2, 4, 8 such that the associated fiber-homogeneous functions $\tilde{\varphi}_i$ on $\mathbf{V}(\mathcal{L}^{-1})$ are algebraically independent. This is done below in Lemma 4.2. Then the functions φ_i , $i = 0, \dots, 3$, generate the free polynomial subalgebra \mathcal{S}' in $\mathcal{S} = S(\mathcal{L}^2)^G$, and comparing the initial segments of their Hilbert functions,

$$\begin{aligned} (h_{\mathcal{S}}(p))_{p \geq 0} &= (1, 2, 4, 6, 10, 14, 20, 27, 36, 46, 58, \dots), \\ (h_{\mathcal{S}'}(p))_{p \geq 0} &= (1, 2, 4, 6, 10, 14, 20, 26, 35, 44, 56, \dots), \end{aligned}$$

we see that the four φ_i generate \mathcal{S} in degrees < 7 , and $h_{\mathcal{S}}(7) = h_{\mathcal{S}'}(7) + 1$, so that one extra generator of degree 7, say φ_4 is needed to generate \mathcal{S} up to degree 7. Furthermore, there are 35 monomials in φ_i , $i = 0, \dots, 3$, of degree 8, and two more monomials $\varphi_0\varphi_4, \varphi_1\varphi_4$ involving φ_4 , and as $h_{\mathcal{S}}(7) = 36$ is one less than the number of monomials of degree 8, there is precisely one linear relation $F_8 = 0$ between the 37 monomials of degree 8. One easily verifies that $h_{\mathcal{S}}(p) = h_R(p) - h_R(p - 8)$ for all $p \in \mathbb{Z}$, where R denotes a free polynomial algebra with generators of degrees 1, 1, 2, 4, 7, so that the natural map $R/F_8R \rightarrow \mathcal{S}$ is an

isomorphism. Thus X is isomorphic to a hypersurface $F_8 = 0$ of degree 8 in the weighted projective space $\mathbb{P}(1, 1, 2, 4, 7) = \text{Proj } R$. \square

For the algebraic independence of $\tilde{\varphi}_0, \dots, \tilde{\varphi}_3$ it is necessary and sufficient that the Jacobian $J = J(\tilde{\varphi}_i)$ is not identically zero. We will present explicitly an ad hoc example of $\varphi_i(v)$ lying in the images of the Reynolds operators $\mathcal{R}_G^{(k_i)}$ and satisfying this property; some motivation for the choice of the example is given in Remark 6.1. We will verify that $J(t_0, v_0) \neq 0$ at a specific point (t_0, v_0) by computing the differentials $d\tilde{\varphi}_i(t_0, v_0)$ approximately as partial sums of their Fourier series, which converge very rapidly. We recall that the period matrix of \mathcal{J} is $Z = \tau B$, where B is given by (4) and $\tau = \frac{3+i\sqrt{7}}{2}$, and we denote $q = e^{2\pi i\tau} = -e^{-\pi\sqrt{7}}$; we also adopt the convention that $q^r = e^{2\pi i r\tau}$ for any $r \in \mathbb{Q}$.

Lemma 4.2. *Let the vectors ξ_i ($i = 0, \dots, 3$) of \mathbb{Z}^3 be defined by*

$$\xi_0 = 0, \quad \xi_1 = \frac{1}{2}(0, 0, 1), \quad \xi_2 = \frac{1}{4}(1, 1, 0), \quad \xi_3 = \frac{1}{8}(2, 1, 1),$$

the first three being equal to the vectors μ_i (10) from Section 6, with the basis (b_i) of M used to identify M with \mathbb{Z}^3 , and $\xi_3 = \frac{7}{8}\mu_3$. Define four G -invariant theta functions on \mathcal{J} by

$$\varphi_0 = \mathcal{R}_G^{(2)}(\theta_{\xi_0,2}), \quad \varphi_1 = \mathcal{R}_G^{(2)}(\theta_{\xi_1,2}), \quad \varphi_2 = \mathcal{R}_G^{(4)}(\theta_{\xi_2,4}), \quad \varphi_3 = \mathcal{R}_G^{(8)}(\theta_{\xi_3,8}),$$

where the operators $\mathcal{R}_G^{(k)}$ are defined in (6). Associate to φ_i the Λ -invariant functions $\tilde{\varphi}_i$ on $\mathbb{C} \times V$, as in (7), and denote by J the Jacobian $J(\tilde{\varphi}_i)$. Set $(t_0, v_0) = (1, (\frac{1}{8}, \frac{1}{16}, \frac{1}{4}))$. Then $J(t_0, v_0) \neq 0$.

Proof. Let us choose $P_k = \{\frac{\nu}{k} \mid \nu \in \{0, 1, \dots, k-1\}\}^3$ for a set of representatives of $\frac{1}{k}\mathbb{Z}^3/\mathbb{Z}^3$. We have

$$\theta_{\xi_i, k_i} = \sum_{u \in \mathbb{Z}^3 + \xi_i} q^{\frac{k_i}{2}B[u]} e^{2k_i\pi i v^\top u} \quad (i = 0, 1, 2, 3, (k_i) = (2, 2, 4, 8)).$$

We compute the G -averages of these four theta-functions, using formula (6). By Proposition 3.1 and Theorem 2.4, the elements $u_{m', m}^{(g, k_i)}$ ($m', m \in P_{k_i}$) of the matrices $U_g^{(k_i)}$ belong to the cyclotomic field $\mathbb{Q}\left(e^{\frac{\pi i}{k_i}}\right)$. We introduce the matrices of the Reynolds operators on the theta functions of degree k_i :

$$\mathcal{R}_G^{(k)} = \left(r_{m', m}^{(k)} \right)_{m', m \in P_k}, \quad r_{m', m}^{(k)} = \frac{1}{336} \sum_{g \in G} u_{m', m}^{(g, k)}, \quad k = 2, 4, 8.$$

The exact values of the elements of the matrices $U_g^{(k_i)}$, belonging to $\mathbb{Q}\left(e^{\frac{\pi i}{8}}\right)$, and the resulting Reynolds matrices were computed with Macaulay2 [M2]. We have:

$$\varphi_i = \sum_{m \in P_{k_i}} r_{\xi_i, m}^{(k_i)} \theta_{m, k_i} = \sum_{u \in \frac{1}{k_i}\mathbb{Z}^3} r_{\xi_i, u}^{(k_i)} q^{\frac{k_i}{2}B[u]} e^{2k_i\pi i v^\top u},$$

where $r_{\xi_i, u}^{(k_i)}$ is defined to be $r_{\xi_i, m}^{(k_i)}$ for the unique $m \in P_{k_i}$ such that $u \equiv m \pmod{\mathbb{Z}^3}$. Replacing further u by $\frac{1}{k_i}u$ with u running over \mathbb{Z}^3 and differentiating, we obtain:

$$\begin{aligned} \partial \tilde{\varphi}_i / \partial t &= 2k_i t^{k_i-1} \sum_{u \in \mathbb{Z}^3} q^{\frac{1}{2k_i}B[u]} r_{\xi_i, \frac{1}{k_i}u}^{(k_i)} e^{2\pi i v^\top u}, \\ \partial \tilde{\varphi}_i / \partial v_j &= 2\pi i t^{k_i} \sum_{u \in \mathbb{Z}^3} q^{\frac{1}{2k_i}B[u]} r_{\xi_i, \frac{1}{k_i}u}^{(k_i)} u_j e^{2\pi i v^\top u}, \quad j = 1, 2, 3. \end{aligned}$$

As B is positive definite, these infinite sums of powers of q contain only a finite number of summands $N_i(c)$ with exponent $\leq c$ for any given constant c , $N_i(c) \sim \frac{4\pi}{3\sqrt{\det B}} (2k_i c)^{\frac{3}{2}}$, and, since $|q| < 1$, converge uniformly on compact subsets of $\mathbb{C} \times V$. The convergence is in fact very rapid. For example, in the case of the slowest convergence, when $i = 3$, $k_i = 8$, if we stop summation at the terms of order $q^{7/2} \approx -2.3211 i \cdot 10^{-13}$, then $N_3(3.5) = 3527$, and all the vectors $u \in \mathbb{Z}^3$ occurring in the truncated sum are in the block $|u_1| \leq 10$, $|u_2| \leq 10$, $|u_3| \leq 14$. Computing the determinant of the thus obtained approximate Jacobian matrix at the point (t_0, v_0) , we obtain: $J(t_0, v_0) \approx 0.000064967853 + 0.000075028580 i$, all the shown decimal digits being exact. \square

5. DEGREE-8 HYPERSURFACES IN $\mathbb{P}(1, 1, 2, 4, 7)$ WITH CORRECT SINGULARITIES

The goal of this section is to show that a hypersurface in $\mathbb{P}(1, 1, 2, 4, 7)$ defined by a degree-8 homogeneous polynomial whose singularities are those of $\mathbb{P}(1, 2, 4, 7)$ is actually isomorphic to $\mathbb{P}(1, 2, 4, 7)$. To achieve this, we shall classify the degree-8 hypersurfaces in $\mathbb{P}(1, 1, 2, 4, 7)$ whose singularities are those of $\mathbb{P}(1, 2, 4, 7)$.

Recall that $\mathbb{P}(1, 2, 4, 7)$ embeds in $\mathbb{P}(1, 1, 2, 4, 7)$ as the degree-8 hypersurface given by the equation

$$F_8^0 = y_3^2 - y_0 y_4.$$

Its singular locus is the union of two irreducible components, $\mathbb{P}^1 = \ell$ and an isolated point p . The singularity at p is of analytic type $\frac{1}{7}(1, 2, 4)$. Here, for a cyclic group μ_d of order d we denote by $\frac{1}{d}(\nu_1, \nu_2, \nu_3)$ the (analytic equivalence class of the) cyclic quotient singularity \mathbb{C}^3 / μ_d , where the generator c_d of μ_d acts by $c_d: (z_1, z_2, z_3) \mapsto (\epsilon^{\nu_1} z_1, \epsilon^{\nu_2} z_2, \epsilon^{\nu_3} z_3)$, $\epsilon = \exp\left(\frac{2\pi i}{d}\right)$. At all but one points of ℓ , the singularity of X is of type $\frac{1}{2}(1, 0, 1)$, that is $\mathbb{C} \times A_1$, the Cartesian product of \mathbb{C} with a surface singularity $\mathbb{C}^2 / \langle -1 \rangle$ of type A_1 . The type of singularity at q , the unique point of ℓ where the type of singularity changes, is $\frac{1}{4}(1, 2, 3)$.

Let us describe in affine charts the hypersurface $\{F_8^0 = 0\}$ in $\mathbb{P}(1, 1, 2, 4, 7)$. Write y_0, \dots, y_4 the coordinates in $\mathbb{P}(1, 1, 2, 4, 7)$ with weights

$$n_0 = 1, \quad n_1 = 1, \quad n_2 = 2, \quad n_3 = 4, \quad n_4 = 7.$$

Set $y_i = 1$, and quotient \mathbb{C}^4 with coordinates $(y_0, \dots, \hat{y}_i, \dots, y_4)$ by the action of the cyclic group μ_{n_i} with weights $(n_0, \dots, \hat{n}_i, \dots, n_4)$ to obtain the affine chart U_i of $\mathbb{P}(1, 1, 2, 4, 7)$. Note that U_0 and U_1 are smooth charts isomorphic to \mathbb{C}^4 since $n_0 = n_1 = 1$.

The restriction of F_8^0 to U_0 is $\{y_3^2 - y_4 = 0\}$. The hypersurface $F_8^0|_{U_0} = 0$ is a smooth hypersurface parameterised by (y_1, y_2, y_3) , isomorphic to \mathbb{C}^3 .

The restriction of F_8^0 to U_1 is $y_3^2 - y_0 y_4$. The singular locus of $F_8^0|_{U_1} = 0$ is the line $y_0 = y_3 = y_4$ with type $\mathbb{C} \times A_1$.

The restriction of F_8^0 to U_2 is $\{y_3^2 - y_0y_4 = 0\}/\frac{1}{2}(1, 1, 0, 1)$. Let us consider the quotient $\mathbb{C}^3/\frac{1}{2}(1, 0, 1)$, where the coordinates of \mathbb{C}^3 are denoted by (u, v, w) . One can identify this quotient with the quadric $y_0y_4 - y_3^2 = 0$ in \mathbb{C}^4 through the mapping $(u, v, w) \mapsto (y_0 = u^2, y_1 = v, y_3 = uw, y_4 = w^2)$. From this we observe that $\{y_0y_4 - y_3^2 = 0\}/\frac{1}{2}(1, 1, 0, 1)$, is the same as $\mathbb{C}^3/\frac{1}{4}(1, 2, 3)$. As a result, the restriction of F_8^0 to U_2 is isomorphic to $\mathbb{C}^3/\frac{1}{4}(1, 2, 3)$. Note that this is the affine chart $x_2 = 1$ in $\mathbb{P}(1, 2, 4, 7)$ with coordinates (x_0, x_1, x_2, x_3) .

The restriction of F_8^0 to U_3 is $\{y_0y_4 = 1\}/\frac{1}{4}(1, 1, 2, 3)$. This chart is contained in $U_0 \cup U_4$.

The restriction of F_8^0 to U_4 is $\{y_3^2 - y_0 = 0\}/\frac{1}{7}(1, 1, 2, 4)$, which is isomorphic to $\mathbb{C}^3/\frac{1}{7}(1, 2, 4)$, that is the affine chart $x_3 = 1$ in $\mathbb{P}(1, 2, 4, 7)$.

Let now consider a hypersurface X in $\mathbb{P}(1, 1, 2, 4, 7)$ defined by any degree-8 homogeneous equation $F_8 = 0$ such that the singularities of X are those of $\mathbb{P}(1, 2, 4, 7)$.

Lemma 5.1. *Write*

$$F_8 = y_4\varphi_1(y_0, y_1) + c_0y_3^2 + c_1y_2^2y_3 + c_2y_2^4 + y_2y_3\varphi_2(y_0, y_1) + y_2^3\psi_2(y_0, y_1) \\ + y_3\varphi_4(y_0, y_1) + y_2^2\psi_4(y_0, y_1) + y_2\varphi_6(y_0, y_1) + \varphi_8(y_0, y_1),$$

with $c_j \in \mathbb{C}$ and φ_i, ψ_i homogeneous polynomials of degree i . Then both c_0 and φ_1 are nonzero.

Proof. Assume first that $c_0 = 0$. Then $Q_3 = (0, 0, 0, 1, 0)$ is in X . It is the origin of the chart $U_3 = \mathbb{C}^4/\frac{1}{4}(1, 1, 2, 3)$ of $\mathbb{P}(1, 1, 2, 4, 7)$. Denote by $\tilde{X}_3 \subseteq \mathbb{C}^4$ the hypersurface defined by $\{F_8|_{y_3=1} = 0\}$ and let $\tilde{Q}_3 \in \tilde{X}_3$ be such that the image of \tilde{Q}_3 in the quotient \mathbb{C}^4/μ_4 is Q_3 . The germ of the canonical sheaf $\omega_{\tilde{X}_3, \tilde{Q}_3}$ is generated by

$$\sigma = \text{res}_{\tilde{X}_3} \left(\frac{dy_0 \wedge dy_1 \wedge dy_2 \wedge dy_4}{F_8|_{y_3=1}} \right),$$

of weight -1 for the μ_4 -action. So the Gorenstein index of $\tilde{X}_3/\frac{1}{4}(1, 1, 2, 3) = X|_{U_3}$ in Q_3 is 4. But X has no singularity of index 4. We conclude that c_0 is nonzero.

Assume now that $\varphi_1 = 0$. Denote $X_4 = X|_{U_4}$. We have $X_4 = \tilde{X}_4/\frac{1}{7}(1, 1, 2, 4)$, where $\tilde{X}_4 \subset \mathbb{C}^4$ is the hypersurface defined by the equation $\{F_8|_{y_4=1} = 0\}$. Under our assumption, F_8 does not contain y_4 , so $F_8|_{y_4=1} = F_8$ defines the affine cone of an octic surface in $\mathbb{P}(1, 1, 2, 4)$ with coordinates y_0, y_1, y_2, y_3 . The cone has to be smooth outside of its vertex O , the origin of the affine space \mathbb{C}^4 with the same coordinates. Indeed, assume the contrary. Then O is a non-isolated singularity, so the image Q_4 of O in X_4 also is a non-isolated singularity. Hence the unique isolated cyclic quotient singularity $p \in X$ of type $\frac{1}{7}(1, 2, 4)$ is different from Q_4 . Moreover, $p \in X$ is not a hypersurface singularity, but all the singularities of $X \setminus Q_4$ are hypersurface, so p cannot be located in the chart U_4 . Then, looking at the singularities of X in the other charts, we see that all of them are hypersurface ones, except possibly points on the axis (y_2, y_3) . By proving that $c_0 \neq 0$, we have excluded the possibility that Q_3 belongs to X , so the only non-hypersurface singularity that may occur in a chart U_i with $i \neq 4$ is of the type (hypersurface in \mathbb{C}^4)/ $\frac{1}{2}(1, 1, 0, 1)$. The embedding dimension of such singularity is at most that of $\mathbb{C}^4/\frac{1}{2}(1, 1, 0, 1)$, equal to 7, but the embedding dimension of p is 12. So no point of a chart U_i for $i \neq 4$ can fit to the role of p , hence this case is impossible, and the singularity of \tilde{X}_4 at O is an isolated quasi-homogeneous singularity.

It remains to see that under this assumption the quotients $\{F_8 = 0\}/\frac{1}{7}(1, 1, 2, 4)$ and $\mathbb{C}^3/\frac{1}{7}(1, 2, 4)$ cannot be isomorphic. This follows from the fact that the first one is non-canonical, whilst the second one is canonical. Indeed, $\tilde{X}_4 = \{F_8 = 0\}$ is non-canonical by Reid's criterion of canonicity (Theorem 4.1 of [Re]) with monomial valuation α defined by $(\alpha(y_i)) = (\frac{1}{8}, \frac{1}{8}, \frac{1}{4}, \frac{1}{2})$; this implies that $X_4 = \tilde{X}_4/\mu_7$ is non-canonical (ibid., Proposition 1.7). The canonicity of the second singularity follows from ibid., Theorem 3.1 and Remark 3.2. We conclude that φ_1 is nonzero. \square

By Lemma 5.1, one can assume that $c_0 \neq 1$ and use the change of coordinates

$$y_3 \mapsto d_0 y_3 + d_1 y_2^2 + y_2 f_2(y_0, y_1) + f_4(y_0, y_1),$$

with $d_j \in \mathbb{C}$, $d_0 \in \mathbb{C}^*$, and f_i homogeneous polynomials of degree i , to normalize the value of c_0 and to kill the terms containing y_3 to the power 1. We obtain for F_8 an expression as in the lemma, but with the constraints

$$c_0 = -1, \quad c_1 = 0, \quad \varphi_2 = 0, \quad \varphi_4 = 0.$$

Next, again by the lemma, $\varphi_1 \neq 0$, and we can change variables y_0, y_1 in such a way that in new variables we have $\varphi_1 = y_0$, so that there is no monomial $y_1 y_4$ in the expression for F_8 . Then, we use the change of coordinates

$$y_4 \mapsto y_4 + y_2^3 g_1(y_0, y_1) + y_2^2 g_3(y_0, y_1) + y_2 g_5(y_0, y_1) + g_7(y_0, y_1),$$

with g_i homogeneous of degree i , to kill all the monomials divisible by y_0 . As a result, we obtain an expression of the form

$$(8) \quad F_8 = y_0 y_4 - y_3^2 + a_1 y_2^4 + a_2 y_2^3 y_1^2 + a_3 y_2^2 y_1^4 + a_4 y_2 y_1^6 + a_5 y_1^8, \quad a_i \in \mathbb{C}.$$

Set $\nu = \min\{i : a_i \neq 0\}$. Finally, using changes of coordinates of the form $y_1 \mapsto \kappa y_1$, $y_2 \mapsto \lambda y_2 + \mu y_1^2$, $\kappa, \lambda \in \mathbb{C}^*$, $\mu \in \mathbb{C}$, we can reduce F_8 to one of the following normal forms:

$$(9) \quad \begin{aligned} \nu = \infty: & \quad F_8 = y_0 y_4 - y_3^2, \\ \nu = 5: & \quad F_8 = y_0 y_4 - y_3^2 + y_1^8, \\ \nu = 4: & \quad F_8 = y_0 y_4 - y_3^2 + y_2 y_1^6, \\ \nu = 3: & \quad F_8 = y_0 y_4 - y_3^2 + (y_2^2 + a_5 y_1^4) y_1^4, \\ \nu = 2: & \quad F_8 = y_0 y_4 - y_3^2 + (y_2^3 + a_4 y_2 y_1^4 + a_5 y_1^6) y_1^2, \\ \nu = 1: & \quad F_8 = y_0 y_4 - y_3^2 + y_2^4 + (a_3 y_2^2 + a_4 y_2 y_1^2 + a_6 y_1^4) y_1^4. \end{aligned}$$

Proposition 5.2. *Except for the case $\nu = \infty$, the degree-8 hypersurface X defined by $F_8 = 0$ in $\mathbb{P}(1, 1, 2, 4, 7)$, with F_8 one of the above normal forms, has a singularity of a type different from the types of singularities of $\mathbb{P}(1, 2, 4, 7)$.*

Proof. Let $\nu = 5$. Then X passes through the origin of the chart U_2 . We have $U_2 = \mathbb{C}^4/\frac{1}{2}(1, 1, 0, 1)$, and $X_2 := X \cap U_2$ is the quotient $\tilde{X}_2/\frac{1}{2}(1, 1, 0, 1)$, where $\tilde{X}_2 = \{y_0 y_4 - y_3^2 + y_1^8 = 0\} \subset \mathbb{C}^4$. Since $y_0 y_4 - y_3^2 + y_1^8$ is μ_2 -invariant and the differential $dy_0 \wedge dy_1 \wedge dy_3 \wedge dy_4$ is not, the generator

$$\text{res}_{\tilde{X}_2} \left(\frac{dy_0 \wedge dy_1 \wedge dy_3 \wedge dy_4}{y_0 y_4 - y_3^2 + y_1^8} \right)$$

of $\omega_{\tilde{X}_2, O}$, where O is the origin of \mathbb{C}^4 , is anti-invariant under μ_2 . So Q_2 , the image of O in U_2 , is a singular point of Gorenstein index 2. But in $\mathbb{P}(1, 2, 4, 7)$, the only point of Gorenstein index 2 is the origin of the chart $x_2 = 1$, with singularity $\frac{1}{4}(1, 2, 3)$. The latter singularity is

non-isolated: the whole image of the coordinate axis with weight 2 consists of singular points of the quotient. Hence it cannot be equivalent to the *isolated* singularity $\{y_0y_4 - y_3^2 + y_1^8 = 0\}/\frac{1}{2}(1, 1, 0, 1)$. Thus the case $\nu = 5$ is impossible.

The cases $2 \leq \nu \leq 5$ are all treated in the same way: in all of them we find an isolated singularity of Gorenstein index 2 at the origin of the chart U_2 , which is impossible.

Let us look now at the case $\nu = 1$. In the chart U_2 , we have

$$X \cap U_2 = \tilde{X}_2/\frac{1}{2}(1, 1, 0, 1),$$

where $\tilde{X}_2 \subset \mathbb{C}^4$ is defined by the equation $\tilde{F} = 0$ with $\tilde{F} = y_0y_4 - y_3^2 + 1 + a_3y_1^4 + a_4y_1^6 + a_5y_1^8$. The singular locus of \tilde{X}_2 is given by $y_0 = y_3 = y_4 = 0$, $y_1^2 = \gamma$, where γ is a double root of $a_5t^4 + a_4t^3 + a_3t^2 + 1 = 0$. But this polynomial cannot have double roots. Indeed, otherwise X would have singular points in the chart U_1 , which are of the type of isolated hypersurface singularities:

$$X \cap U_1 = \{y_0y_4 - y_3^2 + y_2^4 + a_3y_2^2 + a_4y_2 + a_5 = 0\} \subset \mathbb{C}^4.$$

this contradicts the fact that the only isolated Gorenstein singularity of $\mathbb{P}(1, 2, 4, 7)$ is of type $\frac{1}{7}(1, 2, 4)$, and this is not a hypersurface singularity.

Therefore \tilde{X}_2 is smooth and the singularities of $\tilde{X}_2/\mu_2 = X \cap U_2$ can only occur in a subset of the fixed locus of μ_2 . Thus $\text{Sing}(X) \cap U_2$ is the image of the set

$$\{y_0 = y_1 = y_4 = 0\} \cap \tilde{X}_2 = \{y_0 = y_1 = y_4 = -y_3^2 + 1 = 0\}.$$

Hence X has two isolated singular points of type $\frac{1}{2}(1, 1, 1)$ which are $(0, 0, \pm 1, 0)$. But this is impossible. \square

This brings us to the main result of the paper:

Theorem 5.3. *The quotient $X = \mathcal{J}/G$ of the Jacobian \mathcal{J} of the Klein quartic C by its full automorphism group G is isomorphic to the weighted projective space $\mathbb{P}(1, 2, 4, 7)$.*

Proof. This is an obvious consequence of Theorem 4.1, Lemma 5.1, Proposition 5.2 and [MM, Theorem 4.3]. \square

Remark 5.4. By arguments, similar to those used in the reduction of F_8 to normal forms (9), one can easily verify that in the 45-dimensional group of coordinate changes in $\mathbb{P}(1, 1, 2, 4, 7)$, the stabilizer of the octic form $F_8^0 = y_0y_4 - y_3^2$ is 13-dimensional and the connected component of the identity in it is generated by the changes of the form

$$y_0 \mapsto \lambda_0 y_0, \quad y_4 \mapsto \lambda_0^{-1} y_4, \quad y_1 \mapsto \mu y_0 + \lambda_1 y_1, \quad y_2 \mapsto \lambda_2 y_2 + h_2(y_0, y_1),$$

$$y_3 \mapsto y_3 + y_0 y_2 h_1(y_0, y_1) + y_0 h_3(y_0, y_1),$$

$$y_4 \mapsto y_4 + 2(h_1(y_0, y_1)y_2 + h_3(y_0, y_1))y_3 + y_0(h_1(y_0, y_1)y_2 + h_3(y_0, y_1))^2,$$

where $\lambda_i \in \mathbb{C}^*$, $\mu \in \mathbb{C}$ and the h_i are homogeneous of degree i in y_0, y_1 . Thus the dimension of the orbit of F_8^0 is $45 - 13 = 32$. As the vector space of octic forms is of dimension 37, the orbit of F_8^0 is of codimension 5. A transversal slice to the orbit can be given by (8). This family of hypersurfaces in $\mathbb{P}(1, 1, 2, 4, 7)$ represents a versal deformation of the weighted projective space $\mathbb{P}(1, 2, 4, 7)$ inside the space of octics in $\mathbb{P}(1, 1, 2, 4, 7)$. The computation of the infinitesimal deformation space $\text{Ext}^1(\Omega_X^1, \mathcal{O}_X)$ shows that $\dim T_X^1 = 5$, and this implies that (8) is a complete versal deformation of X . It is interesting to note that this family contains

partial smoothings of X with only 3 singular points, of which two are of type $\frac{1}{2}(1, 1, 1)$ (the singular points found in the proof of the case $\nu = 1$ of Lemma 5.2) and the third one is $\frac{1}{7}(1, 2, 4)$. These three singular points are non-smoothable, and even infinitesimally rigid by the result of Schlessinger [Schl].

6. RELATION TO QUOTIENTS BY THE WEYL GROUP FOR THE ROOT SYSTEMS B_3, C_3

As we already noticed, G contains the Weyl group $W = W(B_3) = W(C_3)$ of order 48. We are going to look at the invariants under the action of W . For any $g \in W$, the symplectic matrix $\gamma_g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is bloc-diagonal, that is $b = c = 0$, and $a = (d^\top)^{-1}$ is nothing but the matrix of g^{-1} in the basis given by the columns of $\omega_2 = \alpha C$ (we keep the notation from previous sections). It easily follows from the definitions that $g \cdot \theta_{m,k}$ is just $\theta_{dm,k}$. This implies that the space of W -invariant theta-functions of degree k consists of the functions $\mathcal{R}_W \theta_{m,k}$, m running over $(C^\top)^{-1}(F) \cap P_k$, where F is a fundamental domain for the action of the (real) affine crystallographic group $M \rtimes W$ on the space $M_{\mathbb{R}} = M \otimes \mathbb{R}$ containing the weight lattice M , and \mathcal{R}_W is the Reynolds operator of taking the average over the action of W . Saying that F is a fundamental domain we mean that F is the disjoint union of an open convex polyhedron in $M_{\mathbb{R}}$ with finitely many polyhedra of smaller dimensions contained in its boundary such that every W -orbit in $M_{\mathbb{R}}$ has exactly one representative in F . We are now going to fix the choice of a particular fundamental domain F .

There is a tower of degree two extensions


$$Q = Q(C_3) \subset Q^\vee = Q(B_3) \subset M = Q^*,$$

the three lattices being invariant under W , so taking the semi-direct product with W , we obtain a tower of degree two extensions of the corresponding real affine crystallographic groups:

$$\widetilde{W} = Q \rtimes W \subset \widetilde{W}^\vee = Q^\vee \rtimes W \subset \widetilde{W}^* = M \rtimes W.$$

Both \widetilde{W} and \widetilde{W}^\vee are affine Weyl groups and are generated by affine reflections. By [Bo, VI.2.2], they have fundamental domains which are closed tetrahedra, called alcoves. For \widetilde{W} , the vertices of the standard alcove $\widetilde{C} = C_{\widetilde{W}}$ are $f_0 = 0$, $f_1 = (1, 0, 0)$, $f_2 = \frac{1}{2}(1, 1, 0)$, $f_3 = \frac{1}{2}(1, 1, 1)$. Denoting by x_1, x_2, x_3 the coordinates in the euclidean space \mathbb{R}^3 , in which we place our lattices Q and M , we obtain an alcove $\widetilde{C}^\vee = C_{\widetilde{W}^\vee}$ for \widetilde{W}^\vee as a half of \widetilde{C} cut out by the mirror $x_1 = \frac{1}{2}$ of a reflection contained in \widetilde{W}^\vee but not in \widetilde{W} . Thus \widetilde{C}^\vee is the tetrahedron with vertices $O = f_0$, $A = \frac{1}{2}f_1$, $E = f_2$, $N = f_3$. The subgroup of \widetilde{W}^* leaving invariant the tetrahedron $OAEN$ is of order 2; besides the identity, it contains an element R of order 2, the axial symmetry, or rotation by the angle π with axis KL , where K, L are the middles of AE, ON respectively (see the figure below).

Now we can get a fundamental domain F of \widetilde{W}^* as follows: first choose a plane P containing KL and pick up one of the two halves in which P dissects \widetilde{C}^\vee , say, C_1 . The intersection $D = P \cap \widetilde{C}^\vee$ is a face of C_1 , and it is dissected by KL in two halves D_1, D_2 symmetric to each other by the action of R . Then we obtain the following set of representatives of the orbits of R acting on \widetilde{C}^\vee : the polyhedron C_1 , in which we include all of its faces, except for the face D , from which only the closed part D_1 is included. The representatives of the orbits of R in \widetilde{C}^\vee are at the same time the representatives of the orbits of \widetilde{W}^* in $V_{\mathbb{R}}^*$, so the constructible set we have described is a fundamental domain for \widetilde{W}^* .



To obtain a particular fundamental domain F , we choose in the above construction the plane OKN for P , the tetrahedron $OAKN$ for C_1 , then the triangle KON for the face D , and we keep in F only a half of it, the triangle $D_1 = OKL$. On the picture, F is hatched; the triangle OKL is hatched in a darker gray color as it is included in F and the remaining part of the face OKN is not included. The sides of the three closed triangles of the boundary of F contained in F , as well as the edge AN contained in F , are drawn by thick lines. Only the part OL of the edge ON drawn by a thick line is contained in F , and the edge KN , not contained in F , is drawn by a thin line. The black points O, A, K, L are contained in F , while the vertex N is not in F and is represented by a white point.

Looijenga in [Lo] provides a recipe for the choice of generators of the algebra of W -invariant theta-functions for the lattice of the form $\tau Q + Q$; his generators are the averages of the theta-functions θ_{f_i, k_i} with characteristics f_i , the vertices of the alcove, the degrees k_i being the smallest positive integers such that $k_i f_i \in M$.

Remark 6.1. In the previous sections, we worked with theta-functions $\theta_{m, k}$ defined in Section 1 for a different lattice, $\tau M + Q$, so Looijenga's recipe does not apply $\theta_{m, k}$, but we can use it as a heuristic principle for the choice of theta-functions that would be candidates for generators of the G -invariant subalgebra. We can try the G -averages of the theta-functions whose characteristics are the vertices μ_i of a "pseudo-fundamental domain" $F_G \subset F$ for G , or a "pseudo-alcove" which is just a smaller tetrahedron with vertex O contained in F , having the correct lattice volume with respect to M , equal to $\frac{1}{336}$, and the expected weights k_i . Recall that $b_1 = f_1, b_2 = 2f_2, b_3 = f_3$ are the columns of the matrix $(C^T)^{-1}$, that is they form a basis of M dual to the standard basis of the root system C_3 ; this is the basis in which we write the characteristics of our theta-functions. From this we see, in particular, that the lattice volume of F with respect to M is $\frac{1}{48}$. We can pick up the following ad hoc candidate for the "pseudo-alcove": take the tetrahedron F_G with vertices

$$(10) \quad \mu_0 = 0, \quad \mu_1 = \frac{1}{2}b_1, \quad \mu_2 = \frac{1}{4}(b_1 + b_2), \quad \mu_3 = \frac{1}{7}(2b_1 + b_2 + b_3).$$

F_G obviously has the M -volume $\frac{1}{336}$, and the smallest k_i such that $k_i \mu_i \in M$ coincide with the weights of the expected weighted projective space: $(k_0, k_1, k_2, k_3) = (1, 2, 4, 7)$. We thus may expect that the algebra $S(\mathcal{L}^2)^G$ is generated by the G -averages of the even-degree theta-functions that are obtained as products of theta functions whose characteristics are the

vertices of F_G : $\theta_{\mu_0,1}^2, \theta_{\mu_1,2}, \theta_{\mu_2,4}, \theta_{\mu_0,1}\theta_{\mu_3,7}, \theta_{\mu_3,7}^2$. To present four algebraically independent G -invariant theta functions of degrees 2, 2, 4, 8 in Lemma 4.2, we chose another set, for which the computations are slightly easier: $\theta_{\mu_0,2}, \theta_{\mu_1,2}, \theta_{\mu_2,4}, \theta_{\frac{7}{8}\mu_3,8}$.

We are now returning to the quotient \mathcal{J}/W . As we saw above, the theta-functions $\{\mathcal{R}_W\theta_{m,k}\}_{m \in (C^\top)^{-1}(F) \cap P_k}$ form a basis of the W -invariant theta-functions of degree k on \mathcal{J} . The fact that \mathcal{J}/W is not a complex crystallographic reflection quotient manifests itself in that F is not a closed simplex, but a constructible set, which we can describe as follows:

$$(11) \quad F = OAKN \cup \overline{OAK} \cup \overline{OKL} \cup \overline{OAL} \cup ANK \cup ANL \cup AN.$$

Here we denote by $\overline{A_1 \dots A_n}$ the convex hull of points A_1, \dots, A_n in a real affine space, and by $A_1 \dots A_n$ its relative interior. From this we can deduce the Hilbert function of the invariant algebra $S(\mathcal{L})^W$:

Proposition 6.2. *The Hilbert function $h_{\overline{S}} : k \mapsto \dim H^0(\mathcal{J}, \mathcal{L}^k)^W$ of the algebra $\overline{S} = S(\mathcal{L})^W$ is the sum of a polynomial E and a function $\ell : k \mapsto d_0k + d_1$, where d_0, d_1 are 4-periodic functions of the integer variable k ,*

$$E(k) = \frac{1}{48}k^3 + \frac{3}{16}k^2 + \frac{2}{3}k + 1, \quad d_0 = \begin{cases} 0 & \text{if } k \equiv 0 \text{ or } 2 \\ -\frac{3}{16} & \text{if } k \equiv \pm 1 \end{cases}, \quad d_1 = \begin{cases} 0 & \text{if } k \equiv 0 \\ -\frac{11}{16} & \text{if } k \equiv \pm 1 \pmod{4} \\ -\frac{1}{4} & \text{if } k \equiv 2 \end{cases}.$$

The Hilbert series $\mathcal{H}_{\overline{S}}(t) = \sum_{k=0}^{\infty} h_{\overline{S}}(k)t^k$ of \overline{S} is given by:

$$(12) \quad \mathcal{H}_{\overline{S}}(t) = \frac{1 - t + t^2}{(1 - t)^2(1 - t^2)(1 - t^4)}.$$

Proof. By Ehrhart's Theorem, the number of integer points $h_P(k)$ in the integer multiples kP of an open or closed polytope P in \mathbb{R}^n with rational vertices is a quasi-polynomial in the integer variable k of degree equal to the dimension of P . A quasi-polynomial is a polynomial function whose coefficients are periodic. The common period of the coefficients of the Ehrhart quasi-polynomial of P is the smallest positive integer d such that dP is a lattice polytope, that is, all the coordinates of its vertices are integers (see, for example, [St, Theorem 4.6.25]). By the above, $h_{\overline{S}}$ coincides with the function h_F counting the number of points of M in the multiples kF of F . By (11), $h_F(k)$ can be represented as a linear combination, with coefficients ± 1 , of the numbers $h_\sigma(k)$ for σ running through a finite set of open or closed simplices with rational vertices of dimensions between 0 and 3:

$$h_F = h_{OAKN} + h_{\overline{OAK}} + h_{\overline{OKL}} + h_{\overline{OAL}} + h_{ALN} + h_{ANK} - h_{\overline{OA}} - h_{\overline{OK}} - h_{\overline{OL}} + h_{AN} + h_O.$$

Each of the terms of this linear combination is a quasi-polynomial of degree at most 3 with period $d = 4$, the least common denominator of the coordinates of the vertices of the simplices σ_i , so h_F also is such a quasi-polynomial. Moreover, the dominant coefficient of h_F is constant and is equal to the inverse of the lattice volume of F , that is $\frac{1}{48}$, so it suffices to compute 12 consecutive values of h_F in order to determine the coefficients as solutions to a system of linear equations. Here is the result of computation of $h_{\overline{S}}(k) = h_F(k)$ for $0 \leq k \leq 12$ by Macaulay2 [M2]:

$$1, 1, 3, 4, 8, 10, 16, 20, 29, 35, 47, 56, 72 \dots$$

These values determine completely $h_{\overline{S}}$, and we thus obtain the formulas from the statement of the proposition. \square

Proposition 6.3. *Consider the quotient $X' = \mathbb{P}^3/\mu_2 \times \mu_4$ of the projective space \mathbb{P}^3 by a group of order 8, where μ_n denotes a cyclic group of order n and the generators of μ_2, μ_4 act by diagonal matrices $\text{diag}(1, 1, -1, 1), \text{diag}(i, 1, 1, -1)$ respectively. Then X' and \mathcal{J}/W have the same Hilbert functions.*

Proof. We represent X' as a toric variety, the equivariant compactification $\mathbf{X}_{\Sigma, \mathbf{N}}$ of the 3-dimensional algebraic torus $\mathbb{T} := (\mathbb{C}^*)^3$, defined by a fan Σ in the 3-dimensional \mathbb{R} -vector space $\mathbf{N} \otimes \mathbb{R}$, where $\mathbf{N} \simeq \mathbb{Z}^3$ is the lattice of 1-parametric subgroups in \mathbb{T} (see e.g. [Fu] for definitions and basic properties of toric varieties). Let y_0, \dots, y_3 be the homogeneous coordinates of \mathbb{P}^3 and $\mathbb{T}_0 = (\mathbb{C}^*)^3$ the standard torus in the affine chart \mathbb{C}^3 of \mathbb{P}^3 with affine coordinates $u_1 = y_1/y_0, u_2 = y_2/y_0, u_3 = y_3/y_0$. The simplicial fan Σ defining \mathbb{P}^3 is determined by its 1-dimensional cones, or rays, which are spanned by the four vectors $(1, 0, 0), (0, 1, 0), (0, 0, 1), (-1, -1, -1)$ of the lattice $\mathbf{N}_0 = \mathbb{Z}^3$. The dual to \mathbf{N}_0 is the lattice $\mathbf{M}_0 = \mathbb{Z}^3$ of exponents of monomials in the coordinate algebra $\mathbb{C}[\mathbf{M}_0] = \mathbb{C}[u_1, u_2, u_3]$ of the affine chart of \mathbb{P}^3 that we have chosen. To get the quotient by $\mu_2 \times \mu_4$, we replace \mathbf{M}_0 by the sublattice $\mathbf{M} = \mathbf{M}_0^{\mu_2 \times \mu_4}$ of exponents of $\mu_2 \times \mu_4$ -invariant monomials, which is given by

$$\mathbf{M} = \{(m_1, m_2, m_3) \in \mathbf{M}_0 : m_3 \equiv 0 \pmod{2}, -m_1 - m_2 + m_3 \equiv 0 \pmod{4}\};$$

this is the lattice of exponents of the monomials $\mathbf{u}^{\mathbf{m}} = u_1^{m_1} u_2^{m_2} u_3^{m_3}$ which form a basis of the regular functions on the quotient torus $\mathbb{T} := \mathbb{T}_0/\mu_2 \times \mu_4$. The dual $\mathbf{N} = \mathbf{M}^* = (\mathbf{M}_0^{\mu_2 \times \mu_4})^*$ is the overlattice of $\mathbf{N}_0 = \mathbb{Z}^3$ generated by the vectors $(1, 0, 0), \frac{1}{2}(0, 1, 0), \frac{1}{4}(-1, -1, 1)$, and $X' = \mathbf{X}_{\Sigma, \mathbf{N}}$ is the equivariant compactification of \mathbb{T} defined by the same fan Σ as the original \mathbb{P}^3 , but taken with respect to the lattice \mathbf{N} . The primitive vectors of \mathbf{N} generating the rays of the fan are $v_0 = (1, 0, 0), v_1 = \frac{1}{2}(0, 1, 0), v_2 = -\frac{1}{2}(1, 1, 1), v_3 = (0, 0, 1)$, and we get 4 \mathbb{T} -invariant divisors D_i in X' , defined by $D_i = D_{v_i}$, where D_{v_i} denotes the closure in X' of the kernel of v_i viewed as a 1-parametric subgroup of \mathbb{T} . The Cartier indices of these divisors for $i = 0, 1, 2, 3$ are respectively 4, 2, 2, 4. We omit the routine verification of the following assertion, which makes more precise the statement we are proving:

Proposition 6.3'. *In the above notation, the Hilbert functions $k \mapsto h^0(X', \mathcal{O}_{X'}(kD_i))$ coincide with $h_{\overline{\mathcal{J}}}$ for both divisors D_i of Cartier index 4, that is for $i = 0$ and 3.*

□

It is plausible that the two quotients are indeed isomorphic. We observe that X' belongs to the class of projective toric varieties of dimension n whose fan contains $n + 1$ rays. Such varieties are called weak weighted projective spaces; some authors call them fake weighted projective space (see [Kas]), but we prefer the adjective weak because the class of weak weighted projective spaces contains all the weighted projective spaces. A weak weighted projective space is a weighted projective space if and only if the primitive vectors of \mathbf{N} on the rays $\mathbb{R}_+^* v_i$ of its fan generate the whole of \mathbf{N} . One easily sees that this is not the case for X' . This brings us to the following generalization of the Bernstein–Schwarzman conjecture:

Conjecture 6.4. *If Γ, Γ_1 are commensurable complex crystallographic groups acting on \mathbb{C}^n such that $d\Gamma_1 = d\Gamma$, and if Γ is irreducible and generated by reflections, then the quotient \mathbb{C}^n/Γ_1 is a weak weighted projective space. It is a genuine weighted projective space if and only if Γ_1 is also generated by reflections.*

The cases when Γ is one of the complex crystallographic reflection groups $(Q + \tau Q) \rtimes W$ or $(Q + \tau Q^\vee) \rtimes W$ treated by Looijenga and Bernstein–Schwarzman and $\Gamma_1 = \Lambda \rtimes W$, where $\Lambda = Q + \tau M$, are particular cases of this conjecture: while \mathbb{C}^3/Γ is known to be a weighted projective space, we expect that \mathbb{C}^3/Γ_1 is the weak weighted projective space X' .

REFERENCES

- [BS] Joseph Bernstein and Ossip Schwarzman, Chevalley's theorem for the complex crystallographic groups, *J. Nonlinear Math. Phys.* **13** (2006), n°3, 323–351.
- [BEW] Bruce C. Berndt, Ronald J. Evans and Kenneth S. Williams, Gauss and Jacobi sums, *Canadian Mathematical Society Series of Monographs and Advanced Texts*, John Wiley & Sons, Inc., New York, 1998.
- [BL] Christina Birkenhake and Herbert Lange, Complex abelian varieties. Second edition. Grundlehren der mathematischen Wissenschaften, **302**, Springer-Verlag, Berlin, 2004.
- [Bo] Nicolas Bourbaki, *Éléments de mathématique*, Fasc. XXXIV, Groupes et algèbres de Lie, Chapitres IV–VI, Hermann, Paris, 1968.
- [Co] Arjeh M. Cohen, Finite complex reflection groups. *Ann. Sci. Ecole Norm. Sup.* (4) **9** (1976), n°3, 379–436.
- [DS] Thomas Dedieu and Edoardo Sernesi, Deformations and extensions of Gorenstein weighted projective spaces, to appear in *The Art of Doing Algebraic Geometry (dedicated to Ciro Ciliberto on the occasion of his 70th birthday)*, Springer, Cham, 2022, arXiv:2103.08210 [math.AG].
- [Fu] William Fulton, Introduction to toric varieties, *Annals of Mathematics Studies*, **131**, Princeton University Press, Princeton, NJ, 1993.
- [Gu] Robert Clifford Gunning, Lectures on modular forms, *Annals of Mathematics Studies*, n°48, Princeton University Press, Princeton, N.J. 1962.
- [Ig] Jun-ichi Igusa, Theta functions, *Die Grundlehren der mathematischen Wissenschaften*, Band **194**, Springer-Verlag, New York-Heidelberg, 1972.
- [Kas] Kasprzyk, A. M.: Bounds on fake weighted projective space, *Kodai Math. J.* **32**, 197–208 (2009).
- [KI] Felix Klein, Über die Transformationen siebenter Ordnung der elliptischen Funktionen, *Math. Ann.* **14**, 428–471 (1878). Œuvres, Tome III, 90–136.
- [Lo] Eduard Looijenga, Root systems and elliptic curves. *Invent. Math.* **38** (1976/77), n°1, 17–32.
- [M2] Daniel R. Grayson and Michael Stillman, Macaulay2, A software system for research in algebraic geometry, available at www.math.uiuc.edu/Macaulay2/.
- [MM] Dimitri Markushevich and Anne Moreau, Action of the automorphism group on the Jacobian of Klein's quartic curve, to appear in *Birational Geometry, Kähler–Einstein Metrics and Degenerations*, Springer Proceedings in Mathematics & Statistics, 2022, arXiv:2107.03745 [math.AG].
- [Ma] Barry Mazur, Arithmetic on curves. *Bull. Amer. Math. Soc.*, **14** (1986), n°2, 207–259.
- [Po] Vladimir L. Popov, Discrete complex reflection groups, *Communications of the Mathematical Institute*, Rijksuniversiteit Utrecht, 15. Rijksuniversiteit Utrecht, Mathematical Institute, Utrecht, 1982.
- [Rains] Eric M. Rains, Quotients of abelian varieties by reflection groups, arXiv:2303.04786 [math.AG].
- [Re] Miles Reid, Canonical 3-folds, In: *Journées de Géométrie Algébrique d'Angers, Juillet 1979*, pp. 273–310, Sijthoff & Noordhoff, Alphen aan den Rijn, 1980.
- [Schl] Michael Schlessinger, Rigidity of quotient singularities, *Invent. Math.*, **14** (1971), 17–26.
- [ST] Geoffrey C. Shephard and John Arthur Todd, Finite unitary reflection groups. *Canad. J. Math.*, **6** (1954), 274–304.
- [St] Richard P. Stanley, Enumerative combinatorics, Vol. I, *The Wadsworth & Brooks/Cole Mathematics Series*, Monterey, CA, 1986.

UNIV. LILLE, CNRS, UMR 8524 – LABORATOIRE PAUL PAINLEVÉ, F-59000 LILLE, FRANCE
Email address: dimitri.markouchevitch@univ-lille.fr

UNIVERSITÉ PARIS-SACLAY, CNRS, LABORATOIRE DE MATHÉMATIQUES D'ORSAY, RUE MICHEL MARGAT, BÂT. 307, 91405 ORSAY, FRANCE
Email address: anne.moreau@universite-paris-saclay.fr