



**HAL**  
open science

## **Non-lethal sampling method for the analysis of white muscle fatty acid profiles in European sea bass (*Dicentrarchus labrax*)**

Mickaël Péron, Victor Simon, Fabienne Le Grand, Philippe Soudant, David Mazurais, Marie Vagner

### ► **To cite this version:**

Mickaël Péron, Victor Simon, Fabienne Le Grand, Philippe Soudant, David Mazurais, et al.. Non-lethal sampling method for the analysis of white muscle fatty acid profiles in European sea bass (*Dicentrarchus labrax*). *Fish Physiology and Biochemistry*, 2023, 49 (6), pp.1381-1390. <10.1007/s10695-023-01262-w>. <hal-04394694>

**HAL Id: hal-04394694**

**<https://hal.science/hal-04394694v1>**

Submitted on 22 Aug 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 NON-LETHAL SAMPLING METHOD FOR THE ANALYSIS OF WHITE MUSCLE FATTY ACID  
2 PROFILES IN EUROPEAN SEA BASS (*Dicentrarchus labrax*)

3 Mickael Peron<sup>1</sup>, Victor Simon<sup>1</sup>, Fabienne Le Grand<sup>1</sup>, Philippe Soudant<sup>1</sup>, David Mazurais<sup>1</sup>, Marie Vagner<sup>1</sup>

4 <sup>1</sup> Univ Brest, CNRS, IRD, Ifremer, LEMAR, F-29280 Plouzane, France

5 Abstract :

6 This study presents a novel non-lethal sampling method for assessing fatty acid (FA) composition in juvenile  
7 European sea bass (*Dicentrarchus labrax*) using subcutaneous white muscle biopsies. This research aimed to  
8 evaluate the suitability of the biopsy for FA analysis using two lipid extraction protocols and comparing them to  
9 a lethal routine method. The results showed that a mass of fresh tissue as low as 1.4 mg provided good quality FA  
10 chromatograms for both reserve and membrane lipids. Although the biopsy method displayed high variability in  
11 terms of FA quantity among intra-individual replicates, it showed good FA profile repeatability in both reserve  
12 and membrane lipids. The study highlights the potential of this non-lethal approach for studying FA dynamics in  
13 fish, with its application particularly promising for ecological and experimental studies. However, careful biopsy  
14 implementation is recommended to account for potential lipid droplet and lipid distribution variability within the  
15 tissue.

16 Key words: Fatty acids, European sea bass, Muscle biopsy, Lipid composition, Method comparison

17 1. Introduction

18 Lipids are essential molecules in animals that serve both as energy sources and reserves (non-polar lipids; NL), as  
19 well as key structural components of cell membranes (polar lipids; PL) (Sargent et al. 2003). Due to their structural  
20 function, lipids modulate cell membrane functionality (Singer and Nicholson 1972; Hulbert and Else 1999; Tocher  
21 2003), which in turn influences various biological processes in cells, tissues, organs, and ultimately affects the  
22 overall performance of an individual (Tocher 2010). Fatty acids are also involved in the modulation of gene  
23 expression (réf) and influences the function of membrane proteins thus affecting cellular signaling and  
24 communication (Murphy 1990). Both NL and PL are composed of fatty acids (FA), among which polyunsaturated  
25 fatty acids (PUFA) are the main component of cell membranes and thus play a key role in maintaining vital  
26 functions of most animals (Sargent et al. 2003). Among PUFA, eicosapentaenoic acid (20:5n-3, EPA) and  
27 docosahexaenoic acid (22:6n-3, DHA) have especially been demonstrated to be essential for most marine fish to  
28 ensure growth, reproduction, development (Tocher 2010). However, as for most animals, these FA are weakly  
29 synthesized *de novo* by fish, and must be obtained through the diet. In natural environments, EPA and DHA are  
30 primarily synthesized at the base of the aquatic food web by primary producers like marine microalgae (Arts and  
31 Kohler 2009) and then accumulate through the food chain, ultimately reaching higher consumers (Sargent et al.  
32 1993; Tocher et al. 2006; Li et al. 2010).

33 Fish fatty acid composition analysis is mandatory to meet several objectives, for example, to understand the link  
34 between dietary FA and physiological performance of an individual. Experimental research has explored for  
35 instance the links between dietary FA, fish muscle membrane FA composition, and individual performances, such  
36 as growth, locomotion or metabolism (Tocher 2003; Calder 2012). The effects of dietary FA on fish physiology  
37 have also been studied in the context of global change, in combination with other environmental stressors like

38 hypoxia (McKenzie et al. 2000; McKenzie et al. 2008), warming (Gourtay et al. 2018; Vagner et al. 2019) or  
39 acidification (Hurst et al. 2019). Examining fish FA composition is also valuable for understanding the structure  
40 of trophic food webs in natural ecosystems, as some fatty acids serve as trophic biomarkers that can be used to  
41 identify consumer prey and elucidate trophic interactions within an ecosystem (Dalsgaard et al. 2003; El-Sabaawi  
42 et al. 2010; Mathieu-Resuge et al. 2019; Bertrand et al. 2022).

43 Typically in fish, FA analysis requires the sacrifice of the animal to obtain sufficient amount of tissue from specific  
44 organs (e.g. 50-200 mg, wet weight WW, Meyer et al. 2017), such as the liver, brain or muscle (Couturier et al.  
45 2020; Betancor et al. 2021), to meet the minimal biomass requirements according to the analytical protocol. Aside  
46 from being a lethal method, another drawback of this approach is that it only allows us for the determination of an  
47 individual's FA composition at a single point in time. As a result, monitoring changes in an individual's FA  
48 composition over time becomes challenging, which could otherwise offer valuable insights into the dynamics of  
49 FA incorporation. However, this is only the case for small animals like juvenile fish and larger animals can be  
50 sampled without risking killing the individual.

51 Non-lethal sampling methods have already been used in fish biology to address trophic ecology and ecotoxicology,  
52 using fins and scales (Cano-Rocabayera et al. 2015; Jeffries et al. 2021) or blood (Lawrence et al. 2020). Non-  
53 lethal sampling of muscle has also been realized to study mercury concentrations, with a tissue WW ranging from  
54 47 to 220 mg from fishes of a minimal size of 36 cm and 600 g (Baker et al. 2004). In 2013, Olsen et al. studied  
55 muscle FA composition using biopsy on adult salmon *Salmo salar* with a mean size and weight of 84 cm and 6.6  
56 kg, respectively, and with biopsy samples weighing a minimum of 40 mg WW. However, one of the main  
57 challenges of these non-lethal sampling methods is the necessity to take a small quantity of sampled tissue to  
58 guarantee minimal harm to fish, especially when working with small species or young individuals (Meyer et al.  
59 2017). Given that lipid distribution within a tissue can exhibit heterogeneity (Nanton et al. 2007; Barbosa et al.  
60 2018), the small amount of tissue collected through biopsies may lead to an inaccurate representation of the tissue's  
61 lipid content. This could result in a lack of repeatability, as the variability in FA composition between replicates  
62 obtained from different areas of the same tissue is likely to be high.

63 The objective of our research was to evaluate a novel nonlethal sampling technique for determining the FA  
64 composition of juvenile European sea bass (*Dicentrarchus labrax*; Linnaeus 1758) using subcutaneous white  
65 muscle biopsies. We anticipate that this approach will: i) yield adequate tissue sample quantities for generating  
66 interpretable FA profiles ii) would be repeatable within an individual, with a low intra-individual variability, also  
67 compared to the inter-individual variability in terms of FA quantity and proportions, and iii) produce results  
68 comparable to a traditional but lethal method of sampling using higher tissue biomass. As working with dry  
69 samples is of interest in some particular cases, such as field studies, thanks to its convenience for transport and  
70 storage (Sardenne et al. 2019), we wanted to test whether the conditioning (freeze-drying or fresh) of biopsied  
71 tissues had an effect on FA quantities and proportions and could be compared to our routine tissue analysis.

## 72 2. Material and methods

### 73 2.1. Ethics statement

74 All fish manipulations were performed according to the Animal Care Committee of France (ACCF) and performed  
75 at the Ifremer facilities in Plouzané (France, Agreement number: B29-212-05). All fish manipulations were

76 performed under anesthesia using tricaine (200 mg l<sup>-1</sup>, ethyl 3-aminobenzoate methane sulfonate), and all efforts  
77 were made to minimize animal suffering during manipulation.

## 78 2.2. Animal origin

79 Juvenile European sea bass originating from aquaculture farm have been raised from January 2020 to April 2021  
80 in a single tank. They were maintained at the natural seasonal temperature of the bay of Brest (min-max: 10-19°C)  
81 and were fed ad-libitum with a commercial diet thrice a week (NeoStart, Le Gouessant, Lamballe, France).

## 82 2.3. Muscle sampling and lipid extraction

83 In April 2021, six individuals randomly chosen (15 ± 2 cm; 50 ± 10 g) were euthanized using tricaine (400 mg l<sup>-1</sup>  
84 <sup>1</sup>) in seawater. On each individual, six subcutaneous white muscle biopsies were immediately performed on the  
85 left dorsal side of the fish, just behind the head, using a biopsy punch (2 mm LCH-PUK-20, Kai Medical, Solingen,  
86 Germany) (Fig. S1).

87 Three out of the six biopsies (further named “Fresh”, 8.8 ± 1.0 mg WW) were directly dived into 6 mL of a mix  
88 of chloroform/methanol (2:1, v/v), following a modified method from (Folch, 1957) for lipid extraction. The three  
89 other biopsies (further named “FD” for Freeze-Dried, 9.5 ± 0.8 mg WW) were first flash frozen in liquid nitrogen,  
90 and then freeze-dried for 36 hours before being dived into 6 mL of a mix of chloroform/ methanol (2:1, v/v).

91 After having performed biopsies, a large muscle sample (further named “MC” for Muscle control, 80.0 ± 24.4 mg  
92 WW) was sampled on the same area as where biopsies were performed, in each individual (Fig. S1). MC were  
93 treated following our routine protocol for FA analyses. It was flash frozen, then grounded in liquid nitrogen into a  
94 homogenous powder and finally, 40 to 100 mg WW of muscle powder was dived in a 6 mL mix of  
95 chloroform/methanol (2:1, v/v). To ensure a total lipid extraction, all the lipid extracts (Fresh, FD, and MC) were  
96 sonicated for 10 min and agitated for 20 min before being stored at -20°C under nitrogenous atmosphere prior  
97 further FA analysis.

## 98 2.4. Analysis of the muscle fatty acid composition

### 99 2.4.1. Lipid separation

100 For all samples, lipids were separated into neutral lipids (NL) and polar lipids (PL) fractions following the method  
101 described by Le Grand et al. (2014). An aliquot (one sixth) of the total lipid extract was used for Muscle control  
102 samples, whereas the whole lipid extract was used for biopsies (both Fresh and FD), as the mass of biopsy sampled  
103 tissue ranged from 1.4 to 14.4 mg for Fresh and from 5.5 to 13.6 mg for FD samples. The lipid extract aliquot was  
104 evaporated to dryness under nitrogen, recovered with three washings of 0.5 mL of chloroform/methanol (98:2,  
105 v/v), and deposited at the top of a silica gel column (40 mm × 4 mm, silica gel 60 A 63–200 µm rehydrated with  
106 6% H<sub>2</sub>O (70–230 mesh)). NL were eluted with 10 mL of chloroform/methanol (98:2, v/v) and collected in a glass  
107 vial containing 2.3 µg of internal standard (tricosanoic acid, C23:0). PL were then eluted with 20 mL of methanol  
108 and collected in the same way as NL.

### 109 2.4.2. Neutral lipid and polar lipid transesterification

110 NL and PL fractions were processed following a modified version of Mathieu-Resuge et al. (2019). Briefly, NL  
111 and PL fractions were evaporated to dryness under nitrogen. Fatty Acid Methyl Esters (FAME) were then formed  
112 by adding 800 µL of H<sub>2</sub>SO<sub>4</sub>/methanol (3.4%, v/v) and heating at 100°C for 10 min in a dry bath. After cooling,

113 800  $\mu$ L of hexane and 1.5 mL of hexane-saturated distilled water were added to the mixture. After removal of the  
114 aqueous phase, the hexane phase, containing FAME, was washed two more times with hexane-saturated distilled  
115 water.

#### 116 2.4.3. FAME analysis


117 Analyses of FAME was performed on a Varian CP8400 gas chromatograph using simultaneously two separations  
118 on a polar column (DBWAX: 30 mm  $\times$  0.25 mm ID  $\times$  0.2  $\mu$ m, Agilent) and an apolar column (DB5: 30 m  $\times$  0.25  
119 mm ID  $\times$  0.2  $\mu$ m, Agilent) using hydrogen as carrier gas with a flow rate set at 2 mL.min<sup>-1</sup>, in splitless mode. The  
120 oven temperature was programmed as follows: first, initial heating from 60°C to 150 °C at 50 °C.min<sup>-1</sup>, then to  
121 170 °C at 3.5 °C.min<sup>-1</sup>, to 185 °C at 1.5 °C.min<sup>-1</sup>, to 225 at 2.4 °C.min<sup>-1</sup> and finally to 250 °C at 5.5 °C.min<sup>-1</sup>  
122 and maintained for 25 min. Injector temperature was set at 250°C, detector temperature at 280°C. Identification of  
123 FAME was realized by comparison of their retention time based on those of commercial standards (Supelco 37  
124 Component FAME Mix, PUFA No.1 and No.3, and Bacterial Acid Methyl Ester Mix, Sigma, MO, USA). Internal  
125 standard allowed to calculate FA content ( $\mu$ g.mg<sup>-1</sup> WW). Fatty acid proportion was defined as the mass percentage  
126 of each fatty acid to the total fatty acid content.

#### 127 2.5. Statistical analysis

128 All statistical analyses were conducted using R Studio (v4.2.1). In order to determine if the biopsy method would  
129 be spatially repeatable within one individual, we calculated the intra-individual variation as the CV (standard  
130 deviation/mean) of the studied parameter (FA quantities and proportion), for each individual and for each biopsy  
131 method (n=3 Fresh and n=3 FD for each individual). This was performed on both quantity and proportions of EPA,  
132 DHA, PUFA, monounsaturated fatty acid (MUFA) and saturated fatty acid (SFA) in both NL and PL fractions. In  
133 order to be able to characterize the range of intra-individual variability, we compared these intra-individual CV to  
134 the corresponding inter-individual CV performed on MC samples (n=6).

135 We hypothesized that if the intra-individual CV exceeded either 10% (Sardenne et al. 2019) or the corresponding  
136 inter-individual CV, the biopsy method would not be repeatable enough to be validated and used with confidence.  
137 On the contrary, if the intra-individual variability was lower than 10% or than the inter-individual variability, we  
138 hypothesized that the biopsy method would be repeatable. It is the necessary condition allowing us to calculate the  
139 mean of the biopsy triplicates and to conduct a PERMANOVA test to compare the effect of the method (Fresh,  
140 FD, MC) on the FA proportions. Only the ten main FA were considered in this analysis. To evaluate if there were  
141 any differences in specific FA proportions between the methods, each FA was analyzed using a one-way ANOVA  
142 test. Normality and homoscedasticity were assessed on residuals with the Shapiro-Wilk and Bartlett tests,  
143 respectively. If ANOVA assumptions were not met, a Kruskal-Wallis test was employed instead. When ANOVA  
144 or Kruskal-Wallis tests indicated significant differences, a post-hoc Tukey HSD test was conducted to differentiate  
145 the conditions. Data are presented as mean  $\pm$  SEM.


### 146 3. Results



148 **Fig. 1: Fatty acid quantities (mg.g<sup>-1</sup> WW upper panels) and coefficients of variation (percentage, lower**  
 149 **panels) of FA quantities in neutral lipids (A, C) and polar lipids (B, D) of different method.** The different  
 150 methods are presented as followed: Freeze-dried protocol (FD, red, n=3 intra-individual replicates), fresh protocol  
 151 (Fresh, green, n=3 intra-individual replicates) and muscle control (MC, blue, n=6 inter-individual replicates) in  
 152 white muscle of juvenile European sea bass, *Dicentrarchus labrax*. On the boxplots, each dot represent a replicate.  
 153 On the dot plot, each dot represents the CV of the 3 (for biopsies) or 6 replicates (for MC) and the blue dotted line  
 154 represent the value of MC CV.

### 155 3.1. Adequacy of biopsied tissues for FA analysis.

156 White muscle biopsy samples ranging from 1.5 to 14.4 mg were sufficient to obtain good quality chromatograms.  
 157 In addition, the biopsy method allowed us to quantify over 90% of the FA identified in the Muscle control method.  
 158 The biopsy total lipid extracts contained  $86.1 \pm 13.5 \mu\text{g}$  (min-max: 5.5-409.0  $\mu\text{g}$ ) of FA for NL and  $28.8 \pm 1.7 \mu\text{g}$   
 159 (mix-max: 10.8-45.6  $\mu\text{g}$ ) of FA for PL (not shown). These values were similar to those contained in the 1/6 aliquots



160 of MC total lipid extracts used in the routine method:  $86.1 \pm 13.5 \mu\text{g}$  (min-max: 5.5-409.0  $\mu\text{g}$ ) of FA for NL and  
 161  $28.8 \pm 1.7 \mu\text{g}$  (mix-max: 10.8-45.6  $\mu\text{g}$ ) of FA for PL (not shown).

162 **Fig. 2: Coefficient of variation (percentage) of FA proportions in NL (A) and PL (B) for the different**  
 163 **methods.** Main family and FA are presented: Saturated fatty acids (SFA), monounsaturated fatty acids (MUFA),  
 164 polyunsaturated fatty acids (PUFA), ecosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) for freeze-  
 165 dried protocol (FD, red, n=3 intra-individual replicates), fresh protocol (Fresh, green, n=3 intra-individual  
 166 replicates) and muscle control (MC, blue, n=6, inter-individual replicates) in white muscle of juvenile European  
 167 sea bass, *Dicentrarchus labrax*. Each dot represents the CV of the 3 (for biopsies) or 6 replicates (for MC) and the  
 168 blue dotted line represent the value of MC CV.

### 169 3.2. Repeatability of the biopsy method within an individual

170 Our results indicate that both biopsy methods resulted in high intra-individual variability in total FA quantity in  
 171 NL (Fig. 1 NL A) (Fresh: min-max: 1.0-98.4  $\text{mg g}^{-1}$ , FD: min-max: 0.7-74.4  $\text{mg g}^{-1}$ ) this variability, illustrated by  
 172 the coefficient of variation of the three replicates per individual and biopsy method (CV mean =  $75 \pm 40\%$ , min-  
 173 max = 15-147%) was always above 10% (Fig. 1C). The CV values were higher than those of MC for 5 out of 6  
 174 individuals for both methods. In PL, the intra-individual variability of total FA quantity was lower than in NL (Fig.

175 1B) but still displayed considerable differences between extreme values (Fresh: min-max: 2.2-17.7 mg.g<sup>-1</sup>, FD:  
 176 min-max: 1.4-5.6 mg.g<sup>-1</sup>). The CV was higher than 10% for most individuals and ranged from 3% to 74% for both  
 177 Fresh and FD methods (CV mean = 23 ± 19.8%, Fig. 1D). The CV values were higher than those of MC for 4 out  
 178 of 6 individuals for both methods. In general, no biopsy method distinguished itself from the other in terms of  
 179 variability.

180 Regarding the NL fraction, SFA, MUFA and PUFA proportions (Fig. 2A) varied less within individuals (CV  
 181 <10%), than EPA and DHA (Fig. 2A, E-F, EPA min-max: 3-21.1%; DHA min-max: 2.9-49.6%), whatever the  
 182 sampling method employed, either biopsy or routine. For EPA, the biopsy CV was always lower than the MC CV.  
 183 For DHA, 11 out of 12 CV were lower than the MC CV. Regarding PL, even if the CV varied among individuals,  
 184 they remained under 10% for all individuals and for both methods (Fig. 2B).

### 185 3.3. Comparison between biopsy methods and a traditional but lethal method of sampling

186 Concerning FA quantities, the biopsy results showed a high intra-individual variability for both methods. We then  
 187 considered that biopsy replicates would not be repeatable enough and then could not be pooled to obtain a mean per  
 188 individual, and thus, that testing the effect of method on FA quantities would not be relevant.

189 Regarding biopsy methods, most of the FA proportions in NL and PL had a CV <10% and had lower CV than the  
 190 inter-individual one (Fig. 2). We thus hypothesized that the biopsy triplicates performed within individuals could  
 191 be pooled, and that the effect of method can be tested on FA proportions. When considering the 10 major fatty  
 192 acids (Fig. S2), no significant effect of the method was observed for both NL and PL (PERMANOVA, p > 0.05).  
 193 However, when considering the FA individually, 16:0 and 18:0 FA exhibited a significantly higher proportion in  
 194 the NL fraction when analyzed with the FD method than with the two others (Table 1), and consequently, SFA  
 195 proportions followed this trend. No statistical differences were observed among the three methods for PL FA  
 196 proportions (Table 1).

197 **Table 1: Fatty acid proportion (mass percentage of each FA over total fatty acids) of the 10 major fatty**  
 198 **acids measured in neutral (NL) and polar (PL) lipids with the three methods: Fresh biopsy, freeze-dried**  
 199 **biopsy (FD) and muscle control (MC) in white muscle of juvenile European sea bass, *Dicentrarchus labrax*.**

200 Data are presented as inter-individual means (± SEM, n=6), after pooling intra-individual biopsy triplicates. The  
 201 effect of the method on the FA proportions was tested using a one-way ANOVA or Kruskal-Wallis test, followed  
 202 by Tukey's post hoc test or pairwise Wilcoxon test. Significance was accepted at p < 0.05. Different letters read in  
 203 line indicate significant differences between methods. \*\*\* p < 0.001, \*\* p < 0.01, \* p < 0.05, - NS (not significant).  
 204 SFA: saturated fatty acids; MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids.

NL Fatty Acid	Muscle Biopsies		Muscle Control	Statistics
	Fresh	Freeze-Dried		
14:0	2.9 ± 0.2	3.1 ± 0.2	3.1 ± 0.2	-
16:0	18.1 ± 0.3 <sup>a</sup>	18.2 ± 0.1 <sup>a</sup>	18.2 ± 0.1 <sup>a</sup>	**
18:0	3.7 ± 0.2 <sup>a</sup>	3.6 ± 0.1 <sup>a</sup>	3.6 ± 0.1 <sup>a</sup>	**
16:1n-7	5.7 ± 0.3	5.9 ± 0.3	5.9 ± 0.3	-
18:1n-7	3.0 ± 0.1	3.1 ± 0.1	3.1 ± 0.1	-

18:1n-9	29.0 ± 1.5	28.8 ± 1.2	28.8 ± 1.2	-	205
18:2n-6	13.4 ± 1.1	13.1 ± 0.7	13.1 ± 0.7	-	
18:3n-3	2.5 ± 0.2	2.4 ± 0.1	2.4 ± 0.1	-	
20:5n-3	5.5 ± 0.7	5.5 ± 0.6	5.5 ± 0.6	-	
22:6n-3	2.8 ± 0.4	2.8 ± 0.3	2.8 ± 0.3	-	
SFA	25.6 ± 0.6 <sup>a</sup>	26.0 ± 0.3 <sup>a</sup>	26.0 ± 0.3 <sup>a</sup>	**	
MUFA	42.8 ± 0.9	43.1 ± 0.8	43.1 ± 0.8	-	
PUFA	31.3 ± 0.5	30.7 ± 0.5	30.7 ± 0.5	-	
<b>PL Fatty Acid</b>	<b>Muscle Biopsies</b>			<b>Muscle control</b>	<b>Statistics</b>
	<b>Fresh</b>	<b>Freeze-Dried</b>			
16:0	23.2 ± 0.3	23.7 ± 0.5	22.6 ± 0.3	-	
18:0	5.9 ± 0.1	5.8 ± 0.1	6.1 ± 0.2	-	
16:1n-7	2.5 ± 0.1	2.4 ± 0.1	2.7 ± 0.2	-	
18:1n-7	2.1 ± 0.0	2.1 ± 0.0	2.1 ± 0.1	-	
18:1n-9	13.1 ± 0.4	13.6 ± 0.4	14.0 ± 0.5	-	
18:2n-6	5.8 ± 0.4	5.9 ± 0.4	6.2 ± 0.4	-	
20:4n-6	1.7 ± 0.0	1.7 ± 0.0	1.7 ± 0.1	-	
20:5n-3	14.7 ± 0.2	14.5 ± 0.2	14.2 ± 0.4	-	
22:5n-3	2.4 ± 0.1	2.2 ± 0.2	2.3 ± 0.1	-	
22:6n-3	17.0 ± 0.5	16.8 ± 0.5	16.3 ± 0.7	-	
SFA	31.1 ± 0.4	31.5 ± 0.5	30.6 ± 0.4	-	
MUFA	19.8 ± 0.5	20.2 ± 0.5	21.1 ± 0.6	-	
PUFA	46.8 ± 0.5	46.2 ± 0.5	45.8 ± 0.7	-	

206

## 4. Discussion

207

## 4.1. Adequacy of biopsied tissue for FA analysis

208

Our results show that white muscle biopsy using punch of 2 mm on sea bass juvenile of about 15 cm and 50 g is adequate to perform fatty acid analysis. The sampled biopsied tissues were sufficient to yield at least 5.5 µg of FA for NL and 10.8 µg for PL, which was enough to obtain qualitative chromatograms with FA having a peak area reaching the minimal threshold. The average FA quantity extracted was even much higher than these minimums, which supports that white muscle biopsy method provides an adequate quantity of tissue for FA analysis. Olsen et al. (2013) stated that 5 µg of FA is the minimum amount required to obtain interpretable chromatograms. In the same study, the authors suggested to use a minimum of 60 mg WW of muscle but this value should consider the type of tissue, life stage of the fish, the species, as well as the analytical equipment used. The fish size and weight are crucial considerations when sampling tissue to ensure minimal impact on the animal. In our study, we did not tested the effect of biopsy on fish growth and health condition, as they were sacrificed. But in a related study, Quémeneur et al. (2022) conducted red muscle biopsies on European sea bass of a similar size (around 76.7g in weight) using the same biopsy punch as ours. They observed that the biopsy did not affect the fish's immune system, and the fish continued to gain weight post-procedure. Henderson et al. (2016) used a larger biopsy punch (4mm compared to our 2mm) on Moses perch *Lutjanus russelli* and Grass emperor *Lethrinus laticaudis*. These

221

222 fish had a total length close to those used in our study (approximately 15.8 and 14.0 cm, respectively), and it was  
223 found that the biopsy procedure led to only temporary stress and did not result in any fish mortality.

#### 224 4.2. How repeatable is the FA analysis within an individual using the biopsy method?

225 Quantities of FA obtained with the biopsy methods were highly variable within individuals for both NL (almost  
226 100-time fold between min and max) and PL (almost 5-time fold between min and max). One hypothesis could be  
227 that these variations might be caused by the presence of lipid droplets in the tissue. Lipid droplets are intracellular  
228 structures mainly composed of NL, serving as lipid storage in various tissues, including muscle (Favé et al. 2004;  
229 Granneman et al. 2017; Kiessling et al. 2006), that would hence induce higher FA content variability in NL rather  
230 than in PL. However, lipid droplets are less abundant in fish white muscle compared to red muscle (Granneman et  
231 al. 2017; Kiessling et al. 2006) and may only be present in few samples, leading to FA quantities significantly  
232 higher in samples containing lipid droplets. In addition, the distribution of lipids in white muscle is known to vary  
233 depending on the sampling location (Charette et al. 2021), which may cause FA quantities to differ based on the  
234 sampling location of the biopsy.

#### 235 4.3. Comparison between methods and traditional lethal sampling

236 Contrary to quantities, the analysis of FA proportions using biopsy was repeatable. The low variability for SFA,  
237 MUFA, and PUFA proportions in NL measured within individuals (CV<10%) suggests relative homogeneity of  
238 these proportions within individuals, while higher variability in EPA and DHA proportions could indicate dietary  
239 or metabolic impacts leading to specific heterogeneity in FA profile. Is the biopsy method comparable to a  
240 traditional but lethal method?

241 For NL, the variability measured within individuals and illustrated by the high CV values that exceeded the  
242 targeted 10% threshold as well as the inter-individual variability (Sardenne et al. 2019), makes it difficult to  
243 compare the two-biopsy methods with the control methods. A higher quantity of tissue would have supported less  
244 variability between methods, as shown by Sardenne et al. (2019) who compared the effect of extraction method  
245 from wet and dry muscle tissue (around 100 mg) from gilthead seabream *Sparus aurata* and did not find any  
246 differences between methods when considering total lipids (NL + PL).

247 In terms of proportions, the biopsy method seemed to yield results comparable to the MC for both extraction  
248 protocols. As the intra-individual variability did not reach the threshold of our study, we considered the biopsy  
249 method repeatable and comparable for both extraction protocols. Indeed, no differences between biopsy and  
250 control methods were observed when considering the whole profile of 10 major FA for NL and PL. However,  
251 differences were observed when considering individually certain SFA (16:0 and 18:0) for the FD method despite  
252 being repeatable with our criteria. Nevertheless, we acknowledge that applying a uniform threshold to both  
253 proportions and quantities may not be wholly pertinent, as it involves two distinct aspects. It might be necessary  
254 to calibrate this threshold more appropriately, depending on whether we are investigating proportions or quantities  
255 due to their different nature and analytical variabilities.

#### 256 5. Conclusions

257 This study assessed the feasibility of a novel approach for measuring FA composition in European sea bass using  
258 subcutaneous white muscle biopsies. The method allowed successfully obtaining adequate sample biomass and

259 produced high-quality chromatograms for both analytical protocols. FA quantity obtained with both biopsy  
260 methods was highly variable within individuals, potentially due to factors such as lipid droplets and heterogeneous  
261 lipid distribution in the sampled tissue. We thus recommend to take a special care when realizing the biopsy, and  
262 to ensure that a sufficiently deep piece of tissue is removed, avoiding the muscle directly in contact with the skin  
263 which has different FA composition. The analysis of FA proportions appeared to be achievable with biopsy with  
264 both protocols. Although it is challenging to determine the optimal biopsy method for FA analysis in fish, this  
265 study demonstrates the potential of using subcutaneous white muscle biopsies in European sea bass for examining  
266 FA composition, with a particular focus on FA proportions. The fish species, its fat content and its size must also  
267 be considered to optimize lipid extraction and minimize the effect of the procedure on the animal.

## 268 6. Acknowledgments

269 We thank Fany Sardenne for her help and suggestion on previous versions of the manuscript.

## 270 7. Competing interests

271 The authors declare that they have no known competing financial interests or personal relationships that could  
272 have appeared to influence the work reported in this paper.

## 273 8. Funding

274 This work was supported by the Brittany region to Marie Vagner.

## 275 9. References

- 276 Arts MT, Kohler CC (2009) Health and condition in fish: the influence of lipids on membrane competency and  
277 immune response. In: Kainz M, Brett MT, Arts MT (eds) *Lipids in Aquatic Ecosystems*. Springer, New York,  
278 NY, pp 237–256
- 279 Baker RF, Blanchfield PJ, Paterson MJ, et al (2004) Evaluation of Nonlethal Methods for the Analysis of  
280 Mercury in Fish Tissue. *Transactions of the American Fisheries Society* 133:568–576.  
281 <https://doi.org/10.1577/T03-012.1>
- 282 Barbosa RG, Trigo M, Prego R, et al (2018) The chemical composition of different edible locations (central and  
283 edge muscles) of flat fish (*Lepidorhombus whiffiagonis*). *International Journal of Food Science & Technology*  
284 53:271–281. <https://doi.org/10.1111/ijfs.13583>
- 285 Bell MV, Henderson RJ, Sargent JR (1986) The role of polyunsaturated fatty acids in fish. *Comparative*  
286 *Biochemistry and Physiology Part B: Comparative Biochemistry* 83:711–719. [https://doi.org/10.1016/0305-](https://doi.org/10.1016/0305-0491(86)90135-5)  
287 [0491\(86\)90135-5](https://doi.org/10.1016/0305-0491(86)90135-5)
- 288 Bertrand M, Brosset P, Soudant P, Lebigre C (2022) Spatial and ontogenetic variations in sardine feeding  
289 conditions in the Bay of Biscay through fatty acid composition. *Marine Environmental Research* 173:105514.  
290 <https://doi.org/10.1016/j.marenvres.2021.105514>
- 291 Betancor MB, MacEwan A, Sprague M, et al (2021) Oil from transgenic *Camelina sativa* as a source of EPA and  
292 DHA in feed for European sea bass (*Dicentrarchus labrax* L.). *Aquaculture* 530:735759.  
293 <https://doi.org/10.1016/j.aquaculture.2020.735759>
- 294 Calder PC (2012) Mechanisms of Action of (n-3) Fatty Acids. *The Journal of Nutrition* 142:592S-599S.  
295 <https://doi.org/10.3945/jn.111.155259>
- 296 Cano-Rocabayera O, Maceda-Veiga A, de Sostoa A (2015) Fish fins and scales as non-lethally sampled tissues  
297 for stable isotope analysis in five fish species of north – eastern Spain. *Environ Biol Fish* 98:925–932.  
298 <https://doi.org/10.1007/s10641-014-0328-6>

299 Charette T, Rosabal M, Amyot M (2021) Mapping metal (Hg, As, Se), lipid and protein levels within fish  
300 muscular system in two fish species (Striped Bass and Northern Pike). *Chemosphere* 265:129036.  
301 <https://doi.org/10.1016/j.chemosphere.2020.129036>

302 Couturier LIE, Michel LN, Amaro T, et al (2020) State of art and best practices for fatty acid analysis in aquatic  
303 sciences. *ICES Journal of Marine Science* 77:2375–2395. <https://doi.org/10.1093/icesjms/fsaa121>

304 Dalsgaard J, St. John M, Kattner G, et al (2003) Fatty acid trophic markers in the pelagic marine environment.  
305 In: *Advances in Marine Biology*. Academic Press, pp 225–340

306 El-Sabaawi RW, Sastri AR, Dower JF, Mazumder A (2010) Deciphering the Seasonal Cycle of Copepod  
307 Trophic Dynamics in the Strait of Georgia, Canada, Using Stable Isotopes and Fatty Acids. *Estuaries and Coasts*  
308 33:738–752. <https://doi.org/10.1007/s12237-009-9263-8>

309 Evans RD (2008) Assessment of an underwater biopsy probe for collecting teleost fish tissue samples. *Marine*  
310 *Ecology Progress Series* 368:305–308. <https://doi.org/10.3354/meps07624>

311 Favé G, Coste T, Armand M (2004) Physicochemical properties of lipids: New strategies to manage fatty acid  
312 bioavailability. *Cellular and molecular biology (Noisy-le-Grand, France)* 50:815–31

313 Gourtay C, Chabot D, Audet C, et al (2018) Will global warming affect the functional need for essential fatty  
314 acids in juvenile sea bass (*Dicentrarchus labrax*)? A first overview of the consequences of lower availability of  
315 nutritional fatty acids on growth performance. *Mar Biol* 165:143. <https://doi.org/10.1007/s00227-018-3402-3>

316 Granneman JG, Kimler VA, Zhang H, et al (2017) Lipid droplet biology and evolution illuminated by the  
317 characterization of a novel perilipin in teleost fish. *eLife* 6:e21771. <https://doi.org/10.7554/eLife.21771>

318 Henderson CJ, Stevens TF, Lee SY (2016) Assessing the suitability of a non-lethal biopsy punch for sampling  
319 fish muscle tissue. *Fish Physiol Biochem* 42:1521–1526. <https://doi.org/10.1007/s10695-016-0237-z>

320 Hulbert AJ, Else PL (1999) Membranes as Possible Pacemakers of Metabolism. *Journal of Theoretical Biology*  
321 199:257–274. <https://doi.org/10.1006/jtbi.1999.0955>

322 Hurst TP, Copeman LA, Haines SA, et al (2019) Elevated CO<sub>2</sub> alters behavior, growth, and lipid composition of  
323 Pacific cod larvae. *Marine Environmental Research* 145:52–65. <https://doi.org/10.1016/j.marenvres.2019.02.004>

324 Ibeas C, Izquierdo MS, Lorenzo A (1994) Effect of different levels of n–3 highly unsaturated fatty acids on  
325 growth and fatty acid composition of juvenile gilthead seabream (*Sparus aurata*). *Aquaculture* 127:177–188.  
326 [https://doi.org/10.1016/0044-8486\(94\)90424-3](https://doi.org/10.1016/0044-8486(94)90424-3)

327 Jeffries KM, Teffer A, Michaleski S, et al (2021) The use of non-lethal sampling for transcriptomics to assess the  
328 physiological status of wild fishes. *Comparative Biochemistry and Physiology Part B: Biochemistry and*  
329 *Molecular Biology* 256:110629. <https://doi.org/10.1016/j.cbpb.2021.110629>

330 Kiessling A, Ruohonen K, BJØRNEVIK M Muscle fibre growth and quality in fish

331 Lawrence MJ, Raby GD, Teffer AK, et al (2020) Best practices for non-lethal blood sampling of fish via the  
332 caudal vasculature. *Journal of Fish Biology* 97:4–15. <https://doi.org/10.1111/jfb.14339>

333 Le Grand F, Soudant P, Siah A, et al (2014) Disseminated Neoplasia in the Soft-Shell Clam *Mya arenaria*:  
334 Membrane Lipid Composition and Functional Parameters of Circulating Cells. *Lipids* 49:807–818.  
335 <https://doi.org/10.1007/s11745-014-3917-4>

336 Li Y, Han D, Hu G, et al (2010) Inhibition of starch synthesis results in overproduction of lipids in  
337 *Chlamydomonas reinhardtii*. *Biotechnology and Bioengineering* 107:258–268. <https://doi.org/10.1002/bit.22807>

338 Mathieu-Resuge M, Kraffe E, Le Grand F, et al (2019) Trophic ecology of suspension-feeding bivalves  
339 inhabiting a north-eastern Pacific coastal lagoon: Comparison of different biomarkers. *Marine Environmental*  
340 *Research* 145:155–163. <https://doi.org/10.1016/j.marenvres.2019.02.016>

341 McKenzie DJ, Lund I, Pedersen PB (2008) Essential fatty acids influence metabolic rate and tolerance of  
342 hypoxia in Dover sole (*Solea solea*) larvae and juveniles. *Mar Biol* 154:1041–1051.  
343 <https://doi.org/10.1007/s00227-008-0998-8>

344 McKenzie DJ, Piraccini G, Piccolella M, et al (2000) Effects of dietary fatty acid composition on metabolic rate  
345 and responses to hypoxia in the European eel (*Anguilla anguilla*). *Fish Physiology and Biochemistry* 22:281–  
346 296. <https://doi.org/10.1023/A:1007865327923>

347 McNamara RK, Carlson SE (2006) Role of omega-3 fatty acids in brain development and function: Potential  
348 implications for the pathogenesis and prevention of psychopathology. *Prostaglandins, Leukotrienes and Essential*  
349 *Fatty Acids* 75:329–349. <https://doi.org/10.1016/j.plefa.2006.07.010>

350 Meyer L, Pethybridge H, Nichols PD, et al (2017) Assessing the Functional Limitations of Lipids and Fatty  
351 Acids for Diet Determination: The Importance of Tissue Type, Quantity, and Quality. *Frontiers in Marine*  
352 *Science* 4:

353 Nanton DA, Vegusdal A, Rørå AMB, et al (2007) Muscle lipid storage pattern, composition, and adipocyte  
354 distribution in different parts of Atlantic salmon (*Salmo salar*) fed fish oil and vegetable oil. *Aquaculture*  
355 265:230–243. <https://doi.org/10.1016/j.aquaculture.2006.03.053>

356 Olsen RE, Taranger GL, Svåsand T, Skilbrei OT (2013) Improved method for triacylglycerol-derived fatty acid  
357 profiling by various non-lethal and lethal sampling techniques in Atlantic salmon. *Aquaculture Environment*  
358 *Interactions* 4:251–261. <https://doi.org/10.3354/aei00087>

359 Özogul Y, Özogul F, Çiçek E, et al (2009) Fat content and fatty acid compositions of 34 marine water fish  
360 species from the Mediterranean Sea. *International Journal of Food Sciences and Nutrition* 60:464–475.  
361 <https://doi.org/10.1080/09637480701838175>

362 Quéméneur J-B, Danion M, Cabon J, et al (2022) The relationships between growth rate and mitochondrial  
363 metabolism varies over time. *Sci Rep* 12:16066. <https://doi.org/10.1038/s41598-022-20428-9>

364 Ruxton CHS, Reed SC, Simpson MJA, Millington KJ (2004) The health benefits of omega-3 polyunsaturated  
365 fatty acids: a review of the evidence. *Journal of Human Nutrition and Dietetics* 17:449–459.  
366 <https://doi.org/10.1111/j.1365-277X.2004.00552.x>

367 Sardenne F, Bodin N, Metral L, et al (2019) Effects of extraction method and storage of dry tissue on marine  
368 lipids and fatty acids. *Analytica Chimica Acta* 1051:82–93. <https://doi.org/10.1016/j.aca.2018.11.012>

369 Sargent J r., Bell J g., Bell M v., et al (1993) The Metabolism of Phospholipids and Polyunsaturated Fatty Acids  
370 in Fish. In: *Aquaculture: Fundamental and Applied Research*. American Geophysical Union (AGU), pp 103–124

371 Sargent JR, Tocher DR, Bell JG (2003) 4 - The Lipids. In: Halver JE, Hardy RW (eds) *Fish Nutrition* (Third  
372 Edition). Academic Press, San Diego, pp 181–257

373 Simopoulos AP (1991) Omega-3 fatty acids in health and disease and in growth and development. *The American*  
374 *Journal of Clinical Nutrition* 54:438–463. <https://doi.org/10.1093/ajcn/54.3.438>

375 Singer SJ, Nicolson GL (1972) The Fluid Mosaic Model of the Structure of Cell Membranes | *Science*

376 Skalli A, Robin JH, Le Bayon N, et al (2006) Impact of essential fatty acid deficiency and temperature on  
377 tissues' fatty acid composition of European sea bass (*Dicentrarchus labrax*). *Aquaculture* 255:223–232.  
378 <https://doi.org/10.1016/j.aquaculture.2005.12.006>

379 Tocher DR (2003) Metabolism and Functions of Lipids and Fatty Acids in Teleost Fish. *Reviews in Fisheries*  
380 *Science* 11:107–184. <https://doi.org/10.1080/713610925>

381 Tocher DR (2010) Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquaculture Research*  
382 41:717–732. <https://doi.org/10.1111/j.1365-2109.2008.02150.x>

383 Tocher DR, Zheng X, Schlechtriem C, et al (2006) Highly unsaturated fatty acid synthesis in marine fish:  
384 Cloning, functional characterization, and nutritional regulation of fatty acyl  $\Delta 6$  desaturase of Atlantic cod (*Gadus*  
385 *morhua* L.). *Lipids* 41:1003–1016. <https://doi.org/10.1007/s11745-006-5051-4>

386 Vagner M, Pante E, Viricel A, et al (2019) Ocean warming combined with lower omega-3 nutritional availability  
387 impairs the cardio-respiratory function of a marine fish. *Journal of Experimental Biology* 222:jeb187179.  
388 <https://doi.org/10.1242/jeb.187179>

389

390

