HAL open science

The miracle of integer eigenvalues Dedicated to Professor A.M.Vershik, on the occasion of his 90-th birthday

Richard Kenyon, Maxim Kontsevich, Oleg Ogievetsky, Cosmin Pohoata, Will Sawin, Senya Shlosman

To cite this version:

Richard Kenyon, Maxim Kontsevich, Oleg Ogievetsky, Cosmin Pohoata, Will Sawin, et al.. The miracle of integer eigenvalues Dedicated to Professor A.M.Vershik, on the occasion of his 90-th birthday. 2024. hal-04396050

HAL Id: hal-04396050

https://hal.science/hal-04396050v1

Preprint submitted on 15 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The miracle of integer eigenvalues

Dedicated to Professor A.M.Vershik,
on the occasion of his 90-th birthday

Richard Kenyon[1], Maxim Kontsevich[2], Oleg Ogievetsky[3,4,5],
Cosmin Pohoata[9], Will Sawin[6] and Senya Shlosman[3,5,7,8]

[1] Yale University, USA, [2] IHES, France
[3]Aix Marseille Univ, Universite de Toulon,
CNRS, CPT, Marseille, France
[4]Lebedev Physical Institute, Moscow, Russia,
[5]Inst. of the Information Transmission Problems, Moscow, Russia,
[6]Princeton University, USA, [7]BIMSA, Beijing, China,
[8]Skolkovo Institute of Science and Technology, Moscow, Russia
[9]Emory University, USA

January 10, 2024

Abstract

For partially ordered sets X we consider the square matrices M^X with rows and columns indexed by linear extensions of the partial order on X. Each entry $\left(M^X\right)_{PQ}$ is a formal variable defined by a *pedestal* of the linear order Q with respect to linear order P. We show that all the eigenvalues of any such matrix M^X are $\mathbb{Z}$ -linear combinations of those variables.

1 The statement of the main result

Let $X = \{\alpha_1, ..., \alpha_n\}$ be a partially ordered set with the partial order $\preccurlyeq$. A linear extension P of $\preccurlyeq$ is a bijection $P : X \to [1, ..., n]$, such that for any pair α_i, α_j, satisfying $\alpha_i \preccurlyeq \alpha_j$ we have $P(\alpha_i) \leq P(\alpha_j)$. We denote by Tot_X the set of all such linear extensions (i.e. total orders extending $\preccurlyeq$).

Let P, Q be two linear extensions of $\preccurlyeq$. We call the node $Q^{-1}(k) \in X$ a (P,Q)-disagreement node iff $P(Q^{-1}(k-1)) > P(Q^{-1}(k))$. The index k is called the disagreement index. By definition, the node $Q^{-1}(1)$ is a (P,Q)-agreement node. With every pair P, Q we associate the function $\varepsilon_{PQ} : \{1, ..., n-1\} \to \{0,1\}$, given by

$$\varepsilon_{PQ}(k) = \begin{cases} 1 & \text{if } Q^{-1}(k+1) \text{ is a } (P,Q)\text{-disagreement node,} \\ 0 & \text{otherwise.} \end{cases} \tag{1}$$

Note that for some pairs $(P,Q) \neq (P',Q')$ the functions ε_{PQ}, $\varepsilon_{P'Q'}$ can coincide (see the Examples section).

To formulate our main result we denote by $\mathcal{E} = \{\varepsilon : \{1, ..., n-1\} \to \{0,1\}\}$ the set of all 2^{n-1} different ε functions, and we associate with every ε a corresponding formal variable a_ε. For any poset X consider the square matrix M^X, whose matrix elements are indexed by the pairs (P,Q), and are given by $\left(M^X\right)_{PQ} = a_{\varepsilon_{PQ}}$.

For example, the poset $(X, \preccurlyeq)$ with three elements and one relation: $X = \{\{u,v,w\}, u < v\}$ has three linear extensions of $\preccurlyeq$: $u < v < w$, $u < w < v$, $w < u < v$. Let P be the linear extension $u < v < w$ and Q – the linear extension $u < w < v$. We have $\varepsilon_{PQ} = (0,1)$ since 2 is an agreement index ($u < v$ in both Q and P) and 3 is a disagreement index ($w < v$ in Q but not in P). The matrix M^X is

$$\begin{pmatrix} a_{00} & a_{01} & a_{10} \\ a_{01} & a_{00} & a_{10} \\ a_{01} & a_{10} & a_{00} \end{pmatrix}. \tag{2}$$

The eigenvalues of this matrix are $a_{00} - a_{01}$, $a_{00} - a_{10}$ and $a_{00} + a_{01} + a_{10}$, so they are $\mathbb{Z}$-linear combinations of the letters entering the matrix. One of us (O.O.) conjectured that it holds (the eigenvalues are $\mathbb{Z}$-linear combinations of the letters entering the matrix M^X) for every poset X. Below we present the proof of this conjecture:

Theorem 1 *For every poset X the eigenvalues of the matrix M^X are linear combinations of the variables a_ε with integer coefficients.*

The matrices M^X were introduced in the paper [OS]. It is proven there that the row sums $\sum_Q \left(M^X\right)_{PQ}$ do not depend on the row P, so the matrix M^X is 'stochastic' (up to scale), and $\Pi_X(\{a_\varepsilon\}) = \sum_Q \left(M^X\right)_{PQ}$ is its main

eigenvalue. In [OS] the corresponding sums are called the 'pedestal polynomials'. They enter into the expression for the generating functions of the monotone functions $f : X \to \{0, 1, 2, ...\}$ (e.g. the generating function of the number of plane partitions, spacial partitions, etc.):

$$\sum_{\text{monotone } f:X\to\{0,1,2,...\}} t^{\sum_{x\in X} f(x)} = \Pi_X(t) \prod_{k=1}^{n} \frac{1}{1-t^k}, \tag{3}$$

where the polynomial $\Pi_X(t)$ is obtained from $\Pi_X(\{a_\varepsilon\})$ by the substitution

$$a_\varepsilon \rightsquigarrow t^{\sum_{k=1}^{n-1} k\varepsilon(k)}.$$

We put into the Appendix the relevant combinatorial facts about the pedestals and pedestal polynomials.

Our main tool is the filter semigroup of operators M_F^X, introduced in the next section. They have appeared first in [BHR, BD], where their spectral properties were studied. In fact, part of Theorem 1 can be obtained from Thm 1.2 in [BHR]. We give a shorter and more direct proof.

The next section contains some general facts about posets. It is followed by the section containing proofs.

2 The filter semigroup

At the end of this section we will introduce the filter semigroup. But it is easier to describe it geometrically, as the face semigroup of a hyperplane arrangement, which we do first.

2.1 Faces

Consider the central real hyperplane arrangement A_n consisting of hyperplanes $\{H_{ij} : 1 \le i < j \le n\}$ in $\mathbb{R}^n$ defined by $H_{ij} = \{(x_1, ..., x_n) : x_i = x_j\}$. Every open connected component of the complement $\mathbb{R}^n \setminus \{\cup H_{ij}\}$ is called a chamber. A *cone* is any union of closures of chambers, which is *convex*. Let us introduce the (finite) set $\mathfrak{O}(n)$ of all different cones thus obtained.

Let a poset X of n elements be given, with a binary relation $\preccurlyeq$. To every pair $i, j \in X$ which is in the relation $i \preccurlyeq j$ there corresponds a half-space $K_{ij} = \{x_i \le x_j\} \subset \mathbb{R}^n$ (here we assume that X is identified with $\{1, 2, \ldots, n\}$

as a plain set, ignoring the order). Consider the cone

$$A\left(X,\preccurlyeq\right)=\left\{\bigcap_{i,j:i\preccurlyeq j}K_{ij}\right\}\in\mathfrak{O}\left(n\right)$$

where the intersection is taken over all pairs i,j such that $i\preccurlyeq j$.

The following statements are well-known (and easy to prove), see [B, D, Sa, St].

Claim 2 *The above defined correspondence* $\left(X,\preccurlyeq\right)\rightarrow A\left(X,\preccurlyeq\right)$ *is a one-to-one correspondence between the set of all partial orders on* $\{1,2,...,n\}$ *and the set of all cones* $\mathfrak{O}\left(n\right)$.

We present an illustration of this claim for $n=4$ on Fig.1:



Figure 1: The central real hyperplane arrangement A_4 in $\mathbb{R}^4$, projected to $\mathbb{R}^3$ along the line $x=y=z=t$ and intersected with the sphere $\mathbb{S}^2\subset\mathbb{R}^3$. It is a partition of $\mathbb{S}^2$ into 24 equal triangles, each with the angles $\left(\frac{\pi}{2},\frac{\pi}{3},\frac{\pi}{3}\right)$. The types of convex unions of the triangles are: the sphere, the hemisphere, the region between two great semicircles, an elementary triangle – or e-triangle, a pair of e-riangles with a common side, a triangle made from three e-triangles, a 'square' formed by four e-triangles with a common $\frac{\pi}{2}$-vertex, a triangle made from a 'square' and a fifth adjacent e-triange, a triangle formed by six e-triangles with a common $\frac{\pi}{3}$-vertex. The number of corresponding convex shapes are 1, 12, 60, 24, 36, 48, 6, 24, 8, with total being 219. This is precisely the number of partial orders on the set of four distinct elements, see the sequence A001035 in OEIS [Sl].

Let f',f'' be two faces in $A(X)=A\left(X,\preccurlyeq\right)$. (It is allowed that one or both of them are in fact chambers, i.e. faces of highest dimension). Define the face $f=f''\left(f'\right)\in A\left(X\right)$ – or the *face-product* $f''f'$ – by the following procedure: choose points $x'\in f'$, $x''\in f''$ in general position and let $s_{x'x''}:[0,1]\rightarrow\mathbb{R}^n$ be

a linear segment, $s_{x'x''}(0) = x'$, $s_{x'x''}(1) = x''$. Consider the face $f \in A(X)$ which contains all the points $s_{x'x''}(1-\varepsilon)$ of our segment for $\varepsilon > 0$ small enough. Such a face does exist due to the convexity of $A(X)$. By definition, $f''(f') = f$. Note that if f'' is a chamber then $f''f' = f''$.

The face-product is associative. We mention for completeness that the semigroups $A(X, \preccurlyeq)$ are what are called *left-regular bands,* see [Sa]:

Claim 3 *For every choice of faces $f, g, h \in A(X, \preccurlyeq)$ we have*

$$f(gh) = (fg)h,$$

$$ff = f, \quad fgf = fg.$$

We do not give here the proofs as we are not using these relations.

2.2 Filters

Let F be a filter on X of rank k, i.e. a surjective map $F : X \to \{1, ..., k\}$, preserving the partial order, and let

$$\{b_1, ..., b_{j_1}\}, \{b_{j_1+1}, ..., b_{j_2}\}, ..., \{b_{j_{k-1}+1}, ..., b_{j_k}\} \subset X$$

be its 'floors':

$$\{b_{j_{r-1}+1}, ..., b_{j_r}\} = F^{-1}(r), \ r = 1, ..., k.$$

Consider the face $f_F \in A(X, \preccurlyeq)$, defined by the equations

$$x_{b_{j_{r-1}+1}} = ... = x_{b_{j_r}}, \ r = 1, ..., k$$

and inequalities

$$x_{b_{j_1}} < x_{b_{j_2}} < ... < x_{b_{j_k}}.$$

(More precisely, we write an equation for every floor of F which contains at least two elements of X.) This is a one-to-one correspondence between faces and filters. The filters of the highest rank n, i.e. the linear extensions of $\preccurlyeq$, correspond to the chambers.

The corresponding filter-product looks as follows. For F', F'' being two filters of X, the filter $F = F''F'$ on X is uniquely defined by the following properties:

- For u, v with $F''(u) < F''(v)$ we have $F(u) < F(v)$.
- For u, v with $F''(u) = F''(v)$ we have $F(u) < F(v)$ iff $F'(u) < F'(v)$.

Indeed, let f', f'' be the two faces, corresponding to the filters F', F'', and the general position points x', x'' belong to corresponding faces.

The fact that $F''(u) < F''(v)$ means that $x''_u < x''_v$. But the point $s_{x'x''}(1-\varepsilon)$ is close to the point x'', therefore $[s_{x'x''}(1-\varepsilon)]_u < [s_{x'x''}(1-\varepsilon)]_v$ for all ε small enough.

The fact that $F''(u) = F''(v)$ while $F'(u) < F'(v)$ means that $x''_u = x''_v$ while $x'_u < x'_v$. Since the map $s_{x'x''} : [0,1] \to \mathbb{R}^n$ is linear, for any $t < 1$ we have $[s_{x'x''}(t)]_u < [s_{x'x''}(t)]_v$.

Let F be a filter on X, and P is some filter of rank n, i.e. a linear order on X. Then the filter FP is again a filter of rank n. Consider the square matrix $M_F^X = \left\| \left(M_F^X\right)_{P,Q} \right\|$ where P, Q are linear orders on X :

$$\left(M_F^X\right)_{P,Q} = \begin{cases} 1 & \text{if } Q = FP \\ 0 & \text{if } Q \neq FP \end{cases} .$$

The operators M_F^X play a central role in our proof.

Examples of the operators M_F^X are given in the Example section below.

3 Proof of the main result

The plan of the proof is the following:

1. We will show that the matrix M^X can be written as a linear combination of M_F^X-s with integer coefficients.
2. We will show that all M_F^X-s can be made upper-triangular via conjugation with the **same** matrix, and the resulting upper-triangular matrices have integer entries on the diagonal.

3.1 The filter decomposition

Let us rewrite M^X as the sum over all 2^{n-1} functions $\varepsilon : \{1, ..., n-1\} \to \{0, 1\}$:

$$M^X = \sum_{\varepsilon} a_\varepsilon B_{X,\varepsilon}, \tag{4}$$

where the entries of each matrix $B_{X,\varepsilon}$ are 0 or 1.

For every function ε we define the number $r(\varepsilon) = 1 + \sum_{j=1}^{n-1} \varepsilon(j)$, and we partition the segment $\{1, ..., n\}$ into $r(\varepsilon)$ consecutive segments

$$\begin{aligned}\{1, ..., n\} = \{1, ..., c_1\} \\ &\cup \{c_1 + 1, ..., c_1 + c_2\} \\ &\cup \{c_1 + c_2 + 1, ..., c_1 + c_2 + c_3\} \cup ... \\ &\cup \left\{c_1 + ... + c_{r(\varepsilon)} + 1, ..., n\right\},\end{aligned}$$

where the values $c_1 + 1, c_1 + c_2 + 1, ..., c_1 + ... + c_{r(\varepsilon)} + 1$ are all the points where the function ε takes value 1.

For $c_1, ..., c_r$ being integers summing up to n we denote by $\mathcal{F}_{c_1,...,c_r}$ the set of all filters $F : X \to [1, 2, ..., r]$ such that $|F^{-1}(i)| = c_i$ for all $i = 1, ..., r$.

Lemma 4 *Suppose that the matrix $B_{X,\varepsilon} \neq 0$, and the function ε has the parameters r and $c_1, ..., c_r$. Then the following inclusion-exclusion identity holds:*

$$B_{X,\varepsilon} = \sum_{F \in \mathcal{F}_{c_1,...,c_r}} M_F^X - \left[\sum_{\substack{F \in \mathcal{F}_{c_1+c_2,c_3,...,c_r} \cup \\ \cup \mathcal{F}_{c_1,c_2+c_3,...,c_r} \cup ...}} M_F^X \right] + \left[\sum_{\substack{F \in \mathcal{F}_{c_1+c_2+c_3,c_4,...,c_r} \cup \\ \cup \mathcal{F}_{c_1+c_2,c_3+c_4,...,c_r} \cup ...}} M_F^X \right] - ... \tag{5}$$

where the sums are taken over all possible mergers of neighboring indices c_i, and the signs are $(-1)^{\#mergers}$.

Proof. Indeed, if we take an order Q from the row P which appears in the lhs, then it agrees with P over the first $c_1 - 1$ locations, then it disagrees once, then it agrees again over next $c_2 - 1$ locations, then disagrees once again, etc. But an order Q from the row P which appears in the rhs and corresponds to the first sum in (5), agrees with P over the first $c_1 - 1$ locations, then it **agrees or disagrees** once, then it agrees again over next $c_2 - 1$ locations, then **agrees or disagrees** once again, etc. Therefore we have to remove all these Q-s which agrees with P over the first $c_1 - 1$ locations, then **agrees** once again, then agrees also over next $c_2 - 1$ locations, etc.

See the Examples section for some M_F^X operators. ■

3.2 Conjugation of M_F^X-s to upper-triangular

Our matrices M_F^X are of the size $|\mathsf{Tot}_X| \times |\mathsf{Tot}_X|$. Let us now abolish all order relations on X, getting the poset $\bar{X}$ with $|\mathsf{Tot}_{\bar{X}}| = n!$. Of course, M_F^X is a submatrix of $M_F^{\bar{X}}$. Imagine (after reindexing) that it is an upper-left submatrix. We claim that to the right of this submatrix all matrix elements of $M_F^{\bar{X}}$ are zero, and so M_F^X is a block of $M_F^{\bar{X}}$. Indeed, each row of $M_F^{\bar{X}}$ has exactly one 1, and the rest are 0-s. But each row of M_F^X already has one 1. So it is sufficient to know that the spectrum of $M_F^{\bar{X}}$ consists of integers.

In what follows, the initial poset X will not appear any more, and we will deal only with 'totally unordered' poset $\bar{X}$. The fact that the matrices $M_F^{\bar{X}}$ can be conjugated simultaneously to upper-triangular ones can be deduced from the results of the papers [BHR, BD]. We give a shorter and more direct proof.

Let us consider an even bigger matrix, $N_F^{\bar{X}}$, of size $2^{n(n-1)/2}$. Here F is a filter on X, while the rows and columns of $N_F^{\bar{X}}$ are indexed by *tournaments* between the n entries of $\bar{X}$. A *tournament* is an assignment of an order $\preccurlyeq$ to each pair $i \neq j$ of the elements of the set $\bar{X}$, independently for each pair. If we have a tournament $\preccurlyeq$ and a filter F on $\bar{X}$, then we define a new tournament $\preccurlyeq_F$ by the rule:

1. If $F(i) = F(j)$ then $i \preccurlyeq_F j$ iff $i \preccurlyeq j$,
2. If $F(i) < F(j)$ then $i \preccurlyeq_F j$.

We define $N_F^{\bar{X}}$ by

$$\left(N_F^{\bar{X}}\right)_{\preccurlyeq\preccurlyeq'} = \begin{cases} 1 & \text{if } \preccurlyeq' = \preccurlyeq_F \\ 0 & \text{if } \preccurlyeq' \neq \preccurlyeq_F \end{cases}$$

Any linear order defines a tournament in an obvious way, so our matrices $M_F^{\bar{X}}$ are blocks of $N_F^{\bar{X}}$-s, and it is sufficient to study $N_F^{\bar{X}}$-s.

The key observation now is the fact that $N_F^{\bar{X}}$ is a tensor product of $n(n-1)/2$ two-by-two matrices, corresponding to all pairs (i,j), since the tournament orders $\preccurlyeq$ can be assigned to the pairs independently. And since the tensor product of upper triangular matrices is upper triangular, it is sufficient to check our claim just for the filters and tournaments in the case $n = \left|\bar{X}\right| = 2$.

The two-element no-order set $\bar{X} = \{1, 2\}$ carries three different filters and has two possible tournaments. The three two-by-two matrices $N_F^{\bar{X}}$-s are

$N_1 := \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$, $N_2 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $N_3 := \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}$. Conjugating them by the discrete Fourier transform matrix $U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ brings them to the triple of upper triangular matrices: $UN_1U^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$, $UN_2U^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $UN_3U^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$. Extending the conjugation through the tensor product finishes the proof. ■

Remark 5 *Recall the definition (4) of the matrices B_μ: for a poset X, the set of (0,1)-valued matrices $\{B_{X,\varepsilon}\}_{\varepsilon \in \{0,1\}^{\{1,\dots,n-1\}}}$ is defined by $M^X = \sum_\varepsilon a_\varepsilon B_{X,\varepsilon}$. Let $\mathcal{L}(X)$ be the Lie algebra generated by the matrices $\{B_{X,\varepsilon}\}$. The proof shows that the Lie algebra $\mathcal{L}(X)$ is solvable.*

Let us denote by Φ_T the algebra of functions on the set $\mathsf{Tour}_{\bar{X}}$ of tournaments considered as the set $\{-1,+1\}^{n(n-1)/2} \subset \mathbb{R}^{n(n-1)/2}$ of vertices of the cube. This algebra carries an increasing filtration by subspaces

$$0 \subset \Phi_T^{\leq 0} \subset \Phi_T^{\leq 1} \subset \cdots \subset \Phi_T^{\leq \frac{n(n-1)}{2}} = \Phi_T$$

consisting of restrictions of polynomials of degree $\leq 0, \leq 1, \dots$ to the vertices of the cube. This filtration is **strictly multiplicative** in the sense that

$$\Phi_T^{\leq k} = \underbrace{\Phi_T^{\leq 1} \cdot \dots \cdot \Phi_T^{\leq 1}}_{k \text{ times}} .$$

Our considerations imply that all operators $N_F^{\bar{X}}$ preserve this filtration, and commute with each other on the associated graded space $\oplus_k \Phi_T^{\leq k}/\Phi_T^{\leq k-1}$. Indeed, Φ_T has basis $\prod_{i \in S} x_i$ where S runs through all subsets of $\{1, \dots, \frac{n(n-1)}{2}\}$. For each coordinate x_i we have two *commuting* operators $N_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $(x_i \to -x_i)^* = N_1 + N_3 - N_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ acting on the i-th tensor factor $\mathbb{R}^{\{-1,+1\}} = \mathbb{R}[x_i]/(x_i^2 - 1))$ preserving the filtration by degree, as well as one extra operator $(N_3 - N_1) = \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix}$ which *strictly decreases* the filtration (it acts as $2\partial_{x_i}$ in the basis $(\prod_{i \in S} x_i)_{S \subseteq \{1,\dots,n(n-1)/2\}}$). We see that on the associated graded space $\oplus_k \Phi_T^{\leq k}/\Phi_T^{\leq k-1}$ our operators give the ones coming from the commutative group algebra of the abelian group $(\mathbb{Z}/2\mathbb{Z})^{n(n-1)/2}$ acting by involutions of coordinates.

Restricting functions from Φ_T to the subset $\mathsf{Tot}_X \subset \mathsf{Tour}_{\bar{X}}$ we obtain again a strictly multiplicative filtration on the algebra $\Phi_X := \mathbb{R}^{\mathsf{Tot}_X}$ of functions on Tot_X, preserved by all operators M_F^X where F runs through filters on the poset X.

4 Appendices

4.1 Pedestals

Let again X be a finite poset with the partial order $\preccurlyeq$, and P, Q be a pair of linear orders on X, consistent with $\preccurlyeq$. We define the function q_{PQ} on X by

$$q_{PQ}\left(Q^{-1}(k)\right) = \#\left\{l : l \leq k, Q^{-1}(l) \text{ is a } (P,Q)\text{-disagreement node}\right\}. \quad (6)$$

Clearly, the function q_{PQ} is non-decreasing on X, and $q_{PQ}\left(Q^{-1}(1)\right) = 0$. It is called the pedestal of Q with respect to P.

For example, let X be a 3×2 Young diagram, and

$$P = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}, \; Q = \begin{bmatrix} 1 & 2 & 5 \\ 3 & 4 & 6 \end{bmatrix}$$

be the two standard tableaux. Then

$$q_{PQ} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Let $\mathcal{E}_P$ denotes the set of all pedestals q_{PQ}. Clearly, there is a map $\mathcal{E}_P \to \mathcal{E}$, which to every pedestal q_{PQ} corresponds its 'discrete derivative' ε_{PQ}.

The pedestals were introduced in [S] in the following context. Consider the set $\mathcal{P} = \mathcal{P}_X$ of all non-negative integer-valued non-decreasing functions p on X. Denote by $v(p)$ the 'volume' of p :

$$v(p) = \sum_{\alpha \in X} p(\alpha),$$

and let G be the following generating function:

$$G_X(t) = \sum_{k \geq 0} g_k t^k = \sum_{p \in \mathcal{P}_X} t^{v(p)},$$

i.e. g_k is the number of non-decreasing p-s with $v(p) = k$. For example, if the poset X is in fact the set $X_n = [1, 2, ..., n]$, ordered linearly, then

$$G_{X_n}(t) = \prod_{l=1}^{n} \frac{1}{1 - t^l}$$

is the generating function of the sequence g_k of the number of partitions π of the integer k into at most n parts: $k = \pi(1) + \pi(2) + ... + \pi(n)$, with $\pi(i) \geq 0$, $\pi(i) \leq \pi(i+1)$. Let $\mathcal{Y}_n$ denote the set of all such partitions π (i.e. Young diagrams).

In order to write a formula for G_X for an arbitrary poset X one needs pedestals. Namely, let us fix some ordering P of X, consider all pedestals q_{PQ}, and let

$$\Pi_P(t) = \sum_Q t^{v(q_{PQ})} \tag{7}$$

be the generating function (in fact, generating polynomial) of the sequence of the number of pedestals with a given volume. Then we have the identity:

$$G_X(t) = \Pi_P(t) G_{X_n}(t) \equiv \Pi_P(t) \prod_{l=1}^{n} \frac{1}{1-t^l}, \tag{8}$$

(compare with (3)). In particular, it follows from (8) that the polynomial $\Pi_P(t)$ does not depend on P, and thus can be denoted by $\Pi_X(t)$. The reason for (8) to hold is the existence of the bijection between the set $\mathcal{P}_X$ of nondecreasing functions and the direct product $\mathcal{E}_P \times \mathcal{Y}_n$, respecting the volumes. Namely, to each pedestal q_{PQ} and each partition π it associates the following function p on X :

$$p\left(Q^{-1}(k)\right) = q_{PQ}\left(Q^{-1}(k)\right) + \pi(k), \ k = 1, ..., n.$$

Clearly, the function thus defined is non-decreasing on X. For the check that this is a one-to-one correspondence, see [S].

In the case when X is a (2D) Young diagram, the functions $p \in \mathcal{P}_X$ are called 'reverse plane partitions'. The generating function G_X for these is also given by the famous MacMahon formula,

$$G_X(t) = \prod_{\alpha \in X} \frac{1}{1-t^{h(\alpha)}},$$

where $h(\alpha)$ is the hook length of the cell $\alpha \in X$. That means that for the case of X being a Young diagram nice cancellations happen in the rhs of (8). One can check that for some X being a 3D Young diagram no cancellations happen in (8), and this is the reason why the analog of the MacMahon formula in the 3D case does not exist.

4.2 Pedestal polynomials

The fact that the function $\Pi_P(t)$ (see (7)) does not depend on the order P on X, but only on X, has the following generalization. Instead of characterizing the pedestal q_{PQ} just by its volume let us associate with it the monomial $m_{PQ}(x_1, x_2, x_3, ...) = x_1^{l_1-1} x_2^{l_2-l_1} ... x_r^{l_r-l_{r-1}} x_{r+1}^{n-l_r+1}$, where r is the number of (P,Q)-disagreement nodes, and $l_1, ..., l_r$ are their locations, see (6). Note that $m_{PQ}(1, t, t^2, ...) = t^{v(q_{PQ})}$.

It was shown in [OS] that the polynomial

$$\mathfrak{h}_P(x_1, x_2, x_3, ...) = \sum_{Q \in \mathsf{Tot}_X} m_{PQ}(x_1, x_2, x_3, ...)$$

is also independent of P, so it can be denoted as $\mathfrak{h}_X(x_1, x_2, x_3, ...)$. Another way of expressing this is to say that the matrix $\tilde{M}^X$ of size $|\mathsf{Tot}_X| \times |\mathsf{Tot}_X|$, with entries $\left(\tilde{M}^X\right)_{PQ} = m_{PQ}(x_1, x_2, x_3, ...)$ is *stochastic*, i.e. the vector $(1, 1, ..., 1)$ is the right eigenvector, with the eigenvalue $\mathfrak{h}_X(x_1, x_2, x_3, ...)$.

By replacing the monomials $m_{PQ}(x_1, x_2, x_3, ...)$ with variables $a_{\varepsilon_{PQ}}$ one obtains from $\tilde{M}^X$ our matrix M^X.

Remark 6 *As we just said, we know from [OS] that the rows of the matrix M^X consist of the same matrix elements, permuted. So it is tempting to consider the set of permutations $\pi_{PP'} \in S_{|\mathsf{Tot}_X|}$, which permute the elements of the row P to these of row P'. Unfortunately, rows of the matrix M^X can contain repeated elements, so the permutations $\pi_{PP'}$ are not uniquely defined.*

4.3 Examples

Here we present several examples in which our posets X correspond to partitions; standard Young tableaux are read row by row.

0. In all examples we considered the pedestal matrix is diagonalisable in the generic point. However for special values of variables the pedestal matrix might have non-trivial Jordan blocks. We give a minimal example - partition (3,1). It is essentially the same example as the one before the main theorem, with the pedestal matrix (2), because the box (1,1) comes first in any linear order and can be omitted.
Here it is enough to take a partial evaluation $a_{10} \mapsto -2a_{01}$. Then the Jordan form is

$$\begin{pmatrix} a_{00} - a_{01} & 1 & 0 \\ 0 & a_{00} - a_{01} & 0 \\ 0 & 0 & a_{00} + 2a_{01} \end{pmatrix}.$$

It would be interesting to understand the regimes in which the pedestal matrix is not diagonalisible.

1. Partition (3,2). The standard tableaux are

$$\{1,3,5,2,4\}\ ,\ \{1,2,5,3,4\}\ ,\ \{1,3,4,2,5\}\ ,\ \{1,2,4,3,5\}\ ,\ \{1,2,3,4,5\}.$$

The pedestal matrix $\tilde{M}^X$ is $x_1^2 A_{(3,2)}$, where

$$A_{(3,2)} = \begin{pmatrix} x_1^3 & x_2^3 & x_1^2x_2 & x_2^2x_3 & x_1x_2^2 \\ x_2^3 & x_1^3 & x_2^2x_3 & x_1^2x_2 & x_1x_2^2 \\ x_1^2x_2 & x_2^2x_3 & x_1^3 & x_2^3 & x_1x_2^2 \\ x_2^2x_3 & x_1^2x_2 & x_2^3 & x_1^3 & x_1x_2^2 \\ x_2^2x_3 & x_1^2x_2 & x_2^3 & x_1x_2^2 & x_1^3 \end{pmatrix} .$$

After a replacement

$$\phi : (x_1^3, x_1^2x_2, x_1x_2^2, x_2^3, x_2^2x_3) \to (a_1, a_2, a_3, a_4, a_5)\ , \tag{9}$$

we have

$$A^{\phi}_{(3,2)} = \begin{pmatrix} a_1 & a_4 & a_2 & a_5 & a_3 \\ a_4 & a_1 & a_5 & a_2 & a_3 \\ a_2 & a_5 & a_1 & a_4 & a_3 \\ a_5 & a_2 & a_4 & a_1 & a_3 \\ a_5 & a_2 & a_4 & a_3 & a_1 \end{pmatrix} .$$

The eigenvalues of $A^{\phi}_{(3,2)}$ are

$$a_1-a_3\ ,\ a_1+a_2-a_4-a_5\ ,\ a_1-a_2+a_4-a_5\ ,\ a_1-a_2-a_4+a_5\ ,\ a_1+a_2+a_3+a_4+a_5\ .$$

2. Partition (3,1,1). The standard tableaux are

$$\{1,4,5,2,3\}\ ,\ \{1,3,5,2,4\}\ ,\ \{1,2,5,3,4\}\ ,\ \{1,3,4,2,5\}\ ,\ \{1,2,4,3,5\}, \{1,2,3,4,5\}\ .$$

The pedestal matrix is $x_1^2 A_{(3,1,1)}$ where

$$A_{(3,1,1)} = \begin{pmatrix} x_1^3 & x_1x_2^2 & x_2^3 & x_1^2x_2 & x_2^2x_3 & x_1x_2^2 \\ x_1x_2^2 & x_1^3 & x_2^3 & x_1^2x_2 & x_2^2x_3 & x_1x_2^2 \\ x_1x_2^2 & x_2^3 & x_1^3 & x_2^2x_3 & x_1^2x_2^2 & x_1x_2^2 \\ x_1x_2^2 & x_1^2x_2 & x_2^2x_3 & x_1^3 & x_2^3 & x_1x_2^2 \\ x_1x_2^2 & x_2^2x_3 & x_1^2x_2 & x_2^3 & x_1^3 & x_1x_2^2 \\ x_1x_2^2 & x_2^2x_3 & x_1^2x_2 & x_2^3 & x_1x_2^2 & x_1^3 \end{pmatrix} .$$

After the same replacement (9) (the matrix $A_{(3,1,1)}$ contains the same monomials as the matrix $A_{(3,2)}$) we have

$$A^{\phi}_{(3,1,1)} = \begin{pmatrix} a_1 & a_3 & a_4 & a_2 & a_5 & a_3 \\ a_3 & a_1 & a_4 & a_2 & a_5 & a_3 \\ a_3 & a_4 & a_1 & a_5 & a_2 & a_3 \\ a_3 & a_2 & a_5 & a_1 & a_4 & a_3 \\ a_3 & a_5 & a_2 & a_4 & a_1 & a_3 \\ a_3 & a_5 & a_2 & a_4 & a_3 & a_1 \end{pmatrix} .$$

The eigenvalues of $A^{\phi}_{(3,1,1)}$ are (the notation $(y)_k$ means that the multiplicity of the eigenvalue y is k)

$$(a_1-a_3)_2 \ , \ a_1+a_2-a_4-a_5 \ , \ a_1-a_2+a_4-a_5 \ , \ a_1-a_2-a_4+a_5 \ , \ a_1+a_2+2a_3+a_4+a_5 \ .$$

The example $(3,1,1)$ shows degeneration: the letter a_3 appears twice in every row of $A^{\phi}_{(3,1,1)}$. The corresponding monomial is $x_1^3x_2^2$ so for writing down the decomposition of the matrix B_{a_3} we need filters from $\mathcal{F}_{3,2}$. There are three of them in $\mathcal{F}_{3,2}$ (the notation is like for a matrix; element (i,j) is in the intersection of row i and column j):

- F_1: Floor 1 contains cells (1,1), (1,2) and (2,1);
- F_2: Floor 1 contains cells (1,1), (1,2) and (1,3);
- F_3: Floor 1 contains cells (1,1), (2,1) and (3,1).

The matrices of action of these filters on the linear orders are

$$M_{F_1} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, M_{F_2} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$M_{F_3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} .$$

The family $\mathcal{F}_5$ contains one filter, which acts as the identity I. The matrix B_{a_3} is thus

$$B_{a_3} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{pmatrix} = M_{F_1} + M_{F_2} + M_{F_3} - I \ ,$$

as dictated by the inclusion-exclusion formula.

3. Partition (3,2,1). The standard tableaux are

$$\begin{array}{llll} \{1,4,6,2,5,3\} \ , & \{1,3,6,2,5,4\} \ , & \{1,2,6,3,5,4\} \ , & \{1,3,6,2,4,5\} \ , \\ \{1,2,6,3,4,5\} \ , & \{1,4,5,2,6,3\} \ , & \{1,3,5,2,6,4\} \ , & \{1,2,5,3,6,4\} \ , \\ \{1,3,4,2,6,5\} \ , & \{1,2,4,3,6,5\} \ , & \{1,2,3,4,6,5\} \ , & \{1,3,5,2,4,6\} \ , \\ \{1,2,5,3,4,6\} \ , & \{1,3,4,2,5,6\} \ , & \{1,2,4,3,5,6\} \ , & \{1,2,3,4,5,6\} \ . \end{array}$$

To save the space we write down the pedestal matrix in which the replacement

$$(x_1^6, x_1^5x_2, x_1^4x_2^2, x_1^4x_2x_3, x_1^3x_2^3, x_1^3x_2^2x_3, x_1^2x_2^4, x_1^2x_2^3x_3, x_1^2x_2^2x_3^2, x_1^2x_2^2x_3x_4) \rightarrow$$

$$(a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10})$$

is made. This matrix $A^{\phi}_{(3,2,1)}$ is

$$A^{\phi}_{(3,2,1)} = \begin{pmatrix} a_1 & a_5 & a_7 & a_3 & a_9 & a_2 & a_6 & a_8 & a_3 & a_9 & a_5 & a_2 & a_8 & a_4 & a_{10} & a_6 \\ a_5 & a_1 & a_7 & a_3 & a_9 & a_6 & a_2 & a_8 & a_3 & a_9 & a_5 & a_2 & a_8 & a_4 & a_{10} & a_6 \\ a_5 & a_7 & a_1 & a_9 & a_3 & a_6 & a_8 & a_2 & a_9 & a_3 & a_5 & a_8 & a_2 & a_{10} & a_4 & a_6 \\ a_5 & a_3 & a_9 & a_1 & a_7 & a_6 & a_2 & a_8 & a_4 & a_{10} & a_6 & a_2 & a_8 & a_3 & a_9 & a_5 \\ a_5 & a_9 & a_3 & a_7 & a_1 & a_6 & a_8 & a_2 & a_{10} & a_4 & a_6 & a_8 & a_2 & a_9 & a_3 & a_5 \\ a_2 & a_6 & a_8 & a_3 & a_9 & a_1 & a_5 & a_7 & a_3 & a_9 & a_5 & a_4 & a_{10} & a_2 & a_8 & a_6 \\ a_6 & a_2 & a_8 & a_3 & a_9 & a_5 & a_1 & a_7 & a_3 & a_9 & a_5 & a_4 & a_{10} & a_2 & a_8 & a_6 \\ a_6 & a_8 & a_2 & a_9 & a_3 & a_5 & a_7 & a_1 & a_9 & a_3 & a_5 & a_{10} & a_4 & a_8 & a_2 & a_6 \\ a_6 & a_2 & a_8 & a_4 & a_{10} & a_5 & a_3 & a_9 & a_1 & a_7 & a_5 & a_3 & a_9 & a_2 & a_8 & a_6 \\ a_6 & a_8 & a_2 & a_{10} & a_4 & a_5 & a_9 & a_3 & a_7 & a_1 & a_5 & a_9 & a_3 & a_8 & a_2 & a_6 \\ a_6 & a_8 & a_2 & a_{10} & a_4 & a_5 & a_9 & a_3 & a_7 & a_5 & a_1 & a_9 & a_3 & a_8 & a_6 & a_2 \\ a_5 & a_3 & a_9 & a_2 & a_8 & a_6 & a_4 & a_{10} & a_2 & a_8 & a_6 & a_1 & a_7 & a_3 & a_9 & a_5 \\ a_5 & a_9 & a_3 & a_8 & a_2 & a_6 & a_{10} & a_4 & a_8 & a_2 & a_6 & a_7 & a_1 & a_9 & a_3 & a_5 \\ a_6 & a_4 & a_{10} & a_2 & a_8 & a_5 & a_3 & a_9 & a_2 & a_8 & a_6 & a_3 & a_9 & a_1 & a_7 & a_5 \\ a_6 & a_{10} & a_4 & a_8 & a_2 & a_5 & a_9 & a_3 & a_9 & a_2 & a_6 & a_9 & a_3 & a_7 & a_1 & a_5 \\ a_6 & a_{10} & a_4 & a_8 & a_2 & a_5 & a_9 & a_3 & a_8 & a_6 & a_2 & a_9 & a_3 & a_7 & a_5 & a_1 \end{pmatrix} .$$

The eigenvalues of $A^{\phi}_{(3,2,1)}$ are

$$(a_1-a_4-a_7+a_{10})_3\ ,\ a_1-a_4+a_7-a_{10}\ ,\ (a_1+a_2-a_5-a_6)_2\ ,\ (a_1-a_2-a_5+a_6)_2\ ,$$

$$(a_1-a_2-a_3+a_4+a_7-a_8-a_9+a_{10})_2\ ,\ (a_1-a_2-a_3+a_4-a_7+a_8+a_9-a_{10})_2\ ,$$

$$(a_1-a_4+a_5-a_6+a_7-a_{10})_2\ ,\ a_1+2a_2+2a_3+a_4-a_7-2a_8-2a_9-a_{10}\ ,$$

$$a_1+2a_2+2a_3+2a_5+2a_6+a_7+2a_8+2a_9+a_{10}\ .$$

Acknowledgements. Part of this work was done in IHES during the program dedicated to the 100-th (or rather 102-nd) anniversary of the Ising model, in May – June of 2022. R.K., M.K. and S.S. are grateful to the organizers of the program for the uplifting atmosphere. R.K. was supported by NSF grant DMS-1940932 and the Simons Foundation grant 327929. The work of W.S. was supported by the NSF grant DMS-2101491 and by the Sloan Research Fellowship. The work of S.S. was supported by the RSF under project 23-11-00150.

References

[BHR] Bidigare P., Hanlon P. and Rockmore D., *A combinatorial description of the spectrum for the Tsetlin library and its generalization to hyperplane arrangements*, Duke Math. Journal, 1999, Jul; 99(1):135-74.

[B] Brown K.S., *Semigroups, rings, and Markov chains*, Journal of Theoretical Probability, 2000, Jul; 13:871-938.

[BD] Brown K.S. and Diaconis P., *Random walks and hyperplane arrangements*, Annals of Probability, 1998, Oct; 1:1813-54.

[D] Dorpalen-Barry, G.A., *Cones of hyperplane arrangements* (Doctoral dissertation, 2021, University of Minnesota).

[OS] Ogievetsky O. and Shlosman S., *Plane Partitions and Their Pedestal Polynomials*, Mathematical Notes, 2018, May; 103(5):793-6.

[Sa] Saliola F.V., *The face semigroup algebra of a hyperplane arrangement*, Canadian Journal of Mathematics, 2009, Aug; 61(4):904-29.

[S] Shlosman S., *The Wulff construction in statistical mechanics and combinatorics* Russian Mathematical Surveys, 56(4), p.709-38, 2001.

[Sl] Sloane N.J.A., *The on-line encyclopedia of integer sequences*, http://www.oeis.org/.

[St] Stanley R., *Enumerative Combinatorics*, Cambridge University Press, New York, NY, USA, 2 edition, 2012.