



HAL
open science

Exposure of whales to entanglement risk in Dungeness crab fishing gear in Oregon, USA, reveals distinctive spatio-temporal and climatic patterns

Solène Derville, T.V. Buell, K.C. Corbett, C. Hayslip, L.G. Torres

► To cite this version:

Solène Derville, T.V. Buell, K.C. Corbett, C. Hayslip, L.G. Torres. Exposure of whales to entanglement risk in Dungeness crab fishing gear in Oregon, USA, reveals distinctive spatio-temporal and climatic patterns. *Biological Conservation*, 2023, 281, pp.109989. 10.1016/j.biocon.2023.109989 . hal-04398293

HAL Id: hal-04398293

<https://hal.science/hal-04398293v1>

Submitted on 16 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Exposure of whales to entanglement risk in Dungeness crab fishing gear in Oregon, USA, reveals distinctive spatio-temporal and climatic patterns

Derville, S.^{1,2,*}, Buell, T.³, Corbett, K.³, Hayslip, C.¹, Torres, L. G.¹

¹ GEMM Lab, Marine Mammal Institute, Department of Fisheries, Wildlife and Conservation Sciences, Oregon State University, 2030 SE Marine Science Drive, Newport, OR 97365, United States

² French Institute of Research for Sustainable Development, UMR ENTROPIE, 101 promenade Roger Laroque, 98800 Noumea, New Caledonia

³ Marine Resources Program, Oregon Department of Fish and Wildlife, 2040 Marine Science Drive Newport, OR 97365, United States

* Corresponding Author

s.derville@live.fr +33620799502

Abstract

Entanglement in fishing gear presents a major threat to marine mammals worldwide and a pressing concern for distinct populations of whales off the US West Coast. The lack of understanding of their fine-scale distribution in relation to fishing activity limits management efforts, specifically in Oregon. Based on year-round predictions of orca whale densities and fishing effort compiled from logbooks, we assess co-occurrence between commercial Dungeness crab fishing gear and whales over a decade (2011-2020) as an indicator of exposure to entanglement risk. Generalized Additive Models including temporal, climatic, and ocean upwelling predictors were used to investigate variations in exposure. Exposure peaked in April, at the onset of the upwelling season when whales were predicted to occur in greater numbers and closer to shore. Exposure remained constant until the end of the crab season in nearshore waters <40 fathoms (73 m) and decreased past these depths. Across years, exposure was lower during the marine heatwave (2014-2016) when fishing was more active nearshore and whales were predicted to be less abundant. Exposure was higher before (2011-2013) and after (2017-2020) the heatwave, which correspond to negative phases of the Pacific Decadal Oscillation associated with stronger upwelling, indicating more productive conditions favorable to whales. A recent increase in exposure was also due to a slight shift in fishing effort towards deeper waters. These findings illustrate the use of fine-scale species distribution models to assess space-use conflicts in dynamic marine ecosystems and can be used to guide fisheries management to reduce entanglement risk in Oregon.

Keywords: Entanglement, hindcast predictions, fishing gear, large whales, Dungeness crab, US West Coast

Introduction

Space use conflicts between fishing activities and megafauna is a long-standing marine conservation issue, with entanglement in fishing gear a main concern for large whale population recovery (Clapham, 2016). Entanglement can cause immediate or delayed mortality as it affects whale health, feeding success, and fecundity (van der Hoop, Corkeron and Moore, 2017; Carretta and Henry, 2022). These anthropogenic impacts can accumulate and interact with the effects of changing ocean conditions and prey availability resulting from climate change. Recent research indicates that extreme climatic events, such as marine heatwaves and regime shifts, may result in distribution changes that increase the overlap of large whales with fishing activities (Santora *et al.*, 2020; Meyer-Gutbrod *et al.*, 2021; Samhuri *et al.*, 2021). Hence, improved knowledge of the shifting spatio-temporal distribution of large whales with respect to fishing activities is necessary to anticipate entanglement risk and design appropriate conservation measures in a changing ocean.

Elevated counts of large whale entanglements have occurred in the past decade along the US West Coast (California, Oregon, and Washington), with particularly concerning levels recorded since 2014 (NOAA Fisheries, 2022). In cases where fishing gear involved in entanglements could be identified since 2014, commercial Dungeness crab (*Metacarcinus magister*) pot gear was one of the most frequently identified gear types (Saez, Lawson and Deangelis, 2021). This pot fishery operates in relatively shallow and nearshore waters of the continental shelf, with most landings occurring within the first few months of the crab season from December onwards (Feist *et al.*, 2021). Dungeness crab is the most economically important species harvested along the US West Coast (Rasmuson, 2013) and the fishery is culturally significant to coastal communities in the region (ODFW, 2022). This fishery is managed by states independently (CDFW, 2021; ODFW, 2021; WDFW, 2022) and is subject to local closures or delayed openings as a result of variable crab meat yield and domoic acid contamination, which are influenced by environmental fluctuations (Santora *et al.*, 2020; Feist *et al.*, 2021; Samhuri *et al.*, 2021). The fishery may also be locally closed or reduced due to whale entanglement, creating an inherent management tension between providing fishing opportunities and managing for whale conservation under variable climatic conditions. For example, since 2019 higher rates of whale entanglements that involved commercial Dungeness crab gear caused significant restrictions in the California crab fishery (CDFW, 2019). Consequently, understanding the environmental and social drivers of whale exposure to fishing activities is urgently needed to both protect whale populations and the vibrant, otherwise sustainable, Dungeness crab fishery across the West Coast.

Humpback whales (*Megaptera novaeangliae*) are most impacted by entanglements in commercial Dungeness crab gear, with the total human-caused serious injury and mortality of the California/Oregon/Washington stock (48.3 whales/year; (Carretta *et al.*, 2021) exceeding biologically sustainable levels under the U.S. Marine Mammal Protection Act (Potential Biological Removal level of 29.4 whales/year; (Carretta *et al.*, 2021). To a lesser extent, other rorquals (baleen whales of the family Balaenopteridae) such as the endangered blue whales (*Balaenoptera musculus musculus*) and fin whales (*Balaenoptera physalus*) are also at risk of entanglement while they migrate and forage off the US West Coast. These marine predators are known to shift their distribution in response to seasonal and interannual fluctuations of environmental conditions that drive prey availability (Becker *et al.*, 2017, 2018; Derville *et al.*, 2022). Indeed, rorqual whale distribution in the California Current System (CCS), which extends from California to British Columbia, Canada, is tightly connected to wind-driven upwelling that provides nutrient rich and cool waters that boost primary productivity in surface waters during spring and summer (Hazen *et al.*, 2017; Becker *et al.*, 2018; Abrahms *et al.*, 2019; Derville *et al.*, 2022). Yet, upwelling intensity, timing and duration vary across years (Benoit-Bird, Waluk and Ryan, 2019), notably due to basin-wide forcing reflected by climatic indices such as the Pacific Decadal Oscillation or the El Niño Southern Oscillation Index (PDO, SOI, e.g., Bograd *et al.*, 2009; Peterson *et al.*, 2017). Such variability

was suggested to be a main driver of increased entanglement rates observed in recent years in the southern CCS (Santora *et al.*, 2020). Relatively less is known about the interplay between whale and fishing activities that cause entanglements in the northern CCS (Riekkola *et al.*, 2023). While less entanglements have involved gear confirmed as originating from Oregon and Washington states compared to California (Saez, Lawson and Deangelis, 2021), the issue is of high concern for local coastal communities and state natural resource managers tasked with protecting single stocks of humpback, fin and blue whales that extend along the entire US West Coast. Specificities in fishing practices (ODFW, 2022), and whale ecological relationships (Derville *et al.*, 2022) warrant dedicated research efforts to understand patterns of exposure to entanglement risk in the northern CCS.

We leverage recent research outputs that predict orca whale distribution over the continental shelf off Oregon during the last 10 years (Derville *et al.*, 2022) to retrospectively analyze co-occurrence between commercial Dungeness crab fishing activity and orca whales indicating exposure to entanglement risk. We hypothesize that 1) whale exposure to entanglement risk varies in space and time, 2) spatio-temporal variations of exposure are related to shifting whale habitat and fishing effort distribution within and across years, and 3) variations of exposure are driven by upwelling conditions reflected by basin- and local-scale environmental indicators. Enhanced knowledge about the physical, biological and social drivers of the spatio-temporal variability of whale entanglement risk in Oregon will help design management strategies to address this threat.

MATERIAL & METHODS

Study area

Our study area includes the entire Oregon coast and was divided into six different zones of varying depth and latitude reflecting different fishing contexts (Figure 1). Three latitudinal zones were delineated: south, from Bandon to the southern Oregon border; central, between Bandon and Cascade Head; and north, from Cascade Head to the northern Oregon border. Two depth ranges delineated nearshore and offshore zones using the 40 fathom isobath (73.2 m) as this depth limit was recently adopted in fishery regulations effective May 2021 to restrict commercial Dungeness crab fishing annually to waters within 40 fathoms from May to August (Oregon Secretary of State, 2020) where orca whales generally occur less at this time of the year (Derville *et al.*, 2022). All spatial data were projected in a Universal Transverse Mercator coordinate system (UTM zone 10N) prior to analysis.

Fishing data

Commercial Dungeness crab fishing effort layers were constructed from fishery logbook data from the 2010-2011 (opened February 2011) through 2019-2020 crab seasons (hereafter referred to as “fishing years”). Commercial crab fishing season in Oregon may span from December to August. Fishing effort is reported in logbooks as the number of pot pulls, whereby a pot is lifted out of the water to harvest crab, after which it is usually redeployed. Therefore, pots are typically in the water nearly constantly from the time a vessel enters the fishery until it exits the fishery (ODFW, 2022). By regulation, commercial Dungeness crab pots may not be connected to each other via a groundline, therefore each pot requires a single vertical line connecting it to one or more surface buoys. Logbook data values likely to be outside the range of common fishing practices were excluded using a series of filters. Then, we allocated each vessel's pot limit (the maximum number that may legally be set at one time) proportional to pot pulls across fishing locations that vessel recorded in logbooks, at a monthly time scale. Our approach accounted for sub-sampling of logbooks for data entry in some years, non-compliance (landings with no logbook submitted), and logbook records missing critical data fields (see Supplementary Methods). We

then generated estimates of the number of pots in the water in layers of 5 km resolution, matching whale model outputs, for each month and year the fishery was open.

Whale data

Using the year-round rorqual whale models from (Derville *et al.*, 2022), we hindcasted rorqual densities by month for each year over layers of 5 km resolution. Three different Generalized Additive Models (GAM) were used to predict whale densities in the winter (December-March), spring (April-July), and summer (August-September), in relation to 10 topographic and dynamic ocean circulation variables averaged at a weekly scale (see (Derville *et al.*, 2022). Predictions were spatially limited to waters displaying environmental conditions analogous to those in which the models were trained. Predictions in non-analogous conditions, a practice known as environmental extrapolation, can lead to extreme and unrealistic predictions (Bouchet *et al.*, 2019). We used the Extrapolation Detection (ExDet) tool developed by (Mesgaran, Cousens and Webber, 2014) to evaluate this environmental extrapolation over the weekly layers of environmental variables on which the whale predictions were based (see Supplementary Methods). ExDet was computed with the *dsmextra* R package (version 1.1.5; (Bouchet *et al.*, 2020). ExDet was either considered at the scale of grid cells or averaged by study zone, and removed from further analyses appropriately (see below).

Fishing and whale center of mass

To detect spatial shifts of crab fishing effort and whale distribution, we calculated the depth and latitude of the center of mass for these layers by month for each year. Mean depths of fishing and whale distribution were calculated separately in the north, central and south zones (combining together the nearshore and offshore parts of the same latitudinal zone). Mean latitude of fishing and whale distribution were calculated separately in the offshore and nearshore zones (combining together the northern, central and southern parts of the same depth range). For the purpose of this analysis, we removed all grid cells showing environmental extrapolation with respect to the whale models (i.e., where ExDet < 0 or > 1). GAMs were fitted to the mean depth (models “M_{crab.cmass.depth}”, “M_{whale.cmass.depth}”) or the mean latitude (models “M_{crab.cmass.lat}”, “M_{whale.cmass.lat}”) by zone by month, using a gaussian distribution with identity link and a restricted maximum likelihood method in the *mgcv* R package (Wood, 2011). Explanatory variables included separate penalized thin-plate regression splines of month and year. Smooth basis size was limited to 3 to prevent overfitting. The effect of the explanatory variables on the overall model fit was assessed by examining the percent of deviance explained in comparison to a null model.

Patterns of overlap

First, the Williamson spatial overlap index (Williamson, Stoeckel and Schoeneck, 1989) was calculated per month to assess whether whale densities were uniformly distributed relative to fishing activity. Williamson index values > 1 and < 1 respectively represent overlap greater or less than expected from a uniform distribution:

$$Williamson\ index = \frac{\sum_{z=1}^m N_z \cdot n_z \cdot m}{\sum_{z=1}^m N_z \cdot \sum_{z=1}^m n_z}$$

Where *z* is the spatial unit (grid cells), *m* is the number of spatial units available (number of grid cells where ExDet ∈ 0,1), *N* is the predicted whale density, and *n* is the crab fishing effort.

Second, exposure evaluated co-occurrence between whales and fishing activity at a monthly scale, over layers of 5 km resolution calculated as the product between the fishing (number of pots) layers and the density of rorqual whales layers. Maps of monthly exposure were averaged together to identify spatial areas of elevated entanglement risk throughout the whole study period (2011-2020). For the purpose of this analysis, we removed all grid cells showing environmental extrapolation with respect to the whale models (i.e., where $ExDet < 0$ or > 1). These average maps of exposure were compared to the estimated gear set location of four humpback whale entanglement events with confirmed Oregon Dungeness crab fishing gear (Data provided by NMFS WCR, April 2021).

Variations in exposure

Grid cells of monthly exposure, in each of the six study zones were summed to investigate the spatio-temporal variations of entanglement risk in relation to month, fishing year and proxies of ocean conditions. GAMs were fitted to the log-transformed sum of exposure (models “ $M_{exposure}$ ”) by zone, using a gaussian distribution with an identity link and a restricted maximum likelihood method. Explanatory variables included separate penalized thin-plate regression splines of month, fishing year, and a suite of climate and upwelling indices (see below description). Smooth basis size was limited to 3 to prevent overfitting. Models with alternative sets of predictors were compared with one another using the Akaike Information Criterion (AIC).

In parallel, the same models were produced with the log-transformed summed abundance of whales (models “ M_{whale} ”) by zone and the log-transformed summed fishing effort (models “ M_{crab} ”) by zone to understand whether the changes in exposure could be due to changes in local whale abundance or the amount of fishing. Although the rorqual density models of Derville et al., (2022) were not meant to derive abundance trends for these whale populations as a whole, they allow the relative comparison of local estimates of abundance calculated over different time periods. For the purpose of these models, all zone x month combinations in which the mean $ExDet < 0$ were removed (no zone x month were found with $ExDet > 1$).

Climate and upwelling drivers

Climate indices known to influence upwelling conditions and productivity in the northern CCS were extracted at a monthly resolution: the PDO and the SOI (see Supplementary Methods for data sources). Upwelling conditions were estimated with the Coastal Upwelling Transport Index (Jacox, Edwards, *et al.*, 2018), which provides estimates of model derived vertical transport of seawater (in $m^2 \cdot s^{-1}$) at monthly and daily scales. Monthly CUTI values were incorporated as predictors in the GAMs, while daily values were used to quantify parameters of the upwelling phenology along the Oregon coast (Bograd *et al.*, 2009). Using the algorithms provided by (Oestreich *et al.*, 2022), we used the daily values to calculate the cumulative sum of CUTI by day, for each year that the index was available (1988-2021). The climatological mean, 5th percentile, and 95th percentile across years were calculated and smoothed with a 10-day running mean. Then, we calculated the mean spring transition index (STI), the mean peak upwelling (MAX), and the mean end of positive upwelling accumulation (END) based on the climatological mean of CUTI to describe average upwelling phenology by zone (north, central and south) from 1988 to 2021 (for more details see (Oestreich *et al.*, 2022).

All analyses were performed using R statistical computing (R Core Team, 2021).

RESULTS

Overlap between orca whales and commercial Dungeness crab fishing was assessed over 10 fishing years (83 months). The mean Williamson index was equal to $0.37 \pm \text{SD } 0.23$, hence indicating that overlap between fishing and whales was low and generally less than expected from a uniform distribution.

Spatial variations of exposure

The mean pattern of exposure throughout the study period (2011-2020) revealed several areas of higher risk that were mostly driven by the distribution of fishing effort along the coast of Oregon (Figure 1). Exposure was higher on average in nearshore waters off Astoria, off Garibaldi, north of Newport, north of Charleston, north of Port Orford, and at the southern border of Oregon waters. These areas overlapped with estimated set positions of crab gear involved in three of four confirmed humpback whale entanglements (Data provided by NMFS WCR, April 2021, see Supplementary Table S1) that involved gear estimated to be set in waters off Astoria, Garibaldi, and Charleston (Figure 1c).

Temporal variations of exposure

Local whale abundance increased with month in all zones but was mostly stable across years with a slight decrease in certain zones in the mid study period (M_{whale} , Figure 2a). The M_{whale} that included month and fishing year explained 46 % of the deviance on average, including a marginal 1 to 5 % deviance explained by fishing year depending on the zone considered (Table 1). In comparison, the crab fishing effort was stable across years and tended to decrease across months of a fishing year, particularly in the offshore zones (M_{crab} , Figure 2b). The average deviance explained by month and fishing year in M_{crab} was 36% (Table 1). The resulting temporal variations in exposure showed an increase throughout the first months of the fishing year (December-April), then ended in a plateau (April-August) for the nearshore zones or a decrease in the offshore zones (M_{exposure} , Figure 2c). The average deviance explained by month and fishing year in M_{exposure} was 20%, including a marginal 2 to 10 % deviance explained by fishing year depending on the zone considered (Table 1). Exposure was significantly lower in most zones during the 2014-2016 marine heat wave (Figure 2c).

Whale and fishing distribution shifts

Month and fishing year combined explained 11%, 38%, and 52% of the deviance in the depth of the center of mass for whale distribution ($M_{\text{whale.cmass.depth}}$) in the south, north and central zones respectively. The center of mass did not change across fishing years but it significantly shifted to shallower waters by month throughout the fishing year (min around 400 m; Figure 3a₁). Within the offshore zones, whale distribution also tended to move southward throughout the fishing year ($M_{\text{whale.cmass.lat}}$, deviance explained 51 %, Figure 3a₂) although it was stable in latitude across years. On average, the center of mass for whale distribution shifted south by 77 km in offshore zones between the months of Dec-Mar and May-Aug (t test: $t = -8.1$, $p < 0.001^{***}$).

Month and fishing year combined explained 46%, 45%, and 74% of the deviance in the depth of the center of mass for crab fishing effort ($M_{\text{crab.cmass.depth}}$) in the south, north and central zones respectively. The center of mass moved to shallower waters throughout the fishing year (min around 20 m; Figure 3b₁). Fishing effort also tended to move to shallower waters during the marine heatwave (Figure 3b₁) in the south (Anova: $n = 83$, $F\text{-value} = 13.7$, $p < 0.001^{***}$) and central (Anova: $n = 83$, $F\text{-value} = 3.7$, $p = 0.029^*$) zones while it remained stable in the north (Anova: $n = 83$, $F\text{-value} = 0.5$, $p = 0.6$). Indeed, in the central zone the center of mass for crab fishing effort was on average 5.9 m deeper (Tukey post-hoc test: $p = 0.043^*$) after the marine heat wave (2017-2020) than it was before (2010-2013). In the south zone, the center of mass for fishing effort was significantly deeper before (mean difference 4.1 m; Tukey

post-hoc test: $p = 0.047^*$) and after (mean difference 8.2 m; Tukey post-hoc test: $p < 0.001^{***}$) compared to during the marine heatwave. Due to the relatively flat slope of the continental shelf off the coast of Oregon, a 5 m drop in depth is equivalent to a westward shift of up to 10 km. This shift in fishing behavior in the south of Oregon is corroborated by a significant southward shift of the offshore center of mass throughout the study period ($M_{\text{crab.cmass.lat}}$, deviance explained 42%, Figure 3b₂). On average, the center of mass for crab fishing effort shifted south by 44 km in offshore zones between before and after the marine heatwave (t test: $t = -2.0$, $p = 0.046^*$).

Climatic drivers of exposure

Across the study period, the ecosystem alternated between two negative phases of PDO (2011-2014 and 2017-2021) and a positive phase (2014-2016) corresponding to the marine heat wave (Figure 4a). Within the fishing year, the upwelling conditions off the coast of Oregon varied with latitude (Figure 4b). The upwelling season (from STI to END) is on average longer in the south zone and shorter in the north zone (Figure 4b). Upwelling strength is also stronger in the south, as reflected in higher cumulative sum of CUTI at the peak and at the end of the upwelling season.

The exposure models including month, CUTI and PDO as predictors were systematically selected as the best models across almost all study zones (Table 1), with deviance explained ranging from 40% to 67%. Deviance explained by these predictors was also relatively high in the whale models (56 to 78%). Overall, CUTI and PDO significantly affected local whale abundance (Figure 5a) and exposure (Figure 5c) in almost all study zones but had almost no effect on crab fishing (Figure 5b). Negative PDO phases and strong upwelling (indicated by higher CUTI) were associated with higher local whale abundance and higher exposure overall, particularly in the south. Although the models of crab fishing effort, local whale abundance, and exposure with month, PDO and SOI were sometimes selected as the most parsimonious (Table 1), the effect of SOI was generally very minor (marginal deviance explained in crab fishing - 0.7%, local whale abundance + 0.2%, and exposure + 0.5% on average across zones).

DISCUSSION

We combined long-term fishing effort data and hindcasts of whale densities to identify locations and times with increased entanglement risk, thus providing critical information to managers to reduce space use conflicts between fishers and whales. Variations in exposure rates across months are driven by the timing of whale migration as well as spatial and seasonal patterns of fishing effort. Across years, higher exposure is predicted in the beginning and the end of our study period, due to greater local whale abundance predicted during negative PDO phases and stronger upwelling conditions, and to a lesser extent to a shift of the fishing effort towards deeper waters in recent years.

Co-occurrence indices indicate generally little overlap between the fishing effort and orca whale distribution. Yet, even at these low levels of co-occurrence, interactions between whales and fishing gear are occurring. Entanglements have been shown to contribute to biologically unsustainable rates of total human caused mortality for some humpback and blue whale stocks coast-wide (Carretta *et al.*, 2021). However, entanglements are often unobserved or under-reported (Tackaberry *et al.*, 2022). For example, fresh entanglement scars recorded on humpback whales off the US West Coast revealed that as much as 90% of entanglements may have gone unnoticed between 2009 and 2010 (Robbins, 2012). The likelihood for past entanglements involving Oregon crab gear to be unreported or impossible to trace back to a specific fishing zone limits our ability to estimate the trends of entanglement risk at a local scale. Yet, our analysis of co-occurrence with fishing gear provides a useful tool to locate the areas of elevated risk. Our modeling approach indicates high coincidence between areas of high mean exposure and three of

four entanglement reports involving Oregon crab gear with estimated gear set locations (Figure 1c), suggesting confidence in the spatial patterns of risk we identified.

The seasonality of Dungeness crab fishing and orca whale migration coincides with a peak of exposure around the month of April (Figure 2c). This temporal pattern is driven by the migratory timing of humpback whales (Derville *et al.*, 2022), the main species informing our orca distribution models. Blue and fin whales occur more rarely off the coast of Oregon, and while they were also assessed in this study, their respective phenology and habitat use patterns are likely to limit their interactions with the Dungeness crab fishery; blue whales tend to occur later than humpback whales (encounter rate peaks in September) and fin whales tend to occur further offshore in the winter (over and off the continental slope; (Derville *et al.*, 2022). To date, there have been no confirmed entanglements of either of these two species in Oregon commercial Dungeness crab gear (Saez, Lawson and Deangelis, 2021). Moreover, during the late crab season (May – August) exposure decreases offshore, but remains stable in nearshore waters (Figure 2c), a pattern likely driven by hard shell crab distribution as well as social and economic factors (Davis *et al.*, 2017). Larger fishing vessels typically stop fishing Dungeness crab around April, while smaller vessels tend to continue to fish later in the crab season, consolidating gear in more nearshore waters as the season progresses. We therefore identify marked seasonality and spatial patterns in entanglement risk off Oregon that result from a combination of ecological and human factors.

Across years, we found that variations in exposure are associated with environmental fluctuations that drive ocean productivity in the CCS. Exposure increases at all depths and along the whole Oregon coast (Figure 5c) during negative PDO phases and when upwelling is stronger. On the other hand, positive PDO phases tend to weaken upwelling strength and shift less warm water into the CCS, which results in diminished productivity and influence krill (Peterson *et al.*, 2017), forage fish (Santora *et al.*, 2017) and their predators (Henderson *et al.*, 2014). Between 2014 and 2016/2017, a positive PDO phase coinciding with other climatic drivers generated an anomalously warm water event that had widespread biological impacts throughout the Northeast Pacific (Jacox, Alexander, *et al.*, 2018). Observations of entangled humpback whales off the coast of California increased markedly during that period (Saez, Lawson and Deangelis, 2021), interpreted as the result of habitat compression that increased the overlap of whales with fishing gear (Santora *et al.*, 2020; Feist *et al.*, 2021). In contrast, our model predicted a decrease in exposure of orca whales to commercial Dungeness crab fishing during this marine heatwave in Oregon waters. This pattern was likely caused by decreased productivity and foraging conditions (Peterson *et al.*, 2017) that resulted in lower whale densities in Oregon waters. Given the known distinctiveness of oceanographic conditions and seabed topography of the northern CCS (Hickey and Banas, 2008; Castelao and Luo, 2018), we suggest that physical and biological drivers of whale distribution and therefore entanglement risk vary along the US West Coast. More specifically, the wide continental shelf off the coast of Oregon offers larger extents of suitable habitats for whales and fishing grounds, and our applied models indicate a strong link between local whale abundance and upwelling strength in Oregon (CUTI, restricted to a 43° to 46° latitude range, Figure 5a). While coast-wide management of whale populations that cross state boundaries is necessary, we emphasize the additional need to investigate drivers of whale entanglements at the state level that enable management decisions tailored to local conditions and risks.

Compared to other studies of wildlife-fisheries interaction that used Vessel Monitoring Systems (VMS) data to estimate fishing effort by inferring where fishing events occurred based on vessel location, speed, and bearing reported at pre-determined time intervals (Torres *et al.*, 2013; Breen *et al.*, 2017; Feist *et al.*, 2021), logbook data includes the self-reported position where fishing occurred but may be subject to human error. Also, VMS coverage in Oregon is biased as it is not currently required for all vessels participating in the Dungeness crab fishery, whereas logbook data is required for all participating vessels. While logbook data is an improvement, the data processing requires assumptions, particularly to correct

for the 30% sub-sampling for data entry in some years. We assumed that vessels fished their entire pot limit each month if they made a crab landing that month, potentially overestimating the total amount of gear set. Yet, this bias is likely to be randomly distributed and does not prevent the assessment of the relative distribution of entanglement risk across space and time (see Supplementary Methods). Also, whale density predictions were generated with seasonal models with variable predictive performance. The winter model that covered the peak of the crab season (December- March) had a lower predictive capacity than the spring and summer models (Derville *et al.*, 2022). Finally, the whale models predicted density based on habitat quality and did not account for intrinsic population trends of blue, fin and humpback whales. Indeed, the across-year trend of exposure that we modelled should be interpreted in parallel with the population increase identified in the CCS (Carretta *et al.*, 2021), and encounter rates off the coast of Oregon (Derville *et al.*, 2022). If these population trends persist and fishing efforts remains the same, exposure to entanglement risk is likely to be exacerbated in future PDO negative phases.

CONCLUSIONS

There was no major change in the overall amount of commercial crab gear set off Oregon from 2011 to 2020 (Figure 2b), which is consistent with the time series of fishing distribution acquired through VMS estimates up to 2016 (Feist *et al.*, 2021). Fishing effort slightly shifted to shallower waters during the marine heatwave and then to deeper waters in recent years (Figure 3b₁), potentially following environmentally-driven changes in Dungeness crab distribution. Although this shift in fishing effort contributed to an increase in exposure to entanglement risk in recent years (2017-2020), the main driver of exposure appears to be whale habitat suitability related to climatic fluctuations. These results can inform fishery management measures to reduce the risk of entanglements of orca whales. Temporary area closures and shortening of the crab season are identified as management options with a likely success to reduce whale entanglement, but could burden fishers (Lebon and Kelly, 2019), specifically smaller vessels that continue to fish later into the crab season when more whales use the region. In addition, the effectiveness of any management strategy to balance whale conservation and fishing revenue depends on climatic conditions (Samhuri *et al.*, 2021). To minimize impacts to the industry, one possible approach could be to dynamically fit temporal closures around predicted areas of high whale densities, or only shorten crab fishing seasons during years when upwelling is predicted to be strong (based on negative PDO phase, elevated CUTI and early transition). Nonetheless, managers must consider multiple biological and human factors, such as delayed crab season start due to high levels of domoic acid in crab which can increase co-occurrence of fishing with whales (e.g., California 2016, Santora *et al.*, 2020; Feist *et al.*, 2021). Our models and findings provide scientific grounds for designing management strategies to reduce whale entanglement risk in Oregon. These strategies will require assessment of impacts to the commercial crab industry, management costs and complexity of implementation, and effectiveness to mitigate entanglement.

ACKNOWLEDGMENTS

Financial support from the NOAA Species Recovery Grant (#NA19NMF4720109), the Oregon Dungeness Crab Commission, the Oregon State University Marine Mammal Institute, the Oregon Department of Fish and Wildlife and Oregon Sea Grant. We thank Caren Braby, Michelle K. Jones, Leif K. Rasmuson, and Steven S. Rumrill for reviewing this manuscript and providing valuable feedback. We thank the anonymous reviewers.

AUTHORSHIP CONTRIBUTION STATEMENT

Craig Hayslip, Leigh G Torres, Solène Derville, Troy V Buell, Kelly C Corbett: Data acquisition and curation; Solène Derville: Formal analysis; Leigh G Torres, Troy V Buell, Kelly C Corbett,: Funding acquisition; Solène Derville, Troy V Buell: Methodology; Troy V Buell, Kelly C Corbett, Leigh G

Torres: Project administration; Solène Derville: Visualization; Solène Derville, Leigh G Torres, Troy V Buell, Kelly C Corbett: Writing.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any conflict of interest.

DATA AVAILABILITY STATEMENT

Datasets are made available via the Figshare Digital Repository: https://figshare.com/projects/OPAL_Overlap_Predictions_About_Large_whales/161137

REFERENCES

- Abrahms, B. *et al.* (2019) ‘Dynamic ensemble models to predict distributions and anthropogenic risk exposure for highly mobile species’, *Diversity and Distributions*, 25(8), pp. 1182–1193. Available at: <https://doi.org/10.1111/ddi.12940>.
- Becker, E.A. *et al.* (2017) ‘Habitat-Based Density Models for Three Cetacean Species off Southern California Illustrate Pronounced Seasonal Differences’, *Frontiers in Marine Science*, 4, p. 121. Available at: <https://doi.org/10.3389/fmars.2017.00121>.
- Becker, E.A. *et al.* (2018) ‘Predicting cetacean abundance and distribution in a changing climate’, *Diversity and Distributions*, 25(4), pp. 626–643. Available at: <https://doi.org/10.1111/ddi.12867>.
- Benoit-Bird, K.J., Waluk, C.M. and Ryan, J.P. (2019) ‘Forage Species Swarm in Response to Coastal Upwelling’, *Geophysical Research Letters*, 46(3), pp. 1537–1546. Available at: <https://doi.org/10.1029/2018GL081603>.
- Bograd, S.J. *et al.* (2009) ‘Phenology of coastal upwelling in the California Current’, *Geophysical Research Letters*, 36(1), pp. 1–5. Available at: <https://doi.org/10.1029/2008GL035933>.
- Bouchet, P.J. *et al.* (2019) *From here and now to there and then: Practical recommendations for the extrapolation of cetacean density surface models to novel conditions*. Technical report 2019-01 v1.0, Centre for Research into Ecological & Environmental Modelling (CREEM), University of St Andrews.
- Bouchet, P.J. *et al.* (2020) ‘dsmextra: Extrapolation assessment tools for density surface models’, *Methods in Ecology and Evolution*, 11(11), pp. 1464–1469. Available at: <https://doi.org/10.1111/2041-210X.13469>.
- Breen, P. *et al.* (2017) ‘Where is the risk? Integrating a spatial distribution model and a risk assessment to identify areas of cetacean interaction with fisheries in the northeast Atlantic’, *Ocean and Coastal Management*, 136, pp. 148–155. Available at: <https://doi.org/10.1016/j.ocecoaman.2016.12.001>.
- Carretta, J. V *et al.* (2021) *U.S. Pacific marine mammal stock assessments: 2021*. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-663. Available at: <https://doi.org/10.25923/246k-7589>.
- Carretta, J. V and Henry, A.G. (2022) ‘Risk Assessment of Whale Entanglement and Vessel Strike Injuries From Case Narratives and Classification Trees’, *Frontiers in Marine Science*, 9(June), pp. 1–11. Available at: <https://doi.org/10.3389/fmars.2022.863070>.
- Castelao, R.M. and Luo, H. (2018) ‘Upwelling jet separation in the California Current System’, *Scientific Reports*, 8, p. 16004. Available at: <https://doi.org/10.1038/s41598-018-34401-y>.
- CDFW (2019) *California Department of Fish and Wildlife Declaration of Fishery Closure due to Significant Risk of Marine Life Entanglement in the Dungeness Crab Commercial Fishery*. State of California, Natural Resources Agency, Sacramento, CA, USA.
- CDFW (2021) *Draft Conservation Plan for California’s Commercial Dungeness Crab Fishery December 2021*.
- Clapham, P.J. (2016) ‘Managing leviathan: Conservation challenges for the great whales in a post-whaling world’, *Oceanography*, 29(3), pp. 214–225. Available at: <https://doi.org/10.5670/oceanog.2016.70>.
- Davis, S. *et al.* (2017) *Oregon Dungeness crab fishery bioeconomic model: a fishery interactive simulator learning tool*. Prepared by OSU Coastal Oregon Marine Experiment Station and The Research Group, LLC for the Oregon Dungeness Crab Commission.
- Derville, S. *et al.* (2022) ‘Seasonal, Annual, and Decadal Distribution of Three Rorqual Whale Species Relative to Dynamic Ocean Conditions Off Oregon, USA’, *Frontiers in Marine Science*, 9, pp. 1–19. Available at: <https://doi.org/10.3389/fmars.2022.868566>.
- Feist, B.E. *et al.* (2021) ‘Footprints of fixed-gear fisheries in relation to rising whale entanglements on the U.S. West Coast’, *Fisheries Management and Ecology*, 28(3), pp. 283–294. Available at: <https://doi.org/10.1111/fme.12478>.

- Hazen, E.L. *et al.* (2017) ‘WhaleWatch : a dynamic management tool for predicting blue whale density in the California Current’, *Journal of Applied Ecology*, 54(5), pp. 1415–1428. Available at: <https://doi.org/10.1111/1365-2664.12820>.
- Henderson, E.E. *et al.* (2014) ‘Effects of fluctuations in sea-surface temperature on the occurrence of small cetaceans off Southern California’, *Fishery Bulletin*, 112(2–3), pp. 159–177. Available at: <https://doi.org/10.7755/FB.112.2-3.5>.
- Hickey, B.M. and Banas, N.S. (2008) ‘Why is the northern end of the California Current System so productive?’, *Oceanography*, 21(SPL.ISS. 4), pp. 90–107. Available at: <https://doi.org/10.5670/oceanog.2008.07>.
- van der Hoop, J., Corkeron, P. and Moore, M. (2017) ‘Entanglement is a costly life-history stage in large whales’, *Ecology and Evolution*, 7(1), pp. 92–106. Available at: <https://doi.org/10.1002/ece3.2615>.
- Jacox, M.G., Edwards, C.A., *et al.* (2018) ‘Coastal Upwelling Revisited: Ekman, Bakun, and Improved Upwelling Indices for the U.S. West Coast’, *Journal of Geophysical Research: Oceans*, 123(10), pp. 7332–7350. Available at: <https://doi.org/10.1029/2018JC014187>.
- Jacox, M.G., Alexander, M.A., *et al.* (2018) ‘Forcing of multiyear extreme ocean temperatures that impacted California Current living marine resources in 2016’, *Bulletin of the American Meteorological Society*, 99(1), pp. S1–S157.
- Lebon, K.M. and Kelly, R.P. (2019) ‘Evaluating alternatives to reduce whale entanglements in commercial Dungeness Crab fishing gear’, *Global Ecology and Conservation*, 18, p. e00608. Available at: <https://doi.org/10.1016/j.gecco.2019.e00608>.
- Mesgaran, M.B., Cousens, R.D. and Webber, B.L. (2014) ‘Here be dragons: A tool for quantifying novelty due to covariate range and correlation change when projecting species distribution models’, *Diversity and Distributions*, 20(10), pp. 1147–1159. Available at: <https://doi.org/10.1111/ddi.12209>.
- Meyer-Gutbrod, E.L. *et al.* (2021) ‘Ocean regime shift is driving collapse of the north atlantic right whale population’, *Oceanography*, 34(3), pp. 22–31. Available at: <https://doi.org/10.5670/oceanog.2021.308>.
- NOAA Fisheries (2022) *2021 West Coast Whale Entanglement Summary*.
- ODFW (2021) *Draft Conservation plan for reducing the impact of the Oregon ocean commercial Dungeness crab fishery on ESA-listed species off Oregon*. Available at: https://www.dfw.state.or.us/MRP/shellfish/commercial/crab/docs/2021/Public_CP_DRAFT_8.18.21.pdf (Accessed: 6 February 2023).
- ODFW (2022) *Oregon Dungeness crab fishery management plan*. Oregon Department of Fish and Wildlife Marine Resources Program. Available at: <http://www.dfw.state.or.us/MRP/>.
- Oestreich, W.K. *et al.* (2022) ‘Acoustic signature reveals blue whales tune life-history transitions to oceanographic conditions’, *Functional Ecology*, (January), pp. 1–14. Available at: <https://doi.org/10.1111/1365-2435.14013>.
- Oregon Secretary of State (2020) *The Oregon Bulletin for October 2020, DFW 131-2020*. Available at: <https://secure.sos.state.or.us/oard/displayBulletin.action?bulltnRsn=588>.
- Peterson, W.T. *et al.* (2017) ‘The pelagic ecosystem in the Northern California Current off Oregon during the 2014–2016 warm anomalies within the context of the past 20 years’, *Journal of Geophysical Research: Oceans*, 122, pp. 7267–7290. Available at: <https://doi.org/10.1002/2017JC012952>.
- R Core Team (2021) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rasmuson, L.K. (2013) *The biology, ecology and fishery of the dungeness crab, cancer magister*. 1st edn, *Advances in Marine Biology*. 1st edn. Elsevier Ltd. Available at: <https://doi.org/10.1016/B978-0-12-410498-3.00003-3>.
- Riekkola, L. *et al.* (2023) ‘Retrospective analysis of measures to reduce large whale entanglements in a lucrative commercial fishery’, *Biological Conservation*, 278. Available at: <https://doi.org/10.1016/j.biocon.2022.109880>.

- Robbins, J. (2012) *Scar-Based Inference Into Gulf of Maine Humpback Whale Entanglement : 2010*. Report to the Northeast Fisheries Science Center National Marine Fisheries Service, EA133F09CN0253 Item 0003AB, Task 3.
- Saez, L., Lawson, D. and Deangelis, M. (2021) 'Large whale entanglements off the U.S. West Coast, from 1982-2017', *NOAA Technical Memorandum NMFS-OPR-63*, (February), p. 64.
- Samhour, J.F. *et al.* (2021) 'Marine heatwave challenges solutions to human-wildlife conflict', *Proceedings of the Royal Society B: Biological Sciences*, 288, p. 20211607. Available at: <https://doi.org/10.1098/rspb.2021.1607>.
- Santora, J.A. *et al.* (2017) 'Impacts of ocean climate variability on biodiversity of pelagic forage species in an upwelling ecosystem', *Marine Ecology Progress Series*, 580, pp. 205–220. Available at: <https://doi.org/10.3354/meps12278>.
- Santora, J.A. *et al.* (2020) 'Habitat compression and ecosystem shifts as potential links between marine heatwave and record whale entanglements', *Nature Communications*, 11, p. 536. Available at: <https://doi.org/10.1038/s41467-019-14215-w>.
- Tackaberry, J. *et al.* (2022) 'Low Resighting Rate of Entangled Humpback Whales Within the California , Oregon , and Washington Region Based on Photo-Identification and Long-Term Life History Data', *Frontiers in Marine Science*, 8(January), pp. 1–13. Available at: <https://doi.org/10.3389/fmars.2021.779448>.
- Torres, L.G. *et al.* (2013) 'Scaling down the analysis of seabird-fishery interactions', *Marine Ecology Progress Series*, 473, pp. 275–289. Available at: <https://doi.org/10.3354/meps10071>.
- WDFW (2022) *Marine Life entanglement information and resources*.
- Williamson, C.E., Stoeckel, M.E. and Schoeneck, L.J. (1989) 'Predation Risk and the Structure of Freshwater Zooplankton Communities', *Oecologia*, 79, pp. 76–82. Available at: <https://doi.org/10.1007/BF00378242>.
- Wood, S.N. (2011) 'Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models', *Journal of the Royal Statistical Society (B)*, 73(1), pp. 3–36.

Tables

Table 1: Performance metrics of models of local whale abundance (M_{whale}), crab fishing effort (M_{crab}), and exposure (M_{exposure}) by zone, in relation to month, fishing year, PDO, SOI, and CUTI. Performance metrics include AIC (Akaike Information Criterion) and dev.exp (percent deviance explained). The preferred models (lowest AIC with a delta < 2) are highlighted for each zone.

Zone	Predictors	M_{whale}		M_{crab}		M_{exposure}		
		AIC	dev.exp	AIC	dev.exp	AIC	dev.exp	
north	<40 fathom	month	262	32	262	20	311	12
		month+fishing year	262	33	262	20	311	14
		month+PDO	259	37	262	23	301	25
		month+PDO+SOI	239	36	247	22	279	25
		month+CUTI	234	56	259	23	275	49
		month+CUTI+PDO	235	57	259	27	273	52
	>40 fathom	month	189	26	374	59	260	11
		month+fishing year	187	31	374	59	256	21
		month+PDO	178	45	374	60	244	33
		month+PDO+SOI	165	46	347	59	224	34
		month+CUTI	172	49	374	60	237	45
		month+CUTI+PDO	168	56	373	61	231	51
central	<40 fathom	month	256	55	262	29	304	24
		month+fishing year	252	59	262	29	301	30
		month+PDO	242	64	262	29	288	39
		month+PDO+SOI	225	63	247	27	267	39
		month+CUTI	221	73	257	33	265	58
		month+CUTI+PDO	207	78	257	33	247	67
	>40 fathom	month	233	51	307	55	293	17
		month+fishing year	231	53	307	56	288	24
		month+PDO	222	61	307	55	279	32
		month+PDO+SOI	207	60	288	54	260	30
		month+CUTI	210	67	308	57	265	48
		month+CUTI+PDO	200	73	308	57	254	56
south	<40 fathom	month	270	34	258	12	312	13
		month+fishing year	266	38	258	15	308	19
		month+PDO	266	39	258	12	305	22
		month+PDO+SOI	246	38	243	11	284	23
		month+CUTI	227	66	258	12	274	52
		month+CUTI+PDO	215	71	258	12	259	61
	>40 fathom	month	206	57	325	35	279	7
		month+fishing year	205	59	325	35	278	9
		month+PDO	197	65	325	35	273	20
		month+PDO+SOI	174	69	302	37	248	23
		month+CUTI	192	68	325	37	266	28
		month+CUTI+PDO	181	75	325	38	259	40

Figures




Figure 1: Average whale densities in number of whales predicted by 5 x 5 km grid cell (a), commercial Dungeness crab fishing effort (b), and exposure to entanglement risk (c) calculated over the study period (2011-2020, n = 83 monthly layers). Isobaths (73 m, 200 m, and 1,500 m deep) are shown with grey lines. The 40 fathom (73 m) isobath used in fishing regulations is shown in yellow. In panel c, white triangles indicate the estimated gear set location and year where humpback whales were confirmed to be entangled in commercial Oregon (OR) Dungeness crab fishing gear.




Figure 2: Fitted relationships between predicted local rorqual whale abundance (a; M_{whale}), crab fishing effort (b; M_{crab}) and exposure to entanglement risk (c; $M_{exposure}$) with month (top) and fishing year (bottom) predictors. GAMs were fitted to each response type in each of the six study zones represented on the map. Solid and dashed lines represent the marginal effect of month and fishing year by zone, and shaded areas represent approximate 95 % confidence intervals. The estimated degrees of freedom (edf) and the approximate smooth significance of predictors is indicated in each panel when the p-values were below a 0.05 threshold. Data points are represented in light shades. Y-axes are presented on a log scale.




Figure 3: Fitted relationships between the depth ($a_1 = M_{\text{whale.cmass.depth}}$ and $b_1 = M_{\text{crab.cmass.depth}}$) and the latitude ($a_2 = M_{\text{whale.cmass.lat}}$ and $b_2 = M_{\text{crab.cmass.lat}}$) of the center of mass of the predicted local rorqual whale abundance (a_1 and a_2) and the crab fishing effort (b_1 and b_2). GAMs were fitted to each response type in either the three latitudinal zones (panels a_1 and b_1) or the two depth ranges (panels a_2 and b_2) represented on the maps. Solid and dashed lines represent the marginal effect of month and fishing year by zone, and shaded areas represent approximate 95 % confidence intervals. The estimated degrees of freedom (edf) and the approximate smooth significance of predictors is indicated in each panel when the p-values were below a 0.05 threshold. Data points are represented in light shades.




Figure 4: Temporal evolution of the climatic and upwelling indices affecting ocean conditions in Oregon. a) Pacific Decadal Oscillation (PDO) monthly values from 2011 to 2021. Positive phases are shown in grey and negative phases in black. b) Long-term climatological mean of the cumulative sum of the Coastal Upwelling Transport Index (CUTI) with 5th and 95th percentile intervals. Daily values were averaged from 1988 to 2021, at latitude 43°N (south), 44°N and 45°N (central) and 46°N (north zone). Average timing by zone of the Spring Transition (STI), the peak upwelling (MAX) and the end of the upwelling season (END) are indicated at the top of the plot.




Figure 5: Fitted relationships between predicted local rorqual whale abundance (a; M_{whale}), crab fishing effort (b; M_{crab}) and exposure to entanglement risk (c; $M_{exposure}$) with the Pacific Decadal Oscillation (PDO; top) and the Coastal Upwelling Transport Index (CUTI; bottom) predictors. GAMs including month, PDO and CUTI as predictors were fitted to each response type in each of the six study zones represented on the map. The fitted relationships to month are not represented here. Solid and dashed lines represent the marginal effect of PDO and CUTI by zone, and shaded areas represent approximate 95 % confidence intervals. The estimated degrees of freedom (edf) and the approximate smooth significance of predictors is indicated in each panel when the p-values were below a 0.05 threshold. Data points are represented in light shades. Y-axes are presented on a log scale.