



HAL
open science

Profile Decomposition And Scattering For General Nonlinear Schrödinger Equations

Thomas Duyckaerts, Phan van Tin

► **To cite this version:**

Thomas Duyckaerts, Phan van Tin. Profile Decomposition And Scattering For General Nonlinear Schrödinger Equations. *Journal of Differential Equations*, 2024, 410, pp.113-170. 10.1016/j.jde.2024.07.003 . hal-04398984v2

HAL Id: hal-04398984

<https://hal.science/hal-04398984v2>

Submitted on 10 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROFILE DECOMPOSITION AND SCATTERING FOR GENERAL NONLINEAR SCHRÖDINGER EQUATIONS

THOMAS DUYNCKAERTS AND PHAN VAN TIN

ABSTRACT. We consider a Schrödinger equation with a nonlinearity which is a general perturbation of a power nonlinearity. We construct a profile decomposition adapted to this nonlinearity. We also prove global existence and scattering in a general defocusing setting, assuming that the critical Sobolev norm is bounded in the energy-supercritical case. This generalizes several previous works on double-power nonlinearities.

CONTENTS

1. Introduction	1
1.1. General setting	1
1.2. Well-posedness and profile decomposition	2
1.3. Global well-posedness	2
1.4. Scattering	3
1.5. Previous works	5
1.6. Outline	6
Acknowledgment	6
2. Preliminaries and Cauchy theory	6
2.1. Notations and function spaces	6
2.2. Preliminary nonlinear estimates	7
2.3. Local Lipschitz continuity of the nonlinearity	8
2.4. Cauchy and stability theory for general nonlinearities	13
3. Profile decomposition	16
3.1. Profiles in homogeneous Sobolev spaces	16
3.2. Profiles in non-homogeneous Sobolev spaces	18
3.3. Nonlinear profile decomposition	19
4. Global well-posedness	26
5. General rigidity result	28
6. Scattering	33
Appendix A. Equivalence of Sobolev norms	37
References	37

1. INTRODUCTION

1.1. **General setting.** This article concerns the nonlinear Schrödinger (NLS) equation

$$(1.1) \quad i\partial_t u + \Delta u = g(u),$$

in space dimension $d \geq 1$, where $g(u)$ is a L^2 -supercritical nonlinearity of the form

$$(1.2) \quad g(u) = \iota_0 |u|^{p_0} u + g_1(u), \quad \iota_0 \in \{\pm 1\}$$

and $g_1 : \mathbb{C} \rightarrow \mathbb{C}$ is a L^2 -supercritical lower-order term, i.e. g_1 is C^{k_0} and

$$\exists p_1, p_2, \frac{4}{d} < p_1 \leq p_2 < p_0, \quad \forall u \in \mathbb{C}, \forall k \in \llbracket 0, k_0 \rrbracket, \quad |D^k g_1(u)| \lesssim |u|^{p_2-k} + |u|^{p_1-k},$$

Date: February 10, 2024.

2020 Mathematics Subject Classification. 35Q55, 35A01, 35B40.

Key words and phrases. Nonlinear Schrödinger equations, profile decomposition, stability theory, scattering.

for some k_0 that we will be specified later.

The model case for g is a sum of $k + 1$ powers, $k \geq 1$

$$(1.3) \quad g(u) = \sum_{j=0}^k \iota_j |u|^{p_j} u, \quad \iota_j \in \mathbb{R} \setminus \{0\}, \quad \frac{4}{d} < p_k < \dots < p_1 < p_0.$$

With the regularity assumption:

$$(1.4) \quad \forall j \in \{0, \dots, k\}, p_j \text{ is an even integer, or } \lceil s_0 \rceil < p_j,$$

where $s_0 = \frac{d}{2} - \frac{2}{p_0}$ (see Subsection 2.1 for the notation $\lceil s_0 \rceil$ and other notations that will be used in this introduction). The case of a double-power, energy-subcritical (i.e. $s_0 \leq 1$) nonlinearity was studied in many work. Our goal is give a general setting for the study of (1.1) which includes also the supercritical case $s_0 > 1$ and more general lower order nonlinearity g_1 .

We are interested in the global well-posedness and scattering for solutions of (1.1). Neglecting the lower order term in g , we obtain the usual (NLS) equation with a single power nonlinearity

$$(1.5) \quad i\partial_t u + \Delta u = \iota_0 |u|^{p_0} u,$$

The equation (1.5) is invariant by scaling: if u is a solution of (1.5) and $\lambda > 0$, then $u_\lambda(t, x) = \lambda^{2/p_0} u(\lambda^2 t, \lambda x)$ is also a solution of (1.5). The critical Sobolev exponent s_0 for (1.5) is the unique s_0 such that $\|u\|_{\dot{H}^{s_0}} = \|u_\lambda(0)\|_{\dot{H}^{s_0}}$ for all $\lambda > 0$. The equation (1.5) is well-posed in \dot{H}^{s_0} (with additional technical conditions in high dimensions ensure a minimal regularity of the nonlinearity) see [10].

1.2. Well-posedness and profile decomposition. In Section 2 we prove that (1.1) is locally well-posed in the inhomogenous space H^s , for any $s \geq s_0$, assuming that $g \in C^{\lceil s \rceil + 1}$ and L^2 -supercritical (see Assumption A p.8 for the precise assumptions). We also develop a full stability/long time perturbation theory for (1.1). The existence and uniqueness of solutions yields for all $u_0 \in H^s$ a maximal interval of existence $I_{\max}(u_0) = (T_-(u_0), T_+(u_0))$. Assuming $g(u) = G'(|u|^2)u$ for some C^1 function G , with $G(0) = 0$ we also have conservation of the mass:

$$M(u(t)) = \int_{\mathbb{R}^d} |u(t, x)|^2 dx,$$

and, if $s \geq 1$, of the energy

$$E(u(t)) = \int_{\mathbb{R}^d} |\nabla u(t, x)|^2 dx + \int_{\mathbb{R}^d} G(|u(t, x)|^2) dx.$$

and the momentum

$$P(u(t)) = \mathcal{Im} \int_{\mathbb{R}^d} \nabla u(t, x) \bar{u}(t, x) dx$$

of a solution.

Our first main result is the construction of a profile decomposition adapted to bounded sequences of H^{s_0} solutions of equation (1.1), which builds up on the stability theory developed in Section 2. This amounts to expressing such a sequence as a sum of three distinct types of objects: a dispersive behaving as a solution of the linear Schrödinger equation, concentrating (nonlinear) profiles that are solutions of (1.5) rescaled with a scaling parameter going to 0, and nonconcentrating profiles, that are solutions of (1.1). We refer to Section 3, and in particular Subsection 3.3 and Theorem 3.15 for the detailed statements.

This profile decomposition is valid in the general setting described above, and generalizes various previous constructions on double power nonlinearities (see Subsection 1.5 below for references).

1.3. Global well-posedness. Solutions of (1.1) are not always global. Indeed, in the case of a double power nonlinearity, if $\iota_0 = -1$ (the higher-order nonlinearity is focusing), a standard convexity argument (see [56, 21]) shows that any solution with negative energy and finite variance blows up in finite time (at least in the case of a double-power nonlinearity, (1.3) with $k = 2$). More surprisingly, Merle, Raphaël, Rodnianski and Szeftel [35] have constructed solutions of the homogeneous equation (1.5) with a defocusing, energy-supercritical nonlinearity $s_0 > 1$, $\iota_0 = 1$ that blow up in finite time. It is known however in the defocusing case $\iota_0 = 1$, for many values of p_0 , that solution of (1.5) that remains bounded in the critical Sobolev space are global and scatter.

We will prove that this property implies that solutions of (1.1), with g of the form (1.2) satisfying the same boundedness condition are global. We will thus consider:

Property 1.1. Let $A_0 \in (0, \infty]$. For any solution u of (1.5) with initial data in \dot{H}^{s_0} , if

$$(1.6) \quad \limsup_{t \rightarrow T_+(u)} \|u(t)\|_{\dot{H}^{s_0}} < A_0.$$

Then $T_+(u) = +\infty$ and u scatters for positive times in \dot{H}^{s_0} , i.e. there exists $v_0 \in \dot{H}^{s_0}$ such that

$$\lim_{t \rightarrow \infty} \|u(t) - e^{it\Delta} v_0\|_{\dot{H}^{s_0}} = 0.$$

It is conjectured that Property 1.1 always holds for $A_0 = \infty$ in the defocusing case $\iota_0 = 1$. In the defocusing energy-critical case $d \geq 3$, $p_0 = \frac{4}{d-2}$, Property 1.1 is unconditional (the bound (1.6) is given by conservation of the energy), and was proved in [14], [46] and [51],

The study of Property 1.1 in the defocusing case for other critical exponents was initiated in [25] where it was proved when $d = 3$, $p_0 = 2$ (thus $s_0 = 1/2$), for radial solutions. It was later proved in many other cases: see [29], [37], [33], [16] for supercritical nonlinearities in dimension $d \geq 4$ (with technical restriction if $d \geq 7$), and [53], [41], [42], [19], [55] and [5] for several energy-subcritical nonlinearities.

In the focusing case $\iota_0 = -1$, Property 1.1 is only known to hold when A_0 is small, from the small data theory for equation (1.5). In the focusing energy critical case, $p_0 = \frac{4}{d-2}$, it follows from [24], [30] and [15] that Property 1.1 holds with $A_0 = \|\nabla W\|_{L^2}^2$ in dimension $d \geq 4$, and $d = 3$ in the radial case, where W is the ground state of equation (1.5). This is optimal, since the existence of W shows that Property 1.1 does not hold for $A_0 > \|\nabla W\|_{L^2}^2$.

When $\iota_0 = -1$ and $s_0 \in (0, 1)$, there exists standing wave solutions of (1.5), so that Property 1.1 does not hold for large A_0 . When $\iota_0 = -1$, $s_0 > 1$, travelling wave solutions in \dot{H}^{s_0} do not exist, and the validity of Property 1.1 for large A_0 is an open question. Let us also mention that the analogue of this property was proved for radial focusing nonlinear wave equation in the energy supercritical and subcritical settings (see e.g. [18] for supercritical p_0 in space dimension 3).

Our result on global well-posedness is as follows:

Theorem 1.2. *Let ι_0, s_0, g such that Assumption B holds, and such that Property 1.1 is valid for some $A_0 \in (0, \infty]$. Let u be a solution of (2.1), with initial data in H^{s_0} , such that (1.6) holds. Then $T_+(u) = +\infty$.*

We refer to Definition 2.7 and Section 3, p. 19 for the details of Assumption B. Let us mention that a multi-power non-linearity as in (1.3) with the additional technical assumption (1.4) satisfies this assumption.

If Property 1.1 holds only for radial functions, then the conclusion of Theorem 1.2 is also valid when restricted to radial solutions of (1.1).

Theorem 1.2 is new in the energy supercritical case $s_0 > 1$. In the energy-subcritical and energy-critical cases $0 < s_0 \leq 1$, it was proved in [50] for a double power non-linearity with the stronger assumptions $\iota_0 = 1$, $u_0 \in H^1$, without assuming Property 1.1.

The proof of Theorem 1.2 uses the profile decomposition mentioned above. The Property 1.1 is used to deal with the concentrating profiles.

In the defocusing energy-critical case, by conservation of the energy and the scattering result for the energy-critical Schrödinger defocusing equation, the global well-posedness is unconditional:

Corollary 1.3. *Assume $\iota_0 = 1$. Let $d \in \{3, 4, 5\}$, and g such that Assumption B holds with $p_0 = \frac{4}{d-2}$. Let u be a solution of (2.1), with initial data in H^1 . Then u is global.*

Corollary 1.3 was proved in the case of a double-power nonlinearity in [57] ($d = 3$) and in [50] (for general $d \geq 4$). Corollary 1.3 generalizes these results to more general perturbations of the energy-critical nonlinearity, in low dimension. The restriction on the dimension is due to the regularity assumption $g \in C^2$ in Assumption B. This restriction could be weakened using a refined well-posedness/stability theory as in [7].

1.4. Scattering. Our next goal is to give sufficient conditions for scattering of solutions of (1.1). We recall that a solution of (1.1) with initial data in H^{s_0} is said to scatter (in H^{s_0} , forward in

time) when $T_+(u) = +\infty$ and there exists $v_0 \in H^{s_0}$ such that

$$\lim_{t \rightarrow \infty} \|e^{it\Delta} v_0 - u(t)\|_{H^{s_0}} = 0.$$

We will prove scattering for a general defocusing nonlinearity defined as follows:

Definition 1.4. The nonlinearity g is defocusing when it is of the form $g(u) = G'(|u|^2)u$ for some C^1 function G such that for almost all $a > 0$, $aG'(a) - G(a) > 0$.

Note that any power nonlinearity with a positive coefficient is defocusing in the sense of Definition (1.4). For a multi-power nonlinearity g as in (1.3) the assumption that g is defocusing is equivalent to

$$(1.7) \quad \forall s > 0, \quad \sum_{j=0}^k \iota_j \frac{p_j}{p_j + 2} s^{\frac{p_j}{2} - 1} > 0.$$

Note that (1.7) holds when all the ι_j are positive. Also, (1.7) implies $\iota_0 > 0$, $\iota_k > 0$.

For our scattering results, we distinguish between the energy-supercritical case and the energy-subcritical case:

Theorem 1.5. *Let $d \geq 3$ and assume $s_0 > 1$. Let g be a defocusing nonlinearity that satisfies Assumption B. Assume that Property 1.1 holds for some $A_0 \in (0, \infty]$. Let u be a solution of (1.1) with initial data $u_0 \in H^{s_0}$ and that satisfies (1.6). Then u is global and scatter forward in time.*

When $s_0 \leq 1$, we must further assume that the initial data has finite energy.

Theorem 1.6. *Assume $0 < s_0 \leq 1$. Let g be a defocusing nonlinearity that satisfies Assumption B. Let u be a solution of (1.1) with initial data $u_0 \in H^1$. Then u is global and scatter in both time directions.*

Note that the assumption that g is defocusing together with the fact that $G(a)/a$ goes to 0 as a goes to 0 (which is a consequence of Assumption B) implies that $G \geq 0$. Thus the assumptions of Theorem 1.6 and the conservation of mass and energy imply that any H^1 solution of u is bounded in H^1 .

Theorems 1.5 and Theorem 1.6 show that scattering holds for a multi-power nonlinearity satisfying (1.3), (1.4) and (1.7).

In the case of a double-power nonlinearity the condition (1.7) is equivalent to $\iota_0 > 0$, $\iota_1 > 0$. In this case, Theorem 1.5 is new. Theorem 1.6 is proved in [50, Theorem 1.3] and [57]).

For a double-power nonlinearity with $(\iota_0 > 0, \iota_1 < 0)$ or $(\iota_0 < 1, 0 < s_0 < s_1 < 1)$, there are solitary wave solutions and thus it is impossible to prove an analog of Theorems 1.5 and 1.6. This is an open question for other double power nonlinearities.

When only the main order term of the nonlinearity is defocusing, i.e. when $\iota_0 > 0$, scattering holds for initial data with small mass. We again distinguish between $s_0 > 1$ and $s_0 \in (0, 1]$

Theorem 1.7. *Let $d \geq 3$ and assume $s_0 > 1$. Let g be a nonlinearity that satisfies Assumption B with $\iota_0 > 0$. Assume that Property 1.1 holds for some $A_0 \in (0, \infty]$. There exists $m_c > 0$ such that any solution of (1.1) with initial data in H^{s_0} such that $M(u_0) < m_c$ and that satisfies (1.6) is global and scatters forward in time.*

Theorem 1.8. *Let g be nonlinearity that satisfies Assumption B with $\iota_0 > 0$ and $s_0 \in (0, 1]$. There exists $m_c > 0$ such that any solution of (1.1) with initial data in H^1 and such that $M(u_0) < m_c$ is global and scatters in both time directions.*

Remark 1.9. In Theorem 1.5 and Theorem 1.7, if the analog of (1.6) as t goes to $T_-(u)$ is valid then the conclusions hold backward in time.

Theorem 1.8 is new, even for double power nonlinearities. Theorem 1.8 generalizes [50, Theorem 1.3] which concerns a double power nonlinearity with $\iota_0 > 0$ and $\iota_1 < 0$. In particular the case where $d = 3$, $p_0 = 4$ (thus $s_0 = 1$), $p_1 = 2$ scattering was proved to hold for a larger set of initial data in [28], [27]. In a subsequent work, we will use the material of this article, together with [31] to obtain an improvement of Theorems 1.7 and 1.8 in the same spirit.

The proofs of Theorem 1.5 and Theorem 1.7 follow the by now classical rigidity-compactness roadmap (see [24]), using the profile decomposition constructed in Section 3. This provides, in a contradiction argument, a global critical solution u_c of (1.1) such that there exists $x(t)$ such that

$$K = \{u_c(t, \cdot + x(t)), t \in \mathbb{R}\}$$

has compact closure in H^{s_0} .

To exclude this critical element and obtain a contradiction, we use the virial identity

$$(1.8) \quad \frac{d}{dt} \mathcal{I}m \int x \cdot \nabla u \bar{u} = 2\Phi(u),$$

where

$$\Phi(u) = \int |\nabla u|^2 + \iota_1 \frac{dp_1}{2(p_1 + 2)} |u|^{p_1+2} + \iota_0 \frac{dp_0}{2(p_0 + 2)} |u|^{p_0+2},$$

and the center of mass identity:

$$(1.9) \quad \frac{d}{dt} \int x |u(t)|^2 = 2P(u).$$

which are valid for solutions of u with enough decay at infinity.

Using a localized version of (1.9), one can prove:

$$(1.10) \quad \lim_{t \rightarrow +\infty} \frac{1}{t} |x(t) - X(t)| = 0 \text{ where } X(t) = 2 \frac{P(u_c)}{M(u_c)} t.$$

When the momentum of u_c is zero, this allows to control the growth of $x(t)$. A localized version of (1.8) using the relative compactness K , gives a contradiction if $\inf_{t \in \mathbb{R}} \Phi(u_c(t)) - \frac{|P(u_c)|^2}{M(u_c)} > 0$. In the defocusing/defocusing case (as in Theorem 1.5), this property is true whenever u_c is not identically zero. In the defocusing/focusing case, we show, using the quite complete study of the elliptic problem in [31], and some of the ideas in [28], that Φ is positive in the region described in the assumptions of Theorem 1.7, yielding again the desired contradiction.

One must adapt this argument when the momentum of u_c is not zero. The standard strategy, going back to [17] is to use the Galilean transformation to reduce to the case to a critical solution with zero momentum. However the effect of the Galilean transform on the Sobolev norm \dot{H}^{s_0} of the solution is not explicit, and thus the strategy breaks down in the case $s_0 > 1$, where our proof relies on an induction-type argument on this norm. To tackle this difficulty, we observe that (1.8), (1.9) and the conservation of momentum imply:

$$\frac{d}{dt} \mathcal{I}m \int (x - X(t)) \nabla u \bar{u} = 2\Phi(u(t)) - \frac{2|P(u)|^2}{M(u)},$$

which we localize with a time-dependant localization around $X(t)$. This gives again a contradiction using an improved Cauchy-Schwarz inequality going back to [3] to prove that $\Phi(u) - \frac{|P(u)|^2}{M(u)}$ is still positive for the solutions that we consider.

1.5. Previous works. To our knowledge, the profile decomposition for a general nonlinearity of the form (1.2) was not considered before.

Dynamics of nonlinear Schrödinger equations with general nonlinearities were treated in very few works. In [32], [49], the authors prove that any radial, bounded, global solutions of a nonlinear Schrödinger equation with a general energy-subcritical or energy-critical nonlinearity is asymptotically the sum of a radiation term, solution of the linear Schrödinger equation, and a localized term. We refer to [48] for results in a nonradial setting.

Let us mention a few works on NLS equation with a double-power nonlinearity. The study of this type of equation was initiated in [57], in dimension 3, where the author investigated the global well-posedness, scattering and blow-up phenomena in the case $p_0 = 4$. This includes in particular a scattering result for small mass, in the spirit of Theorem 1.7. Similar results were obtained in [50], in general dimension d in the energy-critical and subcritical setting $s_0 \leq 1$.

The problem with $p_1 = \frac{4}{d}$, $\iota_1 = -1$, $p_0 < \frac{4}{d-2}$ was considered in [11], [43], where the author investigated scattering below or at the mass of the ground-state for the mass-critical homogeneous equation. See also [8] which considers the case $(p_0, p_1, \iota_0, \iota_1) = (4, 2, 1, -1)$ in space dimensions 1, 2 and 3.

The problem with a focusing dominant nonlinearity $\iota_0 = -1$ was considered in many works, always in the energy-critical or subcritical cases. Let us mention in particular [1] where a nine-set theorem in the spirit of [44], [45] was proved. We also refer to [38], [52],[23], [54], [12], [39], [34] for scattering result or scattering blow-up dichotomy in the case $\iota_0 = -1$.

Let us mention that in several of the preceding works, the authors construct and use a profile decomposition adapted to NLS equation with a particular double-power nonlinearity. The profile decomposition in Section 3 generalises these profiles decompositions to the large class of nonlinearities described above.

1.6. Outline. We conclude this introduction by an outline of the article. Section 2 is devoted to some preliminaries, well-posedness and perturbation theory for the NLS equation (1.1) with a general nonlinearity g . In Section 3, we construct a profile decomposition for sequences that are bounded in H^{s_0} adapted to (1.1) with a nonlinearity of the form (1.2). This profile decomposition is based on the linear profile decomposition of Shao [47] (which relies ultimately on the result by Merle-Vega [36]) and the long-time perturbation result proved in Subsection 2.4. In Section 4, we use this profile decomposition to prove our global well-posedness result, Theorem 1.2. In Section 5 we prove a general rigidity result for solutions of (1.1) with a relatively compact trajectory in H^{s_0} . In Section 6, we prove our scattering result Theorems 1.5 and 1.7, using the material of the preceding sections.

ACKNOWLEDGMENT

The second author is supported by Post-doc fellowship from Labex MME-DII: SAIC/2022 No 10078.

2. PRELIMINARIES AND CAUCHY THEORY

This section is concerned with the Cauchy and stability theory for the equation:

$$(2.1) \quad i\partial_t u + \Delta u = g(u).$$

for a general L^2 supercritical nonlinearity $g(u)$ in space dimension $d \geq 1$. Our assumptions on g will of course include the case of double power nonlinearities that we are interested in. We start by introducing some notations and functions spaces (see §2.1) and recalling some nonlinear estimates (see §2.2). In §2.3 we prove estimates on the nonlinearity $g(u)$ that are crucial for the well-posedness theory. In §2.4 we prove our main results.

2.1. Notations and function spaces. For $s \in \mathbb{R}$, $[s]$ is the smallest integer number larger or equal s , and $\lfloor s \rfloor$ is the largest integer smaller or equal s (the integer part of s). If j and k are integers with $j \leq k$, we denote by $\llbracket j, k \rrbracket = \{j, j+1, \dots, k-1, k\}$.

When A and B are two positive quantities depending on some parameters, we denote $A \lesssim B$ when there is a constant $C > 0$ such that $A \leq CB$ and $A \approx B$ when $A \lesssim B$ and $B \lesssim A$.

For each $q \geq 1$, we define q' such that

$$\frac{1}{q} + \frac{1}{q'} = 1.$$

If X is a vector space, $(u, v) \in X^2$, we will make a small abuse of notation, denoting $\|(u, v)\|_X = \|u\|_X + \|v\|_X$.

We fix $d \geq 1$. If m is a complex valued function on \mathbb{R}^d , we define by $m(\nabla)$ the Fourier multiplier with symbol $m(\xi)$, i.e. $\widehat{m(\nabla)u} = m(\xi)\hat{u}(\xi)$, where \hat{u} is the Fourier transform of u . For $s \geq 0$, $p \geq 1$, we define

$$\|u\|_{H^{s,p}} = \|\langle \nabla \rangle^s u\|_{L^p}, \quad \|u\|_{\dot{H}^{s,p}} = \|\ |\nabla|^s u \|_{L^p}$$

where $\langle \xi \rangle = (1 + |\xi|^2)^{s/2}$. It follows from Mihlin multiplier theorem that

$$(2.2) \quad \|u\|_{H^{s,p}} \approx \|u\|_{L^p} + \|\ |\nabla|^s u \|_{L^p}, \quad 1 < p < \infty.$$

(see the proof in the appendix). For a multi-index $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)$, denote

$$D^\alpha = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_d}^{\alpha_d}, \quad |\alpha| = \sum_{i=1}^d |\alpha_i|.$$

For $s > 0$ and $1 < p < \infty$ and $v = s - [s]$, we have (see [2, Lemma 3.2]),

$$(2.3) \quad \sum_{|\alpha|=[s]} \|D^\alpha f\|_{\dot{H}^{v,p}} \approx \|f\|_{\dot{H}^{s,p}}.$$

We recall that a pair (q, r) is *Strichartz-admissible* for the Schrödinger equation when $2 \leq q \leq \infty$, $2 \leq r \leq \infty$, $\frac{2}{q} + \frac{d}{r} = \frac{d}{2}$, and $(q, r, d) \neq (2, \infty, 2)$. We recall that if (q, r) and (a, b) are Strichartz admissible, we have,

$$\|u\|_{L_t^q L_x^r} \lesssim \|u_0\|_{L^2} + \|f\|_{L_t^{q'} L_x^{b'}}$$

for any solution of the Schrödinger equation $i\partial_t u + \Delta u = f$ with initial data u_0 . We denote:

$$S^0(I) = \begin{cases} L^\infty(I, L^2(\mathbb{R}^d)) \cap L^2\left(I, L^{\frac{2d}{d-2}}(\mathbb{R}^d)\right) & \text{if } d \geq 3 \\ L^\infty(I, L^2(\mathbb{R}^2)) \cap L^{q_2}(I, L^{r_2}(\mathbb{R}^2)) & \text{if } d = 2 \\ L^\infty(I, L^2(\mathbb{R})) \cap L^4(I, L^\infty(\mathbb{R})) & \text{if } d = 1, \end{cases}$$

where when $d = 2$, (q_2, r_2) is an admissible pair with $q_2 > 2$ close to 2, and

$$N^0(I) = \begin{cases} L^1(I, L^2(\mathbb{R}^d)) + L^2\left(I, L^{\frac{2d}{d+2}}(\mathbb{R}^d)\right) & \text{if } d \geq 3 \\ L^1(I, L^2(\mathbb{R}^2)) + L^{q_2'}(I, L^{r_2'}(\mathbb{R}^2)) & \text{if } d = 2 \\ L^1(I, L^2(\mathbb{R})) + L^{4/3}(I, L^1(\mathbb{R})) & \text{if } d = 1, \end{cases}$$

Note that the norm of $S^0(I)$ is equivalent to the supremum of all $L^q(I, L^r)$ norms, and that the norm of $N^0(I)$ is smaller than the infimum of all $L^{q'}(I, L^{r'})$ norms, where in each case, (q, r) is taken over all Strichartz admissible pairs (with $q \geq q_2$ if $d = 2$). We also define the following Strichartz spaces, and dual Strichartz spaces:

$$W^0(I) = L^{\frac{2(2+d)}{d}}(I \times \mathbb{R}^d) \quad Z^0(I) = (W^0(I))' = L^{\frac{2(d+2)}{d+4}}(I \times \mathbb{R}^d),$$

so that we have $S^0(I) \subset W^0(I)$ and $Z^0(I) \subset N^0(I)$ (with continuous embedding). We denote

$$X_p(I) = L^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d).$$

For $s \geq 0$, we denote

$$\|u\|_{S^s(I)} = \|\langle \nabla \rangle^s u\|_{S^0(I)}, \quad \|u\|_{\dot{S}^s(I)} = \|\langle \nabla \rangle^s u\|_{S^0(I)}$$

and define similarly $W^s(I)$, $\dot{W}^s(I)$, $N^s(I)$, $\dot{N}^s(I)$, $Z^s(I)$ and $\dot{Z}^s(I)$.

2.2. Preliminary nonlinear estimates.

Lemma 2.1. (*Product Rule 1*) *Let $s \geq 0$ and $1 < r, r_1, r_0, q_1, q_0 < \infty$ such that $\frac{1}{r} = \frac{1}{r_i} + \frac{1}{q_i}$, for $i = 0, 1$. Then,*

$$(2.4) \quad \|\langle \nabla \rangle^s (f\varphi)\|_{L_x^r} \lesssim \|f\|_{L_x^{r_1}} \|\langle \nabla \rangle^s \varphi\|_{L_x^{q_1}} + \|\langle \nabla \rangle^s f\|_{L_x^{r_0}} \|\varphi\|_{L_x^{q_0}}$$

$$(2.5) \quad \|\langle \nabla \rangle^s (f\varphi)\|_{L_x^r} \lesssim \|f\|_{L_x^{r_1}} \|\langle \nabla \rangle^s \varphi\|_{L_x^{q_1}} + \|\langle \nabla \rangle^s f\|_{L_x^{r_0}} \|\varphi\|_{L_x^{q_0}}.$$

Proof. See e.g [13, Proposition 3.3]. For (2.4), see e.g [2, Lemma 2.2] for the statement and [13] for proof in dimension 1 and $s \in (0, 1)$.

By (2.2), (2.4) and Hölder inequality, we have

$$\begin{aligned} \|\langle \nabla \rangle^s (f\varphi)\|_{L^r} &\approx \|f\varphi\|_{L^r} + \|\langle \nabla \rangle^s (f\varphi)\|_{L^r} \lesssim \|f\|_{L^{r_1}} \|\varphi\|_{L^{q_1}} + \|f\|_{L^{r_1}} \|\langle \nabla \rangle^s \varphi\|_{L^{q_1}} + \|\langle \nabla \rangle^s f\|_{L^{r_0}} \|\varphi\|_{L^{q_0}} \\ &\lesssim \|f\|_{r_1} \|\langle \nabla \rangle^s \varphi\|_{q_1} + \|\langle \nabla \rangle^s f\|_{r_0} \|\varphi\|_{q_0}. \end{aligned}$$

Hence (2.5). \square

Lemma 2.2 (*Product rule 2*). (*see e.g [2, Corollary 2.3]*). *Let $s \geq 0$, $n \in \mathbb{N}$ ($n \geq 1$), and, for $i, j \in \llbracket 1, n \rrbracket$, $1 < r, r_k^i < \infty$, such that for all $k \in \llbracket 1, n \rrbracket$, $\frac{1}{r} = \sum_{i=1}^n \frac{1}{r_k^i}$. Then*

$$\left\| \prod_{i=1}^n f_i \right\|_{\dot{H}^{s,r}} \lesssim \sum_{k=1}^n \left(\|f_k\|_{\dot{H}^{s,r_k^k}} \prod_{i \neq k} \|f_i\|_{L^{r_k^i}} \right).$$

Lemma 2.3 (Fractional chain rule). (see e.g [2, Lemma 2.4]). Let $G \in C^1(\mathbb{C}, \mathbb{C})$, $s \in (0, 1)$, $1 < r, r_2 < \infty$, and $1 < r_1 \leq \infty$ satisfying $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$,

$$\|G(u)\|_{\dot{H}^{s,r}} \lesssim \|G'(u)\|_{L^{r_1}} \|u\|_{\dot{H}^{s,r_2}}.$$

Lemma 2.4 (Gagliardo-Nirenberg inequality [6]). Let $s_1 \leq s_2$, $p_2 > 1$, $s = \theta s_1 + (1 - \theta)s_2$, $\frac{1}{p} = \frac{\theta}{p_1} + \frac{1-\theta}{p_2}$. Then

$$\|u\|_{H^{s,p}} \lesssim \|u\|_{H^{s_1,p_1}}^\theta \|u\|_{H^{s_2,p_2}}^{1-\theta}.$$

Lemma 2.5 (Homogeneous Gagliardo-Nirenberg inequality). Let $s_1 \leq s_2$, $p_2 > 1$, $s = \theta s_1 + (1 - \theta)s_2$, $\frac{1}{p} = \frac{\theta}{p_1} + \frac{1-\theta}{p_2}$. Then

$$\|u\|_{\dot{H}^{s,p}} \lesssim \|u\|_{\dot{H}^{s_1,p_1}}^\theta \|u\|_{\dot{H}^{s_2,p_2}}^{1-\theta}.$$

Proof. Let $\varphi \in H^s$ Define $\varphi^\lambda(x) = \varphi(\lambda x)$. Applying Lemma 2.4 for φ^λ , we have

$$\begin{aligned} & \left(\lambda^{-d} \|\varphi\|_{L^p}^p + \lambda^{ps-d} \|\varphi\|_{\dot{H}^{s,p}}^p \right)^{\frac{1}{p}} \\ & \lesssim \left(\lambda^{-d} \|\varphi\|_{L^{p_1}}^{p_1} + \lambda^{p_1 s_1 - d} \|\varphi\|_{\dot{H}^{s_1,p_1}}^{p_1} \right)^{\frac{1}{p_1}} \left(\lambda^{-d} \|\varphi\|_{L^{p_2}}^{p_2} + \lambda^{p_2 s_2 - d} \|\varphi\|_{\dot{H}^{s_2,p_2}}^{p_2} \right)^{\frac{1}{p_2}}. \end{aligned}$$

By dividing both sides by $\lambda^{s-\frac{d}{p}} = \lambda^{\theta s_1 - \frac{d\theta}{p_1}} \lambda^{(1-\theta)s_2 - \frac{d(1-\theta)}{p_2}}$, we have

$$\left(\lambda^{-ps} \|\varphi\|_{L^p}^p + \|\varphi\|_{\dot{H}^{s,p}}^p \right)^{\frac{1}{p}} \lesssim \left(\lambda^{-p_1 s_1} \|\varphi\|_{L^{p_1}}^{p_1} + \|\varphi\|_{\dot{H}^{s_1,p_1}}^{p_1} \right)^{\frac{1}{p_1}} \left(\lambda^{-p_2 s_2} \|\varphi\|_{L^{p_2}}^{p_2} + \|\varphi\|_{\dot{H}^{s_2,p_2}}^{p_2} \right)^{\frac{1}{p_2}}.$$

Let $\lambda \rightarrow +\infty$, we obtain the desired result. \square

Lemma 2.6 (Leibniz rule). Let $f \in C^k(\mathbb{C}, \mathbb{C})$ and $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d) \in \mathbb{N}^d$ such that $|\alpha| \leq k$. Then $D^\alpha(f(u))$ is a linear combinations of terms of the form

$$(\partial_z^{h_1} \partial_{\bar{z}}^{h_2} f)(u) \prod_{k=1}^{h_1} D^{\beta_k} u \prod_{k=1}^{h_2} D^{\gamma_k} \bar{u},$$

where $\sum_{k=1}^{h_1} \beta_k + \sum_{k=1}^{h_2} \gamma_k = \alpha$, $1 \leq h_1 + h_2 \leq |\alpha|$, $|\beta_k| \geq 1$, $|\gamma_k| \geq 1$ for all k .

Proof. The proof is by induction, using the formula:

$$\partial_{x_i} f(u) = \partial_z f(u) \partial_{x_i} u + \partial_{\bar{z}} f(u) \overline{\partial_{x_i} u}.$$

\square

2.3. Local Lipschitz continuity of the nonlinearity. In this subsection we consider the following general classes of nonlinearities:

Definition 2.7. Let $0 < p_1 \leq p_0$, $s \geq 0$ be real numbers. We denote by $\mathcal{N}(s, p_0, p_1)$ the vector space of functions $g \in C^{\lceil s \rceil + 1}(\mathbb{C}, \mathbb{C})$ such that

$$(2.6) \quad \exists C > 0, \forall k \in \llbracket 0, \lceil s \rceil + 1 \rrbracket, \forall z \in \mathbb{C}, \quad |g^{(k)}(z)| \leq C (|z|^{p_0+1-k} + |z|^{p_1+1-k}).$$

In the definition $|g^{(k)}(z)|$ denotes the supremum of all the derivatives (in z, \bar{z}) of order k of g .

Assumption A. $d \geq 1$, $g \in \mathcal{N}(s, p_0, p_1)$ with $0 \leq s, \frac{4}{d} \leq p_1 \leq p_0$, $1 < p_1$ and

$$\lceil s \rceil \leq p_0 \text{ or } g \text{ is a polynomial in } u, \bar{u}.$$

We recall the notation $X_p(I) = L^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)$. We will prove:

Proposition 2.8. If Assumption A is satisfied, we let

$$(2.7) \quad X(I) = X_{p_0}(I) \cap X_{p_1}(I) = L_{t,x}^{p_0(d+2)/2}(I \times \mathbb{R}^d) \cap L_{t,x}^{p_1(d+2)/2}(I \times \mathbb{R}^d).$$

Then

$$(2.8) \quad \|g(u) - g(v)\|_{N^s(I)} \lesssim \left(\|(u, v)\|_{X(I)}^{p_1-1} + \|(u, v)\|_{X(I)}^{p_0-1} \right) \times \left[\|u - v\|_{\dot{W}^s(I)} \|(u, v)\|_{X(I)} + \|u - v\|_{X(I)} \|(u, v)\|_{W^s(I)} \right],$$

In particular

$$(2.9) \quad \|g(u)\|_{N^s(I)} \lesssim \|u\|_{W^s(I)} \left(\|u\|_{X(I)}^{p_0} + \|u\|_{X(I)}^{p_1} \right).$$

Let us insist on the important fact that the first norm of $u - v$ in the second line of (2.8) is the norm in the *homogeneous* space \dot{W}^s .

We start with a few lemmas.

Lemma 2.9. *Let $p \geq 1$ be an integer, and $g(u)$ be a homogeneous polynomial of degree $p+1$ in u , \bar{u} . Let $s \geq 0$, $u, v \in S^s(I)$. Then*

$$(2.10) \quad \begin{aligned} \|\nabla|^s(g(u) - g(v))\|_{N^0(I)} &\lesssim \left(\|u\|_{X_p(I)}^{p-1} + \|v\|_{X_p(I)}^{p-1} \right) \\ &\times \left[\|\nabla|^s(u - v)\|_{W^0(I)} (\|u\|_{X_p(I)} + \|v\|_{X_p(I)}) + \|u - v\|_{X_p(I)} (\|\nabla|^s u\|_{W^0(I)} + \|\nabla|^s v\|_{W^0(I)}) \right], \end{aligned}$$

and

$$(2.11) \quad \|g(u) - g(v)\|_{N^0(I)} \lesssim \|u - v\|_{X_p(I)} \|(u, v)\|_{X_p(I)}^{p-1} \|(u, v)\|_{W^0(I)}.$$

Proof. It is sufficient to prove (2.10) and (2.11) for $g(u) = u^{p+1-j} \bar{u}^j$, where $j \in \llbracket 0, p+1 \rrbracket$. We have

$$(2.12) \quad g(u) - g(v) = (u^{p+1-j} - v^{p+1-j}) \bar{u}^j + v^{p+1-j} (\bar{u}^j - \bar{v}^j).$$

We treat the contribution of the first term in the right-hand side of (2.12). The contribution of the second term is similar. We have

$$(2.13) \quad (u^{p+1-j} - v^{p+1-j}) \bar{u}^j = (u - v) \bar{u}^j \sum_{k=0}^{p-j} u^{p-j-k} v^k.$$

We work on the interval I for all norms. We note the following relations between the exponents defining the W^0 , $Z^0 = (W^0)'$ and X_p norms:

$$(2.14) \quad \frac{d+4}{2(d+2)} = \frac{d}{2(d+2)} + p \times \frac{2}{p(d+2)}.$$

By Hölder inequality, the definitions of X_p and Z_0 and the fact that Z_0 is continuously embedded into \dot{N}^s , we obtain (2.11).

We next prove (2.10). Using (2.13), Lemma 2.1 and the definitions of X_p and Z^0 , we have

$$\begin{aligned} \|(u^{p+1-j} - v^{p+1-j}) \bar{u}^j\|_{\dot{N}^s} &\lesssim \left\| |\nabla|^s \left((u^{p+1-j} - v^{p+1-j}) \bar{u}^j \right) \right\|_{Z^0} \\ &\lesssim \sum_{k=0}^{p-j} \|\nabla|^s(u - v)\|_{W^0} \|u\|_{X_p}^{p-k} \|v\|_{X_p}^k \\ &+ \sum_{k=0}^{p-j} \|u - v\|_{X_p} \left(\|\nabla|^s u\|_{W^0} \|u\|_{X_p}^{p-k-1} \|v\|_{X_p}^k + \|\nabla|^s v\|_{W^0} \|v\|_{X_p}^{k-1} \|u\|_{X_p}^{p-k} \right) \\ &\lesssim \|u - v\|_{W^s} (\|u\|_{X_p}^p + \|v\|_{X_p}^p) + \|u - v\|_{X_p} (\|u\|_{W^s} + \|v\|_{W^s}) (\|u\|_{X_p}^{p-1} + \|v\|_{X_p}^{p-1}). \end{aligned}$$

Combining with the same bound for the second term in (2.12), we obtain (2.10), concluding the proof. \square

Lemma 2.10. *Let s, p be real numbers such that $0 < s$, $\lceil s \rceil \leq p$, $1 < p$ and $\frac{4}{d} \leq p$. Let $g \in C^{\lceil s \rceil + 1}(\mathbb{C}, \mathbb{C})$ such that*

$$(2.15) \quad \forall k \in \llbracket 0, \lceil s \rceil + 1 \rrbracket, \quad \left| g^{(k)}(u) \right| \leq C |u|^{p+1-k}.$$

Then (2.10) and (2.11) hold.

Proof. We have

$$g(u) - g(v) = (u - v) \int_0^1 g_z(v + \theta(u - v)) d\theta + \overline{u - v} \int_0^1 g_{\bar{z}}(v + \theta(u - v)) d\theta,$$

where $g_z = \frac{\partial g}{\partial z}$, $g_{\bar{z}} = \frac{\partial g}{\partial \bar{z}}$. Since $\left| \frac{\partial g}{\partial z} \right|, \left| \frac{\partial g}{\partial \bar{z}} \right| \leq C|z|^p$, we have

$$|g(u) - g(v)| \leq C|u - v|(|u|^p + |v|^p).$$

Thus, by Hölder inequality and (2.14),

$$(2.16) \quad \|g(u) - g(v)\|_{N^0} \lesssim \|g(u) - g(v)\|_{Z^0} \lesssim \|u - v\|_{X_p} \|(u, v)\|_{X_p}^{p-1} \|(u, v)\|_{W^0}.$$

This yields (2.11). We are left with proving (2.10) when $s > 0$. We define a by

$$\frac{1}{a} = \frac{d+4}{2(d+2)} - \frac{2}{p(d+2)}.$$

Note that the assumption $p \geq \frac{4}{d}$ implies $\frac{2(d+2)}{d+4} \leq a \leq \frac{d+2}{2}$. By using product rule Lemma 2.1, we have

$$(2.17) \quad \|g(u) - g(v)\|_{\dot{N}^s} \leq \| |\nabla|^s (g(u) - g(v)) \|_{Z^0}$$

$$(2.18) \quad \lesssim \| |\nabla|^s (u - v) \|_{W^0} \left(\left\| \int_0^1 g_z(v + \theta(u - v)) d\theta \right\|_{L_{t,x}^{\frac{d+2}{2}}} + \left\| \int_0^1 g_{\bar{z}}(v + \theta(u - v)) d\theta \right\|_{L_{t,x}^{\frac{d+2}{2}}} \right)$$

$$(2.19) \quad + \|u - v\|_{X_p} \left(\left\| \int_0^1 |\nabla|^s g_z(v + \theta(u - v)) d\theta \right\|_{L_{t,x}^a} + \left\| \int_0^1 |\nabla|^s g_{\bar{z}}(v + \theta(u - v)) d\theta \right\|_{L_{t,x}^a} \right).$$

By the assumption (2.15), the term (2.18) is bounded as follows

$$(2.20) \quad (2.18) \lesssim \|u - v\|_{\dot{W}^s} (\|u\|_{L_{t,x}^{p(d+2)/2}}^p + \|v\|_{L_{t,x}^{p(d+2)/2}}^p) \approx \|u - v\|_{\dot{W}^s} \|(u, v)\|_{X_p}^p.$$

We now consider the term (2.19). We will prove the following bound

$$(2.21) \quad (2.19) \lesssim \|u - v\|_{X_p} \|(u, v)\|_{\dot{W}^s} \|(u, v)\|_{X_p}^{p-1}$$

Combining (2.20) and (2.21), we obtain the bound (2.10). We are thus left with proving (2.21).

Note that the term (2.19) is bounded by

$$\|u - v\|_{X_p} \left(\sup_{\theta \in [0,1]} \| |\nabla|^s g_z(v + \theta(u - v)) \|_{L_{t,x}^a} + \sup_{\theta \in [0,1]} \| |\nabla|^s g_{\bar{z}}(v + \theta(u - v)) \|_{L_{t,x}^a} \right).$$

Letting $f = g_z$ or $f = g_{\bar{z}}$, we will prove, for a general function $u \in \dot{W}^s \cap X_p$,

$$(2.22) \quad \| |\nabla|^s (f(u)) \|_{L_{t,x}^a} \lesssim \|u\|_{\dot{W}^s} \|u\|_{X_p}^{p-1}.$$

Note that this will conclude the proof of (2.21).

The function f belongs to $C^{\lceil s \rceil}$ and satisfies

$$(2.23) \quad |f^{(k)}(z)| \lesssim |z|^{p-k},$$

for each $0 \leq k \leq \lceil s \rceil$. We use similar argument as in [2, Proof of Lemma 3.3]. Using the equivalence of norms (2.3), we have

$$(2.24) \quad \| |\nabla|^s f(u) \|_{L_{t,x}^a} \lesssim \sum_{|\alpha|=\lceil s \rceil} \|D^\alpha f(u)\|_{L^a \dot{H}^{v,a}},$$

where $v = s - \lceil s \rceil$. By Lemma 2.6, $D^\alpha (f(u))$ is a linear combinations of terms of the form

$$(\partial_z^{h_1} \partial_{\bar{z}}^{h_2} f)(u) \prod_{k=1}^{h_1} D^{\beta_k} u \prod_{k=1}^{h_2} D^{\gamma_k} \bar{u},$$

where $\sum_{k=1}^{h_1} \beta_k + \sum_{k=1}^{h_2} \gamma_k = \alpha$, $1 \leq h_1 + h_2 \leq |\alpha|$, $|\beta_k| \geq 1$, $|\gamma_k| \geq 1$ for all k . To simplify notations we will only consider terms of the form

$$f^{(h)}(u) \prod_{k=1}^h D^{\beta_k} u, \quad \sum_{k=1}^h \beta_k = \alpha, \quad 1 \leq h \leq |\alpha|, \quad |\beta_k| \geq 1,$$

where $f^{(h)} = \left(\frac{\partial}{\partial z}\right)^h f$. The proof is the same for the other terms. We distinguish between the cases $h < p$ and $h = p$.

Case $h < p$.

Using Lemma 2.2 and (2.23), we have

$$(2.25) \quad \left\| f^{(h)}(u) \prod_{i=1}^h D^{\beta_i} u \right\|_{L^a \dot{H}^{v,a}}$$

$$(2.26) \quad \lesssim \left\| f^{(h)}(u) \right\|_{L^{q_1} \dot{H}^{v,q_1}} \prod_{i=1}^h \|D^{\beta_i} u\|_{L_{t,x}^{r_i}}$$

$$(2.27) \quad + \| |u|^{p-h} \|_{L_{t,x}^{q_2}} \sum_{k=1}^h \prod_{i=1, i \neq k}^h \|D^{\beta_i} u\|_{L_{t,x}^{r_i}} \|D^{\beta_k} u\|_{L^{\rho_k} \dot{H}^{v,\rho_k}}.$$

where, for $i \in \llbracket 1, h \rrbracket$,

$$(2.28) \quad \frac{1}{r_i} = \frac{|\beta_i|}{s} \frac{d}{2(d+2)} + \left(1 - \frac{|\beta_i|}{s}\right) \frac{2}{p(d+2)}, \quad \frac{1}{\rho_i} = \frac{|\beta_i| + v}{s} \frac{d}{2(d+2)} + \left(1 - \frac{|\beta_i| + v}{s}\right) \frac{2}{p(d+2)}$$

and

$$(2.29) \quad \frac{1}{q_1} = \frac{v}{s} \frac{d}{2(2+d)} + \frac{2}{p(d+2)} \frac{\lfloor s \rfloor}{s} + \frac{2(p-h-1)}{p(d+2)}, \quad \frac{1}{q_2} = \frac{2(p-h)}{p(d+2)}.$$

Since $\sum_{k=1}^h |\beta_k| = |\alpha| = \lfloor s \rfloor$, we see that the right-hand sides of the equalities in (2.28) and (2.29) are positive, and thus that q_1, q_2 , and, for $i \in \llbracket 1, h \rrbracket$, r_i and ρ_i are finite and positive. Using also that $v = s - \lfloor s \rfloor$, we obtain

$$\frac{1}{a} = \frac{1}{q_1} + \sum_{i=1}^h r_i = \frac{1}{q_2} + \sum_{\substack{1 \leq i \leq h \\ i \neq k}} \frac{1}{r_i} + \frac{1}{\rho_k},$$

which proves that q_1, q_2 , and, for $i \in \llbracket 1, h \rrbracket$, r_i and ρ_i are all greater than a , and that the assumptions of Lemma 2.2 are satisfied.

By Gagliardo-Nirenberg inequality Lemma 2.5 and the definition of r_i, ρ_i , we see that

$$\|D^{\beta_i} u\|_{L_x^{r_i}} \lesssim \| |\nabla|^s u \|_{L_x^{\frac{2(2+d)}{s}}} \|u\|_{L_x^{\frac{p(d+2)}{2}}}^{1 - \frac{|\beta_i|}{s}}, \quad \|D^{\beta_i} u\|_{\dot{H}^{v,\rho_i}} \lesssim \| |\nabla|^s u \|_{L_x^{\frac{2(2+d)}{s}}} \|u\|_{L_x^{\frac{p(d+2)}{2}}}^{1 - \frac{|\beta_i| + v}{s}}.$$

Integrating in time and using Hölder inequality, we obtain

$$(2.30) \quad \|D^{\beta_i} u\|_{L_{t,x}^{r_i}} \lesssim \|u\|_{\dot{W}_s^s}^{\frac{|\beta_i|}{s}} \|u\|_{X_p}^{1 - \frac{|\beta_i|}{s}}, \quad \|D^{\beta_i} u\|_{L_t^{\rho_i} \dot{H}^{v,\rho_i}} \lesssim \|u\|_{\dot{W}_s^s}^{\frac{|\beta_i| + v}{s}} \|u\|_{X_p}^{1 - \frac{|\beta_i| + v}{s}}$$

By (2.30), we obtain:

$$(2.27) \lesssim \|u\|_{X_p}^{p-1} \|u\|_{\dot{W}_s^s}.$$

Using the first inequality in (2.30), (2.26) is estimated by

$$(2.26) \lesssim \left\| f^{(h)}(u) \right\|_{L^{q_1} \dot{H}^{v,q_1}} \prod_{i=1}^h \|u\|_{\dot{W}_s^s}^{\frac{|\beta_i|}{s}} \|u\|_{X_p}^{1 - \frac{|\beta_i|}{s}}$$

$$(2.31) \quad = \left\| f^{(h)}(u) \right\|_{L^{q_1} \dot{H}^{v,q_1}} \|u\|_{\dot{W}_s^s}^{\frac{\lfloor s \rfloor}{s}} \|u\|_{X_p}^{h - \frac{\lfloor s \rfloor}{s}}.$$

In (2.31), we will estimate $\|f^{(h)}(u)\|_{L^{q_1} \dot{H}^{v,q_1}}$ by $\|u\|_{X_p}$ and $\|u\|_{\dot{W}_s^s}$ using fractional chain rule Lemma 2.3 and Gagliardo-Nirenberg inequality Lemma 2.5.

If $s \notin \mathbb{N}$ then $\lfloor s \rfloor \geq h + 1$ (because $h \leq |\alpha| = \lfloor s \rfloor$). Using Fractional chain rule Lemma 2.3, we have

$$\begin{aligned} \left\| f^{(h)}(u) \right\|_{L^{q_1} \dot{H}^{v,r_1}} &\lesssim \left\| f^{(h+1)}(u) \right\|_{L_{t,x}^{q_3}} \|u\|_{L^{q_4} \dot{H}^{v,q_4}} \\ &\lesssim \| |u|^{p-h-1} \|_{L_{t,x}^{q_3}} \|u\|_{L^{q_4} \dot{H}^{v,q_4}} \\ &= \|u\|_{L_{t,x}^{\frac{p-h-1}{q_3}}}^{p-h-1} \|u\|_{L^{q_4} \dot{H}^{v,q_4}}, \end{aligned}$$

where

$$\frac{1}{q_3} = \frac{2(p-h-1)}{p(d+2)}, \quad \frac{1}{q_4} = \frac{1}{q_1} - \frac{1}{q_3} = \frac{v}{s} \frac{d}{2(2+d)} + \frac{\lfloor s \rfloor}{s} \frac{2}{p(d+2)}.$$

For (q_4, r_4) as above, using Gagliardo-Nirenberg inequality Lemma 2.5, we have

$$\|u\|_{L^{q_4} \dot{H}^{v, r_4}} \lesssim \|u\|_{\dot{W}^s}^{\frac{v}{s}} \|u\|_{X_p}^{\frac{[s]}{s}}.$$

Thus, for $s \notin \mathbb{N}$,

$$(2.32) \quad (2.26) \lesssim (2.31) \lesssim \|u\|_{X_p}^{p-h-1} \|u\|_{\dot{W}^s}^{\frac{v}{s}} \|u\|_{X_p}^{\frac{[s]}{s}} \|u\|_{\dot{W}^s}^{\frac{[s]}{s}} \|u\|_{X_p}^{h-\frac{[s]}{s}} = \|u\|_{\dot{W}^s} \|u\|_{X_p}^{p-1},$$

yielding (2.22). If $s \in \mathbb{N}$ then $v = 0$, $\sum_{i=1}^k |\beta_i| = s$, and $\frac{1}{q_1} = \frac{p-h}{p(d+2)/2}$. Using (2.30), we obtain

$$(2.26) \lesssim (2.31) \lesssim \|u\|_{L^{q_1} \dot{H}^{v, a}}^{p-h} \|u\|_{\dot{W}^s} \|u\|_{X_p}^{h-1} = \|u\|_{X_p}^{p-1} \|u\|_{\dot{W}^s},$$

which proves (2.22) in this case also.

Case $h = p$.

In this case, we have $p = |\alpha|$, and thus, since $|\alpha| = [s] \leq [s] \leq p$, s is an integer, $v = 0$ and $s = p$.

By the assumptions on f , we have $|f^{(h)}(u)| \lesssim 1$. Thus

$$(2.25) \lesssim \left\| f^{(h)}(u) \prod_{i=1}^h D^{\beta_i} u \right\|_{L^a \dot{H}^{v, a}} \lesssim \left\| \prod_{i=1}^h D^{\beta_i} u \right\|_{L^a \dot{H}^{v, a}} \lesssim \prod_{i=1}^h \|D^{\beta_i} u\|_{L^{r_i}},$$

where the r_i are defined as above. Using (2.30) and since $\sum_{i=1}^h |\beta_i| = s$, and $h = p$, we obtain (2.22), which concludes the proof of the Lemma. \square

Proof of Proposition 2.8. Proposition 2.8 follows easily from Lemmas 2.9 and 2.10. We fix $g \in \mathcal{N}(s, p_0, p_1)$ with $0 \leq s$, $p_0 \geq q_d$ and $p_0 \geq [s]$. All the norms used are over the interval I .

The estimate (2.9) is exactly (2.8) with $v = 0$. To prove (2.8), we decompose $g(u)$ as follows:

$$g(u) = P(u) + \tilde{g}(u).$$

Where $P(u)$ is the Taylor expansion of g at $u = 0$ up to order $[s]$. As a consequence P is a polynomial of the form

$$P(u) = \sum_{p_1+1 \leq k_1+k_2 \leq [s]} a_{k_1, k_2} u^{k_1} \bar{u}^{k_2}.$$

By Lemma 2.9, using that by the definition of X and the assumptions $p_1 \geq \frac{4}{d}$ and $p_0 \geq [s]$ we have

$$p_1 \leq q \leq [s] - 1 \implies \|u\|_{X_q} \leq \|u\|_X,$$

we obtain

$$(2.33) \quad \|P(u) - P(v)\|_{N^s} \lesssim \left(\|u\|_X^{p_0-1} + \|v\|_X^{p_0-1} + \|u\|_X^{p_1-1} + \|v\|_X^{p_1-1} \right) \\ \times \left[\|u - v\|_{\dot{W}^s} (\|u\|_X + \|v\|_X) + \|u - v\|_X (\|u\|_{W^s} + \|v\|_{W^s}) \right].$$

Next, we notice that since $g \in \mathcal{N}(s, p_0, p_1)$, the definition of P and the inequalities $p_1 < p_0$ and $[s] \leq p_0$ imply, for $k \in \llbracket 0, [s] + 1 \rrbracket$, $z \in \mathbb{C}$,

$$|z| \geq 1 \implies \left| \tilde{g}^{(k)}(z) \right| \leq C|z|^{p_0+1-k}, \quad |z| \leq 1 \implies \left| \tilde{g}^{(k)}(z) \right| \leq C|z|^{[s]+1-k}.$$

We let χ be a smooth function such that

$$\chi = \begin{cases} 1 & \text{if } |x| < 1, \\ 0 & \text{if } |x| > 2. \end{cases}$$

Set $g_0(u) = (1 - \chi(u))\tilde{g}(u)$, $g_1(u) = \chi(u)\tilde{g}(u)$, so that g_0 satisfies the assumptions of Lemma 2.10 with $p = \max(p_0, [s]) = p_0$, and g_1 the same assumptions with $p = [s]$. Combining the conclusion of this Lemma for g_0, g_1 with the estimate (2.33), we obtain the conclusion of the proposition. \square

2.4. Cauchy and stability theory for general nonlinearities. In all this subsection, we fix g, p_0, p_1, s such that Assumption A holds. We assume furthermore

$$(2.34) \quad s \geq s_0 = \frac{d}{2} - \frac{2}{p_0}.$$

Note that these assumptions are satisfied for a sum of powers:

$$g(u) = \sum_{j=0}^k \lambda_j |u|^{p_j} u,$$

where $k \geq 0$, $\frac{4}{d} \leq p_1 < \dots < p_{k-1} < p_0$, $\lambda_j \in \mathbb{R}$ for all j , provided $g \in C^{\lceil s \rceil + 1}$ (which is the case, for example, when all the p_j are even integers), $s \geq \frac{d}{2} - \frac{2}{p_0}$, and $\lceil s \rceil \leq p_0$ if g is not a polynomial.

Definition 2.11. Let $0 \in I$ be an interval. By definition, a solution u to (1.1) on I , with initial data in H^s ($s \geq s_0$) is a function $u \in C(I, H^s)$ such that for all $K \subset I$ compact, $u \in S^s(K)$ and u satisfies the following Duhamel formula

$$(2.35) \quad u(t) = e^{it\Delta} u_0 - i \int_0^t e^{i(t-\tau)\Delta} g(u)(\tau) d\tau,$$

for all $t \in I$.

We recall from (2.7) the definition of $X(I)$. Noting that the assumption $p_0 \geq \frac{4}{d}$ implies $p_0(d+2)/2 > 2$ (and $p_0(d+2)/2 \geq 6$ if $d=1$), we can choose q_0 such that $(p_0(d+2)/2, q_0)$ is an admissible pair. By Sobolev inequality and the definitions of s_0, q_0 , one can check

$$(2.36) \quad \|u\|_{L_x^{\frac{p_0(d+2)}{2}}} \lesssim \| |\nabla|^{s_0} u \|_{L^{q_0}}$$

and thus

$$(2.37) \quad \|u\|_{L^{\frac{p_0(d+2)}{2}}(I \times \mathbb{R}^d)} \lesssim \| |\nabla|^{s_0} u \|_{S^0(I)}, \quad \|u\|_{X(I)} \lesssim \|u\|_{S^{s_0}(I)}.$$

Proposition 2.12. Let $u_0 \in H^s$. Let g, p_0, p_1, s such that Assumption A holds and $s \geq \frac{d}{2} - \frac{2}{p_0}$. Let $0 \in I$ be an interval of \mathbb{R} , and $A > 0$. Assume that $\|u_0\|_{H^s} \leq A$ and

$$\|e^{i\cdot\Delta} u_0\|_{X(I)} = \delta \leq \delta_0(A) \text{ small.}$$

Then there exists a unique solution u to (1.1) such that

$$u(0) = u_0, \quad \|u\|_{S^s(I)} \lesssim \|u_0\|_{H^s}, \quad \|u\|_{X(I)} \leq 2\delta.$$

Moreover, if $u_{0,k} \rightarrow u_0$ in H^s (so that, for k large, $\|e^{it\Delta} u_{0,k}\|_{X(I)} < \delta$) then the corresponding solution $u_k \rightarrow u$ in $C(I, H^s)$.

Remark 2.13. By Strichartz estimates, (2.37) and Proposition 2.12, if $\|u_0\|_{H^s}$ is small then u is global and

$$\|u\|_{S^s} \lesssim \|u_0\|_{H^s}.$$

Proof. We use similar argument as in [24, Lemma 2.5]. We work on the interval I for all norms.

Consider

$$B = \left\{ u : \|u\|_X \leq 2\delta, \|u\|_{S^s} \leq MA \right\},$$

for some large universal constant M , with the topology induced by the norm in $S^s(I)$. We denote by $\Psi(u)$ the right hand side of (2.35). We show that if $\delta \leq \delta_0(A)$ small enough, and M large enough (independently of A), Ψ is a contraction map on B .

For $u \in B$, we have, by Strichartz estimates and Proposition 2.8

$$\begin{aligned} \|\Psi(u)\|_{X(I)} &\leq \|e^{it\Delta} u_0\|_{X(I)} + \left\| \int_0^t e^{i(t-\tau)\Delta} g(u)(\tau) d\tau \right\|_{X(I)} \\ &\leq \delta + C \left\| \int_0^t e^{i(t-\tau)\Delta} g(u)(\tau) d\tau \right\|_{S^s(I)} \leq \delta + C \|g(u)\|_{N^s(I)} \leq \delta + C \|u\|_{S^s} (\|u\|_X^{p_1} + \|u\|_X^{p_0}) \end{aligned}$$

Thus,

$$\|\Psi(u)\|_{X(I)} \leq \delta + CMA\delta^{p_1} \leq 2\delta,$$

if δ is small enough (so that $CMA\delta^{p_1-1} \leq 1$). Similarly,

$$\|\Psi(u)\|_{S^s} \leq \|e^{i\cdot\Delta}u_0\|_{S^s} + C\|g(u)\|_{N^s} \leq CA + CMA\delta^{p_1} \leq MA,$$

choosing $M \geq 2C$, and $\delta \leq \delta_0(A)$ small enough. Thus $\Psi(u) \in B$.

Now, let $u, v \in B$. We have, by Strichartz estimates and Proposition 2.8,

$$\begin{aligned} \|\Psi(u) - \Psi(v)\|_{S^s} &\leq C\|g(u) - g(v)\|_{N^s} \leq \|u - v\|_{S^s} (\|u\|_{S^s} + \|v\|_{S^s}) \left(\|u\|_{X^s}^{p_1-1} + \|v\|_{X^s}^{p_1-1} \right) \\ &\leq C\delta^{p_1-1}MA\|u - v\|_{S^s} \end{aligned}$$

Thus, taking δ small enough, we obtain that Ψ is a contraction map on B . By the fixed point theorem, there exists a unique $u \in B$ such that $u = \Psi(u)$. Thus, u is a solution to (1.1). Also, since $u \in B$ (and since we can take $A = \|u_0\|_{H^s}$) we have as announced $\|u\|_{S(I)} \lesssim \|u_0\|_{H^s}$, $\|u\|_{X(I)} \leq 2\delta$. If $\|u_0\|_{H^s}$ is small then $\|e^{i\cdot\Delta}u_0\|_{X(\mathbb{R})} \leq C\|u_0\|_{H^s}$ is small. Thus, u is global and $\|u\|_{S^s(\mathbb{R})} \leq M\|u_0\|_{H^s}$. \square

Remark 2.14. We have $\|e^{i\cdot\Delta}u_0\|_{X(\mathbb{R})} \leq C\|u_0\|_{H^s} < \infty$. Thus, for each $\varepsilon > 0$, there exists a small interval I around 0 such that $\|e^{i\cdot\Delta}u_0\|_{X(I)} < \varepsilon$. This implies local existence of solution.

Remark 2.15. If u, v are two solutions to (1.1) on $0 \in I$ such that $u(0) = v(0)$ then $u \equiv v$. We first show this assertion when $I' \subset I$ is a small interval around 0. Shrinking I' if necessary, we assume

$$\|u\|_{X(I')} \leq \delta_0(A), \quad \|v\|_{X(I')} \leq \delta_0(A),$$

with $\delta_0(A)$ as in Proposition 2.12, and

$$A = \max_{t \in I'} \|u(t)\|_{H^s} + \|v(t)\|_{H^s}.$$

Thus, $u, v \in B$ (where B is as in the preceding proof), which shows that $u = v$ on I' . Repeating this argument, we deduce that $u \equiv v$ on all I . This allows us to define a maximal interval $I_{\max} = (T_-, T_+)$ with $T_- < 0, T_+ > 0$.

Lemma 2.16. *Let u be a maximal solution on I_{\max} as in Proposition 2.12, Remark 2.15. Assume that*

$$u \in X([0, T_+)).$$

Then $T_+ = \infty$, and u scatters in the future in H^s . A similar result holds in the past, if $u \in X((T_-, 0])$.

Remark 2.17. Lemma 2.16 implies the blow-up criterion $T_+ < \infty \implies \|u\|_{X(0, T_+)} = \infty$ (and similarly for T_-). In the case $s > \frac{d}{2} - \frac{2}{p_0}$ we can also show (see e.g. [10]) that if $T_+(u)$ is the maximum time of existence of u and $T_+(u) < +\infty$ then

$$\|u(t)\|_{H^s} \rightarrow \infty, \text{ as } t \rightarrow T_+(u).$$

Proof. We only work on $[0, T_+)$. In $(T_-, 0]$, we use similar argument. We only need to prove that

$$(2.38) \quad \|u\|_{S^s(0, T_+)} < \infty.$$

Indeed, if (2.38) holds and T_+ is finite, then Proposition 2.12 shows that one can extend the solution u beyond T_+ , a contradiction with the definition of T_+ . Thus $T_+ = \infty$, and Proposition 2.8 implies $\|g(u)\|_{N([0, \infty))} < \infty$, which implies by standard arguments scattering in H^s .

We divide $[0, T_+)$ into finite intervals I_k such that $\|u\|_{X(I_k)} < \varepsilon$. On each $I_k = [t_k, t_{k+1}]$, we have

$$\begin{aligned} \|u\|_{S^s(I_k)} &\leq C\|u(t_k)\|_{H^s} + C\|u\|_{S^s(I_k)} (\|u\|_{X(I_k)}^{p_1} + \|u\|_{X(I_k)}^{p_0}) \\ &\leq C\|u(t_k)\|_{H^s} + C\|u\|_{S^s(I_k)} (\varepsilon^{p_1} + \varepsilon^{p_0}). \end{aligned}$$

Choosing ε small and using that $\|u(t_{k+1})\|_{H^s} \lesssim \|u\|_{S^s(I_k)}$, we obtain by induction on k that $\|u\|_{L^\infty([0, T_{k_{\max}}], H^s)} < \infty$ then (2.38), concluding the proof. \square

One can also prove the existence of wave operators for equation (1.1):

Proposition 2.18. *Let $u_0 \in H^s$. Let g, p_0, p_1, s such that Assumption A holds and $s \geq \frac{d}{2} - \frac{2}{p_0}$. Let $v_0 \in H^s$ and $v_L(t) = e^{it\Delta}v_0$. Then there exist a unique solution $u \in C^0((T_-(u), \infty))$ of (1.1) such that*

$$\lim_{t \rightarrow \infty} \|u(t) - v_L(t)\|_{H^s} = 0.$$

We omit the proof which is similar to the proof of Proposition 2.12.

Theorem 2.19. *(Long time perturbation theory) Let $A > 0, s \geq s_0$. There exists constants $\varepsilon(A, s) \in (0, 1], C(A, s) > 0$ with the following properties. Let $0 \in I$ be an compact interval of \mathbb{R} and w be a solution of the following equation*

$$Lw = g(w) + e,$$

and $u_0 \in H^s$ such that

$$\|w_0\|_{H^s} + \|w\|_{X(I)} \leq A, \quad \|e\|_{N^s(I)} + \|u_0 - w(0)\|_{H^s} = \varepsilon \leq \varepsilon(A, s).$$

Then the solution u of (1.1) with initial data u_0 is defined on I and satisfies

$$(2.39) \quad \|u - w\|_{S^s(I)} \leq C(A, s)\varepsilon, \quad \|u\|_{S^s(I)} \leq C(A, s).$$

Proof. Without loss of generality, using the reversibility of the equation, we can assume $I = [0, T)$. We can also assume that $I \subset I_{\max}(u)$ (indeed, if $T_+(u) < T$, the proof below will show that (2.39) holds with I replaced by $[0, T_+)$, contradicting the blow-up criterion).

Divide I into $J = J(A, s)$ subintervals $I_j = [t_j, t_{j+1}]$, $j \in [0, J - 1]$, with $t_0 = 0$, such that

$$(2.40) \quad \|w\|_{X(I_j)} \leq \delta,$$

where δ is a small constant (depending only on s), to be specified. On each I_j , by Strichartz and Proposition 2.8, we have

$$\begin{aligned} \|w\|_{S^s(I_j)} &\leq C \|w(t_j)\|_{H^s} + \|g(w)\|_{N^s(I_j)} + \|e\|_{N^s(I_j)} \\ &\lesssim \|w(t_j)\|_{H^s} + \|w\|_{S^s(I_j)} (\|w\|_{X(I_j)}^{p_1} + \|w\|_{X(I_j)}^{p_0}) + \varepsilon \\ &\lesssim \|w(t_j)\|_{H^s} + \varepsilon + \|w\|_{S^s(I_j)} (\delta^{p_1} + \delta^{p_0}). \end{aligned}$$

Thus, for δ small enough, $\|w\|_{S^s(I_j)} \leq \|w(t_j)\|_{H^s} + \varepsilon$. Since $\|w(t_{j+1})\|_{H^s} \leq \|w\|_{S^s(I_j)}$, $\|w(0)\|_{H^s} \leq A$, we obtain by a finite induction $\|w(t_j)\|_{H^s} \leq C(A, s)$, and thus

$$(2.41) \quad \|w\|_{S^s(I)} \leq C(A, s).$$

Thus, we may divide I into $J = J(A, s)$ subintervals, which we still denote by $I_j = [t_j, t_{j+1}]$, $j \in [0, J - 1]$, such that for all j ,

$$(2.42) \quad \|w\|_{W^s(I_j)} + \|w\|_{X(I_j)} \leq \delta.$$

Taking δ small (independently of A and ε), and $\varepsilon \leq \varepsilon(A, s)$ small, we prove by induction

$$(2.43) \quad \forall j \in [0, J - 1], \quad \|u - w\|_{S^s(I_j)} \leq (Kj + K + 1)\varepsilon,$$

for some large constant K independent of A and ε . This will imply the desired conclusion (2.39). More precisely, we will prove, for $j \in [0, J - 1]$,

$$(2.44) \quad \|u(t_j) - w(t_j)\|_{H^s} \leq (Kj + 1)\varepsilon \implies \|u - w\|_{S^s(I_j)} < (Kj + K + 1)\varepsilon,$$

which will yield (2.43), since $\|u(t_0) - w(t_0)\|_{H^s} \leq \varepsilon$.

Assuming $j \in [0, J - 1]$, $\|u(t_j) - w(t_j)\|_{H^s} \leq (Kj + 1)\varepsilon$, we argue by contradiction, assuming also that there exists $T \in [t_j, t_{j+1}]$ such that $\|u - w\|_{S^s([t_j, T])} = (Kj + K + 1)\varepsilon$. Then by the equation satisfied by $u - w$ and Strichartz estimates, we obtain

$$(2.45) \quad \|u - w\|_{S^s([t_j, T])} \leq \|u(t_j) - w(t_j)\|_{H^s} + C\|e\|_{N^s([t_j, T])} + C\|g(u - w)\|_{N^s([t_j, T])}$$

We have $\|u\|_{W^s([t_j, T])} \leq \|u - w\|_{W^s([t_j, T])} + \|w\|_{W^s([t_j, T])} \leq (Kj + K + 1)\varepsilon + \delta$, and similarly $\|u\|_{X^s([t_j, T])} \leq C(Kj + K + 1)\varepsilon + \delta$. Thus by Proposition 2.8, taking ε small enough, so that $(Kj + K + 1)\varepsilon \leq 1$.

$$\|g(u - w)\|_{N^s([t_j, T])} \leq C(Kj + K + 1) \left((Kj + K + 1)^{p_1} \varepsilon^{p_1} + \delta^{p_1} \right).$$

Going back to (2.45), we deduce

$$(Kj + K + 1)\varepsilon = \|u - w\|_{S^s([t_j, T])} \leq C\varepsilon + (Kj + 1)\varepsilon + C(Kj + K + 1)\varepsilon((Kj + K + 1)^{p_1}\varepsilon^{p_1} + \delta^{p_1}).$$

Taking K large enough, $\varepsilon \leq \varepsilon(s, A)$ small (so that $(Kj + K + 1)\varepsilon \leq \delta$), and δ small, we obtain.

$$(Kj + K + 1)\varepsilon \leq (Kj + K/2 + 1)\varepsilon,$$

a contradiction which concludes the proof. \square

Remark 2.20. Assume $d \geq 1$, $g(u) = |u|^{p_0}u$, $s \geq s_0 = \frac{d}{2} - \frac{2}{p_0}$, $p_0 \geq \frac{4}{d}$ and $g \in C^{\lceil s \rceil + 1}$ (i.e. p_0 is an even integer or $p_0 > \lceil s \rceil$). Then analogs of Proposition 2.12 and Theorem 2.19 where all the spaces are replaced by homogeneous spaces hold. Precisely, in the statement of Proposition 2.12, one can replace in this case $X(I)$ by $X_{p_0}(I) = L^{\frac{p_0(d+2)}{2}}(I \times \mathbb{R}^d)$, H^s by \dot{H}^s , $S^s(I)$ by $\dot{S}^s(I)$, and similarly for Proposition 2.12. The proof is the same, replacing Proposition 2.8 by Lemma 2.9. See also e.g. [24].

Proposition 2.21 (Conservation laws). *Let g such that Assumption A holds and $g(z) = G'(|z|^2)z$ for a C^1 function $G : [0, \infty) \rightarrow \mathbb{R}$ with $G(0) = 0$. Let $u \in C^0(I, H^s(\mathbb{R}^d))$ be a solution of (2.1), where $s \geq \frac{d}{2} - \frac{2}{p_0}$. Then the mass*

$$M(u) := \int |u(t, x)|^2 dx$$

is conserved on I . Furthermore, if $s \geq 1$, then the momentum:

$$P(u) := \text{Im} \int \nabla u(t, x) \bar{u}(t, x) dx$$

and the energy

$$E(u) := \int |\nabla u(t, x)|^2 dx + \int G(|u(t, x)|^2) dx.$$

are well-defined and conserved on I .

We omit the classical proof. See e.g. [9]. Observe that the assumptions of Proposition 2.21 imply

$$(2.46) \quad G(|u|^2) \lesssim |u|^{p_0+2} + |u|^{p_1+2},$$

so that by Sobolev inequalities, $E(u)$ is well-defined if $s \geq 1$.

3. PROFILE DECOMPOSITION

3.1. Profiles in homogeneous Sobolev spaces. Let $0 < s < \frac{d}{2}$. We denote by $u_L(t) = e^{it\Delta}u_0$ the solution to the linear Schrödinger equation on $\mathbb{R} \times \mathbb{R}^d$

$$(3.1) \quad i\partial_t u_L + \Delta u_L = 0,$$

with initial data

$$(3.2) \quad u_L|_{t=0} = u_0 \in \dot{H}^s.$$

Definition 3.1. Let $p > \frac{4}{d}$ such that $s = \frac{d}{2} - \frac{2}{p}$. A linear \dot{H}^s -*profile*, in short *profile*, is a sequence $(\varphi_{L_n})_n$, of solutions of (3.1), of the form

$$(3.3) \quad \varphi_{L_n}(t, x) = \frac{1}{\lambda_n^{\frac{p}{2}}} \varphi_L \left(\frac{t - t_n}{\lambda_n^2}, \frac{x - x_n}{\lambda_n} \right),$$

where φ_L is a fixed solution of (3.1), (3.2) and $\Lambda_n = (\lambda_n, t_n, x_n)_n$ is a sequence in $(0, \infty) \times \mathbb{R} \times \mathbb{R}^d$ (called sequence of transformations) such that

$$(3.4) \quad \lim_{n \rightarrow \infty} \frac{-t_n}{\lambda_n^2} = \tau \in \mathbb{R} \cup \{\pm\infty\}.$$

Definition 3.2. We say that two sequence of transformations $\Lambda_n = (\lambda_n, t_n, x_n)$ and $M_n = (\mu_n, s_n, y_n)$ are *orthogonal* when they satisfy

$$\lim_{n \rightarrow \infty} \frac{|t_n - s_n|}{\lambda_n^2} + \frac{|x_n - y_n|}{\lambda_n} + \left| \log \left(\frac{\lambda_n}{\mu_n} \right) \right| = \infty.$$

We say that two \dot{H}^s -profiles φ_{L_n} and ψ_{L_n} are *equivalent* (in \dot{H}^s) when

$$\lim_{n \rightarrow \infty} \|\varphi_{L_n}(0) - \psi_{L_n}(0)\|_{\dot{H}^s} = 0.$$

We say that they are *orthogonal* when one of the two profiles is identically 0 or when the corresponding sequence of transformations are orthogonal.

Remark 3.3. For a given \dot{H}^s -profile φ_{L_n} , the choice of the solution φ_L and the sequence of transformations $(\Lambda_n)_n$ are not unique. However the definitions of equivalent and orthogonal profiles do not depend on these choices. Also, two equivalent \dot{H}^s -profiles are orthogonal to the same \dot{H}^s -profiles.

Definition 3.4. Let $(u_{0,n})_n$ be a bounded sequence in \dot{H}^s , and $u_{L_n} = e^{it\Delta}u_{0,n}$. We say that the sequence $\left((\varphi_{L_n}^j)_n \right)_{j \geq 1}$ of \dot{H}^s -profiles is a *profile decomposition* of $(u_{0,n})_n$ if these profiles are pairwise orthogonal, and satisfy

$$(3.5) \quad \lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|w_{L_n}^J\|_{X_p(\mathbb{R})} + \|\nabla^s w_{L_n}^J\|_{W^0(\mathbb{R})} = 0,$$

where

$$(3.6) \quad w_{L_n}^J = u_{L_n} - \sum_{j=1}^J \varphi_{L_n}^j.$$

Proposition 3.5. *For any bounded sequence $(u_{0,n})_n$ in \dot{H}^s , there exists a subsequence (that we still denote by $(u_{0,n})_n$) that admits a profile decomposition $\left((\varphi_{L_n}^j)_n \right)_j$. Furthermore, we have the Pythagorean expansion:*

$$(3.7) \quad \forall J \geq 1, \quad \|u_{0,n}\|_{\dot{H}^s}^2 = \sum_{j=1}^J \left\| \varphi_{L_n}^j(0) \right\|_{\dot{H}^s}^2 + \|w_{L_n}^J(0)\|_{\dot{H}^s}^2 + o(1), \quad n \rightarrow \infty,$$

where w_n^J is as in Definition 3.4.

Proposition 3.5 is proved by Shao in [47], using the L^2 -critical profile decomposition of Merle and Vega [36], generalized to higher dimension by Bégout and Vargas [4]. See also [26] for the energy-critical case $p = \frac{4}{d-2}$.

Remark 3.6. In the notations of Definition 3.4, if $(u_{0,n})_n$ admits a profile decomposition, then

$$(3.8) \quad (\lambda_n^j)^{\frac{2}{p}} \varphi_{L_n}^k(t_{j,n}, x_{j,n} + \lambda_n^j \cdot) \xrightarrow{n \rightarrow \infty} 0 \text{ in } \dot{H}^s, \quad j \neq k$$

$$(3.9) \quad (\lambda_n^j)^{\frac{2}{p}} w_{L_n}^J(t_{j,n}, x_{j,n} + \lambda_n^j \cdot) \xrightarrow{n \rightarrow \infty} 0 \text{ in } \dot{H}^s, \quad J \geq j$$

$$(3.10) \quad (\lambda_n^j)^{\frac{2}{p}} u_{L_n}(t_{j,n}, x_{j,n} + \lambda_n^j \cdot) \xrightarrow{n \rightarrow \infty} \varphi_{L_n}^j(0) \text{ in } \dot{H}^s.$$

Indeed, (3.8) follows from the orthogonality of the profiles. The property (3.9) follows from (3.5) and the orthogonality of the profiles. Finally, (3.10) follows easily from the two other properties. By (3.9), (3.10), one sees that the initial data of the nonzero profiles are exactly the nonzero weak limits of the form (3.10). This implies in particular that the profile decomposition is unique, up to reordering and equivalent profiles, if one ignore the null profiles.

Let $q = \frac{2d}{d-2s} = \frac{d}{2}p$ be the Lebesgue exponent such that the Sobolev embedding $\dot{H}^s \subset L^q$ holds. Let $(u_{0,n})_n$ be a sequence that has a profile decomposition as above. We next prove the analog of the property (3.5) of the remainder w_n^J in the space $L^\infty(\mathbb{R}, L^q)$ and obtain a Pythagorean expansion of the L^q norm of $u_{0,n}$. Extracting subsequences, we can assume that for all j , the following limit exists in $\mathbb{R} \cup \{\pm\infty\}$:

$$(3.11) \quad \tau^j = \lim_{n \rightarrow \infty} \frac{-t_n^j}{(\lambda_n^j)^2}.$$

We can also assume (translating the profiles in time if necessary), $\tau^j \in \{0, -\infty, +\infty\}$. If $\tau^j \in \{\pm\infty\}$, we have

$$(3.12) \quad \lim_{n \rightarrow \infty} \|\varphi_{L_n}^j(0)\|_{L^q} = 0.$$

Lemma 3.7. *Let $(u_{0,n})_n$ be as in Proposition 3.5. Then*

$$(3.13) \quad \lim_{J \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \left\| w_{L_n}^j \right\|_{L^\infty(\mathbb{R}, L^q)} = 0$$

$$(3.14) \quad \lim_{n \rightarrow \infty} \|u_{0,n}\|_{L^q}^q = \sum_{\substack{j \geq 1 \\ \tau^j = 0}} \|\varphi_{L_n}^j(0)\|_{L^q}^q.$$

Proof. This follows from the elliptic profile decomposition of Patrick Gérard [20]. We first prove (3.13) by contradiction, assuming that there exists $\eta > 0$ and a sequence $J_k \rightarrow \infty$ such that

$$(3.15) \quad \forall k, \quad \lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|w_{L_n}^{J_k}\|_{L^\infty(\mathbb{R}, L^q)} \geq \eta.$$

Let

$$\varepsilon_k = \limsup_{n \rightarrow \infty} \|w_{L_n}^{J_k}\|_{X_p(\mathbb{R})} \xrightarrow{k \rightarrow \infty} 0,$$

where the convergence to 0 follows from (3.5). For all k , there exists N_k such that

$$n \geq N_k \implies \|w_{L_n}^{J_k}\|_{X_p(\mathbb{R})} \leq 2\varepsilon_k.$$

By (3.15), we can find $n_k \geq N_k$, $t_k \in \mathbb{R}$ such that

$$\|w_{L_{n_k}}^{J_k}(t_k)\|_{L^q} \geq \frac{1}{2}\eta.$$

As a consequence, by the main result of [20], we can find $\varphi \in \dot{H}^s \setminus \{0\}$ such that

$$w_{L_{n_k}}^{J_k}(t_k) \xrightarrow{k \rightarrow \infty} \varphi.$$

By Strichartz estimates, this implies that the sequence $\left(w_{L_{n_k}}^{J_k}(\cdot + t_k)\right)_k$ converges weakly to $e^{i\cdot\Delta}\varphi$ in $X_p(\mathbb{R})$, contradicting (3.5). This concludes the proof of (3.13). As a consequence (using also (3.12)), we have the ‘‘elliptic’’ profile decomposition at fixed time:

$$u_{0,n} = \sum_{\substack{1 \leq j \leq J \\ \tau^j = 0}} \varphi_{L_n}^j(0) + \tilde{w}_{L_n}^J(0), \quad \lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|\tilde{w}_{L_n}^J(0)\|_{L^q} = 0.$$

The pseudo-Pythagorean expansion (3.14) then follows from [20]: see the expansion (1.11) there. \square

3.2. Profiles in non-homogeneous Sobolev spaces. In all this subsection, we fix $s_0 > 0$ and $p_0 > \frac{4}{d}$ with $s_0 = \frac{d}{2} - \frac{2}{p_0}$. We consider a sequence $(u_{0,n})$, which is bounded in the *inhomogeneous* Sobolev space $H^{s_0}(\mathbb{R}^d)$. We assume that $u_{0,n}$ admits a \dot{H}^{s_0} profile decomposition $(\varphi_{L_n}^j)_{n,j}$, where

$$\varphi_{L_n}^j(t, x) = \frac{1}{(\lambda_n^j)^{\frac{2}{p_0}}} \varphi_L^j \left(\frac{t - t_n^j}{\lambda_n^j}, \frac{x - x_n^j}{\lambda_n^j} \right).$$

Claim 3.8. For all j , the sequence $(\lambda_n^j)_n$ is bounded.

Proof. We have

$$(3.16) \quad (\lambda_n^j)^{\frac{2}{p_0}} u_{L_n}(t_n^j, \lambda_n^j \cdot + x_n^j) \xrightarrow{n \rightarrow \infty} \varphi_L^j(0) \text{ weakly in } \dot{H}^{s_0}.$$

Furthermore,

$$\left\| (\lambda_n^j)^{\frac{2}{p_0}} u_{L_n}(t_n^j, \lambda_n^j \cdot + x_n^j) \right\|_{L^2} = (\lambda_n^j)^{\frac{2}{p_0} - \frac{N}{2}} \|u_{0,n}\|_{L^2}.$$

Assume that $\lambda_n^j \rightarrow \infty$ along a subsequence in n . As a consequence, since $\|u_{0,n}\|_{L^2}$ is bounded, we obtain, along the same subsequence, that $(\lambda_n^j)^{\frac{2}{p_0}} u_{L_n}(t_n^j, \lambda_n^j \cdot + x_n^j)$ converges strongly to 0 in L^2 . By uniqueness of the distributional limit, $\varphi_L^j(0) = 0$ as announced. \square

Using the claim and extracting subsequences we obtain that for all j , one of the following holds:

- $\varphi_L^j \equiv 0$. In this case we say that $(\varphi_{L_n}^j)_n$ is a *null profile*, and denote $j \in \mathcal{J}_0$.
- $\varphi_L^j \not\equiv 0$ and $\lim_{n \rightarrow \infty} \lambda_n^j = \lambda^j \in (0, \infty)$. In this case we say that $(\varphi_{L_n}^j)_n$ is a *non-concentrating profile*, and denote $j \in \mathcal{J}_{NC}$. The weak limit (3.16) and the fact that the sequence $(u_{0,n})$ is bounded in H^{s_0} proves that $\varphi_0^j = \varphi_L^j(0) \in H^{s_0}$. Replacing φ_0^j by $\frac{1}{(\lambda^j)^{\frac{N}{2}}} \varphi_0^j \left(\frac{x}{\lambda^j}\right)$ and λ_n^j by λ_n^j/λ^j , we see that we can assume $\lambda^j = 1$. As a consequence, we can assume $\lambda_n^j = 1$ for all n (this will modify the profile $\varphi_{L_n}^j$ only by a term which goes to 0 in H^{s_0} as $n \rightarrow \infty$).
- $\varphi_L^j \not\equiv 0$ and $\lim_{n \rightarrow \infty} \lambda_n^j = 0$. In this case we say that $(\varphi_{L_n}^j)_n$ is a *concentrating profile*, and denote $j \in \mathcal{J}_C$.

Remark 3.9. Assume that the sequence $(u_{0,n})$ is bounded in H^{s_2} for some $s_2 > s_0$. Then it is easy to see that \mathcal{J}_C is empty, i.e. that there is no concentrating profile.

Remark 3.10. Let s such that $0 < s < s_0$. Then $\left((\varphi_{L_n}^j)_n\right)_{j \in \mathcal{J}_{NC}}$ is a \dot{H}^s profile decomposition of $(u_{0,n})_n$. Indeed, by Remark 3.9 the \dot{H}^s -profile decomposition of this sequence has no concentrating profile. By the preceding subsection, we have the Pythagorean expansion

$$(3.17) \quad \forall J \geq 1, \quad \|u_{0,n}\|_{\dot{H}^s}^2 = \sum_{\substack{j \in \mathcal{J}_{NC} \\ 1 \leq j \leq J}} \left\| \varphi_L^j(0) \right\|_{\dot{H}^s}^2 + \|\bar{w}_{L_n}^J(0)\|_{\dot{H}^s}^2 + o(1), \quad n \rightarrow \infty,$$

where $\bar{w}_{L_n}^J = u_{L_n} - \sum_{\substack{j \in \mathcal{J}_{NC} \\ 1 \leq j \leq J}} \varphi_{L_n}^j$. One can also prove, as a consequence of the orthogonality of the profiles,

$$(3.18) \quad \forall J \geq 1, \quad \|u_{0,n}\|_{L^2}^2 = \sum_{\substack{j \in \mathcal{J}_{NC} \\ 1 \leq j \leq J}} \left\| \varphi_L^j(0) \right\|_{L^2}^2 + \|\bar{w}_{L_n}^J(0)\|_{L^2}^2 + o(1), \quad n \rightarrow \infty.$$

Remark 3.11. Let $2 < q < q_0 = \frac{2d}{d-2s_0}$, and τ^j be defined by (3.11). Assume as before $\tau^j \in \{0, \pm\infty\}$. Then if $\tau_j \in \{\pm\infty\}$, one has, by standard properties of the linear Schrödinger equation,

$$(3.19) \quad \lim_{n \rightarrow \infty} \|\varphi_{L_n}^j(0)\|_{L^q} = 0.$$

Moreover, if $j \in \mathcal{J}_C$, then (3.19) holds by a simple scaling argument. Finally, using Lemma 3.7 and the same argument as in Remark 3.10, one obtain

$$\lim_{n \rightarrow \infty} \|u_{0,n}\|_{L^q}^q = \sum_{\substack{j \in \mathcal{J}_{NC} \\ \tau^j = 0}} \|\varphi_L^j(0)\|_{L^q}^q.$$

3.3. Nonlinear profile decomposition. We now construct a nonlinear profile decomposition, based on the preceding linear profile decomposition, and adapted to the equation (2.1), where g satisfy the following assumptions.

Assumption B.

$$(3.20) \quad g(u) = g_0(u) + g_1(u), \quad g_0(u) = \iota_0 |u|^{p_0} u, \quad \iota_0 \in \{\pm 1\}$$

and $g_1(u)$ is a nonlinearity of lower order. Precisely, we will assume

$$(3.21) \quad g_1 \in \mathcal{N}(s_0, p_2, p_1), \quad \frac{4}{d} < p_1 \leq p_2 < p_0, \quad 1 < p_1,$$

where as usual $s_0 = \frac{d}{2} - \frac{2}{p_0}$, and

$$(3.22) \quad [s_0] \leq p_0, \quad \text{or } g \text{ is a polynomial in } u, \bar{u}.$$

Furthermore, the nonlinearity is of the form $g(z) = G(|z|^2)z$, $G \in C^1([0, \infty), \mathbb{R})$ with $G(0) = 0$.

We note that the last condition on g implies conservation of the mass, and, if $s_0 \geq 1$, of the energy and the momentum. Note that Assumption B implies Assumption A with $s = s_0$.

We will also consider the equation (2.1) with $g = g_0$, that is the homogeneous nonlinear Schrödinger equation (1.5) with initial data in \dot{H}^{s_0} .

To each (linear) profile $\varphi_{L_n}^j$, we associate a *nonlinear profile* φ_n^j and a *modified nonlinear profile* $\tilde{\varphi}_n^j$ in the following way:

- If $j \in \mathcal{J}_0$, $\tilde{\varphi}_n^j$ and φ_n^j are both equal to the constant null function.
- If $j \in \mathcal{J}_{NC}$, the modified nonlinear profile and the nonlinear profile are equal, and defined by

$$\tilde{\varphi}_n^j(t, x) = \varphi_n^j(t, x) = \varphi^j(t - t_n^j, x - x_n^j),$$

where φ^j is the unique solution of (2.1) such that

$$(3.23) \quad \lim_{t \rightarrow \tau^j} \left\| \varphi^j(t) - \varphi_L^j(t) \right\|_{H^{s_0}} = 0, \quad \tau^j = \lim_{n \rightarrow \infty} -t_n^j.$$

This solution is given by the well-posedness theory, Proposition 2.12 (if τ^j is finite) or by the existence of wave operators, Proposition 2.18 (if $\tau^j \in \{\pm\infty\}$).

- If $j \in \mathcal{J}_C$, the nonlinear profile φ_n^j is defined by

$$(3.24) \quad \varphi_n^j(t, x) = \frac{1}{(\lambda_n^j)^{\frac{2}{p_0}}} \varphi^j \left(\frac{t - t_n^j}{(\lambda_n^j)^2}, \frac{x - x_n^j}{\lambda_n^j} \right)$$

where φ^j is the unique solution of the homogeneous equation (1.5) such that

$$\lim_{t \rightarrow \tau^j} \left\| \varphi^j(t) - \varphi_L^j(t) \right\|_{\dot{H}^{s_0}} = 0, \quad \tau^j = \lim_{n \rightarrow \infty} \frac{-t_{j,n}}{(\lambda_n^j)^2}.$$

By definition of φ^j , we see that $\varphi^j(\tau)$ is in \dot{H}^{s_0} for all τ in the domain of existence of φ^j . However it is not necessarily in H^{s_0} . To tackle with this difficulty, we fix σ^j in the maximal interval of existence of φ^j (if τ^j is finite, we can take $\sigma^j = \tau^j$, if $\tau^j = \pm\infty$, $|\sigma^j|$ large and with the same sign than τ^j). We let

$$s_n^j = (\lambda_n^j)^2 \sigma^j + t_n^j$$

and denote by $\tilde{\varphi}_n^j$ the solution of (1.5) such that

$$(3.25) \quad \tilde{\varphi}_n^j(s_n^j) = \chi(x - x_n^j) \varphi_n^j(s_n^j) = \chi(x - x_n^j) \frac{1}{(\lambda_n^j)^{\frac{2}{p_0}}} \varphi^j \left(\sigma^j, \frac{x - x_n^j}{\lambda_n^j} \right),$$

where $\chi \in C_0^\infty(\mathbb{R}^N)$ is radially symmetric, $\chi(x) = 1$ for $|x| < 1$, $\chi(x) = 0$ for $|x| > 2$.

Lemma 3.12. *Let $j \in \mathcal{J}_C$. Then*

$$(3.26) \quad \lim_{n \rightarrow \infty} \left\| \tilde{\varphi}_n^j(0) - \varphi_n^j(0) \right\|_{\dot{H}^{s_0}} = 0.$$

More precisely, let $0 \in J_n$ be a sequence of interval. Let $I_n^j = \left\{ \frac{t - t_n^j}{(\lambda_n^j)^2}, t \in J_n \right\}$. Assume that there exists an interval I in the domain of existence of φ^j such that $\|\varphi^j\|_{X_{p_0}(I)} < \infty$ and for large n , $I_n^j \subset I$. Then for large n , J_n is included in the domain of existence of $\tilde{\varphi}_n^j$ and

$$(3.27) \quad \sup_{t \in J_n} \left\| \varphi_n^j(t) - \tilde{\varphi}_n^j(t) \right\|_{\dot{H}^{s_0}} + \left\| \varphi_n^j - \tilde{\varphi}_n^j \right\|_{\dot{S}^{s_0}(J_n)} \xrightarrow{n \rightarrow \infty} 0$$

$$(3.28) \quad \forall s \in [0, s_0), \quad \sup_{t \in J_n} \left\| \tilde{\varphi}_n^j(t) \right\|_{H^s} + \left\| \tilde{\varphi}_n^j \right\|_{S^s(J_n)} \xrightarrow{n \rightarrow \infty} 0$$

Proof. We first prove (3.27). Without loss of generality, we can assume $\sigma_j \in I$. Let

$$\tilde{\Phi}_n^j(\tau, y) = (\lambda_n^j)^{2/p} \tilde{\varphi}_n^j(t_n^j + (\lambda_n^j)^2 \tau, x_n^j + \lambda_n^j y).$$

Then

$$\tilde{\Phi}_n^j(\sigma_j, y) = (\lambda_n^j)^{\frac{2}{p_0}} \tilde{\varphi}_n^j(s_n^j, x_n^j + \lambda_n^j y) = \chi(\lambda_n^j y) \varphi^j(\sigma^j, y).$$

Thus, using that $\lambda_n^j \rightarrow 0$ as $n \rightarrow \infty$, we obtain that I is included in the domain of existence of $\tilde{\Phi}_n^j$ and

$$\lim_{n \rightarrow \infty} \left\| \tilde{\Phi}_n^j(\sigma^j) - \varphi^j(\sigma^j) \right\|_{\dot{H}^{s_0}} = 0.$$

Using the long time perturbation theory for equation (1.5) (see Theorem 2.19 and Remark 2.20), we obtain

$$\sup_{\tau \in I} \left\| \varphi^j(\tau) - \tilde{\Phi}_n^j(\tau) \right\|_{\dot{H}^{s_0}} + \left\| \varphi^j - \tilde{\Phi}_n^j \right\|_{\dot{S}^{s_0}(I)} \xrightarrow{n \rightarrow \infty} 0.$$

By the change of variable $\tau = \frac{t - t_{j,n}}{(\lambda_n^j)^2}$, $y = \frac{x - x_{j,n}}{\lambda_n^j}$, we obtain (3.27). Applying (3.27), we obtain (3.26).

To prove (3.28), we first notice that using (3.25) and $\lambda_n^j \rightarrow 0$,

$$\lim_{n \rightarrow \infty} \|\tilde{\varphi}_n^j(s_n^j)\|_{L^2} = 0,$$

By conservation of the L^2 norm, and interpolation with the bound of the \dot{H}^s norm which follows from (3.27)

$$(3.29) \quad \forall s \in [0, s_0), \quad \lim_{n \rightarrow \infty} \sup_{t \in J_n} \|\tilde{\varphi}_n^j\|_{\dot{H}^s} = 0.$$

It remains to prove the second limit in (3.28). We let $s \in [0, s_0)$. To any $J \subset J_n$ such that $\|\tilde{\varphi}_n^j\|_{X(J)} \leq \varepsilon$ (where ε is a small constant), and $a_n \in J$, we have by Strichartz estimates and Proposition 2.8,

$$\|\tilde{\varphi}_n^j\|_{S^s(J)} \lesssim \|\tilde{\varphi}_n^j(a_n)\|_{\dot{H}^s} + \|g_0(\tilde{\varphi}_n^j)\|_{N^s(J)} \lesssim \|\tilde{\varphi}_n^j(a_n)\|_{\dot{H}^s} + \|\tilde{\varphi}_n^j\|_{S^s(J)} \|\tilde{\varphi}_n^j\|_{X(J)}^{p_0},$$

and thus (if ε is small enough),

$$\|\tilde{\varphi}_n^j\|_{S^s(J)} \lesssim \|\tilde{\varphi}_n^j(a_n)\|_{\dot{H}^s}.$$

Since

$$\limsup_{n \rightarrow \infty} \|\tilde{\varphi}_n^j\|_{S^{s_0}(J_n)} < \infty,$$

and $S^{s_0}(J)$ is continuously embedded in $X(J)$, we can divide the interval J_n in N subintervals J_n^k , $k \in \llbracket 1, N \rrbracket$ (N independent of n), such that $\|\tilde{\varphi}_n^j\|_{X(J_n^k)} \leq \varepsilon$. Arguing as in the proof of the long-time perturbation theory result (Theorem 2.19), and using (3.29), we obtain (3.28). \square

When $j \in \mathcal{J}_C$, the modified profiles $\tilde{\varphi}_n^j$ are approximate solutions of (2.1):

Lemma 3.13. *Let $j \in \mathcal{J}_C$, and J_n be as in Lemma 3.12. Let*

$$(3.30) \quad \tilde{e}_n^j = i\partial_t \tilde{\varphi}_n^j + \Delta \tilde{\varphi}_n^j - g(\tilde{\varphi}_n^j) = -g_1(\tilde{\varphi}_n^j).$$

Then

$$\lim_{n \rightarrow \infty} \|\tilde{e}_n^j\|_{N^{s_0}(J_n)} = 0.$$

Proof. Since $\tilde{\varphi}_n^j$ is a solution of (1.5) with $p = p_0$, we indeed have $\tilde{e}_n^j = -g_1(\tilde{\varphi}_n^j)$. As a consequence, by Proposition 2.8, using that $g_1 \in \mathcal{N}(s_0, p_2, p_1)$,

$$(3.31) \quad \|\tilde{e}_n^j\|_{N^{s_0}(J_n)} \lesssim \|\tilde{\varphi}_n^j\|_{S^{s_0}(J_n)} \left(\|\tilde{\varphi}_n^j\|_{L^{p_1(d+2)/2}(J_n \times \mathbb{R}^d)}^{p_1} + \|\tilde{\varphi}_n^j\|_{L^{p_2(d+2)/2}(J_n \times \mathbb{R}^d)}^{p_2} \right) \\ \lesssim \|\tilde{\varphi}_n^j\|_{S^{s_0}(J_n)} \left(\|\tilde{\varphi}_n^j\|_{S^{s_1}(J_n)}^{p_1} + \|\tilde{\varphi}_n^j\|_{S^{s_2}(J_n)}^{p_2} \right),$$

where $s_k = \frac{d}{2} - \frac{2}{p_k}$, $k \in \{1, 2\}$. Since $s_1 < s_2 < s_0$, the conclusion of the Lemma follows from (3.28). \square

We next give the announced approximation result. For this we must also modify the linear remainder $w_{L,n}^J$: we let $\tilde{w}_{L,n}^J$ be the solution of the linear wave equation with initial data

$$(3.32) \quad \tilde{w}_{L,n}^J(0) = u_{0,n} - \sum_{j=1}^J \tilde{\varphi}_n^j(0).$$

Claim 3.14. For all s with $0 < s \leq s_0$,

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|\tilde{w}_{L,n}^J\|_{X(\mathbb{R})} + \|\tilde{w}_{L,n}^J\|_{\dot{W}^s(\mathbb{R})} = 0.$$

Proof. We have

$$\tilde{w}_{L,n}^J(0) = w_{L,n}^J(0) + \sum_{1 \leq j \leq J} (\varphi_{L,n}^j(0) - \tilde{\varphi}_n^j(0)).$$

By Lemma 3.12 and linear profile decomposition, we have

$$\lim_{n \rightarrow \infty} \left\| \varphi_{L,n}^j(0) - \tilde{\varphi}_n^j(0) \right\|_{\dot{H}^{s_0}} \leq \lim_{n \rightarrow \infty} \left\| \varphi_{L,n}^j(0) - \varphi_n^j(0) \right\|_{\dot{H}^{s_0}} + \lim_{n \rightarrow \infty} \left\| \varphi_n^j(0) - \tilde{\varphi}_n^j(0) \right\|_{\dot{H}^{s_0}} \rightarrow 0.$$

Combining with (3.5), we obtain

$$(3.33) \quad \lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|\tilde{w}_{L,n}^J\|_{X_{p_0}(\mathbb{R})} + \|\nabla^{s_0} \tilde{w}_{L,n}^J\|_{W^0(\mathbb{R})} = 0.$$

By Lemma 3.12, we also have

$$\tilde{w}_{L,n}^J(0) = u_{0,n} - \sum_{\substack{j \in \mathcal{J}_{NC} \\ 1 \leq j \leq J}} \varphi_n^j(0) + o_n(1), \text{ in } H^s, 0 \leq s < s_0,$$

which shows by the Pythagorean expansion (3.18) that

$$\sup_J \limsup_{n \rightarrow \infty} \|\tilde{w}_{L,n}^J(0)\|_{L^2} < \infty.$$

Using Strichartz estimates, we obtain

$$(3.34) \quad \lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|\tilde{w}_{L,n}^J\|_{W^0(\mathbb{R})} < \infty,$$

which yields, combining with (3.33), the conclusion of the claim. \square

Theorem 3.15 (Approximation by profiles). *Let $(u_{0,n})_n$ be a sequence bounded in H^{s_0} that admits a profile decomposition $(\varphi_{L_n}^j)_j$. Define as above the nonlinear profiles φ_n^j , the modified nonlinear profiles $\tilde{\varphi}_n^j$ and the modified remainder $\tilde{w}_{L,n}^J$. Let I_n be a sequence of intervals such that $0 \in I_n$, and assume that for each $j \geq 1$, for large n , $0 \in I_n \subset I_{\max}(\varphi_n^j)$,*

$$(3.35) \quad j \in \mathcal{J}_C \implies \limsup_{n \rightarrow \infty} \|\nabla^{s_0} \varphi_n^j\|_{S^0(I_n)} < \infty,$$

$$(3.36) \quad j \in \mathcal{J}_{NC} \implies \limsup_{n \rightarrow \infty} \|\varphi_n^j\|_{S^{s_0}(I_n)} < \infty.$$

Let u_n be the solution of (2.1) with initial data $u_{0,n}$. Then for large n , $I_n \subset I_{\max}(u_n)$,

$$u_n(t) = \sum_{1 \leq j \leq J} \tilde{\varphi}_n^j(t, x) + \tilde{w}_{L,n}^J(t) + r_n^J(t), \quad t \in I_n,$$

with

$$\limsup_{n \rightarrow \infty} \|u_n\|_{S^{s_0}(I_n)} < \infty$$

and

$$(3.37) \quad \lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|r_n^J\|_{S^{s_0}(I_n)} = 0.$$

We will also need the fact that Pythagorean expansions of the Sobolev norms hold in the setting of the preceding Theorem:

Lemma 3.16. *With the same assumptions and notations as in Theorem 3.15, if $(t_n)_n$ is a sequence of time with $t_n \in I_n$ for all time, then for all $J \geq 1$,*

$$\|u_n(t_n)\|_{\dot{H}^{s_0}}^2 = \sum_{j=1}^J \|\tilde{\varphi}_n^j(t_n)\|_{\dot{H}^{s_0}}^2 + \|\tilde{w}_n^J(t_n)\|_{\dot{H}^{s_0}}^2 + o_{J,n}(1),$$

where $\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} o_{J,n}(1) = 0$. Furthermore, for all s with $0 \leq s < s_0$,

$$\|u_n(t_n)\|_{\dot{H}^s}^2 = \sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} \|\tilde{\varphi}_n^j(t_n)\|_{\dot{H}^s}^2 + \|\tilde{w}_n^J(t_n)\|_{\dot{H}^s}^2 + o_{J,n}(1),$$

Before proving Theorem 3.15 and Lemma 3.16, we need two technical lemmas.

Lemma 3.17. *Let (q, r) be a Schrödinger admissible pair with q, r finite, φ such that $|D|^{s_0} \varphi \in L^q L^r(\mathbb{R} \times \mathbb{R}^d)$, $\psi \in C_0^\infty(\mathbb{R} \times \mathbb{R}^d)$. Let $\Lambda_n = (\lambda_n, t_n, x_n)$ and $M_n = (\mu_n, s_n, y_n)$ be two sequences of transformations that are orthogonal in the sense of Definition 3.2. Let*

$$\varphi_n(t, x) = \frac{1}{\lambda_n^{p_0}} \varphi\left(\frac{t-t_n}{\lambda_n^2}, \frac{x-x_n}{\lambda_n}\right), \quad \psi_n(t, x) = \psi\left(\frac{t-s_n}{\mu_n^2}, \frac{x-y_n}{\mu_n}\right).$$

Assume that $(\mu_n/\lambda_n)_n$ is bounded. Then,

$$\lim_{n \rightarrow \infty} \left\| |\nabla|^{s_0} (\varphi_n \psi_n) \right\|_{L^q L^r} = 0.$$

Proof. By density (since (q, r) are finite), we can assume $\varphi \in C_0^\infty(\mathbb{R} \times \mathbb{R}^d)$. Rescaling and translating $\varphi_n \psi_n$, we can assume (since μ_n/λ_n is bounded) that one of the following holds:

- $\lim_{n \rightarrow \infty} \lambda_n = \lim_{n \rightarrow \infty} \mu_n = 1$. In this case, the fact that the two sequences are orthogonal implies that $\varphi_n \psi_n = 0$ for large n .
- $\lim_n \lambda_n = \infty$ and $\forall n, \mu_n = 1, s_n = 0, y_n = 0$. In this case, we have

$$|\varphi_n \psi_n(t, x)| \leq \frac{1}{\lambda_n^{2/p_0}} \|\varphi\|_\infty |\psi(t, x)|$$

and thus $\varphi_n \psi_n$ goes to 0 in $L^q L^r$ as $n \rightarrow \infty$. The same argument proves that for all $\alpha \in \mathbb{N}, \beta \in \mathbb{N}^d, \partial_t^\alpha \partial_x^\beta (\varphi_n \psi_n)$ goes to 0 in $L^q L^r$ as $n \rightarrow \infty$. Interpolating we obtain the conclusion of the lemma. \square

Lemma 3.18. *Let $(\Lambda_n^j)_n = (\lambda_n^j, t_n^j, x_n^j)_n, 1 \leq j \leq J$ be a family of sequences of transformations that are pairwise orthogonal. For $j \in \llbracket 1, J \rrbracket$, we let φ^j such that $|\nabla|^{s_0} \varphi^j \in S^0(\mathbb{R})$, and let φ_n^j be defined by (3.24). Then*

$$(3.38) \quad \lim_{n \rightarrow \infty} \left\| \sum_{j=1}^J \varphi_n^j \right\|_{X_{p_0}(\mathbb{R})}^{\frac{p_0(d+2)}{2}} = \sum_{j=1}^J \left\| \varphi^j \right\|_{X_{p_0}(\mathbb{R})}^{\frac{p_0(d+2)}{2}}$$

$$(3.39) \quad \lim_{n \rightarrow \infty} \left\| \sum_{j=1}^J |\nabla|^{s_0} \varphi_n^j \right\|_{W^0(\mathbb{R})}^{\frac{2(d+2)}{d}} = \sum_{j=1}^J \left\| |\nabla|^{s_0} \varphi^j \right\|_{W^0(\mathbb{R})}^{\frac{2(d+2)}{d}}$$

$$(3.40) \quad \lim_{n \rightarrow \infty} \left\| g_0 \left(\sum_{j=1}^J \varphi_n^j \right) - \sum_{j=1}^J g_0(\varphi_n^j) \right\|_{\dot{N}^{s_0}(\mathbb{R})} = 0.$$

Furthermore, assuming that $\lambda_n^j = 1$ and that $\varphi^j \in S^{s_0}(\mathbb{R})$ for all n , one has

$$(3.41) \quad \lim_{n \rightarrow \infty} \left\| \sum_{j=1}^J \varphi_n^j \right\|_{X_{p_1}(I)}^{\frac{p_1(d+2)}{2}} - \sum_{j=1}^J \left\| \varphi_n^j \right\|_{X_{p_1}(I)}^{\frac{p_1(d+2)}{2}} = 0$$

$$(3.42) \quad \lim_{n \rightarrow \infty} \left\| \sum_{j=1}^J \varphi_n^j \right\|_{W^0(\mathbb{R})}^{\frac{2(d+2)}{d}} - \sum_{j=1}^J \left\| \varphi_n^j \right\|_{W^0(\mathbb{R})}^{\frac{2(d+2)}{d}} = 0$$

$$(3.43) \quad \lim_{n \rightarrow \infty} \left\| g \left(\sum_{j=1}^J \varphi_n^j \right) - \sum_{j=1}^J g(\varphi_n^j) \right\|_{N^{s_0}(\mathbb{R})} = 0.$$

Proof. Proof of (3.41), (3.42) and (3.43). By a density argument (and Proposition 2.8 for the proof of (3.43)), we can assume that $\varphi^j \in C_0^\infty(\mathbb{R} \times \mathbb{R}^d)$. As a consequence of the orthogonality of the sequences Λ_n^j , we deduce that for large n , the supports of the functions $\varphi_n^j, j \in \llbracket 1, J \rrbracket$ are two-by-two disjoint. The three estimates follow immediately.

Proof of (3.40). By a density argument and (2.10), we can assume $\varphi^j \in C_0^\infty(\mathbb{R} \times \mathbb{R}^d)$. We argue by induction on J . We fix $J \geq 2$. Arguing by contradiction, reordering the profiles and extracting subsequences, we see that we can assume

$$\forall n, \quad \lambda_n^J = \min_{j \in \llbracket 1, J \rrbracket} \lambda_n^j.$$

It is sufficient to prove:

$$(3.44) \quad \lim_{n \rightarrow \infty} \left\| g_0 \left(\sum_{j=1}^J \varphi_n^j \right) - g_0 \left(\sum_{j=1}^{J-1} \varphi_n^j \right) - g_0(\varphi_n^J) \right\|_{\dot{N}^{s_0}(\mathbb{R})} = 0.$$

We let $\chi \in C_0^\infty(\mathbb{R} \times \mathbb{R}^d)$, with $\chi = 1$ on the support of φ^J . We let $\chi_n(t, x) = \chi\left(\frac{t-t_n^J}{(\lambda_n^J)^2}, \frac{x-x_n^J}{\lambda_n^J}\right)$. By Lemma 3.17, and Lemma 2.9 or Lemma 2.10 (see (2.10)),

$$\begin{aligned} g_0\left(\sum_{j=1}^J \varphi_n^j\right) &= g_0\left((1-\chi_n)\sum_{j=1}^{J-1} \varphi_n^j + \varphi_n^J\right) + o_n(1) \text{ in } \dot{N}^{s_0} \\ g_0\left(\sum_{j=1}^{J-1} \varphi_n^j\right) &= g_0\left((1-\chi_n)\sum_{j=1}^{J-1} \varphi_n^j\right) + o_n(1) \text{ in } \dot{N}^{s_0}. \end{aligned}$$

Next, we note that, since $g_0(0) = 0$ and the supports of $(1-\chi_n)\sum_{j=1}^{J-1} \varphi_n^j$ and φ_n^J are disjoint, one has

$$g_0\left((1-\chi_n)\sum_{j=1}^{J-1} \varphi_n^j + \varphi_n^J\right) = g_0\left((1-\chi_n)\sum_{j=1}^{J-1} \varphi_n^j\right) + g_0(\varphi_n^J).$$

Combining the preceding estimates, we obtain (3.44), and hence (3.40). We omit the similar proofs of (3.38) and (3.39). \square

Proof of Theorem 3.15. We let, for $t \in I_n$,

$$(3.45) \quad v_n^J(t) = \sum_{1 \leq j \leq J} \tilde{\varphi}_n^J(t, x) + \tilde{w}_{L,n}^J(t, x), \quad e_n^J = Lv_n^J - g(v_n^J).$$

We will prove that there exists a constant $C > 0$ such that

$$(3.46) \quad \forall J \geq 1, \quad \limsup_{n \rightarrow \infty} \|v_n^J\|_{X(I_n)} + \|v_n^J\|_{W^{s_0}(I_n)} \leq C,$$

and that for all $\delta > 0$, there exists J_δ such that

$$(3.47) \quad \forall J \geq J_\delta, \quad \limsup_{n \rightarrow \infty} \|e_n^J\|_{N^{s_0}(I_n)} \leq \delta.$$

By the definitions (3.6) and (3.32) of $w_{L,n}^J$ and $\tilde{w}_{L,n}^J$, we have $v_n^J(0) = u_{0,n}$. This implies, together with (3.46) and the assumption that $(u_{0,n})_n$ is bounded in H^{s_0} that there exists a constant $C > 0$ independent of J such that

$$\forall J, \quad \limsup_{n \rightarrow \infty} \|v_n^J(0)\|_{H^{s_0}} \leq C.$$

Thus we see that for $J \geq J_\delta$, and n large enough, the assumptions of Theorem 2.19 are satisfied. The conclusion of Theorem 3.15 follows. We are left with proving (3.46) and (3.47). In all the proof, we will denote by C a large positive constant that may change from line to line, might depend on the sequence $(u_n)_n$, but is *independent of J* .

Proof of (3.46). We note that it is sufficient to prove the bound in (3.46) for J large.

By the Pythagorean expansion (3.7) we have,

$$\forall J \geq 1, \quad \sum_{j=1}^J \|\varphi_L^j(0)\|_{\dot{H}^{s_0}}^2 \leq \limsup_{n \rightarrow \infty} \|u_n(0)\|_{\dot{H}^{s_0}}^2 \leq C.$$

Hence $\sum_{j=1}^\infty \|\varphi_L^j(0)\|_{\dot{H}^{s_0}}^2 < \infty$. Letting ε_0 be a small positive number, we see that there exists $J_0 \geq 1$ such that

$$(3.48) \quad \sum_{j=J_0}^\infty \|\varphi_L^j(0)\|_{\dot{H}^{s_0}}^2 \leq \varepsilon_0.$$

Using the small data theory for equation (1.5) (see Proposition 2.12 and Remark 2.20), we deduce that if $j \in \mathcal{J}_C$ and $j \geq J_0$, φ^j is global and

$$(3.49) \quad \sum_{\substack{j \geq J_0 \\ j \in \mathcal{J}_C}} \|\varphi^j\|_{\dot{S}^{s_0}(\mathbb{R})}^2 < \infty$$

Arguing similarly with the Pythagorean expansion of the H^{s_0} norm given by (3.7), (3.18), we obtain (taking a larger J_0 if necessary) that if $j \in \mathcal{J}_{NC}$ and $j \geq J_0$, φ^j is global and

$$(3.50) \quad \sum_{\substack{j \geq J_0 \\ j \in \mathcal{J}_{NC}}} \|\varphi^j\|_{S^{s_0}(\mathbb{R})}^2 < \infty.$$

Next, we see that the assumptions of the theorem implies that for all $1 \leq j \leq J_0 - 1$, there exists an interval $I^j \subset I_{\max}(\varphi^j)$ such that for large n , $\left\{ \frac{t-t_n^j}{(\lambda_n^j)^2}, t \in I_n \right\} \subset I^j$ and

$$\begin{cases} \|\varphi^j\|_{\dot{S}^{s_0}(I^j)} < \infty & \text{if } j \in \mathcal{J}_C \\ \|\varphi^j\|_{S^{s_0}(I^j)} < \infty & \text{if } j \in \mathcal{J}_{NC}. \end{cases}$$

We let $\bar{\varphi}^j(t, x) = \varphi^j(t, x)$, if $t \in I^j$, $\bar{\varphi}^j(t, x) = 0$ if $t \in \mathbb{R} \setminus I^j$. We denote by $\bar{\varphi}_n^j$ the corresponding modulated profiles, defined similarly as in (3.24). By (3.49), (3.50), the definition of $\bar{\varphi}_n^j$ and Lemma 3.18, we obtain, for $J \geq J_0$.

$$(3.51) \quad \limsup_{n \rightarrow \infty} \left\| \sum_{j=1}^J \bar{\varphi}_n^j \right\|_{\dot{W}^{s_0}(I_n)}^{\frac{2(2+d)}{d}} = \limsup_{n \rightarrow \infty} \left\| \sum_{j=1}^J \varphi_n^j \right\|_{\dot{W}^{s_0}(I_n)}^{\frac{2(2+d)}{d}} \leq \limsup_{n \rightarrow \infty} \left\| \sum_{j=1}^{J_0-1} \bar{\varphi}_n^j + \sum_{j=J_0}^J \varphi_n^j \right\|_{\dot{W}^{s_0}(\mathbb{R})}^{\frac{2(2+d)}{d}} \\ = \sum_{j=1}^{J_0-1} \|\bar{\varphi}^j\|_{\dot{W}^{s_0}(\mathbb{R})}^{\frac{2(2+d)}{d}} + \sum_{j=J_0}^J \|\varphi^j\|_{\dot{W}^{s_0}(\mathbb{R})}^{\frac{2(2+d)}{d}} \leq C.$$

A similar argument yields $\limsup_{n \rightarrow \infty} \left\| \sum_{j=1}^J \bar{\varphi}_n^j \right\|_{X_{p_0}(I_n)} \leq C$. We also have, for $0 \leq s < s_0$, by (3.28) in Lemma 3.12,

$$(3.52) \quad \lim_{n \rightarrow \infty} \left\| \sum_{j=1}^J \bar{\varphi}_n^j \right\|_{W^s(I_n)}^{\frac{2(2+d)}{d}} = \lim_{n \rightarrow \infty} \left\| \sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} \varphi_n^j \right\|_{W^s(I_n)}^{\frac{2(2+d)}{d}},$$

and the same argument as above, using (3.50) and Lemma 3.18, yields $\limsup_{n \rightarrow \infty} \left\| \sum_{j=1}^J \bar{\varphi}_n^j \right\|_{W^s(I_n)} \leq C$. Combining the estimates above with Claim 3.14 and (3.34), we obtain (3.46).

Proof of (3.47). We have, by (3.45),

$$e_n^J = \overbrace{\sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} L\bar{\varphi}_n^j - g(\bar{\varphi}_n^j)}^0 + \overbrace{\sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_C}} L\bar{\varphi}_n^j - g_0(\bar{\varphi}_n^j)}^0 + \overbrace{L\tilde{w}_{L,n}^J - g(v_n^J)}^0 + \sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_C}} g_0(\bar{\varphi}_n^j) + \sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} g(\bar{\varphi}_n^j)$$

By Proposition 2.8, (3.34) and (3.46),

$$\left\| g(v_n^J) - g\left(\sum_{j=1}^J \bar{\varphi}_n^j\right) \right\|_{N^{s_0}(I_n)} \leq C \left(\|\tilde{w}_{L,n}^J\|_{X(I_n)} + \|\tilde{w}_{L,n}^J\|_{\dot{W}^{s_0}(I_n)} \right)$$

By Claim 3.14, there exists J_δ such that, for all $J \geq J_\delta$,

$$(3.53) \quad \forall J \geq J_\delta, \quad \limsup_{n \rightarrow \infty} \left\| g(v_n^J) - g\left(\sum_{j=1}^J \bar{\varphi}_n^j\right) \right\|_{N^{s_0}(I_n)} \leq \frac{\delta}{2}.$$

Using Lemma 2.9 or Lemma 2.10 together with Lemma 3.18:

$$(3.54) \quad g_0\left(\sum_{j=1}^J \bar{\varphi}_n^j\right) = g_0\left(\sum_{j=1}^J \varphi_n^j\right) + o_n(1) = \sum_{j=1}^J g_0(\varphi_n^j) + o_n(1) \text{ in } \dot{N}^{s_0}(I_n),$$

and similarly (using also Lemma 3.12),

$$(3.55) \quad g_0\left(\sum_{j=1}^J \bar{\varphi}_n^j\right) = g_0\left(\sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} \bar{\varphi}_n^j\right) + o_n(1) = \sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} g_0(\bar{\varphi}_n^j) + o_n(1) \\ = \sum_{1 \leq j \leq J} g_0(\bar{\varphi}_n^j) + o_n(1) \text{ in } N^0(I_n)$$

By Proposition 2.8 and Lemma 3.12, using that $g_1 \in \mathcal{N}(s_0, p_0, p_1)$, we obtain

$$\begin{aligned} & \left\| g_1 \left(\sum_{j=1}^J \tilde{\varphi}_n^j \right) - g_1 \left(\sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} \tilde{\varphi}_n^j \right) \right\|_{N^0(I_n)} \\ & \lesssim \sum_{j \in \mathcal{J}_C, 1 \leq j \leq J} \|\tilde{\varphi}_n^j\|_{X_{p_1}} \left(\sum_{1 \leq j \leq J} \|\tilde{\varphi}_n^j\|_{X_{p_1}} \right)^{p_1-1} \sum_{1 \leq j \leq J} \|\tilde{\varphi}_n^j\|_{W^0} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Combining with Lemma 3.13 and Lemma 3.18, we obtain

$$(3.56) \quad g_1 \left(\sum_{j=1}^J \tilde{\varphi}_n^j \right) = g_1 \left(\sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} \tilde{\varphi}_n^j \right) + o_n(1) = \sum_{\substack{1 \leq j \leq J \\ j \in \mathcal{J}_{NC}}} g_1 \left(\tilde{\varphi}_n^j \right) + o_n(1) \text{ in } N^{s_0}(I_n).$$

By (3.53), (3.54), (3.55) and (3.56), we obtain (3.47), which concludes the proof. \square

Sketch of proof of Lemma 3.16. By Theorem 3.15, $(u_n(t_n))_n$ is bounded in $H^{s_0}(\mathbb{R}^d)$, and

$$u_n(t_n) = \sum_{j=1}^J \tilde{\varphi}_n^j(t_n) + \tilde{w}_{L,n}^J(t_n) + r_n^J(t_n).$$

We will interpret this expansion as a profile decomposition of $u_n(t_n)$. Using the property of $\tilde{w}_{L,n}^J$ given by Claim 3.14, the property (3.37) of r_n^J and the bound of $\varphi_n^j - \tilde{\varphi}_n^j$, $j \in \mathcal{J}_C$ given by Lemma 3.12, we obtain

$$u_n(t_n, x) = \sum_{j=1}^J \varphi_n^j(t_n, x) + R_n^J(x),$$

where

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|e^{i \cdot \Delta} R_n^J\|_{\dot{W}^{s_0}(\mathbb{R}) \cap X_{p_0}(\mathbb{R})} = 0.$$

We have

$$\varphi_n^j(t_n, x) = \frac{1}{(\lambda_n^j)^{\frac{2}{p_0}}} \varphi^j \left(\frac{t_n - t_n^j}{(\lambda_n^j)^2}, \frac{x - x_n^j}{\lambda_n^j} \right).$$

Extracting subsequences, we can assume that $\frac{t_n - t_n^j}{(\lambda_n^j)^2}$ has a limit σ^j as $n \rightarrow \infty$. We define a new linear profile $\psi_{L,n}^j$ by

$$\psi_{L,n}^j(t, x) = \frac{1}{(\lambda_n^j)^{\frac{2}{p_0}}} \psi_L^j \left(\frac{t + t_n - t_n^j}{(\lambda_n^j)^2}, \frac{x - x_n^j}{\lambda_n^j} \right),$$

where ψ_L^j is the solution of the linear Schrödinger equation such that

$$\lim_{t \rightarrow \sigma^j} \left\| \psi_L^j(t) - \varphi^j(t) \right\|_{\dot{H}^{s_0}} = 0.$$

With these choice of ψ_L^j , we see that $\left((\psi_{L,n}^j)_n \right)_{j \geq 1}$ is a \dot{H}^{s_0} profile decomposition for the sequence $(u_n(t_n))_n$. The conclusion of the lemma follows from the Pythagorean expansions (3.7), (3.17) and (3.18). \square

4. GLOBAL WELL-POSEDNESS

In this section we prove our theorem on global well-posedness, Theorem 1.2.

We first observe that Property 1.1 is equivalent to the existence of uniform space-time bound for solutions of equation (1.5) that are bounded in critical norm.

Proposition 4.1. *Let $d \geq 2$, $p_0 > \frac{4}{d}$, $s_0 = \frac{d}{2} - \frac{2}{p_0}$. Assume Property 1.1. Then for all $A \in (0, A_0)$, there exists $\mathcal{F}(A) > 0$ such that for all interval $0 \in I$, for all solution $u \in C^0(I, \dot{H}^{s_0})$ of (1.5) such that*

$$(4.1) \quad \sup_{t \in I} \|u(t)\|_{\dot{H}^{s_0}} \leq A$$

we have

$$(4.2) \quad \|u\|_{\dot{S}^{s_0}(I)} \leq \mathcal{F}(A).$$

Sketch of proof. The proof is by contradiction. Let us denote by $\mathcal{P}(A)$ the property that there exists $\mathcal{F}(A) \in (0, \infty)$ such that (4.1) implies (4.2). By the small data theory for equation (1.5), $\mathcal{P}(A)$ holds for small $A > 0$. Assuming that it does not hold for all $A \in (0, A_0)$, we obtain the existence of a critical $A_c \in (0, A_0)$ such that $\mathcal{P}(A)$ holds for $A < A_c$, but $\mathcal{P}(A_c)$ does not hold. Thus there exists a sequence of intervals $I_n = (a_n, b_n) \ni 0$, and of solutions $u_n \in C^0((a_n, b_n), \dot{H}^{s_0})$ such

$$\sup_{a_n < t < b_n} \|u_n(t)\|_{\dot{H}^{s_0}} \xrightarrow{n \rightarrow \infty} A_c,$$

and

$$\lim_{n \rightarrow \infty} \|u_n\|_{\dot{S}^{s_0}(a_n, 0)} = \lim_{n \rightarrow \infty} \|u_n\|_{\dot{S}^{s_0}(0, b_n)} = +\infty.$$

By a standard compactness argument, using the homogeneous profile decomposition of Subsection 3.1, with the analog, for the homogeneous equation, of Theorem 3.15, we obtain, after extraction of subsequences, that there exists $x_n \in \mathbb{R}^N$, $\lambda_n > 0$ and $\varphi_0 \in \dot{H}^{s_0} \setminus \{0\}$ such that

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{\lambda_n^{\frac{p_0}{2}}} u_n \left(0, \frac{\cdot - x_n}{\lambda_n} \right) - \varphi_0 \right\|_{\dot{H}^{s_0}} = 0,$$

and the solution φ of (1.5) with initial data φ_0 satisfies that for all $t \in \mathbb{R}$ there exist $x(t) \in \mathbb{R}^N$, $\lambda(t) > 0$ such that

$$\left\{ \frac{1}{\lambda(t)^{\frac{p_0}{2}}} \varphi \left(t, \frac{\cdot - x(t)}{\lambda(t)} \right), t \in I_{\max}(\varphi) \right\}$$

has compact closure in \dot{H}^{s_0} and

$$\sup_{t \in I_{\max}} \|\varphi(t)\|_{\dot{H}^{s_0}} \leq A_c$$

(see the similar proof of Theorem 6.5 below). Then φ is global by Property 1.1. Since φ is not the zero solution, the preceding compactness property implies $\liminf_{t \rightarrow \infty} \|\varphi(t)\|_{L^{\frac{d_{p_0}}{2}}} > 0$, a contradiction with the fact that by Property 1.1, φ must be scattering. The proof is complete. \square

Proof of Theorem 1.2. We argue by contradiction, assuming that there exists a solution u of (2.1) such that $T_+(u) < \infty$ and

$$\limsup_{n \rightarrow \infty} \|u(t)\|_{\dot{H}^{s_0}} < m \in (0, A_0).$$

Let ε_0 be such that $m + 3\varepsilon_0 < A_0$. By conservation of mass, we have indeed,

$$(4.3) \quad \limsup_{t \rightarrow T^+(u)} \|u(t)\|_{\dot{H}^{s_0}} < \infty.$$

Let $t_n = T_+(u) - 1/2^n$. Extracting subsequences, we can assume by Proposition 3.5 that $(u(t_n))_n$ admits a profile decomposition $((\varphi_{L,n}^j)_n)_{j \geq 1}$. By the Pythagorean expansion (3.7) of the \dot{H}^{s_0} norm, we have

$$(4.4) \quad \sum_{j \geq 1} \|\varphi_{L,n}^j(0)\|_{\dot{H}^{s_0}}^2 \leq m.$$

By the Pythagorean expansion of the L^2 norm,

$$(4.5) \quad \sum_{j \geq 1} \|\varphi_{L,n}^j(0)\|_{L^2}^2 < \infty.$$

We denote by φ_n^j the nonlinear profiles associated to the preceding profile decomposition. We will prove that for every $j \in \mathcal{J}_c$,

$$(4.6) \quad \sup_{0 \leq \tau < T_+ - t_n} \|\varphi_n^j(\tau)\|_{\dot{H}^{s_0}} \leq m + 2\varepsilon_0, \quad n \gg_j 1.$$

Let $\varepsilon > 0$ be a small constant. By (4.4), (4.5) and the small data theory for equations (1.5) and (2.1) there exists $J_0 \geq 1$ such that for $j \geq J_0 + 1$, φ^j is global and

$$(4.7) \quad \varepsilon \geq \begin{cases} \|\varphi^j\|_{S^{s_0}(\mathbb{R})} & \text{if } j \in \mathcal{J}_{NC} \\ \|\varphi^j\|_{\dot{S}^{s_0}(\mathbb{R})} & \text{if } j \in \mathcal{J}_C. \end{cases}$$

In particular (4.6) is satisfied for $j \in \mathcal{J}_C$, $j \geq J_0 + 1$. We next prove by contradiction that (4.6) holds for $j \in \mathcal{J}_C \cap \llbracket 1, J_0 \rrbracket$. If not, by (4.4), there exists $\tau'_n \in [0, T_+ - t_n)$ such that

$$(4.8) \quad \sup_{j \in \llbracket 1, J_0 \rrbracket \cap \mathcal{J}_C} \sup_{0 \leq \tau \leq \tau'_n} \|\varphi_n^j(\tau)\|_{\dot{H}^{s_0}} = m + \varepsilon_0 \in (0, A_0).$$

By the local well-posedness theory for equation (2.1) and the fact that $\lambda_n^j = 1$ for $j \in \mathcal{J}_{NC}$, there exists $\tau_0 > 0$ such that

$$(4.9) \quad \limsup_{n \rightarrow \infty} \sup_{j \in \llbracket 1, J_0 \rrbracket \cap \mathcal{J}_{NC}} \|\varphi_n^j\|_{S^{s_0}(0, \tau_0)} < \infty.$$

Since $m + \varepsilon_0 < A_0$, by (4.8) and Proposition 4.1, we obtain

$$\forall j \in \llbracket 1, J_0 \rrbracket \cap \mathcal{J}_C, \quad \limsup_{n \rightarrow \infty} \|\varphi_n^j\|_{\dot{S}^{s_0}(0, \tau'_n)} < \infty$$

Note that since $\tau'_n \rightarrow 0$ as $n \rightarrow \infty$, we have $\tau_0 > \tau'_n$ for large n . Combining with (4.7) and (4.9), we see that the assumptions of Theorem 3.15 are satisfied on the interval $I_n = [0, \tau'_n)$. By Lemma 3.16, for all sequence $(\sigma_n)_n$ with $0 \leq \sigma_n \leq \tau'_n$, for all J ,

$$\limsup_n \sum_{j \in \llbracket 1, J \rrbracket \cap \mathcal{J}_C} \|\varphi_n^j(\sigma_n)\|_{\dot{H}^{s_2}} \leq m.$$

This clearly contradicts (4.8), proving (4.6).

Next, we observe that (4.6) implies by Proposition 4.1

$$\forall j \in \llbracket 1, J_0 \rrbracket \cap \mathcal{J}_C, \quad \limsup_{n \rightarrow \infty} \|\varphi_n^j\|_{\dot{S}^{s_0}(0, T_+ - t_n)} < \infty$$

Combining this information with (4.7) and (4.9), we see that the Assumptions of Theorem 3.15 are satisfied on the interval $I_n = [0, T_+ - t_n)$. By the conclusion of the theorem, we obtain that for large n , $u(\cdot + t_n) \in S^{s_0}((0, T_+ - t_n))$. This implies $u \in S^{s_0}((0, T_+))$, contradicting the blow-up criterion for equation (2.1). The proof is complete. \square

5. GENERAL RIGIDITY RESULT

In this section, we consider equation (2.1), where g satisfies Assumption A p. 8, and is of the form $g(u) = G'(|u|^2)u$ for some C^1 function G . We recall that with these assumptions, the mass $M(u)$, the energy $E(u)$ and the momentum $P(u)$ are conserved for $H^1 \cap H^{s_0}$ solutions of (2.1), where as usual $s_0 = \frac{d}{2} - \frac{2}{p_0}$. We will also consider the virial functional:

$$(5.1) \quad \Phi(u) = \int |\nabla u|^2 + \frac{d}{2} \int (G'(|u|^2)|u|^2 - G(|u|^2)).$$

Since the assumptions on g imply $G'(|u|^2)|u|^2 + G(|u|^2) \lesssim |u|^{p_0+2} + |u|^{p_1+2}$, one easily checks, using Sobolev inequalities, that Φ is well-defined if $u \in H^{s_0} \cap H^1$.

We prove the following result:

Proposition 5.1. *With the assumptions above, let u be a solution of (2.1) defined on $[0, \infty)$ such that there exists $x(t)$, $t \in [0, \infty)$ with*

$$(5.2) \quad K = \{u(t, x + x(t)); t \geq 0\}$$

has compact closure in $H^{s_0} \cap H^1$. Then

$$(5.3) \quad \min_{t \geq 0} \left| \Phi(u(t)) - \frac{|P(u)|^2}{M(u)} \right| = 0.$$

Define

$$(5.4) \quad X(t) = 2 \frac{P(u)}{M(u)} t,$$

where $P(u)$ is momentum. The proof of Proposition 5.1 relies on an asymptotic estimate of $x(t)$ and a localized virial argument. We start with two lemmas.

Lemma 5.2. *With the assumptions above, let u be a solution (2.1) such that there exists $x(t)$, $t \geq 0$ such that $K = \{u(t, x + x(t)); t \geq 0\}$ has compact closure in H^1 . Then*

$$(5.5) \quad \lim_{t \rightarrow +\infty} \frac{|x(t) - X(t)|}{t} = 0.$$

Proof. We can assume that x is continuous (see e.g [17, Proposition 3.2]).

We argue by contradiction, assuming that there exists a sequence $t_n \rightarrow +\infty$, $\varepsilon_0 > 0$ such that

$$(5.6) \quad \frac{|x(t_n) - X(t_n)|}{t_n} \geq \varepsilon_0.$$

Without loss of generality we may assume $x(0) = 0$.

For $R > 0$, we let

$$(5.7) \quad t_0(R) = \inf\{t \geq 0; |x(t) - X(t)| \geq R\}.$$

Since $x(0) = 0 = X(0)$ and $x(t) - X(t)$ is continuous, we have $t_0(R) > 0$.

We define $R_n = |x(t_n) - X(t_n)|$ and $\tilde{t}_n = t_0(R_n)$ so that $t_n \geq \tilde{t}_n$. Thus we have the following properties:

$$(5.8) \quad \forall 0 \leq t < \tilde{t}_n \Rightarrow |x(t) - X(t)| < R_n,$$

$$(5.9) \quad |x(\tilde{t}_n) - X(\tilde{t}_n)| = R_n,$$

$$(5.10) \quad \frac{R_n}{\tilde{t}_n} \geq \varepsilon_0,$$

where (5.10) follows from (5.6) and $t_n \geq \tilde{t}_n$.

By precompactness of K , for any $\varepsilon > 0$, there exists $R_0(\varepsilon) > 0$ such that for all $t \geq 0$:

$$(5.11) \quad \int_{|x-x(t)| \geq R_0(\varepsilon)} (|u|^2 + |\nabla u|^2) dx \leq \varepsilon.$$

Let $\tilde{R}_n = R_n + R_0(\varepsilon)$ (for ε small enough).

Let $\theta \in \mathcal{D}(\mathbb{R})$ be such that $\theta(x) = x$ for $|x| \leq 1$, $\theta(x) = 0$ for $|x| \geq 2$ and $\|\theta\|_{L^\infty} < 2$. We write $\varphi(x) = (\theta(x_1), \theta(x_2), \cdot, \cdot, \cdot, \theta(x_d))$. Thus, $\varphi(x) = x$ for $|x| \leq 1$ and $\|\varphi\|_{L^\infty} < 2d$.

We define

$$(5.12) \quad z_{\tilde{R}_n}(t) = \int_{\mathbb{R}^d} \tilde{R}_n \varphi \left(\frac{x - X(t)}{\tilde{R}_n} \right) |u(t, x)|^2 dx.$$

Let $t \in [0, \tilde{t}_n]$. We have $z'_{\tilde{R}_n}(t) = ([z'_{\tilde{R}_n}(t)]_1, [z'_{\tilde{R}_n}(t)]_2, \cdot, \cdot, \cdot, [z'_{\tilde{R}_n}(t)]_d)$, where

$$\begin{aligned} [z'_{\tilde{R}_n}(t)]_j &= -X'_j(t) \int_{\mathbb{R}^d} \theta' \left(\frac{x_j - X_j(t)}{\tilde{R}_n} \right) |u(t, x)|^2 dx \\ &\quad + 2\mathcal{I}m \int_{\mathbb{R}^d} \theta' \left(\frac{x_j - X_j(t)}{\tilde{R}_n} \right) \bar{u} u_j dx, \quad (\text{where } u_j = \partial_j u = \partial_{x_j} u). \end{aligned}$$

For $|x_j - X_j(t)| \geq \tilde{R}_n$, we have $|x(t) - X(t)| \geq \tilde{R}_n$ then $|x - x(t)| \geq \tilde{R}_n - R_n$ (by the definition of R_n and triangle inequality). Thus, $|x - x(t)| \geq R_0(\varepsilon)$. For $|x_j - X_j(t)| \leq \tilde{R}_n$, $\theta' \left(\frac{x_j - X_j(t)}{\tilde{R}_n} \right) = 1$. By (5.11) we deduce, for $t \in [0, \tilde{t}_n]$,

$$(5.13) \quad [z'_{\tilde{R}_n}(t)]_j = -\frac{2P_j(u)}{M(u)} M(u) + 2P_j(u) + O(\varepsilon) = O(\varepsilon).$$

Furthermore, we have

$$\begin{aligned} z_{\tilde{R}_n}(0) &= \int_{\mathbb{R}^d} \tilde{R}_n \varphi \left(\frac{x}{\tilde{R}_n} \right) |u_0(x)|^2 dx \\ &= \int_{|x| < R_0(\varepsilon)} \tilde{R}_n \varphi \left(\frac{x}{\tilde{R}_n} \right) |u_0(x)|^2 dx + \int_{|x| > R_0(\varepsilon)} \tilde{R}_n \varphi \left(\frac{x}{\tilde{R}_n} \right) |u_0(x)|^2 dx. \end{aligned}$$

Thus, for some constant $C > 0$,

$$(5.14) \quad |z_{\tilde{R}_n}(0)| \leq R_0(\varepsilon)M(u) + C\tilde{R}_n\varepsilon \leq 2R_0(\varepsilon)M(u) + CR_n\varepsilon,$$

where we have used for the first bound that $|\varphi(x)| \lesssim |x|$ and for the second bound that $\varphi \in L^\infty$. Furthermore,

$$\begin{aligned} z_{\tilde{R}_n}(\tilde{t}_n) &= \int_{|x-x(\tilde{t}_n)| \geq R_0(\varepsilon)} \tilde{R}_n \varphi \left(\frac{x-X(\tilde{t}_n)}{\tilde{R}_n} \right) |u(\tilde{t}_n, x)|^2 dx \\ &\quad + \int_{|x-x(\tilde{t}_n)| \leq R_0(\varepsilon)} \tilde{R}_n \varphi \left(\frac{x-X(\tilde{t}_n)}{\tilde{R}_n} \right) |u(\tilde{t}_n, x)|^2 dx \\ (5.15) \quad &= I + II. \end{aligned}$$

Using (5.11), we have

$$(5.16) \quad |I| \lesssim \tilde{R}_n\varepsilon.$$

Furthermore, in the integral defining II , we have:

$$\begin{aligned} |x - X(\tilde{t}_n)| &\leq |x - x(\tilde{t}_n)| + |x(\tilde{t}_n) - X(\tilde{t}_n)| \\ &\leq R_0(\varepsilon) + R_n = \tilde{R}_n, \end{aligned}$$

where we have used the definition of \tilde{R}_n .

We next write

$$\begin{aligned} II &= \int_{|x-x(\tilde{t}_n)| \leq R_0(\varepsilon)} (x - X(\tilde{t}_n)) |u(\tilde{t}_n, x)|^2 dx \\ &= (x(\tilde{t}_n) - X(\tilde{t}_n)) \int_{\mathbb{R}^d} |u(\tilde{t}_n, x)|^2 dx \\ &\quad - (x(\tilde{t}_n) - X(\tilde{t}_n)) \int_{|x-x(\tilde{t}_n)| \geq R_0(\varepsilon)} |u(\tilde{t}_n, x)|^2 dx + \int_{|x-x(\tilde{t}_n)| \leq R_0(\varepsilon)} (x - x(\tilde{t}_n)) |u(\tilde{t}_n, x)|^2 dx. \end{aligned}$$

By (5.9), (5.11), we have

$$(5.17) \quad |II| \geq R_n(M(u) - C\varepsilon) - R_0(\varepsilon)M(u).$$

By (5.15), (5.16) and (5.17), we obtain

$$|z_{\tilde{R}_n}(\tilde{t}_n)| \geq R_n(M(u) - C\varepsilon) - R_0(\varepsilon)M(u) - \tilde{R}_n\varepsilon.$$

Thus,

$$(5.18) \quad |z_{\tilde{R}_n}(\tilde{t}_n)| \geq R_n(M(u) - C\varepsilon) - 2R_0(\varepsilon)M(u),$$

where we chose $\varepsilon \ll M(u)$.

By (5.13), (5.14) and (5.18), we have

$$R_n M(u) \lesssim R_0(\varepsilon)M(u) + \varepsilon \tilde{t}_n.$$

By (5.10), we deduce:

$$R_n M(u) \lesssim R_0(\varepsilon)M(u) + \varepsilon \frac{R_n}{\varepsilon_0}.$$

Choosing $\varepsilon \ll \varepsilon_0 M(u)$, we obtain $R_n \lesssim R_0(\varepsilon)$.

Letting $n \rightarrow +\infty$, we obtain a contradiction. This completes the proof. \square

The second lemma concerns the derivative of the localized virial functional. We consider

$$W_R(t) = R\mathcal{I}m \int_{\mathbb{R}^d} \varphi \left(\frac{x - X(t)}{R} \right) \nabla u \bar{u} \, dx,$$

where $X(t)$ is defined by (5.4) and φ is as in the proof of Lemma 5.2. By the relative compactness of K and the continuous embedding of $H^1 \cap H^{s_0}$ into $L^{p_0+2} \cap L^{p_1+2}$, we have that for any $\varepsilon > 0$, there exists $R_1(\varepsilon)$ such that

$$(5.19) \quad \int_{|x-x(t)| \geq R_1(\varepsilon)} |\nabla u|^2 + |u|^2 + |G'(|u|^2)||u|^2 + |G(|u|^2)| \, dx \leq \varepsilon.$$

By Cauchy-Schwarz inequality, we have

$$(5.20) \quad |W_R(t)| \leq CR \|\nabla u(t)\|_2 \|u(t)\|_2 \lesssim R.$$

For convenience, we denote $\sum_{j=1}^d$ by \sum_j , $\partial_{x_j} u = \partial_j u$ by u_j , u_t by $\partial_t u$ and $\int_{\mathbb{R}^d}$ by \int . We have

$$W'_R(t) = \underbrace{\sum_j R\mathcal{I}m \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) \frac{-X'_j(t)}{R} u_j \bar{u} \, dx}_{A_1} + \underbrace{\sum_j R\mathcal{I}m \int \theta \left(\frac{x_j - X_j(t)}{R} \right) \partial_t (u_j \bar{u}) \, dx}_{A_2}.$$

We will prove the following result on the terms A_1 and A_2 :

Lemma 5.3. *Let $\varepsilon, \tilde{\varepsilon} > 0$ be small. There exist $L(\tilde{\varepsilon})$ depending on $\tilde{\varepsilon}$ such that for $R \geq 2R_1(\tilde{\varepsilon})$, $L(\tilde{\varepsilon}) < t < \frac{R}{2\tilde{\varepsilon}}$,*

$$A_1 = \frac{-2|P(u)|^2}{M(u)} + O(\varepsilon), \quad A_2 = 2\Phi(u) + O(\varepsilon).$$

Proof. Writing $\partial_t (u_j \bar{u}) = (u_t \bar{u} + u_j \bar{u}_t)$ and integrating by part, we obtain

$$\begin{aligned} A_2 &= \sum_j -2R\mathcal{I}m \int \theta \left(\frac{x_j - X_j(t)}{R} \right) \bar{u}_j u_t - \sum_j \mathcal{I}m \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) \bar{u} u_t \\ &= \sum_j C_j + \sum_j D_j = I + II. \end{aligned}$$

Using the equation (2.1), we obtain

$$\begin{aligned} C_j &= 2R\mathcal{R}e \int \theta \left(\frac{x_j - X_j(t)}{R} \right) \bar{u}_j (-\Delta u + G'(|u|^2)u) \, dx \\ &= 2R\mathcal{R}e \int \nabla \left(\theta \left(\frac{x_j - X_j(t)}{R} \right) \bar{u}_j \right) \nabla u + \theta \left(\frac{x_j - X_j(t)}{R} \right) \frac{1}{2} \partial_j G(|u|^2) \\ &= 2R \int \theta \left(\frac{x_j - X_j(t)}{R} \right) \partial_j \frac{|\nabla u|^2}{2} \, dx + 2 \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) |u_j|^2 \, dx \\ &\quad + 2R \int \theta \left(\frac{x_j - X_j(t)}{R} \right) \frac{1}{2} \partial_j G(|u|^2) \, dx. \end{aligned}$$

Summing up, we obtain

$$\begin{aligned} I &= -2 \int \sum_j \theta' \left(\frac{x_j - X_j(t)}{R} \right) \left(\frac{|\nabla u|^2}{2} + \frac{1}{2} G(|u|^2) \right) \, dx + 2 \sum_j \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) |u_j|^2 \, dx \\ &= -d \int |\nabla u|^2 + G(|u|^2) \, dx - 2 \sum_j \int \left(\theta' \left(\frac{x_j - X_j(t)}{R} \right) - 1 \right) \left(\frac{|\nabla u|^2}{2} + \frac{1}{2} G(|u|^2) \right) \, dx \\ &\quad + 2 \int |\nabla u|^2 \, dx + 2 \sum_j \int \left(\theta' \left(\frac{x_j - X_j(t)}{R} \right) - 1 \right) |u_j|^2 \, dx. \end{aligned}$$

Moreover

$$\begin{aligned}
D_j &= \mathcal{R}e \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) \bar{u} i u_t dx \\
&= \mathcal{R}e \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) \bar{u} (-\Delta u + G'(|u|^2)u) dx \\
&= \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) G'(|u|^2)|u|^2 dx - \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) \left(\Delta \frac{|u|^2}{2} - |\nabla u|^2 \right) dx \\
&= \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) (G'(|u|^2)|u|^2 + |\nabla u|^2) dx - \frac{1}{2R^2} \int \theta''' \left(\frac{x_j - X_j(t)}{R} \right) |u|^2 dx.
\end{aligned}$$

where we have used $\int \theta' \left(\frac{x_j - X_j(t)}{R} \right) \partial_{kk} \frac{|u|^2}{2} = 0$ for all $k \neq j$. Thus,

$$\begin{aligned}
II &= d \int |\nabla u|^2 + G'(|u|^2)|u|^2 dx \\
&\quad + \sum_j \int \left(\theta' \left(\frac{x_j - X_j(t)}{R} \right) - 1 \right) (|\nabla u|^2 + G'(|u|^2)|u|^2) dx - \frac{1}{2R^2} \int \theta''' \left(\frac{x_j - X_j(t)}{R} \right) |u|^2 dx.
\end{aligned}$$

Combining the above, we have

$$\begin{aligned}
(5.21) \quad A_2 &= 2\Phi(u) \\
&\quad - \sum_j \frac{1}{2R^2} \int \theta''' \left(\frac{x_j - X_j(t)}{R} \right) |u|^2 dx
\end{aligned}$$

$$(5.22) \quad + \sum_j \int \left(\theta' \left(\frac{x_j - X_j(t)}{R} \right) - 1 \right) (2|u_j|^2 + G'(|u|^2)|u|^2 - G(|u|^2)).$$

Applying Lemma 5.2, for each $\tilde{\varepsilon}$, there exists $L(\tilde{\varepsilon})$ such that

$$(5.23) \quad |x(t) - X(t)| \leq \tilde{\varepsilon}t, \quad \forall t \geq L(\tilde{\varepsilon}).$$

Assume $L(\tilde{\varepsilon}) < t < \frac{R}{2\tilde{\varepsilon}}$ and $R \geq 2R_1(\varepsilon)$. Then

$$|(5.21)| + |(5.22)| \lesssim \int_{|x-X(t)| \geq R} \frac{1}{2R^2} |u|^2 + 2|\nabla u|^2 + |G'(|u|^2)||u|^2 + |G(|u|^2)| dx,$$

where we have used the fact that for $|x - X(t)| \leq R$ then $|x_j - X_j(t)| \leq R$, for each $1 \leq j \leq d$.

Thus, $\theta' \left(\frac{x_j - X_j(t)}{R} \right) = 1$ and $\theta''' \left(\frac{x_j - X_j(t)}{R} \right) = 0$ for each $1 \leq j \leq d$.

Moreover, for $|x - X(t)| \geq R$ and $L(\tilde{\varepsilon}) < t < \frac{R}{2\tilde{\varepsilon}}$, we have

$$|x - x(t)| \geq R - |x(t) - X(t)| \geq R - \tilde{\varepsilon}t > R/2 \geq R_1(\varepsilon).$$

This implies that

$$A_2 = 2\Phi(u) + O(\varepsilon).$$

Similarly, for $L(\tilde{\varepsilon}) < t < \frac{R}{2\tilde{\varepsilon}}$, $R \geq 2R_1(\varepsilon)$, we have

$$\begin{aligned}
A_1 &= \sum_j -X_j'(t) \mathcal{I}m \int \theta' \left(\frac{x_j - X_j(t)}{R} \right) u_j \bar{u} dx \\
&= \sum_j \frac{-2P_j(u)}{M(u)} \mathcal{I}m \int u_j \bar{u} dx - \frac{2P_j(u)}{M(u)} \mathcal{I}m \int \left(\theta' \left(\frac{x_j - X_j(t)}{R} \right) - 1 \right) u_j \bar{u} dx \\
&= \sum_j \frac{-2P_j(u)}{M(u)} P_j(u) + O(\varepsilon) = \frac{-2|P(u)|^2}{M(u)} + O(\varepsilon).
\end{aligned}$$

This completes the proof of Lemma 5.3. \square

We are now ready to prove Proposition 5.1.

Proof of Proposition 5.1. Assume that $\inf_{t \geq 0} \left(\Phi(u) - \frac{|P(u)|^2}{M(u)} \right) = \delta > 0$. We fix small parameters $\tilde{\varepsilon} \ll \delta$ and $\varepsilon \ll \delta$.

From Lemma 5.3, we have

$$W'_R(t) = A_1 + A_2 = 2 \left(\Phi(u) - \frac{|P(u)|^2}{M(u)} \right) + O(\varepsilon),$$

for $L(\tilde{\varepsilon}) < t < \frac{R}{2\tilde{\varepsilon}}$, $R \geq 2R_1(\tilde{\varepsilon})$.

Let $T_1 = L(\tilde{\varepsilon})$, $T_2 = \frac{R}{2\tilde{\varepsilon}}$, where R is large (and in particular $R > 2\tilde{\varepsilon}L(\tilde{\varepsilon})$). We have

$$\begin{aligned} CR &\geq |W_R(T_2) - W_R(T_1)| = (T_2 - T_1)|W'_R(t_0)|, \quad \text{for some } T_2 > t_0 > T_1 \\ &\geq \delta \left(\frac{R}{2\varepsilon} - L(\tilde{\varepsilon}) \right), \end{aligned}$$

(where we have used $\varepsilon \ll \delta$ for the last inequality). This gives a contradiction letting $R \rightarrow \infty$, since $\tilde{\varepsilon} \ll \delta$. \square

6. SCATTERING

This section is dedicated to the proof of Theorems 1.5, 1.6, 1.7 and 1.8. Recall that $M(\varphi) = \int |\varphi|^2$. We consider the following property:

Property 6.1. There exists $m_c > 0$ such that $\forall \varphi \in (H^{s_0} \cap H^1) \setminus \{0\}$, if $M(\varphi) < m_c$ then $\Phi(\varphi) \geq \frac{1}{2} \int |\nabla u|^2$ and $E(\varphi) \geq \frac{1}{2} \int |\nabla u|^2$.

Property 6.1 holds for all m_c when g is defocusing. In the case where $\iota_0 > 0$, we have the following:

Lemma 6.2. *Let g be a nonlinearity that satisfies Assumption B with $\iota_0 > 0$. Then Property 6.1 is true.*

Proof. Using that $\iota_0 = 1$, we have

$$\begin{aligned} \Phi(u) &= \int |\nabla u|^2 + \frac{d}{2} \int (G'(|u|^2)|u|^2 - G(|u|^2)) \\ &= \int |\nabla u|^2 + \frac{dp_0}{2(2+p_0)} \int |u|^{p_0+2} + \frac{d}{2} \int (G'_1(|u|^2)|u|^2 - G_1(|u|^2)), \end{aligned}$$

where $G_1(s) = G(s) - s^{p_0/2}$ is such that $g_1(u) = G'_1(|u|^2)u$. By Assumption (B), we have $g_1 \in \mathcal{N}(s_0, p_2, p_1)$, for some $\frac{4}{d} < p_1 < p_2 < p_0$. Using the definition of \mathcal{N} , this implies

$$(6.1) \quad |G_1(|u|^2)| + |G'_1(|u|^2)|u|^2 \lesssim |u|^{p_2+2} + |u|^{p_1+2}.$$

The claim that there exists m_c such that $\Phi(u)$ is larger than $\frac{1}{2} \int |\nabla u|^2$ when $0 < M(u) < m_c$ follows easily from (6.1) and the generalized Gagliardo-Nirenberg inequalities (See (3.25) in [31])

$$\|u\|_{L^{p_j+2}}^{p_j+2} \lesssim \|u\|_{L^2}^{p_j-\theta p_0} \|\nabla u\|_{L^2}^{2(1-\theta)} \|u\|_{L^{p_0+2}}^{\theta(p_0+2)}.$$

The lower bound for the energy is obtained by the same proof. \square

We have the following result.

Lemma 6.3. *If Property 6.1 holds, for each $\varphi \in H^{s_0} \cap H^1$ such that $M(\varphi) < m_c$, we have*

$$\Phi(\varphi) - \frac{|P(\varphi)|^2}{M(\varphi)} > 0.$$

We next prove Theorems 1.5 and 1.7. The proofs of Theorems 1.6 and 1.8 are similar, but simpler and we omit them. Theorems 1.5 and 1.7 are an immediate consequence of the following result:

Theorem 6.4. *Let g satisfy Assumption B. Assume that $s_0 \geq 1$, Properties 1.1 and 6.1 hold. Let u be a solution of (2.1) with $M(u) < m_c$ and u satisfies (1.6). Then u scatters forward in time.*

We will prove Theorem 6.4 as a consequence of

Theorem 6.5. *Let g satisfy Assumption B. Assume that $s_0 \geq 1$, Properties 1.1 and 6.1 hold. Then for all $A \in (0, A_0)$ there exists $\mathcal{F}(A, \eta) > 0$ such that for any interval I , for any solution $u \in C^0(I, H^{s_0})$ of (2.1) such that*

$$(6.2) \quad M(u) \leq m_c - \eta \quad \text{and} \quad \sup_{t \in I} \|u(t)\|_{H^{s_0}}^2 + \eta \|u(t)\|_2^2 \leq A^2,$$

one has $u \in S^{s_0}(I)$ and $\|u\|_{S^{s_0}(I)} \leq \mathcal{F}(A, \eta)$.

Theorem 6.5 implies Theorem 6.4 by the scattering criterion (Lemma 2.16) and letting η go to zero.

Proof of Theorem 6.5. We argue by contradiction, following the compactness/rigidity scheme as in [24]. We fix $\eta > 0$ throughout the argument. In all the proof, we will endow H^{s_0} with the norm defined by

$$(6.3) \quad \|u\|_{H^{s_0}}^2 = \|u\|_{H^{s_0}}^2 + \eta \|u\|_2^2.$$

We will denote by $\mathcal{P}(A)$ the property that there exists $\mathcal{F}(A)$ such that for any interval I , for any solution $u \in C^0(I, H^{s_0})$ such that (6.2) holds, one has $u \in S^{s_0}$ and $\|u\|_{S^{s_0}(I)} \leq \mathcal{F}(A)$.

By the small data theory for (2.1) (see Proposition 2.12), if $A > 0$ is small and $\|u(t)\|_{H^{s_0}} \leq A$ for some $t \in I_{\max}(u)$, then u is globally defined, scatters and $\|u\|_{S^{s_0}(\mathbb{R})} \lesssim A$. This implies that $\mathcal{P}(A)$ holds for small $A > 0$.

Thus if the conclusion of Theorem 6.5 does not hold, there exists $A_c \in (0, A_0)$ such that for all $A < A_c$, $\mathcal{P}(A)$ holds, and $\mathcal{P}(A_c)$ does not hold, i.e. there exists a sequence of intervals $((a_n, b_n))_n$, a sequence $(u_n)_n$ of solutions of (2.1) on (a_n, b_n) ,

$$(6.4) \quad u_n \in C^0((a_n, b_n), H^{s_0}), \quad M(u_n) \leq m_c - \eta, \quad \lim_{n \rightarrow \infty} \sup_{a_n < t < b_n} \|u_n(t)\|_{H^{s_0}} = A_c.$$

and $\lim_{n \rightarrow \infty} \|u_n\|_{S^{s_0}((a_n, b_n))} = \infty$. Time translating u_n , we can assume

$$(6.5) \quad a_n < 0 < b_n, \quad \lim_{n \rightarrow \infty} \|u_n\|_{S^{s_0}((a_n, 0))} = \lim_{n \rightarrow \infty} \|u_n\|_{S^{s_0}((0, b_n))} = +\infty.$$

We will prove

Claim 6.6. For any sequences $(a_n)_n, (b_n)_n$ with $a_n < 0 < b_n$, for any sequence $(u_n)_n$ of solutions of (2.1) satisfying (6.4), (6.5), there exist, after extraction of subsequences, a sequence $(x_n)_n \in (\mathbb{R}^d)^{\mathbb{N}}$ and $\varphi \in H^{s_0}$ such that

$$(6.6) \quad \lim_{n \rightarrow \infty} \|u_n(\cdot - x_n) - \varphi\|_{H^{s_0}} = 0.$$

We first assume the claim and conclude the proof of Theorem 6.5. By the claim, there exist (after extraction of subsequences) $\varphi \in H^{s_0}$ and $(x_n)_n$ such that (6.6) holds. Let u be the solution of (2.1) such that $u(0) = \varphi$. Since \mathcal{R}_η is closed in H^{s_0} , we have $\varphi \in \mathcal{R}_\eta$.

We next prove by contradiction

$$(6.7) \quad \lim_{n \rightarrow \infty} a_n = -\infty, \quad \lim_{n \rightarrow \infty} b_n = +\infty.$$

Assume to fix ideas, and after extraction of subsequences $\lim_{n \rightarrow \infty} b_n = b \in [0, \infty)$. Using that $\lim_n \|u_n\|_{S^{s_0}((0, b_n))} = \infty$, we must have $T_+(u) < \infty$ and $b \geq T_+(u)$. By the last assertion of (6.4), we obtain

$$\sup_{0 \leq t < T_+(u)} \|u(t)\|_{H^{s_0}} < \infty.$$

This implies by Theorem 1.2 that $T_+(u) = +\infty$, a contradiction. Hence (6.7). Next, we see that (6.6), perturbation theory for equation (2.1) and the last assertion in (6.4) implies that for any compact interval $I \subset I_{\max}(u)$,

$$\sup_{t \in I} \|u(t)\|_{H^{s_0}} \leq A_c.$$

This implies by Theorem 1.2 that u is global and

$$\sup_{t \in \mathbb{R}} \|u(t)\|_{H^{s_0}} \leq A_c.$$

By (6.5) and stability theory for equation (2.1), one has

$$\|u\|_{S^{s_0}((-\infty, 0))} = \|u\|_{S^{s_0}((0, +\infty))} = +\infty.$$

If $(t_n)_n$ is any sequence of times, Claim 6.6 and the preceding properties imply that one can extract subsequence such that $u(t_n, \cdot - x_n)$ converges in H^{s_0} for some sequence $(x_n)_n \in (\mathbb{R}^d)^{\mathbb{N}}$. This is classical that it implies that one can find a function $x(t)$, $t \in \mathbb{R}$ such that K defined by (5.2) has compact closure in H^{s_0} . We give a sketch of proof of this fact. Using the compactness and the fact that the solution u is not identically 0, we first notice that there exists $R_0 > 0$, $\eta > 0$ such that

$$\inf_{t \in \mathbb{R}} \left(\sup_{X \in \mathbb{R}^d} \int_{|x| < R_0} |u(t, x - X)|^2 dx \right) \geq \eta.$$

Thus for all t , there exists $x(t) \in \mathbb{R}^d$ such that

$$\int_{|x| < R_0} |u(t, x - x(t))|^2 dx \geq \eta/2.$$

For this choice of $x(t)$, one can check that K defined by (5.2) is compact.

By Proposition 5.1, we have

$$(6.8) \quad \min_{t \geq 0} \left| \Phi(u(t)) - \frac{|P(u)|^2}{M(u)} \right| = 0.$$

By (6.8), there exists a sequence of times $(t_n)_n$ such that

$$(6.9) \quad \lim_{t_n \rightarrow \infty} \Phi(u(t_n)) - \frac{|P(u(t_n))|^2}{M(u(t_n))} = 0.$$

By the claim, extracting subsequences, there exists $(x_n)_n$ such that $u(t_n, \cdot - x_n)$ converges to φ_0 in H^{s_0} (up to extract subsequence). By (6.9),

$$(6.10) \quad \Phi(\varphi_0) - \frac{|P(\varphi_0)|^2}{M(\varphi_0)} = 0.$$

Since \mathcal{R}_η is closed in H^{s_0} , $\varphi_0 \in \mathcal{R}_\eta$. This implies, by Lemma 6.3, that

$$\Phi(\varphi_0) - \frac{|P(\varphi_0)|^2}{M(\varphi_0)} > 0.$$

This contradicts to (6.10). This completes the proof. \square

We are left with proving Claim 6.6.

Proof of Claim 6.6. Step 1. Profile decomposition. Extracting subsequences, we can assume that $u_{0,n} = u_n(0)$ has a profile decomposition as in Section 3:

$$u_{0,n} = \sum_{j=1}^J \varphi_{L_n}^j(0) + w_n^J,$$

where $\varphi_{L_n}^j = \frac{1}{\lambda_n^j} \varphi_L^j \left(\frac{t-t_n^j}{\lambda_n^j}, \frac{x-x_n^j}{\lambda_n^j} \right)$. We denote by φ_n^j the corresponding nonlinear profiles, and $\tilde{\varphi}_n^j$ the modified nonlinear profiles.

Our goal is to prove that there is a unique $j_0 \geq 1$ such that φ^{j_0} is not identically zero, and that $j_0 \in \mathcal{J}_c$. We first note that there is at least one j such that φ^j is not identically zero. If not, by the small data local well-posedness, we would have

$$\lim_{n \rightarrow \infty} \|u_n\|_{S^{s_0}(I_n)} = 0,$$

a contradiction with our assumptions.

In the remaining step, we will prove that there is at most one nonzero profile. Arguing by contradiction, we assume that there is at least two nonzero profiles, say φ^1 and φ^2 . By the small data theory, there exists $\varepsilon_0 > 0$ such that

$$(6.11) \quad \inf_{t \in I_n} \|\tilde{\varphi}_n^j\|_{H^{s_0}} \geq \varepsilon_0, \quad j \in \{1, 2\}.$$

Step 2. Bound of the H^{s_0} norm. We prove that for all $j \geq 1$ we have $I_n \subset I_{\max}(\tilde{\varphi}_n^j)$ and, for large n ,

$$(6.12) \quad \sup_{t \in I_n} \|\tilde{\varphi}_n^j(t)\|_{H^{s_0}} \leq \sqrt{A_c^2 - \frac{1}{4}\varepsilon_0^2}.$$

By (6.4), and the Pythagorean expansions (3.7), (3.18), we obtain the bounds, for $J \geq 1$

$$\sum_{j \geq 1} \|\varphi_L^j(0)\|_{\dot{H}^{s_0}}^2 \leq A_c^2, \quad \sum_{j \in \mathcal{J}_{NC}} \|\varphi_L^j(0)\|_{H^{s_0}}^2 \leq A_c^2$$

Fixing a small $\varepsilon > 0$, we obtain, by the small data theory for equations (2.1) and (1.5) and Lemma 3.12, that there exists $J_0 \geq 1$ such that, for $j \geq J_0 + 1$, $I_{\max}(\varphi^j) = \mathbb{R}$,

$$(6.13) \quad \forall j \geq J_0 + 1, \quad \begin{cases} \|\varphi^j\|_{\dot{S}^{s_0}(\mathbb{R})} < \infty & \text{if } j \in \mathcal{J}_C \\ \|\varphi^j\|_{S^{s_0}(\mathbb{R})} < \infty & \text{if } j \in \mathcal{J}_{NC}. \end{cases}$$

and

$$(6.14) \quad \forall j \geq J_0 + 1, \quad \limsup_{n \rightarrow \infty} \sup_{t \in \mathbb{R}} \|\tilde{\varphi}_n^j(t)\|_{H^{s_0}} \leq \varepsilon.$$

We next prove by contradiction that for $j \in \llbracket 1, J_0 \rrbracket$, $I_n \subset I_{\max}(\tilde{\varphi}_n^j)$ and (6.12) holds. If not, we can assume (inverting time if necessary, and using Theorem 1.2) that for large n , there exists $b'_n \in (0, b_n]$ such that $[0, b'_n] \subset \bigcap_{1 \leq j \leq J_0} I_{\max}(\tilde{\varphi}_n^j)$ and

$$(6.15) \quad \sup_{1 \leq j \leq J_0} \sup_{0 \leq t \leq b'_n} \|\tilde{\varphi}_n^j(t)\|_{H^{s_0}}^2 = A_c^2 - \frac{1}{2}\varepsilon_0^2.$$

This implies that for large n

$$\forall j \in \mathcal{J}_C \cap \llbracket 1, J_0 \rrbracket, \quad \sup_{0 \leq t \leq b'_n} \|\varphi_n^j(t)\|_{\dot{H}^{s_0}} < A_c < A_0.$$

Thus by Proposition 4.1, we obtain a constant $C > 0$ such that for large n ,

$$(6.16) \quad \sup_{j \in \mathcal{J}_C \cap \llbracket 1, J_0 \rrbracket} \|\varphi_n^j\|_{\dot{S}^{s_0}([0, b'_n])} \leq C.$$

Going back to (6.15), we see also that for large n

$$\forall j \in \llbracket 1, J_0 \rrbracket \cap \mathcal{J}_{NC}, \quad \sup_{0 \leq t \leq b'_n} \|\varphi_n^j(t)\|_{H^{s_0}} \leq \sqrt{A_c^2 - \frac{1}{4}\varepsilon_0^2}.$$

Also, using the Pythagorean expansion of the mass we see that

$$\forall j \in \mathcal{J}_{NC}, \quad M(\varphi^j) \leq m_c - \eta.$$

Using that $\mathcal{P}(A)$ holds for $A = \sqrt{A_c^2 - \frac{1}{4}\varepsilon_0^2}$ we obtain that there exists a constant $C > 0$ such that for large n

$$(6.17) \quad \sup_{j \in \mathcal{J}_{NC} \cap \llbracket 1, J_0 \rrbracket} \|\varphi_n^j\|_{S^{s_0}([0, b'_n])} \leq C.$$

Combining (6.13), (6.16) and (6.17), we obtain that the assumptions of Theorem 3.15 are satisfied on $[0, b'_n]$. Using the Pythagorean expansion of Lemma 3.16 together with the limit in (6.4), we obtain

$$\limsup_{n \rightarrow \infty} \sup_{0 \leq t \leq b'_n} \sum_{j=1}^{J_0} \|\tilde{\varphi}_n^j(t)\|_{H^{s_0}}^2 \leq A_c^2.$$

By (6.11), we deduce

$$\forall j \in \llbracket 1, J_0 \rrbracket, \quad \limsup_{n \rightarrow \infty} \sup_{0 \leq t \leq b'_n} \|\tilde{\varphi}_n^j(t)\|_{H^{s_0}}^2 \leq A_c^2 - \varepsilon_0^2,$$

contradicting (6.15). This proves that (6.12) holds for all $j \geq 1$, for large n .

Step 3. Uniqueness of the nonzero profile.

In this step we still assume that φ^1 and φ^2 are nonzero profiles. Using (6.4) and (6.12), and arguing as in Step 2, we see that the assumptions of Theorem 3.15 are satisfied on $[a_n, b_n]$. This

proves that u_n scatters for large n , contradicting (6.5). This concludes the proof that there is only one nonzero profile.

Step 4. End of the proof.

We assume that φ^1 is the only nonzero profile. By the same argument as before, we obtain that for large n , $I_n \subset I_{\max}(\tilde{\varphi}_n^1)$ and

$$\limsup_{n \rightarrow \infty} \sup_{t \in I_n} \|\tilde{\varphi}_n^1\|_{H^{s_0}} \leq A_C.$$

If $1 \in \mathcal{J}_C$, we obtain by Proposition 4.1 that $\limsup_{n \rightarrow \infty} \|\varphi_n^1\|_{\dot{S}^{s_0}(I_n)} < \infty$. Thus the assumptions of Theorem 3.15 are satisfied on I_n , a contradiction with (6.5). Thus $1 \in \mathcal{J}_{NC}$. By the same argument, we obtain $\limsup_{n \rightarrow \infty} \sup_{t \in I_n} \|\tilde{w}_{L_n}^1(t)\|_{H^{s_0}} = 0$. Indeed, if not, we would have by the conservation of the H^{s_0} norm for the linear Schrödinger equation (and after extraction of a subsequence) $\lim_{n \rightarrow \infty} \sup_{t \in I_n} \|\tilde{w}_{L_n}^1(t)\|_{H^{s_0}} = \varepsilon_0 > 0$, and the same strategy as in Steps 2,3 would yield that u_n scatters for large n , a contradiction. We have proved

$$u_n(0, x) = \varphi_L^1(-t_{1,n}, x - x_{1,n}) + o(1) \text{ in } H^{s_0}.$$

By (6.5), $t_{1,n}$ must be bounded, and we can assume $t_{1,n} = 0$ for all n , i.e.

$$(6.18) \quad u_n(0, x) = \varphi^1(0, x - x_{1,n}) + o(1) \text{ in } H^{s_0},$$

which concludes the proof of the claim. \square

APPENDIX A. EQUIVALENCE OF SOBOLEV NORMS

In this appendix we prove (2.2). We recall Mihklin multiplier theorem [40], [22, Theorem 2.5]: if $m \in C^\infty(\mathbb{R}^d \setminus \{0\})$ is such that $|\xi|^\alpha \partial_\xi^\alpha m(\xi)$ is bounded for all multi-indices α such that $|\alpha| \leq 1 + \frac{n}{2}$, then $m(D)$ is a bounded operator from L^p to L^p for $1 < p < \infty$.

Using Mihklin multiplier theorem with $m(\xi) = |\xi|^s(1+|\xi|^2)^{-s/2}$ then with $m(\xi) = (1+|\xi|^2)^{-s/2}$, we see that

$$\|u\|_{L^p} + \| |\nabla|^s u \|_{L^p} \lesssim \| \langle \nabla \rangle^s u \|_{L^p}.$$

By Mihklin theorem with $m(\xi) = \frac{(1+|\xi|^2)^{s/2}}{1+|\xi|^s}$, we obtain

$$\| \langle \nabla \rangle^s u \|_{L^p} \lesssim \| (1 + |\nabla|^s) u \|_{L^p} \lesssim \| u \|_{L^p} + \| |\nabla|^s u \|_{L^p},$$

which concludes the proof of (2.2).

REFERENCES

- [1] T. Akahori, S. Ibrahim, H. Kikuchi, and H. Nawa. *Global dynamics above the ground state energy for the combined power-type nonlinear Schrödinger equations with energy-critical growth at low frequencies*, volume 1331 of *Mem. Am. Math. Soc.* Providence, RI: American Mathematical Society (AMS), 2021.
- [2] J. An and J. Kim. Local well-posedness for the inhomogeneous nonlinear Schrödinger equation in $H^s(\mathbb{R}^n)$. *Nonlinear Anal. Real World Appl.*, 59:Paper No. 103268, 21, 2021.
- [3] V. Banica. Remarks on the blow-up for the Schrödinger equation with critical mass on a plane domain. *Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5)*, 3(1):139–170, 2004.
- [4] P. Bégout and A. Vargas. Mass concentration phenomena for the L^2 -critical nonlinear Schrödinger equation. *Trans. Amer. Math. Soc.*, 359(11):5257–5282, 2007.
- [5] J. Bellazzini, V. D. Dinh, and L. Forcella. Scattering for non-radial 3d nls with combined nonlinearities, 2022.
- [6] H. Brezis and P. Mironescu. Gagliardo-Nirenberg inequalities and non-inequalities: the full story. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire*, 35(5):1355–1376, 2018.
- [7] A. Bulut, M. Czubak, D. Li, N. Pavlović, and X. Zhang. Stability and unconditional uniqueness of solutions for energy critical wave equations in high dimensions. *Comm. Partial Differential Equations*, 38(4):575–607, 2013.
- [8] R. Carles and C. Sparber. Orbital stability vs. scattering in the cubic-quintic Schrödinger equation. *Rev. Math. Phys.*, 33(3):27, 2021. Id/No 2150004.
- [9] T. Cazenave. *Semilinear Schrödinger equations*, volume 10 of *Courant Lecture Notes in Mathematics*. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.
- [10] T. Cazenave and F. B. Weissler. The Cauchy problem for the critical nonlinear Schrödinger equation in H^s . *Nonlinear Anal.*, 14(10):807–836, 1990.
- [11] X. Cheng. Scattering for the mass super-critical perturbations of the mass critical nonlinear Schrödinger equations. *Ill. J. Math.*, 64(1):21–48, 2020.
- [12] X. Cheng, C. Miao, and L. Zhao. Global well-posedness and scattering for nonlinear Schrödinger equations with combined nonlinearities in the radial case. *J. Differ. Equations*, 261(6):2881–2934, 2016.
- [13] F. M. Christ and M. I. Weinstein. Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation. *J. Funct. Anal.*, 100(1):87–109, 1991.

- [14] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global well-posedness and scattering for the energy-critical Schrödinger equation in \mathbb{R}^3 . *Ann. Math. (2)*, 167(3):767–865, 2008.
- [15] B. Dodson. Global well-posedness and scattering for the focusing, cubic Schrödinger equation in dimension $d = 4$. *Ann. Sci. Éc. Norm. Supér. (4)*, 52(1):139–180, 2019.
- [16] B. Dodson, C. Miao, J. Murphy, and J. Zheng. The defocusing quintic NLS in four space dimensions. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire*, 34(3):759–787, 2017.
- [17] T. Duyckaerts, J. Holmer, and S. Roudenko. Scattering for the non-radial 3D cubic nonlinear Schrödinger equation. *Math. Res. Lett.*, 15(6):1233–1250, 2008.
- [18] T. Duyckaerts, C. Kenig, and F. Merle. Scattering for radial, bounded solutions of focusing supercritical wave equations. *IMRN*, 2012.
- [19] C. Gao and Z. Zhao. On scattering for the defocusing high dimensional inter-critical NLS. *J. Differ. Equations*, 267(11):6198–6215, 2019.
- [20] P. Gérard. Description du défaut de compacité de l’injection de Sobolev. *ESAIM Control Optim. Calc. Var.*, 3:213–233, 1998.
- [21] R. T. Glassey. On the blowing up of solutions to the Cauchy problem for nonlinear Schrödinger equations. *J. Math. Phys.*, 18(9):1794–1797, 1977.
- [22] L. Hörmander. Estimates for translation invariant operators in L^p spaces. *Acta Math.*, 104:93–140, 1960.
- [23] J. Huang and J. Zhang. Sharp conditions of global existence and scattering for a focusing energy-critical Schrödinger equation. *J. Math. Anal. Appl.*, 410(2):561–576, 2014.
- [24] C. E. Kenig and F. Merle. Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case. *Invent. Math.*, 166(3):645–675, 2006.
- [25] C. E. Kenig and F. Merle. Scattering for $H^{1/2}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions. *Trans. Amer. Math. Soc.*, 362(4):1937–1962, 2010.
- [26] S. Keraani. On the defect of compactness for the Strichartz estimates of the Schrödinger equations. *J. Differential Equations*, 175(2):353–392, 2001.
- [27] R. Killip, J. Murphy, and M. Visan. Scattering for the cubic-quintic NLS: crossing the virial threshold. *SIAM J. Math. Anal.*, 53(5):5803–5812, 2021.
- [28] R. Killip, T. Oh, O. Pocovnicu, and M. Vişan. Solitons and scattering for the cubic-quintic nonlinear Schrödinger equation on \mathbb{R}^3 . *Arch. Ration. Mech. Anal.*, 225(1):469–548, 2017.
- [29] R. Killip and M. Visan. Energy-supercritical NLS: critical \dot{H}^s -bounds imply scattering. *Commun. Partial Differ. Equations*, 35(6):945–987, 2010.
- [30] R. Killip and M. Visan. The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher. *Am. J. Math.*, 132(2):361–424, 2010.
- [31] M. Lewin and S. Rota Nodari. The double-power nonlinear Schrödinger equation and its generalizations: uniqueness, non-degeneracy and applications. *Calc. Var. Partial Differ. Equ.*, 59(6):48, 2020. Id/No 197.
- [32] B. Liu and A. Soffer. The large time asymptotic solutions of nonlinear Schrödinger type equations. *Applied Numerical Mathematics*, 2023.
- [33] C. Lu and J. Zheng. The radial defocusing energy-supercritical NLS in dimension four. *J. Differ. Equations*, 262(8):4390–4414, 2017.
- [34] Y. Luo. Sharp scattering for the cubic-quintic nonlinear Schrödinger equation in the focusing-focusing regime. *J. Funct. Anal.*, 283(1):34, 2022. Id/No 109489.
- [35] F. Merle, P. Raphaël, I. Rodnianski, and J. Szeftel. On blow up for the energy super critical defocusing nonlinear Schrödinger equations. *Invent. Math.*, 227(1):247–413, 2022.
- [36] F. Merle and L. Vega. Compactness at blow-up time for L^2 solutions of the critical nonlinear Schrödinger equation in 2D. *Internat. Math. Res. Notices*, (8):399–425, 1998.
- [37] C. Miao, J. Murphy, and J. Zheng. The defocusing energy-supercritical NLS in four space dimensions. *J. Funct. Anal.*, 267(6):1662–1724, 2014.
- [38] C. Miao, G. Xu, and L. Zhao. The dynamics of the 3d radial NLS with the combined terms. *Commun. Math. Phys.*, 318(3):767–808, 2013.
- [39] C. Miao, T. Zhao, and J. Zheng. On the 4d nonlinear Schrödinger equation with combined terms under the energy threshold. *Calc. Var. Partial Differ. Equ.*, 56(6):39, 2017. Id/No 179.
- [40] S. G. Mikhlin. On the multipliers of Fourier integrals. *Dokl. Akad. Nauk SSSR*, 109:701–703, 1956.
- [41] J. Murphy. The defocusing $\dot{H}^{1/2}$ -critical NLS in high dimensions. *Discrete Contin. Dyn. Syst.*, 34(2):733–748, 2014.
- [42] J. Murphy. The radial defocusing nonlinear Schrödinger equation in three space dimensions. *Commun. Partial Differ. Equations*, 40(2):265–308, 2015.
- [43] J. Murphy. Threshold scattering for the 2d radial cubic-quintic NLS. *Commun. Partial Differ. Equations*, 46(11):2213–2234, 2021.
- [44] K. Nakanishi and W. Schlag. Global dynamics above the ground state energy for the focusing nonlinear Klein-Gordon equation. *Journal of Differential Equations*, 250(5):2299–2333, 2011.
- [45] K. Nakanishi and W. Schlag. Global dynamics above the ground state energy for the cubic NLS equation in 3D. *Calc. Var. Partial Differential Equations*, 44(1-2):1–45, 2012.
- [46] E. Ryckman and M. Visan. Global well-posedness and scattering for the defocusing energy-critical nonlinear Schrödinger equation in \mathbb{R}^{1+4} . *Am. J. Math.*, 129(1):1–60, 2007.
- [47] S. Shao. Maximizers for the Strichartz and the Sobolev-Strichartz inequalities for the Schrödinger equation. *Electron. J. Differ. Equ.*, 2009:13, 2009. Id/No 03.

- [48] A. Soffer and X. Wu. On the large time asymptotics of schrödinger type equations with general data, arXiv:2203.00724,2023.
- [49] A. Soffer and X. Wu. Soliton resolution for nonlinear schrödinger type equations in the radial case, arXiv:2304.04245,2023.
- [50] T. Tao, M. Visan, and X. Zhang. The nonlinear Schrödinger equation with combined power-type nonlinearities. *Commun. Partial Differ. Equations*, 32(8):1281–1343, 2007.
- [51] M. Visan. The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions. *Duke Math. J.*, 138(2):281–374, 2007.
- [52] J. Xie. Scattering for focusing combined power-type NLS. *Acta Math. Sin., Engl. Ser.*, 30(5):805–826, 2014.
- [53] J. Xie and D. Fang. Global well-posedness and scattering for the defocusing \dot{H}^s -critical NLS. *Chin. Ann. Math., Ser. B*, 34(6):801–842, 2013.
- [54] G. X. Xu and J. W. Yang. Long time dynamics of the 3d radial NLS with the combined terms. *Acta Math. Sin., Engl. Ser.*, 32(5):521–540, 2016.
- [55] X. Yu. Global well-posedness and scattering for the defocusing $\dot{H}^{1/2}$ -critical nonlinear schrödinger equation in \mathbb{R}^2 . *Anal. PDE*, 14(7):2225–2268, 2021.
- [56] V. Zakharov. Collapse of langmuir waves. *Soviet Physics JETP*, 35:908–914, 1972.
- [57] X. Zhang. On the Cauchy problem of 3d energy-critical Schrödinger equations with subcritical perturbations. *J. Differ. Equations*, 230(2):422–445, 2006.

(Thomas Duyckaerts) LAGA (UMR 7539),
 INSTITUT GALILÉE, UNIVERSITÉ SORBONNE PARIS NORD,
 99 AVENUE JEAN-BAPTISTE CLÉMENT,
 93430 VILLETANEUSE, FRANCE
 AND DÉPARTEMENT MATHÉMATIQUES ET APPLICATION,
 ÉCOLE NORMALE SUPÉRIEURE
 45 RUE D'ULM
 75005 PARIS, FRANCE
Email address, Thomas Duyckaerts: `duyckaer@math.univ-paris13.fr`

(Phan Van Tin) LAGA (UMR 7539),
 INSTITUT GALILÉE, UNIVERSITÉ SORBONNE PARIS NORD,
 99 AVENUE JEAN-BAPTISTE CLÉMENT,
 93430 VILLETANEUSE, FRANCE
Email address, Phan Van Tin: `vantin.phan@math.univ-paris13.fr`