



**HAL**  
open science

## The consequences of COVID-19 pandemic on patients with monoclonal gammopathy–associated systemic capillary leak syndrome (Clarkson disease)

Marc Pineton de Chambrun, Quentin Moyon, Stanislas Faguer, Geoffrey Urbanski, Alexis Mathian, Noémie Zucman, Marie Werner, Charles-Edouard Luyt, Franco Verlicchi, Zahir Amoura, et al.

### ► To cite this version:

Marc Pineton de Chambrun, Quentin Moyon, Stanislas Faguer, Geoffrey Urbanski, Alexis Mathian, et al.. The consequences of COVID-19 pandemic on patients with monoclonal gammopathy–associated systemic capillary leak syndrome (Clarkson disease). *Journal of Allergy and Clinical Immunology: In Practice*, 2022, 10 (2), pp.626-629. 10.1016/j.jaip.2021.11.023 . hal-04399103

**HAL Id: hal-04399103**

**<https://hal.science/hal-04399103v1>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **TITLE:** The Consequences Of COVID-19 Pandemic On Patients With Monoclonal Gammopathy  
2 Associated Systemic Capillary-Leak Syndrome (Clarkson's disease)

3

4 **RUNNING TITLE:** Consequences Of COVID-19 Pandemic In Clarkson's disease

5

6 **AUTHORS:** Marc Pineton de Chambrun<sup>1,2,3</sup>, MD, MSc; Quentin Moyon<sup>1,3</sup>, MD, MSc; Stanislas Faguer<sup>4</sup>,  
7 MD, PhD; Geoffrey Urbanski<sup>5</sup>, MD, MSc; Alexis Mathian<sup>3</sup>, MD, PhD; Noémie Zucman<sup>6</sup>, MD; Marie  
8 Werner<sup>7</sup>, MD; Charles-Edouard Luyt<sup>1,2</sup>, MD, PhD; Franco Verlicchi<sup>8</sup>, MD; Zahir Amoura<sup>3</sup>, MD, MSc for  
9 the EurêClark Study Group.

10

11 **AUTHOR AFFILIATIONS:** <sup>1</sup>Sorbonne Université, Assistance Publique-Hôpitaux de Paris (APHP), Hôpital La Pitié-  
12 Salpêtrière, Service de Médecine Intensive-Réanimation, Paris, France; <sup>2</sup>Sorbonne Université, INSERM,  
13 UMRS\_1166-ICAN, Institut de Cardiométabolisme et Nutrition (ICAN), Paris, France ; <sup>3</sup>Sorbonne Université,  
14 APHP, Hôpital La Pitié-Salpêtrière, Institut E3M, Service de Médecine Interne 2; Centre de Référence National  
15 Lupus Systémique, Syndrome des Anticorps Anti-phospholipides et Autres Maladies Auto-Immunes  
16 Systémiques Rares, Paris, France; <sup>4</sup>Département de Néphrologie Et Transplantation D'organes, Centre de  
17 Référence Des Maladies Rénales Rares, INSERM U1297 (I2MC, équipe 12), Centre Hospitalier Universitaire de  
18 Toulouse, Toulouse, France; <sup>5</sup>Service de Médecine Interne et d'Immunologie Clinique, Centre Hospitalier  
19 Universitaire, Angers, France; <sup>6</sup>Université de Paris, APHP, DMU ESPRIT, Service de Médecine Intensive  
20 Réanimation, Hôpital Louis Mourier, Colombes, France; <sup>7</sup>Université Paris Saclay, AP-HP, Hôpital Bicêtre, Service  
21 de réanimation chirurgicale adulte, Le Kremlin-Bicêtre, France; <sup>8</sup>Transfusion Service, Romagna Local Health  
22 Unit, Ravenna, Italy.

23

24 Correspondence and requests for reprints should be addressed to Marc Pineton de Chambrun, MD,  
25 MSc, Service de Médecine Intensive-Réanimation, Hôpital La Pitié-Salpêtrière, 47-83, boulevard de  
26 l'Hôpital, 75651 Paris Cedex 13, France; E-mail: [marc.dechambrun@gmail.com](mailto:marc.dechambrun@gmail.com);

27 [marc.pinetondechambrun@aphp.fr](mailto:marc.pinetondechambrun@aphp.fr) Tel: +33 (0)1 84 82 76 32, +33 (0)1 42 16 38 16

1 **Word count:** 979/1000

2 **Table count:** 2

3 **References:** 7

4

5 **Key words:** systemic capillary-leak syndrome; COVID-19; Clarkson’s disease; vaccination

6

7 **Competing interest:** all authors certify they have no conflict of interest to disclose

8

9 **Funding:** none

10

11 **Acknowledgment:** none

12

13 **Patients and Public involvement in the research:** none

14

15 **Clinical Implications box:** SARS-CoV-2 infection and COVID-19 vaccination can trigger severe relapse  
16 of systemic capillary-leak syndrome (SCLS, Clarkson’s disease).

17

18

19

20

21

22

23

24

25

1           The systemic capillary-leak syndrome (SCLS), also known as Clarkson’s disease, is a rare  
2 condition characterized by recurrent life-threatening episodes of capillary hyperpermeability in the  
3 presence of a monoclonal gammopathy. Viral infections, especially influenza virus, are known to  
4 reveal the disease and to elicit its recurrence<sup>1-3</sup>. Patients are asymptomatic between episodes.  
5 Monoclonal gammopathy of undetermined significance occurs in 3.2% of persons 50 years of age or  
6 older and in 5.3% of those 70 years of age or older<sup>4</sup>. However, the incidence and prevalence of  
7 Clarkson’s disease in this population is unknown, certainly very low<sup>1</sup>. The prevention of severe  
8 episodes rely on chronic perfusion of high-dose intravenous immunoglobulins (IVIg)<sup>5</sup>. Aside IVIg and a  
9 modest effect of terbutaline, no other treatment have shown to prevent recurrence. Moreover,  
10 despite some promising drugs, none have been proven effective in reducing capillary-leak during  
11 severe episodes. Several pathophysiological pathways, such as phosphodiesterase inhibition, are  
12 candidate for the treatment of severe episodes.

13           The coronavirus disease 2019 (COVID-19) and its preventive vaccines have been recently  
14 reported to trigger Clarkson’s disease first episodes and relapses<sup>6,7</sup>. We aimed to describe the  
15 outcome of European patients with Clarkson’s disease from the EurêClark Registry during the COVID-  
16 19 pandemic. All patients with a diagnosis of monoclonal gammopathy-associated SCLS included in  
17 the EurêClark Registry and alive at the start of COVID-19 pandemic (February 1<sup>st</sup> 2020), were included  
18 and evaluated until July 10<sup>th</sup> 2021 (**Table 1**). Thirty patients were included, with a female-to-male  
19 ratio 1.3 and a mean±SD age 58±14 years. Every patient had an IgG gammopathy with kappa (n=24)  
20 or lambda light chain (n=7). Most patients were under long-term treatment with intravenous  
21 immunoglobulins (n=27, 90%). Five patients (17%) experienced a relapse related to a proven (n=3) or  
22 highly probable (see Table 1 footnotes) severe acute respiratory syndrome coronavirus 2 (SARS-CoV-  
23 2) infection, with a fatal outcome in 4 patients. None had evidence of COVID-19 pneumonia and all  
24 experienced typical flare of Clarkson’s disease with severe hypovolemic shock and refractory multiple  
25 organ failure. Twenty patients underwent COVID-19 vaccination with: BNT162b2 (Pfizer-BioNTech)  
26 n=17; Ad26.COV2.S (Janssen) n=1; and mRNA-1273 (Moderna) n=2. Vaccination was uneventful in 18

1 patients, including two not receiving IVIg. Two patients treated with IVIg had a relapse after a second  
2 dose mRNA vaccine, with favorable outcome in both cases. During the time of the study, 5 patients  
3 had a new diagnosis Clarkson's disease and were reported to the EurêClark Registry (**Table 2**). All  
4 required intensive care unit management, and one died during this opening episode. Four had their  
5 first flare triggered by a PCR-confirmed COVID-19 infection. The last patient, previously known to  
6 have a monoclonal gammopathy, had a typical opening flare of SCLS 3 days after the first injection of  
7 ChAdOx1 (AstraZeneca).

8           Our results highlight the burden of COVID-19 pandemic for patients with Clarkson's disease.  
9 They face the threat of both the infection and its preventive vaccine. Moreover, IVIg seems to  
10 protect them imperfectly against these foes. Several lessons can be drawn from our results jointly  
11 with previous reports. First, COVID-19 infection seems to induce very frequently a relapse of  
12 Clarkson's. Every patient from our cohort with a proven or suspected SARS-CoV-2 infection had a  
13 severe flare, fatal in 80% of cases. To the best of our knowledge and in the published literature, there  
14 are no report of uneventful SARS-CoV-2 infection in these patients. As previously reported, viral  
15 infection frequently elicit relapse in Clarkson's disease patients. However, vascular leakage seems to  
16 play a major role in COVID-19 pathophysiology and may explain the high relapse risk during SARS-  
17 CoV-2 infection. While IVIg have been shown to prevent severe episodes<sup>8</sup> and to improve survival<sup>5</sup>,  
18 they failed to prevent the COVID-19-related relapses in 4 of our patients, including one treated at full  
19 dose (2g/kg/4 weeks). The tapered dose in the 3 others might have lowered their preventive effect.  
20 The lack of SARS-CoV-2 specific immunoglobulins in available preparations of IVIg could explain this  
21 lower efficacy<sup>6</sup>. A recent study revealed the apparition of anti-SARS-CoV-2 NP and S-RBD antibodies  
22 in IVIg preparation containing plasma collected in Italy after March 2020<sup>9</sup>. Yet, it cannot be denied  
23 that some patients from our cohort might have had asymptomatic COVID-19 during the time of the  
24 study. Second, while the incidence of Clarkson's disease is unknown, being probably very low, this  
25 pandemic has been associated with an elevated number of new diagnosis in a short period of time.  
26 This findings highlights the critical role of viral triggers in the onset of SCLS. Third, while we usually

1 recommended vaccination in our patients (especially against influenzae virus), COVID-19 vaccines  
2 have been shown to trigger Clarkson's disease episodes. All four vaccines authorised by the European  
3 Medicines Agency (EMA, BNT162b2, Ad26.COVS.2.S, mRNA-1273 and ChAdOx1) have be incriminated  
4 and not solely the ChAdOx1 (AstraZeneca) for which this adverse effect was recently pointed out by  
5 the EMA's safety committee. Conversely with cases reported by Kirk M. Druey's team<sup>7</sup>, our 2 patients  
6 relapsing after vaccination were given IVIg regularly. Furthermore, 18 patients, including 2 not  
7 receiving IVIg had no adverse event after COVID-19 vaccination.

8           This study has several strength and limitations. First, the small sample size of this series  
9 should be put in perspective with the rarity of the disease. Second, while the retrospective nature of  
10 this work is inevitably associated with selection bias, most informations originate from prospectively  
11 collected data. Third, only symptomatic episodes after vaccination were monitored but not any other  
12 vaccine adverse reactions. Last, we did not evaluate the anti-spike antibody response to confirm that  
13 vaccination participates in Clarkson's disease relapse prevention. In conclusion, COVID-19 pandemic  
14 has serious consequences in SCLS patients. SARS-CoV-2 infection is associated with a high risk of  
15 relapse and all COVID-19 vaccines can trigger episodes. High-dose IVIg remains the only effective  
16 preventive treatment and should not be stopped during the pandemic. In our opinion, the  
17 benefit/risk ratio favors COVID-19 vaccination in our patients under IVIg, but further data is needed  
18 to determine its safest modalities.

19  
20  
21  
22  
23  
24  
25  
26

## 1 REFERENCES

- 2 1. Druey KM, Parikh SM. Idiopathic systemic capillary leak syndrome (Clarkson disease). *J Allergy*  
3 *Clin Immunol.* 2017;140:663–70.
- 4 2. Pineton de Chambrun M, Luyt C-E, Beloncle F, Gousseff M, Mauhin W, Argaud L, et al. The  
5 Clinical Picture of Severe Systemic Capillary-Leak Syndrome Episodes Requiring ICU Admission.  
6 *Crit Care Med.* 2017;45:1216–23.
- 7 3. Pineton de Chambrun M, Mathian A, Luyt C-E, Combes A, Amoura Z, EurêClark Study Group.  
8 Myocardial dysfunction is frequent in systemic capillary-leak syndrome (Clarkson disease) severe  
9 episodes. *J Allergy Clin Immunol.* 2018;141:1539–40.
- 10 4. Kyle RA, Larson DR, Therneau TM, Dispenzieri A, Kumar S, Cerhan JR, et al. Long-Term Follow-up  
11 of Monoclonal Gammopathy of Undetermined Significance. *N Engl J Med.* 2018;378:241–9.
- 12 5. Pineton de Chambrun M, Gousseff M, Mauhin W, Lega J-C, Lambert M, Rivière S, et al.  
13 Intravenous Immunoglobulins Improve Survival in Monoclonal Gammopathy-Associated  
14 Systemic Capillary-Leak Syndrome. *Am J Med.* 2017;130:1219.e19-1219.e27.
- 15 6. Pineton de Chambrun M, Cohen-Aubart F, Donker DW, Cariou P-L, Luyt C-E, Combes A, et al.  
16 SARS-CoV-2 Induces Acute and Refractory Relapse of Systemic Capillary Leak Syndrome  
17 (Clarkson’s Disease). *The American Journal of Medicine [Internet].* 2020; Available from:  
18 <http://www.sciencedirect.com/science/article/pii/S0002934320303739>
- 19 7. Matheny M, Maleque N, Channell N, Eisch AR, Auld SC, Banerji A, et al. Severe Exacerbations of  
20 Systemic Capillary Leak Syndrome After COVID-19 Vaccination: A Case Series. *Ann Intern Med.*  
21 2021;
- 22 8. Gousseff M, Arnaud L, Lambert M, Hot A, Hamidou M, Duhaut P, et al. The systemic capillary leak  
23 syndrome: a case series of 28 patients from a European registry. *Ann Intern Med.* 2011;154:464–  
24 71.
- 25 9. Pisani G, Cristiano K, Simeoni M, Martina A, Pati I, Carocci A, et al. Detection of antibodies  
26 against SARS-CoV-2 both in plasma pools for fractionation and in commercial intravenous

1 immunoglobulins produced from plasma collected in Italy during the pandemic. Blood Transfus.

2 2021;

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24



1 **EURÊCLARK STUDY GROUP**

2 PINETON de CHAMBRUN Marc<sup>1,49</sup>, GOUSSEFF Marie<sup>2</sup>, MAUHIN Wladimir<sup>1</sup>, HOT Arnaud<sup>3</sup>, LEGA Jean-  
3 Christophe<sup>5</sup>, LAMBERT Marc<sup>4</sup>, RIVIERE Sophie<sup>6</sup>, DOSSIER Antoine<sup>7</sup>, RUIVARD Marc<sup>8</sup>, LHOTE François<sup>9</sup>,  
4 BLAISON Gilles<sup>10</sup>, MERCERON Sybille<sup>11</sup>, ZAPPELLA Nathalie<sup>11</sup>, ALRIC Laurent<sup>12</sup>, AGARD Christian<sup>13</sup>,  
5 Mathieu LACOUT<sup>13</sup>, SAADOUN David<sup>14</sup>, GRAVELEAU Julie<sup>15</sup>, SOUBRIER Martin<sup>16</sup>, HAROCHE Julien<sup>1</sup>,  
6 BOILEAU Julien<sup>17</sup>, LUCCHINI LECOMTE Marie-Josee<sup>18</sup>, HANSLIK Thomas<sup>19</sup>, CHRISTIDES Christine<sup>20</sup>,  
7 LEVESQUE Hervé<sup>21</sup>, TALASCZKA Aline<sup>22</sup>, BULTE Caroline<sup>21</sup>, HACHULLA Eric<sup>4</sup>, DECAUX Olivier<sup>23</sup>,  
8 SONNEVILLE Romain<sup>7</sup>, IBOUANGA Florent<sup>24</sup>, ARNULF Bertrand<sup>25</sup>, BENEDIT Marcel<sup>26</sup>, VIALARD Jean  
9 François<sup>27</sup>, TIEULIE Nathalie<sup>28</sup>, HADDAD Fadi<sup>29</sup>, MOULIN Bruno<sup>30</sup>, COHEN-AUBERT Fleur<sup>1</sup>, LOVEY  
10 Pierre-Yves<sup>31</sup>, LE MOAL Sylvie<sup>32</sup>, BIBES Béatrice<sup>33</sup>, RIVARD Georges-Etienne<sup>34</sup>, RONDEAU Eric<sup>35</sup>,  
11 MALIZIA Giuseppe<sup>36</sup>, DEBOURDEAU Philippe<sup>37</sup>, ABGUEGUEN Pierre<sup>38</sup>, BOSSERAY Annick<sup>39</sup>, DEVAQUET  
12 Jérôme<sup>40</sup>, PRESNE Claire<sup>41</sup>, LIFERMAN François<sup>42</sup>, LIMAL Nicolas<sup>43</sup> ARGAUD Laurent<sup>44</sup>, HERNU  
13 Romain<sup>44</sup>, DE LA SALLE Sylvie<sup>44</sup>, FAGUER Stanislas<sup>45</sup>, URBANSKI Geoffrey<sup>46</sup>, ZUCMAN Noémie<sup>47</sup>,  
14 WERNER Marie<sup>48</sup>, LUYT Charles-Edouard<sup>49</sup>, MOYON Quentin<sup>1,49</sup>, VERLICCHI Franco<sup>50</sup>, TRONCOSO  
15 Jorge Álvarez<sup>51</sup>, HARTY John<sup>52</sup>, GODMER Pascal<sup>2</sup>, HIE Miguel<sup>1</sup>, PAPO Thomas<sup>7</sup>, HATRON Pierre-Yves<sup>4</sup>,  
16 MATHIAN Alexis<sup>1</sup>, AMOURA Zahir<sup>1</sup>

17

18 <sup>1</sup>Service de médecine interne 2, CHU La Pitié-Salpêtrière, APHP, Paris, France

19 <sup>2</sup>Service de médecine interne, CH Bretagne Atlantique, Vannes, France

20 <sup>3</sup>Service de médecine interne, CHU Edouard Herriot, Lyon, France

21 <sup>4</sup>Service de médecine interne, Hôpital Claude Huriez, CHRU Lille, France

22 <sup>5</sup>Service de médecine interne et vasculaire, CHU Lyon Sud, France

23 <sup>6</sup>Service de médecine interne, Hôpital S<sup>t</sup>-Eloi, CHRU Montpellier, France

24 <sup>7</sup>Service de médecine interne, CHU Bichat, Paris, APHP, France

25 <sup>8</sup>Service de médecine interne, Hôpital d'Estaing, CHU de Clermont-Ferrand, France

26 <sup>9</sup>Service de médecine interne, Hôpital Delafontaine, Saint-Denis, France

- 1 <sup>10</sup>Service de médecine interne et rhumatologie, Hôpital Pasteur, Colmar, France
- 2 <sup>11</sup>Service de réanimation polyvalente, CH André Mignot, Versailles, France
- 3 <sup>12</sup>Service de médecine interne, Pôle digestif, CHU Purpan, Toulouse, France
- 4 <sup>13</sup>Service de médecine interne, CHU Hôtel-Dieu, Nantes, France
- 5 <sup>14</sup>Service de médecine interne et immunologie clinique, CHU La Pitié-Salpêtrière, APHP, Paris, France
- 6 <sup>15</sup>Service de médecine polyvalente, CH S<sup>t</sup>-Nazaire, France
- 7 <sup>16</sup>Service de rhumatologie, Hôpital Gabriel-Montpied, CHU Clermont-Ferrand, France
- 8 <sup>17</sup>Service de médecine interne, CH Morlaix, France
- 9 <sup>18</sup>Service de médecine Interne, CH Notre-Dame de la Miséricorde, Ajaccio, France
- 10 <sup>19</sup>Service de médecine interne, CHU Ambroise Paré, Boulogne-Billancourt, France
- 11 <sup>20</sup>Service de médecine interne, CH Avignon, France
- 12 <sup>21</sup>Service de médecine interne, Hôpital de Bois-Guillaume, CHU de Rouen, France
- 13 <sup>22</sup>Service de médecine interne, CH Roubaix, France
- 14 <sup>23</sup>Service de médecine interne, Hôpital Pontchaillou, CHU Rennes, France
- 15 <sup>24</sup>Service de médecine Interne, CH S<sup>t</sup>-Omer, France
- 16 <sup>25</sup>Service d'immunologie clinique, CHU S<sup>t</sup>-Louis, Paris, APHP, France
- 17 <sup>26</sup>Service de réanimation polyvalente, CH Moulins, France
- 18 <sup>27</sup>Service de médecine interne, Hôpital Haut-Lévêque, CHU Bordeaux Sud, Pessac, France
- 19 <sup>28</sup>Service de rhumatologie, Hôpital Pasteur 2, CHU Nice, France
- 20 <sup>29</sup>Service de médecine interne, Hôtel-Dieu de France, Beyrouth, Liban
- 21 <sup>30</sup>Service de néphrologie dialyse transplantation rénale, Hôpital Civil, CHU de Strasbourg, France
- 22 <sup>31</sup> Service d'hématologie, CH Sion, Suisse
- 23 <sup>32</sup>Service de médecine interne, CH de S<sup>t</sup>-Brieuc, France
- 24 <sup>33</sup>Service de médecine interne, CHP S<sup>t</sup>-Grégoire, France
- 25 <sup>34</sup>Service d'hématologie, CHU S<sup>te</sup>-Justine, Montréal, Canada
- 26 <sup>35</sup>Service d'urgences néphrologiques et de transplantation rénale, CHU Tenon, France

- 1 <sup>36</sup>Divisione di Medicina, Ospedale V. Cervello, Palermo, Italy
- 2 <sup>37</sup>Institut Sainte-Catherine, Avignon, France
- 3 <sup>38</sup>Service des maladies infectieuses et tropicales, CHU d'Angers, Angers, France
- 4 <sup>39</sup>Service de médecine interne, CHU Grenoble Alpes, France
- 5 <sup>40</sup>Service de réanimation polyvalente, CH Foch, Suresnes, France
- 6 <sup>41</sup>Service de néphrologie, Hôpital sud, CHU Amiens, Salouël, France
- 7 <sup>42</sup>Service de médecine interne, CH Dax, France
- 8 <sup>43</sup>Service de médecine interne, CHU Henri Mondor, APHP, Créteil, France
- 9 <sup>44</sup>Service de réanimation médicale, CHU Edouard Herriot, Lyon, France
- 10 <sup>45</sup>Département de Néphrologie Et Transplantation D'organes, CHU de Toulouse, Toulouse, France
- 11 <sup>46</sup>Service de Médecine Interne et d'Immunologie Clinique, CHU d'Angers, Angers, France
- 12 <sup>47</sup>Service de Médecine Intensive Réanimation, Hôpital Louis Mourier, Colombes, France
- 13 <sup>48</sup>Service de réanimation chirurgicale adulte, Le Kremlin-Bicêtre, France
- 14 <sup>49</sup>Service de Médecine Intensive-Réanimation, Hôpital La Pitié-Salpêtrière, Paris, France
- 15 <sup>50</sup>Transfusion Service, Romagna Local Health Unit, Ravenna, Italy
- 16 <sup>51</sup>Medicina Interna, Unidad de Enfermedades Autoinmunes Sistémicas, Hospital Universitario La Paz-Cantoblanco-Carlos III, Madrid, Espana
- 17 <sup>52</sup>Daisy Hill Hospital, Newry, Northern Ireland

**Table 1. Consequences and outcome of patients with known Clarkson's disease during the COVID-19 pandemic (2020-2021)**

Patients	Sex	Age	Diagnosis	Monoclonal Gammopathy	IVIg	Date	Dose	Covid-19	Covid-19 vaccine	Flare	Interval <sup>1</sup>	ICU <sup>2</sup>	Outcome
1	F	45	2013	IgG K	1	2013	0.5g/kg/4w	PCR +	-	Yes	Concomitant	Yes	Dead
2	M	62	2015	IgG K	1	2015	2g/kg/4w	-	BNT162b2	No	-	-	Alive
3	M	53	2010	IgG K	1	2011	0.5g/kg/4w	-	BNT162b2	No	-	-	Alive
4	F	81	2012	IgG K & L	1	2012	0.5g/kg/12w	-	BNT162b2	No	-	-	Alive
5	M	76	2004	IgG K	1	2021	1g/kg/4w	-	-	No	-	-	Alive
6	M	75	2007	IgG L	1	2009	0.5g/kg/16w	-	BNT162b2	No	-	-	Alive
7	M	56	2010	IgG K	1	2010	1g/kg/4w	PCR +	-	Yes	Concomitant	Yes	Alive
8	F	54	2008	IgG L	1	2008	0.5g/kg/6w	Probable <sup>3</sup>	-	Yes	Concomitant	Yes	Dead
9	M	61	2011	IgG K	1	2012	0.7g/kg/6w	-	BNT162b2	No	-	-	Alive
10	M	67	2003	IgG K	1	2005	2g/kg/4w	-	BNT162b2	No	-	-	Alive
11	F	81	2013	IgG K	1	2014	0.5g/kg/6w	-	BNT162b2	No	-	-	Alive
12	F	80	2013	IgG K	1	2014	1g/kg/4w	-	BNT162b2	No	-	-	Alive
13	F	59	2009	IgG K	0	-	-	-	BNT162b2	No	-	-	Alive
14	F	62	2002	IgG K	1	2007	1g/kg/8w	-	mRNA-1273	No	-	-	Alive
15	F	73	2011	IgG L	1	2011	2g/kg/6w	-	BNT162b2	No	-	-	Alive

16	M	73	2015	IgG L	1	2015	0.5g/kg/8w	-	Ad26.COVS.S	No	-	-	Alive
17	M	51	2008	IgG K	1	2008	1g/kg/4w	-	-	No	-	-	Alive
18	F	68	2016	IgG L	1	2016	0.5g/kg/4w	-	-	No	-	-	Alive
19	F	59	2008	IgG K	1	2008	0.5g/kg/4w	-	BNT162b2	No	-	-	Alive
20	M	59	2012	IgG K	1	2012	2g/kg/4w	PCR +	-	Yes	Concomitant	Yes	Dead
21	F	40	2017	IgG K	1	2018	1g/kg/4w	-	BNT162b2	No	-	-	Alive
22	F	46	2016	IgG K	1	2019	0.5g/kg/4w	-	BNT162b2	Yes	2d after 2 <sup>nd</sup> dose	No	Alive
23	F	52	2018	IgG K	0	-	-	-	BNT162b2	No	-	-	Alive
24	F	61	2003	IgG K	0	-	-	Probable <sup>3</sup>	-	Yes	Concomitant	Yes	Dead
25	M	43	2021	IgG L	1	2021	2g/kg/4w	-	BNT162b2	No	-	-	Alive
26	M	45	2020	IgG K	1	2021	2g/kg/4w	-	BNT162b2	No	-	-	Alive
27	M	60	2019	IgG K	1	2019	1g/kg/3w	-	BNT162b2	No	-	-	Alive
28	F	27	2021	IgG K	1	2021	2g/kg/4w	-	-	No	-	-	Alive
29	F	37	2016	IgG K	1	2016	1g/kg/4w	-	mRNA-1273	Yes	3d after 2 <sup>nd</sup> dose	Yes	Alive
30	F	43	2011	IgG K	1	2011	1g/kg/6w	-	-	No	-	-	Alive

EurêClark Registry was approved by local review boards and by the Commission Nationale de l'Informatique et des Libertés n°1001704; no AP-HP 14 in 1997.

Abbreviation: COVID-19, the coronavirus disease 2019; M, male; F, female; IgG, G immunoglobulin; K, kappa light chain; L, lambda light chain; g/kg/w, gram per kilogram of body weight delivered every x weeks; PCR, polymerase-chain reaction; d, days; ICU, intensive care unit.

<sup>1</sup>Interval between COVID-19 infection or COVID-19 vaccination and Clarkson's disease flare.

<sup>2</sup>Admission to the ICU.

<sup>3</sup>Two patients died during severe flare of Clarkson's disease complicated by refractory cardiac arrest probably related to an undiagnosed COVID-19 infection. Both had fever and viral symptoms upon emergency department admission which happened respectively during the epidemic peak of the first wave in France (March 23, 2020) and of the third wave in Italy (March 5, 2021). As cardiac arrest occurred very early in both patients, deep airway COVID-19 PCR could not be taken.

**Table 2. New diagnosis of Clarkson’s disease during the COVID-19 pandemic (2020-2021)**

Patients	Sex	Age, years	Monoclonal Gammopathy	Covid-19	Covid-19 vaccine	Flare	Interval <sup>1</sup>	ICU <sup>2</sup>	Hb <sup>3</sup> , g/dL	MV	RRT	Compartment syndrome	Outcome
A	M	44	? <sup>4</sup>	PCR +	-	Yes	Concomitant	Yes	23	Yes	Yes	4 limbs	Alive
B	F	47	? <sup>5</sup>	PCR +	-	Yes	Concomitant	Yes	19	Yes	Yes	No	Dead
C	M	56	IgG K	-	ChAdOx1	Yes	3d after 1 <sup>st</sup> dose	Yes	23	No	No	No	Alive
D	M	35	IgG L	PCR +	-	Yes	Concomitant	Yes	25	Yes	Yes	4 limbs	Alive
E	M	38	IgG K	PCR +	-	Yes	Concomitant	Yes	26	Yes	No	Lower limbs	Alive

Abbreviation: COVID-19, the coronavirus disease 2019; M, male; F, female; IgG, G immunoglobulin; K, kappa light chain; L, lambda light chain; PCR, polymerase-chain reaction; d, days; ICU, intensive care unit ; Hb, hemoglobin; MV, mechanical ventilation ; RRT, renal replacement therapy.

<sup>1</sup>Interval between COVID-19 infection or COVID-19 vaccination and Clarkson’s disease flare

<sup>2</sup>Admission to the ICU

<sup>3</sup>Hemoglobin highest value during the episode

<sup>4</sup>Monoclonal gammopathy could not be found during the acute episode and will be tested few weeks/month after the flare

<sup>5</sup>Monoclonal gammopathy could not be found during the acute episode and the patient died