



HAL
open science

Ab Initio Molecular Dynamics calculations on NO oxidation over oxygen functionalized High Oriented Pyrolytic Graphite

Alejandro Rivero Santamaría, Gilberto Alou, Céline Toubin, Maurice Monnerville

► **To cite this version:**

Alejandro Rivero Santamaría, Gilberto Alou, Céline Toubin, Maurice Monnerville. Ab Initio Molecular Dynamics calculations on NO oxidation over oxygen functionalized High Oriented Pyrolytic Graphite. Workshop on Quo vadis theoretical modeling of gas-phase reactions: applications in atmospheric chemistry and climate change, Université de Montpellier, Jun 2023, Montpellier, France. hal-04400362

HAL Id: hal-04400362

<https://hal.science/hal-04400362>

Submitted on 17 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ab Initio Molecular Dynamics calculations on NO oxidation over oxygen functionalized High Oriented Pyrolytic Graphite.

A. Rivero Santamaría¹, G. Alou Angulo¹, C. Toubin¹ and M. Monnerville¹

¹Univ. Lille, CNRS, UMR 8523 – PhLAM – Physique des Lasers Atomes et Molécules, F-59000
Lille, France

alejandro.rivero@univ-lille.fr

Nitrogen monoxide (NO) belongs to the group of the most polluting gases in the atmosphere. It serves as a critical precursor for the formation of smog and acid rain, which currently pose significant environmental and human health challenges. Carbonaceous materials have been identified as promising support to carry out the catalytic oxidation of NO to NO₂ at room temperature. After NO oxidation the NO₂ can be easily removed from the reaction chamber by the water scrubbing method allowing an efficient control of NO emissions. The understanding of the reaction mechanism involved in this complex process is fundamental to developing practical applications for this decontamination method.

We presented here ab initio molecular dynamics (AIMD) calculations on the NO oxidation over oxygen functionalized High Oriented Pyrolytic Graphite (O-HOPG) surface. Four different NO incidence energies were studied and the initial conditions were selected in order to facilitate the collision between the incident molecule and the epoxy group present at the surface. Our simulation results indicate that the reaction probability (HOPG-O + NO_(g) -> HOPG + NO_{2(g)}) increase with the NO incidence energy. The correlations obtained between the initial molecule orientation and the reactivity shed light on the factors that facilitate the catalytic oxidation of NO over activated carbonaceous surfaces.