

Development of a standardized questionnaire to report mechanisms of fall-related injuries in sport climbing

Maxime Luiggi, Cécile Martha

▶ To cite this version:

Maxime Luiggi, Cécile Martha. Development of a standardized questionnaire to report mechanisms of fall-related injuries in sport climbing. ECSS Paris 2023 – the 28th Annual Congress of the European College of Sport Science, Jul 2023, Paris, France. hal-04401015

HAL Id: hal-04401015

https://hal.science/hal-04401015

Submitted on 17 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Development of a standardized questionnaire to report mechanisms of fall-related injuries in sport climbing


M. Luiggi¹, C. Martha²







Introduction



Sport climbing is rapidly growing in popularity. While falls are frequent and mostly harmless given the recommended positioning during a fall (Image 1), the most severe injuries in sport climbing are caused by a fall¹. However, the fall mechanisms leading to fall-related injury (FRI) in sport climbing are under studied. No standardized tool exists to report them. The purpose of this study is to develop a standardized self-report tool of fall mechanisms.

Method (1): Participants, material & analyses

Participants: 107 sport climbers attending a sport climbing gym (mean age 31 ± 14), 60% men.

Material: A total of 40 fall mechanisms were compiled from a previous empirical study⁽²⁾ and composed both an observational tool and a self-administered questionnaire. Fall mechanisms were divided into three fall temporalities dimensions: 1) the reasons that lead to the fall (fall reason, n=12); 2) the climber's or belayer's behaviors during the fall (fall behaviors; n=20); 3) the details concerning the end of the fall (End; n=9).

Method (2): Procedure and analyses


Step 1: Observation of climbers who may fall

Step 2: Analysis of the fall by the observer using his observation grid


Step 3: Analysis of the fall by the climber and the belayer using the self-reporting questionnaire

Analyses: Use of Gwett's $AC1^{(3)}$ to measure the degree of agreement between the climber and the observer. AC1 was interpreted as (4): Very good > 0.8; Good between 0.6 and 0.8.

Results (1): Frequency of fall reasons and fall behaviors (fall number = 79)



Results (2): Degree of agreement between the climber and the observer



Conclusion & perspective

 The degree of agreement between the fall mechanisms reported by the observer and the climber is good even very good (Gwet's coefficient > 0.77).

Observer Climber

The self-report tool of fall mechanisms can be used in the event of a fall-related injury, to measure the contribution of each mechanism to the occurrence of a fall leading to injury.

Short reference list: 1 Rauch et al. (2019). Int J Environ Res Public Health, 17(1):203; 2 Luiggi, M., Lafaye, P., & Martha, C. (2023). Journal of Sports Medicine and Physical Fitness; ³Gwet K. (2001). Handbook of Inter-Rater Reliability: How to Estimate the Level of Agreement Between Two or Multiple Raters. Gaithersburg, MD: STATAXIS Publishing Company. Landis J.R, & Koch G.G. (1977). Biometrics, 33:159-74.