



HAL
open science

Is the Faber-Krahn inequality true for the Stokes operator?

Antoine Henrot, Idriss Mazari, Yannick Privat

► **To cite this version:**

Antoine Henrot, Idriss Mazari, Yannick Privat. Is the Faber-Krahn inequality true for the Stokes operator?. 2024. hal-04402162v1

HAL Id: hal-04402162

<https://hal.science/hal-04402162v1>

Preprint submitted on 18 Jan 2024 (v1), last revised 31 Aug 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

Is the Faber-Krahn inequality true for the Stokes operator?

Antoine Henrot*

Idriss Mazari-Fouquer†

Yannick Privat‡§

January 18, 2024

Abstract

The goal of this paper is to investigate the minimisation of the first eigenvalue of the (vectorial) incompressible Dirichlet-Stokes operator. After providing an existence result, we investigate optimality conditions and we prove the following surprising result: while the ball satisfies first and second-order optimality conditions in dimension 2, it does not in dimension 3, so that the Faber-Krahn inequality for the Stokes operator is probably true in \mathbb{R}^2 , but does not hold in \mathbb{R}^3 . The multiplicity of the first eigenvalue of the Dirichlet-Stokes operator in the ball in \mathbb{R}^3 plays a crucial role in the proof of that claim.

1 Introduction

1.1 Setting

Scope of the paper. The optimisation of spectral quantities with respect to a domain is a central question in shape optimisation and in the calculus of variations. Of particular importance is the paradigmatic question of minimising (with respect to the domain Ω) the first eigenvalue of a differential operator under a volume constraint on Ω . The celebrated Faber-Krahn inequality asserts that, when the operator is the (scalar) Dirichlet-Laplacian, the ball minimises the first eigenvalue with a volume constraint. In the present paper, we investigate the minimisation of the first eigenvalue of the Dirichlet-Stokes operator. Namely, for a bounded open set Ω consider the first eigenvalue

$$\lambda_1(\Omega) := \min_{\substack{\mathbf{u} \in W_0^{1,2}(\Omega)^3 \\ \nabla \cdot \mathbf{u} = 0 \text{ in } \Omega, \mathbf{u} \neq 0}} \frac{\int_{\Omega} \|\nabla \mathbf{u}\|^2}{\int_{\Omega} \|\mathbf{u}\|^2}, \quad (1)$$

where $\nabla \cdot$ stands for the divergence operator. This eigenvalue is associated with the eigen-equation

$$\begin{cases} -\Delta \mathbf{u} + \nabla p = \lambda_1(\Omega) \mathbf{u} & \text{in } \Omega, \\ \nabla \cdot \mathbf{u} = 0 & \text{in } \Omega, \\ \mathbf{u} = 0 & \text{on } \Omega. \end{cases} \quad (2)$$

*Université de Lorraine, CNRS, Institut Elie Cartan de Lorraine, BP 70239 54506 Vandœuvre-lès-Nancy Cedex, France (antoine.henrot@univ-lorraine.fr).

†CEREMADE, UMR CNRS 7534, Université Paris-Dauphine, Université PSL, Place du Maréchal De Lattre De Tassigny, 75775 Paris cedex 16, France, (mazari@ceremade.dauphine.fr)

‡Université de Lorraine, CNRS, Institut Elie Cartan de Lorraine, Inria, BP 70239 54506 Vandœuvre-lès-Nancy Cedex, France. (yannick.privat@univ-lorraine.fr).

§Institut Universitaire de France (IUF)

In (2) the function p (unique up to an additive constant) is the pressure, which can be interpreted as the Lagrange multiplier associated with the incompressibility constraint.

The problem under consideration in this article is the following:

$$\boxed{\inf_{\Omega \subset \mathbb{R}^d, \Omega \text{ bounded}, |\Omega| \leq V_0} \lambda_1(\Omega)}. \quad (3)$$

The constant $V_0 > 0$ is a given volume constraint (which will be immaterial as we will be led to work with a scale invariant formulation of (3)).

The main contributions of the article are the following:

1. First, we obtain an **existence result** in the class of quasi-open sets (this is the natural framework for such existence results; we refer to Definition 1) This is Theorem 1. The method of proof relies on the concentration-compactness principle of Lions [37], which was adapted by Bucur [3] to the setting of shape optimisation. The main difficulty here is the incompressibility condition.
2. Second, we investigate the **local optimality** of the ball by checking first and second-order optimality conditions for Hadamard variations. In Theorem 2, we prove that the ball satisfies these optimality conditions in \mathbb{R}^2 . On the other hand, we prove in Theorem 3 that the ball does not satisfy first-order optimality conditions in \mathbb{R}^3 . The proof of the non-optimality of the ball in \mathbb{R}^3 relies on the fact that λ_1 is, in this case, a multiple eigenvalue.
3. We then derive precise **necessary optimality conditions** for the minimality of a set in \mathbb{R}^3 ; this is Theorem 4; this theorem and its proof are linked to some recent results in the optimisation of the first **curl** eigenvalue [25].

We discuss related works in details in section 1.3 but let us already highlight some aspects of our problem. One of the first results in spectral shape optimisation is the celebrated Faber-Krahn inequality [23, 33], which asserts that, under a volume constraint, the ball minimises the first eigenvalue of the Dirichlet-Laplacian in any dimension. This inequality can be derived from numerous methods, some of which can be generalised to the minimisation of other spectral quantities and the ball is very often the minimiser (or at least a local minimiser) for the first eigenvalue of several scalar differential operators. Thus, Theorem 3 hints at a deeper discrepancy between the scalar and the vectorial case. In general, let us observe that the literature devoted to spectral optimisation problems in the vectorial case is scarce. To the best of our knowledge, the results closest to ours were derived very recently by Enciso, Gerner & Peraltas-Sala [22, 21, 25] in the case of the **curl** operator, a problem which was also investigated by Cantarella, DeTurck, Gluck & Teytel in the early 2000's [11].

Notations. Throughout the paper we use the following conventions:

1. For any $k \in \mathbb{N}$, $p \in [1; +\infty)$, $W^{k,p}(\Omega)$ (resp. $W_0^{k,p}(\Omega)$) denotes the usual Sobolev space of order k and index p (resp. the functions of $W^{k,p}(\Omega)$ whose trace on $\partial\Omega$ is zero).
2. $\mathcal{C}_c^\infty(\mathbb{R}^d)$ is the set of compactly supported \mathcal{C}^∞ functions.
3. ∇ is the gradient operator, $\nabla \cdot$ the divergence operator and $\nabla \wedge$ the curl operator.
4. All bold letters will be used to designate a \mathbb{R}^2 or \mathbb{R}^3 vector or vector field.

5. The double-dot product of two matrices $A = (a_{ij})_{1 \leq i, j \leq N}$ and $B = (b_{ij})_{1 \leq i, j \leq N}$ is the real number $A : B$ given by $A : B = \sum_{i, j=1}^N a_{ij} b_{ij}$.
6. $\mathbb{B}_d = \{x \in \mathbb{R}^d \mid \|x\| < 1\}$, the Euclidean unit ball of \mathbb{R}^d .
7. $\mathcal{C}^\infty(\Omega)$ denotes the space of infinitely differentiable real-valued functions in Ω .
8. $\text{Per}(\Omega)$, for Ω bounded connected, denotes the perimeter of Ω in the sense of De Giorgi

Precise statement of the problem. To set our problem in an appropriate (from the point of view of existence properties) framework, we recall the definition of quasi-open sets [30, Chapter 3]:

Definition 1. A subset Ω of \mathbb{R}^d is called **quasi-open** if there exists a non-increasing sequence of open sets $\{\omega_k\}_{k \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow \infty} \text{cap}(\omega_k) = 0 \quad \text{and} \quad \forall k \in \mathbb{N}, \quad \Omega \cup \omega_k \text{ is open,}$$

where cap denotes the capacity of an open set. Recall that it is defined as

$$\text{cap}(\omega) := \sup_{\substack{K \text{ compact} \\ K \subset \omega}} \inf_{\substack{v \in \mathcal{C}_c^\infty(\mathbb{R}^d) \\ v \geq 1 \text{ on } K}} \int_K |\nabla v|^2.$$

We define

$$\mathcal{O} := \{\Omega \subset \mathbb{R}^d, \Omega \text{ quasi-open}, |\Omega| > 0\}.$$

Quasi-open sets are the natural framework in which to consider spectral shape optimisation problems [10, 49]. For any given $\Omega \in \mathcal{O}$ we can define [30, Chapter 3] the Sobolev space $W_0^{1,2}(\Omega)$ as the set of functions $v \in W^{1,2}(\mathbb{R}^d)$ that are equal to 0 on $\mathbb{R}^d \setminus \Omega$ up to a set of zero capacity. For any $\Omega \in \mathcal{O}$ we set

$$\lambda_1(\Omega) := \min_{\substack{\mathbf{u} \in W_0^{1,2}(\Omega)^3 \\ \nabla \cdot \mathbf{u} = 0 \text{ in } \Omega, \mathbf{u} \neq 0}} \frac{\int_\Omega \|\nabla \mathbf{u}\|^2}{\int_\Omega \|\mathbf{u}\|^2}. \quad (4)$$

It is clear that for any $t > 0$ and any $\Omega \in \mathcal{O}$ we have $\lambda_1(t\Omega) = \frac{1}{t^2} \lambda_1(\Omega)$, and we thus introduce the following scale invariant functional:

$$\mathcal{F} : \mathcal{O} \ni \Omega \mapsto |\Omega|^{\frac{2}{d}} \lambda_1(\Omega).$$

The shape optimisation problem under consideration is

$$\boxed{\inf_{\Omega \in \mathcal{O}} \left(\mathcal{F}(\Omega) := |\Omega|^{\frac{2}{d}} \lambda_1(\Omega) \right)}. \quad (5)$$

1.2 Main results of the paper

Existence of an optimal shape. We begin with the following:

Theorem 1. *The variational problem (5) has a solution Ω^* .*

The proof of this theorem relies, in this vectorial setting, of the approach to the Faber-Krahn inequality of Bucur & Freitas [5, Proposition 3.1] or Bucur & Varchon [6], which in turn relies on the concentration-compactness principle of Lions [37].

Of course, if \mathcal{O} were to be replaced with $\mathcal{O}_D := \{\Omega \in \mathcal{O}, \Omega \subset D\}$ for a given compact set D (“box constraint”), the existence of a minimiser would follow from an adaptation of the Buttazzo-Dal Maso theorem [10], although one should be cautious when handling the zero-divergence constraint. This fact is actually used in the proof of our existence result. We refer to section 1.3 for more references on shape optimisation without box constraints.

Remark 1. When we work in the two-dimensional case, if the domain Ω is simply connected, we can introduce a potential ψ such that the velocity \mathbf{u} can be written $\mathbf{u} = (-\partial_y \psi, \partial_x \psi)^\top$ in such a way that the first eigenvalue of the Stokes operator coincides with the first eigenvalue of the buckling problem [27, Chapter 11]. Our existence result is thus linked to the theorem of Ashbaugh & Bucur [1], which asserts the existence of an optimal domain for the buckling problem in the class of simply connected domains. We will comment more on this aspect of the problem in section 1.3.

Optimality conditions and (semi-)differentials of eigenvalues. The local optimality of a shape will be investigated by means of Hadamard perturbations [30, Chapter 5]. This already requires some regularity of the shape under consideration. To be more precise: if Ω is a bounded domain with \mathcal{C}^2 boundary, if $\Phi \in W^{3,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ we define, for any $t \in (-1; 1)$ small enough,

$$\Omega_{t\Phi} := (\text{Id} + t\Phi)\Omega.$$

This definition is meaningful as, for any t small enough, $(\text{Id} + t\Phi)$ is a smooth diffeomorphism. For a given shape functional F and a given shape Ω , we say that F is differentiable at Ω if, for any $\Phi \in W^{3,\infty}(\mathbb{R}^d, \mathbb{R}^d)$, the limit

$$\langle dF(\Omega), \Phi \rangle := \lim_{t \rightarrow 0} \frac{F(\Omega_{t\Phi}) - F(\Omega)}{t}$$

exists and if it is a linear form in Φ . In this case, this limit is called the first-order shape derivative of F at Ω in the direction Φ .

The functional F might not be differentiable (typically, when considering the shape derivative of a multiple eigenvalue, as can be the case here), but we can usually define the **semi-differential** of F at Ω in the direction Φ as

$$\langle \partial F(\Omega), \Phi \rangle := \lim_{t \searrow 0} \frac{F(\Omega_{t\Phi}) - F(\Omega)}{t}.$$

Such semi-differentials are of particular importance when dealing with multiple eigenvalues. A systematic approach was developed by several authors to handle such situations; while we refer to section 1.3 for precise references, let us point to [13], which is the most related to our work as it deals with semi-differentials for the linear elasticity system.

The first-order optimality conditions for (5) read: if Ω^* is a smooth optimal set for any smooth vector field Φ , there holds

$$\langle dF(\Omega), \Phi \rangle \geq 0 \text{ if } F \text{ is differentiable,} \quad \langle \partial F(\Omega), \Phi \rangle \geq 0 \text{ if } F \text{ is semi-differentiable.} \quad (6)$$

We say that F is twice-differentiable at Ω if it is differentiable and if, for any $\Phi \in W^{3,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ the limit

$$\langle d^2 F(\Omega)\Phi, \Phi \rangle = \lim_{t \rightarrow 0} \frac{F(\Omega_{t\Phi}) - F(\Omega) - \langle dF(\Omega), \Phi \rangle}{t^2}$$

exists.

Local optimality of \mathbb{B}_2 in \mathbb{R}^2 . The main result here is the following:

Theorem 2. *The first eigenvalue $\lambda_1(\mathbb{B}_2)$ is simple. Consequently, \mathcal{F} is twice differentiable at \mathbb{B}_2 and there holds:*

1. For any $\Phi \in W^{3,\infty}(\mathbb{R}^d, \mathbb{R}^d)$,

$$\langle d\mathcal{F}(\mathbb{B}_2), \Phi \rangle = 0.$$

2. There exists a constant $c_0 > 0$ such that, for any $\Phi \in W^{3,\infty}(\mathbb{R}^d, \mathbb{R}^d)$,

$$\langle d^2\mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle \geq c_0 \|\langle \Phi, \nu \rangle\|_{W^{\frac{1}{2},2}(\partial\mathbb{B}_2)}^2$$

where ν is the normal vector on $\partial\mathbb{B}_2$.

This theorem is to be expected as it is a generalisation of the classical Faber-Krahn inequality to the case of the Stokes operator.

Remark 2. The differentiability of \mathcal{F} at \mathbb{B}_2 is immediate if $\lambda_1(\mathbb{B}_2)$ is a simple eigenvalue. We refer to [30, Chapter 5], or to the implicit function theorem of Mignot, Murat & Puel [39] (for its application in the context of shape derivatives, see *e.g.* [19]).

The Faber-Krahn inequality is not true for the Stokes operator in \mathbb{R}^3 . A much more surprising result is that *the Faber-Krahn inequality is not true for the Stokes operator in \mathbb{R}^3* . To the best of our knowledge, this is the first result of this kind, where the dimension has an influence on the optimality of the unit ball for the lowest eigenvalue of a differential operator. More precisely, we have:

Theorem 3. *The first eigenvalue $\lambda_1(\mathbb{B}_3)$ has multiplicity 3. \mathcal{F} is semi-differentiable at \mathbb{B}_2 , but does not satisfy the first-order optimality conditions (6): there exists a vector field Φ such that*

$$\langle \partial\mathcal{F}(\mathbb{B}_3)\Phi, \Phi \rangle < 0.$$

In particular, \mathbb{B}_3 does not solve (5).

Necessary optimality conditions in \mathbb{R}^3 . In fact, Theorem 3, which we singled out, is an easy consequence of the following optimality conditions:

Theorem 4. *Let Ω^* be a solution of (5). If Ω^* has a \mathcal{C}^2 boundary then $\lambda_1(\Omega^*)$ is a simple eigenvalue. Furthermore, if \mathbf{u} is an associated first eigenfunction, then $\|(\nabla\mathbf{u})\nu\|$ is constant on $\partial\Omega$.*

Remark 3. The eigenvalues of the Stokes operator are generically simple with respect to the domain; we refer to Ortega & Zuazua and Chitour, Kateb & Long [17, 41]. Regarding the multiplicity of eigenvalues of the Stokes operator let us also mention the recent work of Falocchi & Gazzola [24]; it deals with Navier boundary conditions rather than with Dirichlet ones. Nevertheless, for an optimal domain (for another eigenvalue than the first one), multiplicity is very often expected [28, chapter 11].

Remark 4. It is actually easy to prove that, at an optimiser, $\lambda_1(\Omega^*)$ has multiplicity at most 2. This will follow from considering the semi-differential of the eigenvalue. However, in order to lower the possible multiplicity to 1, we will draw inspiration from the work of Gerner [25, Proof of Theorem 2].

Although it is highly unlikely that these necessary optimality conditions will provide a full characterisation of the optimiser, the Poincaré-Hopf theorem (also known as the hairy ball theorem, see [16, Theorem 34.1] and [46, Theorem 39.7]) easily implies the following corollary:

Corollary 1. *Assume Ω^* has a \mathcal{C}^3 boundary. If Σ is a connected component of $\partial\Omega^*$, Σ has Euler characteristic 1, and is thus homeomorphic to a torus.*

Indeed, it suffices to observe that, as for any eigenfunction \mathbf{u} in Ω , as $\mathbf{u} = 0, \nabla \cdot \mathbf{u} = 0$ in Ω , the vector field $(\nabla \mathbf{u})\nu$ is tangential to $\partial\Omega$ (see Remark 5). Thus, Theorem 4 yields the applicability of the hairy ball theorem. It is notable that this line of reasoning was previously used in [21, 22].

Remark 5. [A consequence of choosing Dirichlet conditions] An elementary yet important observation that will be used several times in the paper is the following fact: for any \mathcal{C}^2 domain Ω , if \mathbf{u}_Ω is a first eigenfunction of the Dirichlet-Stokes operator, then the incompressibility condition and the boundary conditions imply that

$$\langle (\nabla \mathbf{u}_\Omega)\nu, \nu \rangle = 0 \quad \text{on } \partial\Omega.$$

Indeed, writing, in coordinates, $\nu = (\nu_i)_{1 \leq i \leq d}$ and $\mathbf{u}_\Omega = (u_i)_{1 \leq i \leq d}$, we have

$$\langle (\nabla \mathbf{u}_\Omega)\nu, \nu \rangle = \sum_{i=1}^d \nu_i \left(\sum_{j=1}^d \frac{\partial u_i}{\partial x_j} \nu_j \right) \quad \text{a.e. on } \partial\Omega.$$

Since $\mathbf{u}_\Omega \equiv 0$ on $\partial\Omega$ there holds

$$\forall i, j, \quad \frac{\partial u_i}{\partial x_j} \nu_i = \frac{\partial u_j}{\partial x_i} \nu_j \quad \text{a.e. on } \partial\Omega.$$

Consequently, one has

$$\langle (\nabla \mathbf{u}_\Omega)\nu, \nu \rangle = \sum_{i=1}^d \nu_i \left(\sum_{j=1}^d \frac{\partial u_i}{\partial x_j} \nu_j \right) = \sum_{i=1}^d \frac{\partial u_i}{\partial x_i} \left(\sum_{j=1}^d \nu_j^2 \right) = \nabla \cdot \mathbf{u} = 0$$

a.e. on $\partial\Omega$.

1.3 Bibliographical references

The problem under consideration in this article fits in several active lines of research. Let us describe some of the main ones.

Spectral optimisation problems for scalar operators. Due to the wealth of information they provide on the interplay between the geometry and the analytic properties (*i.e.* regarding functions defined on the domain) of domains, spectral optimisation problems have become a tenet of applied mathematics. The basic question in spectral optimisation is the following: given a certain spectral quantities, defined through a differential operator on a domain, what is the domain minimising or maximising this spectral quantity? It would be pointless to try and give an exhaustive bibliographical account of the developments of the field, but let us highlight some of the key contributions. The seminal works of Faber [23] and Krahn [33], investigating the minimisation of the first eigenvalue of the (scalar) Dirichlet-Laplacian bolstered the development of new approaches blending geometric and analytic tools. Typical questions include the existence of optimisers, their

geometry and their stability. Regarding the existence of optimisers, when no direct comparison principles can yield the explicit description of an optimiser, the first general theorem is due to Buttazzo & Dal Maso [10], when additional box constraints are enforced on the set of admissible domains. Bucur [3] developed a framework designed to handle the unbounded case. This was done using the concentration compactness-principle of Lions [37], and later used by Bucur & Varchon [6] and Bucur & Freitas [5] to derive existence results for the optimisation of eigenvalues of scalar operators. Let us also mention some more general results due, independently, to Bucur and Mazzoleni & Pratelli [4, 38]. We refer to [28, chapter 2] as a reference for existence problems. Regarding the geometry of optimisers, these are usually very delicate questions. While symmetrisation techniques-which can be used to derive direct comparison results-can be available for specific problems, there are no general tools that can be employed for generic spectral optimisation problems. Regarding rearrangements, we refer to the monograph [32]. Another possibility to obtain information regarding the optimal domains is to derive tractable optimality conditions; as the latter are often expressed in terms of an overdetermined boundary value problem having a solution on the optimal domain, results similar to the Serrin theorem [45] (see also [40]) yield the result. For an introduction to the geometric aspects of spectral optimisation we refer to [29].

Optimality conditions for multiple eigenvalues. A tenet of the Stokes problem in dimension 3 is that the first eigenvalue of the Dirichlet-Stokes operator has multiplicity 3 at the ball. While seemingly innocuous this remark actually implies the non-minimality of the ball in \mathbb{R}^3 . Naturally, when deriving optimality conditions for multiple eigenvalues, one needs to be very careful in handling shape derivatives, as they do not exist. The approach used in this article is based on the apparatus that was set up by Cox [18], Chenais & Rousselet [43] and Chatelain & Choulli [15] to handle these difficulties. The relevant formulae will be recalled in the course of this article. In a recent paper, Caubet, Dambrine & Mahadevan [13] developed this semi-differential approach to accomodate the linear elasticity system.

Shape optimisation for vectorial operators. The literature devoted to shape optimisation problems for vectorial operators is scarce, but growing. Of particular importance to us in this regard are the different contributions to the study of isoperimetric inequalities for the **curl** operator, that is, the optimisation of the first positive eigenvalue of

$$\begin{cases} \mathbf{curl}(\mathbf{u}) = \xi \mathbf{u} & \text{in } \Omega, \\ \langle u, \nu \rangle = 0 & \text{on } \partial\Omega. \end{cases}$$

This problem is of paramount importance in the study of magnetic fields, and has a strong connection [14] to the Stokes eigenvalue problem with a tangential flow condition, rather than a Dirichlet condition. To the best of our knowledge, the main works on this **curl**-isoperimetric problems can be found, one the one hand, in the works of Cantarella, DeTurck, Gluck and Teytel [11, 12] and, on the other hand, in the research of Enciso, Gerner & Peraltas-Sala [21, 22, 25]. Let us briefly review their finding: in [11, 12], the authors, using explicit computations of eigenfunctions, observe the optimality of the ball among all concentric spherical shells with a given volume constraint, but also conjecture that the minimiser of the **curl**-isoperimetric problem is a “spheromak”, that is, a sphere where the south and north poles are glued together. In their paper, they also give optimality conditions for the optimiser, which lead to the same type of conclusion as Corollary 1. It is interesting to note that their analysis recovers parts of the semi-differentiability results of [15]. In [21, 22], another take on the problem is introduced, and the authors investigate possible symmetries of optimisers. In [22], Enciso & Peraltas-Sala investigate whether a solution of the **curl**-isoperimetric problem can have axial symmetry; their conclusion is no, provided some regularity of the minimiser is assumed *a priori*. Their results hinge on a variational analysis (*à la* Hadamard). In [21] on the other hand, they seek optimisers in

the class of convex domains, and they prove that one can not have “too regular” optimisers. Gerner, in [25], went much further in the fine characterisation of possible optimisers, both in the euclidean and riemannian settings. Although the adaptation of some of his results is not immediate, one of the necessary optimality conditions we derive is inspired by [25, Theorem 2]. Finally, let us mention the recent work of Lamberti & Zaccaron [35], in which, also using a semi-differential approach, the authors investigate the optimal shape of an electromagnetic cavity.

The buckling problem. Let us conclude this bibliographical paragraph by mentioning a closely related problem in the two-dimensional case, the buckling problem. Indeed, in dimension 2, assume Ω is simply connected, so that we might write the first Dirichlet-Stokes eigenfunction \mathbf{u} as

$$\mathbf{u} = \begin{pmatrix} -\partial_y \psi \\ \partial_x \psi \end{pmatrix}$$

for some function ψ . Taking the (2-dimensional) **curl** of the equation, it appears that $(\psi, \lambda_1(\Omega))$ actually solves

$$\begin{cases} -\Delta^2 \psi = \lambda_1(\Omega) \Delta \psi & \text{in } \Omega, \\ \psi = \partial_\nu \psi = 0 & \text{on } \partial\Omega. \end{cases}$$

In fact, with a bit more work, one can see that $\lambda_1(\Omega)$ coincides with the eigenvalue

$$\Lambda(\Omega) := \min_{v \in W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega)} \frac{\int_{\Omega} (\Delta v)^2}{\int_{\Omega} |\nabla v|^2}.$$

This is the well-known *buckled plate eigenvalue problem*, leading to the spectral optimisation problem

$$\inf_{\Omega \subset \mathbb{R}^2, |\Omega| \leq c} \Lambda(\Omega). \quad (7)$$

We refer to [27, Chapter 11] for more details, but let us underline the following aspects of the problem: first of all, the existence of an optimal domain remained open until the contribution of Ashbaugh & Bucur [1] and, later, of Stollenwerk [48, 47]. Second, this problem is related to a long-standing conjecture by Pólya & Szegő that asserts that the ball is a solution of this variational problem. A very active line of research has focused on the computation of shape-derivatives for this problem and, more generally, for polyharmonic problems. Regarding the Hadamard type shape derivatives, we refer to [26]. For applications of this calculus to polyharmonic problems, let us point to the numerous works of Buoso & Lamberti [7, 8, 9]. To the best of our knowledge, the second-order derivative of the first eigenvalue at the ball was not known.

2 Proof of Theorem 1

We pick a minimising sequence $\{\Omega_k\}_{k \in \mathbb{N}}$ for (5). By scaling invariance of \mathcal{F} we might assume that $|\Omega_k|$ does not depend on k : there exists $V_0 > 0$ such that $|\Omega_k| = V_0$ for any $k \in \mathbb{N}$. We consider the auxiliary problem

$$\inf_{\Omega \in \mathcal{O}, |\Omega| \leq V_0} \lambda_1(\Omega). \quad (8)$$

Our proof is in two steps. We first apply the concentration-compactness principle in order to show that one might, in (8), further assume $\Omega \subset D$ for a fixed compact set $D \subset \mathbb{R}^d$. We then investigate the existence of an optimiser with this additional box constraint.

Step 1: Using the. Let us consider, for any $k \in \mathbb{N}$, the function w_k defined as

$$w_k := \|\mathbf{u}_k\|$$

where for any k the function \mathbf{u}_k is an¹ eigenfunction of Ω_k . The function w_k is extended by 0 outside of Ω_k .

By definition and elementary computations, we have

$$\int_{\mathbb{R}^d} w_k^2 = 1, \quad \int_{\mathbb{R}^d} |\nabla w_k|^2 \leq \int_{\mathbb{R}^d} \|\nabla \mathbf{u}_k\|^2. \quad (9)$$

By the concentration-compactness principle of Lions [37] we know that one of the following occurs:

(i) **Concentration:** there exists a function $w \in L^2(\mathbb{R}^d; \mathbb{R}_+)$ and a sequence $\{y_k\}_{k \in \mathbb{N}} \in (\mathbb{R}^d)^\mathbb{N}$ such that $w_k(\cdot + y_k) \xrightarrow[k \rightarrow \infty]{} w$ in $L^2(\mathbb{R}^d)$.

(ii) **Dichotomy:** There exists $\alpha_1 \in (0, 1)$, $\{y_k\}_{k \in \mathbb{N}} \in (\mathbb{R}^d)^\mathbb{N}$, two sequences $\{R_k, R'_k\}_{k \in \mathbb{N}}$ such that

$$R_k - R'_k \xrightarrow[k \rightarrow +\infty]{} +\infty, \quad R_k, R'_k \xrightarrow[k \rightarrow +\infty]{} +\infty$$

and such that

$$\int_{\mathbb{B}(y_k, R_k)} w_k^2 \xrightarrow[k \rightarrow +\infty]{} \alpha_1, \quad \int_{\mathbb{B}(y_k, R'_k)^c} w_k^2 \xrightarrow[k \rightarrow +\infty]{} 1 - \alpha_1$$

and

$$\liminf_{k \rightarrow \infty} \left(\int_{\mathbb{R}^d} |\nabla w_k|^2 - \int_{\mathbb{B}(y_k, R_k)} |\nabla w_k|^2 - \int_{\mathbb{B}(y_k, R'_k)^c} |\nabla w_k|^2 \right) \geq 0.$$

(iii) **Vanishing:** for any $r > 0$,

$$\lim_{k \rightarrow \infty} \sup_{y \in \mathbb{R}^d} \int_{\mathbb{B}(y, r)} w_k^2 = 0.$$

In this first step of the proof, we rule out vanishing and dichotomy.

Vanishing does not occur. Let us exclude vanishing. By the exact same arguments as in [6, Lemma 3.3, Proof of Theorem 3.2] and (9), if the sequence $\{w_k\}_{k \in \mathbb{N}}$ vanishes, we have

$$\limsup_{k \rightarrow +\infty} \lambda_1(\Omega_k) = +\infty.$$

This contradicts the fact that the sequence is minimising.

¹We do not rule out the possibility of Ω_k having a multiple eigenvalue. Although one could easily discard this case by invoking the fact that the first eigenvalue is generically simple, we do not require this.

Dichotomy does not occur. Argue by contradiction and assume dichotomy holds. In that case, define

$$\eta_k := \frac{R'_k - R_k}{4}.$$

Introduce $\{\psi_{k,1}, \psi_{k,2}\}_{k \in \mathbb{N}}$ as two non-negative, radially (with respect to y_k) symmetric, non-increasing functions such that

$$\psi_{k,1}(x) = \begin{cases} 1 & \text{in } \mathbb{B}(y_k, R_k), \\ 0 & \text{in } \mathbb{B}(y_k, R_k + 2\eta_k)^c, \end{cases} \quad \psi_{k,2}(x) = \begin{cases} 1 & \text{in } \mathbb{B}(y_k, R'_k - \eta_k)^c, \\ 0 & \text{in } \mathbb{B}(y_k, R'_k - 2\eta_k) \end{cases}$$

and

$$\|\nabla \psi_{k,i}\|_{L^\infty} \leq \frac{1}{\eta_k}, \quad k \in \mathbb{N}, \quad i \in \{1, 2\}.$$

Now, let, for any $k \in \mathbb{N}$ and any $i \in \{1, 2\}$

$$\mathbf{v}_{k,i} := \psi_{k,i} \mathbf{u}_k.$$

In particular,

$$\int_{\mathbb{R}^d} \|\mathbf{v}_{k,1}\|^2 \xrightarrow{k \rightarrow \infty} \alpha_1, \quad \int_{\mathbb{R}^d} \|\mathbf{v}_{k,2}\|^2 \xrightarrow{k \rightarrow \infty} 1 - \alpha_1$$

and

$$\liminf_{k \rightarrow \infty} \left(\int_{\mathbb{R}^d} \|\nabla \mathbf{u}_k\|^2 - \int_{\mathbb{R}^d} \|\nabla \mathbf{v}_{k,1}\|^2 - \int_{\mathbb{R}^d} \|\nabla \mathbf{v}_{k,2}\|^2 \right) \geq 0.$$

Up to iterating that construction, we can assume that $\{\|\mathbf{v}_{k,1}\|\}_{k \in \mathbb{N}}$ is in a situation of concentration. In particular, there exists \mathbf{v}_1 such that, up to a translation by a vector $z_k \in \mathbb{R}^d$ (which we take equal to 0 without loss of generality),

$$\mathbf{v}_{k,1} \xrightarrow[k \rightarrow +\infty]{} \mathbf{v}_1 \text{ strongly in } L^2(\mathbb{R}^d), \text{ weakly in } W^{1,2}(\mathbb{R}^d)^3.$$

Let us now observe that

$$\nabla \cdot \mathbf{v}_1 = 0$$

in the sense of distributions. To prove this fact, let $\phi \in \mathcal{C}_c^\infty(\mathbb{R}^d)$ and consider the quantity

$$\int_{\mathbb{R}^d} \langle \mathbf{v}_{k,1}, \nabla \phi \rangle.$$

As ϕ is compactly supported, $\text{supp}(\phi) \cap \{\nabla \cdot \mathbf{v}_{k,1} \neq 0\} = \emptyset$ for k large enough. Indeed, should this not be the case, as $\nabla \cdot \mathbf{v}_{k,1} = 0$ in $\mathbb{B}(y_k, R_k + \eta_k)$, it would follow that

$$\text{dist}(\text{supp}(\phi), \mathbb{B}(y_k, R_k)) \geq \eta_k \xrightarrow[k \rightarrow +\infty]{} \infty.$$

This contradicts the strong L^2 convergence of $\{\mathbf{v}_{k,1}\}_{k \in \mathbb{N}}$, thereby leading to a contradiction. Thus,

$$\int_{\mathbb{R}^d} \langle \mathbf{v}_{k,1}, \nabla \phi \rangle = 0$$

for any k large enough. Passing to the limit provides the result.

Now, we know that

$$\lambda^* := \inf_{\Omega \in \mathcal{O}, |\Omega| \leq V_0} \lambda_1(\Omega) \geq \min_{i=1,2} \liminf_{k \rightarrow \infty} \left\{ \frac{\int_{\mathbb{R}^d} \|\nabla \mathbf{v}_{k,i}\|^2}{\int_{\mathbb{R}^d} \|\mathbf{v}_{k,i}\|^2} \right\}.$$

If this minimum is reached for $i = 1$, then $\Omega^* := \{\mathbf{v}_1 \neq 0\}$ is an optimal domain. Else, we apply the procedure once more (as in [6]) to $\{\mathbf{v}_{k,2}\}_{k \in \mathbb{N}}$. Either this sequence is in a situation of compactness, in which case we are done, or it is once again in a situation of dichotomy, giving rise to two new sequences $\{\mathbf{v}_{k,2}^1, \mathbf{v}_{k,2}^2\}_{k \in \mathbb{N}}$, with $\{\mathbf{v}_{k,2}^1\}_{k \in \mathbb{N}}$ in a situation of compactness. We then iterate the construction on $\mathbf{v}_{k,2}^2$ if necessary. Either this process stops, in which case existence follows, or we obtain a decreasing sequence $\{\alpha_j\}_{j \in \mathbb{N}, j \geq 2}$ such that

$$\sum_{j=1}^{+\infty} \alpha_j \leq 1$$

and, for any $j \in \mathbb{N}$, we have a sequence $\{\mathbf{v}_{k,2}^j\}_{k \in \mathbb{N}, j \leq J_k}$ where $J_k \xrightarrow[k \rightarrow +\infty]{} \infty$ such that for any j , $\{\mathbf{v}_{k,2}^j\}_{k \in \mathbb{N}}$ is in a situation of concentration, and $\{\{\mathbf{v}_{k,2}^{J_k} \neq 0\}\}_{k \in \mathbb{N}}$ is a minimising sequence.

Observe that if $\sum_{j=1}^{\infty} \alpha_j < 1$, then vanishing holds for $\{\mathbf{v}_{k,2}\}_{k \in \mathbb{N}}$, which leads to a contradiction, so that we can work under the assumption that

$$\sum_{j=1}^{\infty} \alpha_j = 1.$$

Consequently, we deduce that

$$\lambda^* \geq \liminf_{k \rightarrow \infty} \frac{\int_{\mathbb{R}^d} \|\nabla \mathbf{v}_{k,1}\|^2 + \sum_{j=1}^{J_k} \int_{\mathbb{R}^d} \|\nabla \mathbf{v}_{k,2}^j\|^2}{\sum_{j=1}^{J_k} \alpha_j} > \lambda^*$$

by convexity since we have, by construction,

$$\liminf_{k \rightarrow \infty} \frac{\int_{\mathbb{R}^d} \|\nabla \mathbf{v}_{k,1}\|^2}{\alpha_1} > \lambda^*.$$

This is a contradiction, and the result follows.

We deduce that we have concentration of the sequence. Upon examination of the truncation procedure, one can see that the condition $R_k \xrightarrow[k \rightarrow \infty]{} +\infty$ can be weakened, and that $\{R_k\}_{k \in \mathbb{N}}$ might be chosen to be constant, as is the case in the classical statement of the concentration-compactness lemma [37]. This allows to introduce a compact set $D \subset \mathbb{R}^d$, and to only consider sequences $\{\Omega_k\}_{k \in \mathbb{N}} \subset D$, that is, we can now simply deal with an optimisation problem with box constraints.

Step 2: Existence under box-constraints. We consider a fixed compact domain $D \subset \mathbb{R}^d$. Let us define

$$\widehat{\mathcal{O}}_{V_0} = \{\Omega \text{ quasi-open included in } D, |\Omega| \leq V_0\}$$

and let us introduce, for $\Omega \in \widehat{\mathcal{O}}_V$, the quantity $\widehat{\lambda}_1(V)$ given by

$$\widehat{\lambda}_1(V_0) = \inf_{\substack{\mathbf{u} \in W_0^{1,2}(D)^3 \\ \int_D \mathbf{u} \cdot \nabla v = 0 \quad \forall v \in H^1(D) \\ |\Omega_{\mathbf{u}}| \leq V_0 \\ \int_{\Omega} \|\mathbf{u}\|^2 = 1}} \int_D \|\nabla \mathbf{u}\|^2 \quad (10)$$

where $\Omega_{\mathbf{u}}$ denotes the quasi open set $\{\mathbf{u} \neq 0\}$.

Let $\{\mathbf{u}_k\}_{k \in \mathbb{N}}$ be minimising for Problem (10). For any $k \in \mathbb{N}$, let $\Omega_k := \{\mathbf{u}_k \neq 0\}$. In view of the Poincaré inequality [30, Lemma 4.5.3], $\{\mathbf{u}_k\}_{k \in \mathbb{N}}$ is bounded in $W_0^{1,2}(D)^3$. It follows that there exists $\mathbf{u}^* \in W_0^{1,2}(D)^3$ such that $\{\mathbf{u}_k\}_{k \in \mathbb{N}}$ converges to \mathbf{u}^* up to a (not relabelled) subsequence, weakly in $W_0^{1,2}$ and strongly in L^2 . Thus, $\{u_k\}_{k \in \mathbb{N}}$ converges almost everywhere to \mathbf{u}^* , and therefore

$$|\Omega_{\mathbf{u}^*}| \leq \liminf_{k \rightarrow +\infty} |\Omega_k| \leq V_0.$$

where $\Omega_{\mathbf{u}^*}$ denotes the quasi-open set $\{\mathbf{u}^* \neq 0\}$. We infer that $\Omega_{\mathbf{u}^*}$ belongs to $\widehat{\mathcal{O}}_V$. Furthermore, since $\mathbf{u}^* = 0$ quasi-everywhere on $D \setminus \Omega_{\mathbf{u}^*}$, by weak H^1 -convergence, one has

$$\int_{\Omega_{\mathbf{u}^*}} \|\nabla \mathbf{u}^*\|^2 \leq \liminf_{k \rightarrow +\infty} \int_{\Omega_k} \|\nabla \mathbf{u}_k\|^2 = \widehat{\lambda}_1(V_0) \leq \int_{\Omega_{\mathbf{u}^*}} \|\nabla \mathbf{u}^*\|^2$$

by minimality, whence the equality of these quantities. By strong convergence in $L^2(D; \mathbb{R}^3)$ and weak convergence in $W_0^{1,2}(D)^3$, there holds

$$\int_D \|\mathbf{u}^*\|^2 = 1$$

and, for any $v \in W^{1,2}(D)$,

$$0 = \lim_{k \rightarrow +\infty} \int_D \langle \mathbf{u}_k, \nabla v \rangle = \int_D \langle \mathbf{u}^*, \nabla v \rangle.$$

Thus, $\Omega_{\mathbf{u}^*}$ is a solution of (10). The proof is now concluded.

3 Proof of Theorem 2

Throughout our analysis, we will be using some basic facts about the Stokes eigenfunctions in \mathbb{B}_2 , which we thus first recall.

3.1 Preliminaries

The first eigenpair in \mathbb{B}_2 . We let $E_1(\mathbb{B}_2)$ be the first eigenspace associated with $\lambda_1(\mathbb{B}_2)$. We begin with the following fact (see Remark 6):

Lemma 1. *The eigenvalue $\lambda_1(\mathbb{B}_2)$ is simple: $\dim(E_1(\mathbb{B}_2)) = 1$. Furthermore,*

$$\lambda_1(\mathbb{B}_2) = j_{1,1}^2,$$

where $j_{1,1}$ is the first positive root of J_1 , the Bessel function of the first kind of order 1. Finally, $E_1(\mathbb{B}_2)$ is spanned by the (eigenfunction

$$\mathbf{u}_{\mathbb{B}_2} := \frac{J_1(j_{1,1}r)}{\sqrt{\pi}j_{1,1}|J_0(j_{1,1})|} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}.$$

and the pressure is constant.

While we single out this result for later reference, the following spectral decomposition of the Dirichlet-Stokes operator in the ball is well-known:

Lemma 2. *A Hilbert basis of eigenfunctions of the Dirichlet-Stokes operator is given, in polar coordinates, by*

$$\phi_{0,k}(r, \theta) = \frac{-J'_0(\sqrt{\lambda_{0,k}}r)}{\sqrt{\pi}\sqrt{\lambda_{0,k}}|J_0(\sqrt{\lambda_{0,k}})|} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}, \quad (11)$$

and

$$\begin{aligned} \phi_{j,k,m}(r, \theta) &= \frac{J_j(\sqrt{\lambda_{j,k}}r) - J_j(\sqrt{\lambda_{j,k}}r)^j}{\lambda_{j,k}|J_j(\sqrt{\lambda_{j,k}})|r} j(-1)^{m+1} Y_{j,m}(\theta) \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \\ &+ \frac{-\sqrt{\lambda_{j,k}}J'_j(\sqrt{\lambda_{j,k}}r) + jJ_j(\sqrt{\lambda_{j,k}}r)^{j-1}}{\lambda_{j,k}|J_j(\sqrt{\lambda_{j,k}})|} Y_{j,m+1}(\theta) \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \end{aligned} \quad (12)$$

for $j \in \mathbb{Z}^*$, $k \in \mathbb{N}^*$ and $m = 1, 2$

Here, (r, θ) are the usual polar coordinates (see [31, 36]). The functions $Y_{j,m}(\theta)$ are defined by $Y_{j,1}(\theta) = \frac{1}{\sqrt{\pi}} \cos(j\theta)$ and $Y_{j,2}(\theta) = \frac{1}{\sqrt{\pi}} \sin(j\theta)$, with the agreement that $Y_{j,3} = Y_{j,1}$, and J_j is the Bessel function of the first kind of order j . Denoting by $j_{j,k} > 0$ the k^{th} positive zero of J_j , the eigenvalues of the Dirichlet-Stokes operator are the doubly indexed sequence $(\lambda_{j,k})_{j \in \mathbb{Z}, k \in \mathbb{N}^*}$, where $\lambda_{j,k} = j_{|j|+1,k}^2$ is of multiplicity 1 if $j = 0$, and 2 if $j \neq 0$.

Remark 6. [Regarding Lemmata 1-2] Lemmata 1-2 are essentially contained in [31, 36], which contain explicit computations of the eigen-elements of the Dirichlet-Stokes operator in special geometries. While we refer to these articles for detailed computations, let us briefly indicate how they might be derived in a straightforward manner: consider $(\mathbf{u}_k, \lambda_k(\mathbb{B}_2))$ an eigenpair of the Dirichlet-Stokes operator. As $\nabla \cdot \mathbf{u}_k = 0$, there exists a scalar function ψ_k such that

$$\mathbf{u}_k = \mathbf{curl}(\psi_k) := \begin{pmatrix} -\partial_y \psi_k \\ \partial_x \psi_k \end{pmatrix}$$

and, taking the (2 dimensional) **curl** of (2), it appears that ψ_k solves

$$\begin{cases} -\Delta^2 \psi_k = \lambda_k(\mathbb{B}_2) \Delta \psi_k & \text{in } \mathbb{B}_2, \\ \psi_k = \partial_\nu \psi_k = 0 & \text{on } \partial \mathbb{B}_2. \end{cases} \quad (13)$$

In other words, ψ_k is an eigenvalue of the aforementioned buckling problem (see section 1.3). In other words, knowing the eigenspaces of the buckling problem leads to determining the eigenspaces of the Dirichlet-Stokes operator. However, the eigenspaces of the buckling operator can be easily computed in the usual radial coordinates. It should be observed that the three-dimensional case, although it follows a similar pattern (boiling it down to a scalar operator) is much more involved. We refer to Appendix A.

Basic results about the shape differentiability of eigenfunctions and eigenvalues. Since $\lambda_1(\mathbb{B}_2)$ is simple, the following differentiability result follows from the implicit function theorem of Mignot, Murat & Puel [39]:

Lemma 3. *Let*

$$\mathcal{X} := \{\Phi \in W^{3,\infty}(\mathbb{R}^2; \mathbb{R}^2), \Phi \text{ compactly supported}\}.$$

For any \mathcal{C}^2 domain Ω such that $\lambda_1(\Omega)$ is simple, the eigenvalue mapping $\Omega \mapsto \lambda_1(\Omega)$ is twice shape differentiable at Ω in the following sense: for any $\Phi \in \mathcal{X}$, the map $f_\Phi : t \mapsto \lambda_1((\text{Id} + t\Phi)\Omega)$ is twice differentiable at $t = 0$. We will use the notations

$$\langle d\lambda_1(\Omega), \Phi \rangle := f'_\Phi(0), \langle d^2\lambda_1(\Omega)\Phi, \Phi \rangle := f''_\Phi(0).$$

Similarly, the mapping $\Omega \mapsto \mathbf{u}_\Omega$ is twice shape-differentiable at Ω , where \mathbf{u}_Ω is the first Dirichlet-Stokes normalized eigenfunction of Ω , in the sense that the mapping $g_\Phi : t \mapsto \mathbf{u}_{(\text{Id}+t\Phi)\Omega}$ is twice differentiable at $t = 0$. We let \mathbf{u}'_Φ be its derivative at $t = 0$.

For a detailed introduction to the Hadamard shape calculus, we refer to [30, Chapter 5] and to [26]. To proceed with the proof of Theorem 2, we need tractable expressions for the first and second-order shape derivatives of the eigenvalue at a domain Ω .

Proposition 1. *For any \mathcal{C}^2 domain Ω such that $\lambda_1(\Omega)$ is simple, let \mathbf{u}_Ω be its associated first eigenfunction. For any $\Phi \in \mathcal{X}$, the shape derivative \mathbf{u}'_Φ solves*

$$\begin{cases} -\Delta \mathbf{u}'_\Phi + \nabla p' = \lambda_1(\Omega) \mathbf{u}'_\Phi + \langle d\lambda_1(\Omega), \Phi \rangle \mathbf{u}_\Omega & \text{in } \Omega \\ \nabla \cdot \mathbf{u}'_\Phi = 0 & \text{in } \Omega \\ \mathbf{u}'_\Phi = -\nabla \mathbf{u}_\Omega \nu \langle \Phi, \nu \rangle & \text{on } \partial\Omega, \\ \int_\Omega \langle \mathbf{u}_\Omega, \mathbf{u}'_\Phi \rangle = 0. \end{cases} \quad (14)$$

The first order derivative of λ_1 is

$$\langle d\lambda_1(\Omega), \Phi \rangle = - \int_{\partial\Omega} \|(\nabla \mathbf{u}_\Omega) \nu\|^2 \langle \Phi, \nu \rangle. \quad (15)$$

If, in addition, the vector field Φ is normal to $\partial\Omega$, the second-order shape derivative of λ_1 at Ω is given by

$$\langle d^2\lambda_1(\Omega)\Phi, \Phi \rangle = 2 \int_{\partial\Omega} \langle \mathbf{u}'_\Phi, (\nabla \mathbf{u}'_\Phi) \nu \rangle + \int_{\partial\Omega} H \|(\nabla \mathbf{u}_\Omega) \nu\|^2 \langle \Phi, \nu \rangle^2 - 2 \int_{\partial\Omega} \langle (\nabla \mathbf{u}_\Omega) \nu, \nabla p_\Omega \rangle \quad (16)$$

where H is the mean curvature of $\partial\Omega$ and p_Ω is the pressure field associated with \mathbf{u}_Ω .

Proof of Proposition 1. That \mathbf{u}'_Φ solves (14) is a standard consequence of general formulae for the shape differentiation of Dirichlet boundary value problem, and we refer to [30, Chapter 5] for the detailed computations. Once (14) is established, it suffices to multiply the equation by \mathbf{u}_Ω and to integrate by parts to obtain

$$\langle d\lambda_1(\Omega), \Phi \rangle \int_\Omega \|\mathbf{u}_\Omega\|^2 = - \int_{\partial\Omega} \|(\nabla \mathbf{u}_\Omega) \nu\|^2 \langle \Phi, \nu \rangle.$$

As

$$\int_\Omega \|\mathbf{u}_\Omega\|^2 = 1,$$

the conclusion follows.

To obtain the expression for $\langle d^2\lambda_1(\Omega)\Phi, \Phi \rangle$ for normal vector fields, it suffices to apply the Hadamard formula for integrals on variable boundaries [30, Proposition 5.4.18]. This yields

$$\begin{aligned} \langle d^2\lambda_1(\Omega)\Phi, \Phi \rangle &= -2 \int_{\partial\Omega} \langle (\nabla \mathbf{u}'_\Phi) \nu, \nabla \mathbf{u}_\Omega \nu \rangle \langle \Phi, \nu \rangle \\ &\quad - \int_{\partial\Omega} \left(H \|(\nabla \mathbf{u}_\Omega) \nu\|^2 + \frac{\partial \|(\nabla \mathbf{u}_\Omega) \nu\|^2}{\partial \nu} \right) \langle \Phi, \nu \rangle^2. \end{aligned}$$

However, recall that, introducing Δ_τ the tangential laplacian on $\partial\Omega$, we have

$$\Delta \mathbf{u} = \Delta_\tau \mathbf{u} + H(\nabla \mathbf{u}) \nu + \frac{\partial((\nabla \mathbf{u}) \nu)}{\partial \nu}.$$

Consequently,

$$\begin{aligned}
\langle d^2 \lambda_1(\Omega) \Phi, \Phi \rangle &= -2 \int_{\partial\Omega} \langle (\nabla \mathbf{u}'_{\Phi}) \nu, \nabla \mathbf{u}_{\Omega} \nu \rangle \langle \Phi, \nu \rangle \\
&\quad - \int_{\partial\Omega} \left(H \|(\nabla \mathbf{u}_{\Omega}) \nu\|^2 + \frac{\partial \|(\nabla \mathbf{u}_{\Omega}) \nu\|^2}{\partial \nu} \right) \langle \Phi, \nu \rangle^2 \\
&= -2 \int_{\partial\Omega} \langle (\nabla \mathbf{u}'_{\Phi}) \nu, \nabla \mathbf{u}_{\Omega} \nu \rangle \langle \Phi, \nu \rangle \\
&\quad + \int_{\partial\Omega} H \|(\nabla \mathbf{u}_{\Omega}) \nu\|^2 \langle \Phi, \nu \rangle^2 - 2 \int_{\partial\Omega} \langle (\nabla \mathbf{u}_{\Omega}) \nu, \nabla p_{\Omega} \rangle \langle \Phi, \nu \rangle^2 \\
&= 2 \int_{\partial\Omega} \langle \mathbf{u}'_{\Phi}, (\nabla \mathbf{u}'_{\Phi}) \nu \rangle + \int_{\partial\Omega} H \|(\nabla \mathbf{u}_{\Omega}) \nu\|^2 \langle \Phi, \nu \rangle^2 - 2 \int_{\partial\Omega} \langle (\nabla \mathbf{u}_{\Omega}) \nu, \nabla p_{\Omega} \rangle \langle \Phi, \nu \rangle^2
\end{aligned}$$

as claimed. \square

3.2 The ball is a critical point for Problem (5) (in dimension two)

We first prove that \mathbb{B}_2 is a critical point for \mathcal{F} (recall that $\mathcal{F}(\Omega) = |\Omega| \lambda_1(\Omega)$).

Proposition 2. *The ball \mathbb{B}_2 is a critical point for the problem (5): for any $\Phi \in \mathcal{X}$, there holds*

$$\langle d\mathcal{F}(\mathbb{B}_2), \Phi \rangle = 0.$$

Proof of Proposition 2. Fix $\Phi \in \mathcal{X}$. The derivative of the volume $\text{Vol} : \Omega \mapsto |\Omega|$ at \mathbb{B}_2 in the direction Φ is given by

$$\langle d\text{Vol}(\mathbb{B}_2), \Phi \rangle = \int_{\partial\mathbb{B}_2} \langle \Phi, \nu \rangle. \quad (17)$$

For the sake of notational convenience, introduce

$$\varphi := \langle \Phi, \nu \rangle.$$

Then, (17) and Proposition 1 yield

$$\langle d\mathcal{F}(\mathbb{B}_2), \Phi \rangle = \lambda_1(\mathbb{B}_2) \int_{\partial\mathbb{B}_2} \varphi - |\mathbb{B}_2| \int_{\partial\mathbb{B}_2} \|(\nabla \mathbf{u}_{\mathbb{B}_2}) \nu\|^2 \varphi.$$

Consequently, \mathbb{B}_2 is a critical point for \mathcal{F} if, and only if,

$$|\mathbb{B}_2| \cdot \|(\nabla \mathbf{u}_{\mathbb{B}_2}) \nu\|^2 = \lambda_1(\mathbb{B}_2). \quad (18)$$

However, this relation holds because of the explicit expression given in Lemma 1. \square

3.3 The ball satisfies strong second-order conditions for Problem (5)

In this section, we turn to second-order optimality conditions for \mathcal{F} at \mathbb{B}_2 .

Expression of the second order derivative of \mathcal{F} at \mathbb{B}_2 . By the Hadamard structure theorem, since \mathbb{B}_2 is a critical point of \mathcal{F} , the second order derivative $\langle d^2\mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle$ only depends on the normal trace $\varphi = \langle \Phi, \nu \rangle$ of Φ . Let us show the following:

Lemma 4. *For any $\Phi \in \mathcal{X}$, there holds*

$$\langle d^2\mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle = 2\lambda_1(\mathbb{B}_2) \left(\int_{\partial\mathbb{B}_2} \varphi^2 \right) - \frac{2\lambda_1(\mathbb{B}_2)}{\pi} \left(\int_{\partial\mathbb{B}_2} \varphi \right)^2 + 2\pi I[\mathbf{u}'_\Phi]$$

where I is a quadratic form defined as

$$I[\mathbf{u}'_\Phi] = \int_{\mathbb{B}_2} \|\nabla \mathbf{u}'_\Phi\|^2 - \lambda_1(\mathbb{B}_2) \int_{\mathbb{B}_2} \|\mathbf{u}'_\Phi\|^2.$$

Proof of Lemma 4. Observe that we have

$$\langle d^2\mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle = \langle d^2\text{Vol}(\mathbb{B}_2)\Phi, \Phi \rangle \lambda_1(\mathbb{B}_2) + 2\langle d\text{Vol}(\mathbb{B}_2), \Phi \rangle \langle d\lambda_1(\mathbb{B}_1), \Phi \rangle + |\mathbb{B}_2| \cdot \langle d^2\lambda_1(\mathbb{B}_2)\Phi, \Phi \rangle.$$

However, we have

$$\langle d^2\text{Vol}(\mathbb{B}_2)\Phi, \Phi \rangle = \int_{\partial\mathbb{B}_2} H\varphi^2.$$

Using Proposition 1 (and noting first that $p_{\mathbb{B}_2}$, the pressure, is constant here and, second that $H = 1$) we obtain

$$\begin{aligned} \langle d^2\mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle &= \lambda_1(\mathbb{B}_2) \int_{\partial\mathbb{B}_2} \varphi^2 - 2 \int_{\partial\Omega} \varphi \int_{\partial\mathbb{B}_2} \|(\nabla \mathbf{u}_{\mathbb{B}_2})\nu\|^2 \varphi \\ &\quad + 2|\mathbb{B}_2| \cdot \int_{\partial\Omega} \langle (\nabla \mathbf{u}'_\Phi)\nu, \mathbf{u}'_\Phi \rangle + |\mathbb{B}_2| \cdot \int_{\partial\mathbb{B}_2} \|(\nabla \mathbf{u}_{\mathbb{B}_2})\nu\|^2 \varphi^2. \end{aligned}$$

From (18) this equation simplifies as

$$\begin{aligned} \langle d^2\mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle &= 2\lambda_1(\mathbb{B}_2) \left(\int_{\partial\mathbb{B}_2} \varphi^2 \right) - \frac{2\lambda_1(\mathbb{B}_2)}{|\mathbb{B}_2|} \left(\int_{\partial\mathbb{B}_2} \varphi \right)^2 + 2|\mathbb{B}_2| \cdot \int_{\partial\Omega} \langle (\nabla \mathbf{u}'_\Phi)\nu, \mathbf{u}'_\Phi \rangle \\ &= 2\lambda_1(\mathbb{B}_2) \left(\int_{\partial\mathbb{B}_2} \varphi^2 \right) - \frac{2\lambda_1(\mathbb{B}_2)}{\pi} \left(\int_{\partial\mathbb{B}_2} \varphi \right)^2 + 2\pi \cdot \int_{\partial\mathbb{B}_2} \langle (\nabla \mathbf{u}'_\Phi)\nu, \mathbf{u}'_\Phi \rangle. \end{aligned}$$

Now, multiply (14) by \mathbf{u}'_Φ and integrate by parts. We obtain

$$\begin{aligned} \int_{\mathbb{B}_2} \langle \mathbf{u}'_\Phi, \Delta \mathbf{u}'_\Phi \rangle &= \int_{\mathbb{B}_2} \langle \mathbf{u}'_\Phi, \nabla p' \rangle - \lambda_1(\mathbb{B}_2) \int_{\mathbb{B}_2} \|\mathbf{u}'_\Phi\|^2 - \langle d\lambda_1(\mathbb{B}_1), \Phi \rangle \int_{\mathbb{B}_2} \langle \mathbf{u}, \mathbf{u}'_\Phi \rangle \\ &= -\lambda_1(\mathbb{B}_2) \int_{\mathbb{B}_2} \|\mathbf{u}'_\Phi\|^2. \end{aligned}$$

To go from the first to the second line, we used the orthogonality of $\mathbf{u}_{\mathbb{B}_2}$ and of \mathbf{u}'_Φ , as well as the following fact:

$$\begin{aligned} \int_{\mathbb{B}_2} \langle \mathbf{u}'_\Phi, \nabla p' \rangle &= - \int_{\mathbb{B}_2} p' \text{div} \mathbf{u}'_\Phi + \int_{\partial\mathbb{B}_2} p' \langle \mathbf{u}'_\Phi, \nu \rangle \\ &= \int_{\partial\mathbb{B}_2} p' \langle \mathbf{u}'_\Phi, \nu \rangle \\ &= - \int_{\partial\mathbb{B}_2} p' \langle (\nabla \mathbf{u}_{\mathbb{B}_2})\nu, \nu \rangle \varphi \\ &= 0. \end{aligned}$$

We used again, in this last part, the fact that $\langle (\nabla \mathbf{u}_{\mathbb{B}_2})\nu, \nu = 0 \rangle$ (see Remark 5). Consequently, we obtain

$$\int_{\partial \mathbb{B}_2} \langle (\nabla \mathbf{u}'_{\Phi})\nu, \mathbf{u}'_{\Phi} \rangle = \int_{\mathbb{B}_2} \|\nabla \mathbf{u}'_{\Phi}\|^2 + \int_{\mathbb{B}_2} \langle \mathbf{u}'_{\Phi}, \Delta \mathbf{u}'_{\Phi} \rangle = \int_{\mathbb{B}_2} \|\nabla \mathbf{u}'_{\Phi}\|^2 - \lambda_1(\mathbb{B}_2) \int_{\mathbb{B}_2} \|\mathbf{u}_{\Phi}\|^2.$$

□

Our main lemma is the following:

Lemma 5. *There exists $c_0 > 0$ such that, for any $\Phi \in \mathcal{X}$ such that $\varphi = \langle \Phi, \nu \rangle \in \{1, \cos, \sin\}^\perp$ (for the L^2 scalar product on $L^2(\partial \mathbb{B}_2)$) there holds*

$$\langle d^2 \mathcal{F}(\mathbb{B}_2)\Phi, \Phi \rangle \geq c_0 \|\varphi\|_{W^{\frac{1}{2},2}(\partial \mathbb{B}_2)}^2.$$

The proof of this lemma relies on a diagonalisation of the quadratic form I .

Remark 7. The vector fields Φ such that $\varphi \in \text{Span}\{\cos, \sin, 1\}$ correspond to translation and dilations; of course, \mathcal{F} is constant along such deformations, whence the need to assume $\varphi \in \text{Span}\{1, \sin, \cos\}^\perp$ to obtain coercivity.

Diagonalisation of the quadratic form I . Write $\mathbf{u}_{\mathbb{B}_2} = (u_1, u_2)^\top$, and $\mathbf{u}_{\Phi'} = (u'_{\Phi,1}, u'_{\Phi,2})^\top$. With a slight abuse of notation, we write, in polar coordinates, $\mathbf{u}_{\mathbb{B}_2}(x) = \mathbf{u}(r, \theta) = \phi_{0,1}(r, \theta)\vec{e}_r$, where $\phi_{0,1}$ is given by (11). Similarly, as we know that we can assume the perturbation Φ to be normal, we identify Φ with φ , and we decompose $\varphi : \partial \mathbb{B}^2 \rightarrow \mathbb{R}$ as a Fourier series by considering the two sequences $(\alpha_n)_{n \in \mathbb{N}}$ and $(\beta_n)_{n \in \mathbb{N}}$ in $\ell^2(\mathbb{N})$ such that

$$\varphi(\theta) = \sum_{n=0}^{+\infty} (\alpha_n \cos(n\theta) + \beta_n \sin(n\theta)). \quad (19)$$

Note that, since $J_1 = -J'_0$, one has in particular

$$u_1(r, \theta) = c_1 J_1(j_{1,1}r) \sin \theta \quad \text{and} \quad u_2(r, \theta) = c_1 J_1(j_{1,1}r) \cos \theta \quad (20)$$

where

$$c_1 = \frac{1}{\sqrt{\pi} z_{1,1} |J_0(z_{1,1})|}. \quad (21)$$

As $\text{div}(\mathbf{u}'_{\Phi}) = 0$, we can write

$$\mathbf{u}'_{\Phi} = \begin{pmatrix} -\partial_y \psi'_{\Phi} \\ \partial_x \psi'_{\Phi} \end{pmatrix}$$

where $\psi_{\Phi'} \in W^{2,3}(\mathbb{B}_2)$. Taking the **curl** of (14) it appears that ψ'_{Φ} satisfies

$$\Delta^2 \psi'_{\Phi} + \lambda_1(\mathbb{B}_2) \Delta \psi'_{\Phi} = 0 \quad \text{in } B_2.$$

As $\Delta \psi'_{\Phi} + \lambda_1(\mathbb{B}_2) \psi'_{\Phi}$ is harmonic, there exist $(c_n)_{n \in \mathbb{N}}, (d_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N})$ such that

$$\Delta \psi'_{\Phi} + \lambda_1(\mathbb{B}_2) \psi'_{\Phi} = \sum_{n=0}^{+\infty} (c_n \cos(n\theta) + d_n \sin(n\theta)) r^n.$$

Solving this equation in ψ'_{Φ} and using the Bessel functions, we thus deduce that there exist two sequences $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N})$ such that

$$\psi'_{\Phi}(r, \theta) = \sum_{n=0}^{+\infty} ((a_n \cos(n\theta) + b_n \sin(n\theta))J_n(j_{1,1}r) + (c_n \cos(n\theta) + d_n \sin(n\theta))r^n). \quad (22)$$

Let us now identify all coefficients.

Lemma 6. *For every $n \geq 2$, one has*

$$a_n = \frac{j_{1,1}}{\sqrt{\pi} (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1}))} \alpha_n, \quad (23)$$

$$b_n = \frac{j_{1,1}}{\sqrt{\pi} (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1}))} \beta_n, \quad (24)$$

$$c_n = -\frac{j_{1,1}J_n(j_{1,1})}{\sqrt{\pi} (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1}))} \alpha_n, \quad (25)$$

$$d_n = -\frac{j_{1,1}J_n(j_{1,1})}{\sqrt{\pi} (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1}))} \beta_n. \quad (26)$$

Proof of Lemma 6. In what follows, to determine the coefficients a_n, b_n, c_n, d_n , we will exploit (22) and the boundary conditions

$$u'_1 = \frac{\partial \psi'_{\Phi}}{\partial x_2} = -\frac{\partial u_1}{\partial \nu} \varphi \quad \text{on } \partial \mathbb{B}_2 \quad \text{and} \quad u'_2 = -\frac{\partial \psi'_{\Phi}}{\partial x_1} = -\frac{\partial u_2}{\partial \nu} \varphi \quad \text{on } \partial \mathbb{B}_2.$$

Recall that

$$\begin{pmatrix} u'_1 \\ u'_2 \end{pmatrix} = \frac{j_{1,1}}{\sqrt{\pi}} \begin{pmatrix} \sin \theta \\ -\cos \theta \end{pmatrix} \sum_{n=0}^{+\infty} (\alpha_n \cos(n\theta) + \beta_n \sin(n\theta)) \quad \text{on } \partial \mathbb{B}_2.$$

Using that $u'_1 = \partial \psi'_{\Phi} / \partial x_2$, it follows that

$$\begin{aligned} u'_1 &= \sum_{n=0}^{+\infty} \sin \theta j_{1,1} J'_n(j_{1,1}) (a_n \cos(n\theta) + b_n \sin(n\theta)) + n \sin \theta (c_n \cos(n\theta) + d_n \sin(n\theta)) \\ &\quad + \sum_{n=0}^{+\infty} n \cos \theta J_n(j_{1,1}) (b_n \cos(n\theta) - a_n \sin(n\theta)) + n \cos \theta (d_n \cos(n\theta) - c_n \sin(n\theta)) \quad \text{on } \partial \mathbb{B}_2. \end{aligned}$$

Identifying the cosine and sine terms in the relation above yields

- in front of the constant term:

$$j_{1,1} J'_1(j_{1,1}) b_1 + d_1 = -\frac{j_{1,1}}{\sqrt{\pi}} \beta_1,$$

- in front of the term $\cos \theta$:

$$(j_{1,1} J'_2(j_{1,1}) + 2J_2(j_{1,1})) b_2 + 2d_2 = -\frac{j_{1,1}}{\sqrt{\pi}} \beta_2,$$

- in front of the term $\sin \theta$:

$$-(j_{1,1} J'_2(j_{1,1}) + 2J_2(j_{1,1})) a_2 - 2c_2 = -\frac{j_{1,1}}{\sqrt{\pi}} (2\alpha_0 - \alpha_2),$$

- in front of the term $\cos((n+1)\theta)$:

$$\begin{aligned} & (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1})) b_n + ((n+2)J_{n+2}(j_{1,1}) + j_{1,1}J'_{n+2}(j_{1,1})) b_{n+2} + (n+2)d_{n+2} \\ &= -\frac{j_{1,1}}{\sqrt{\pi}}(\beta_{n+2} - \beta_n), \end{aligned}$$

- in front of the term $\sin((n+1)\theta)$:

$$\begin{aligned} & (j_{1,1}J'_n(j_{1,1}) - nJ_n(j_{1,1})) a_n - ((n+2)J_{n+2}(j_{1,1}) + j_{1,1}J'_{n+2}(j_{1,1})) a_{n+2} - (n+2)c_{n+2} \\ &= -\frac{j_{1,1}}{\sqrt{\pi}}(-\alpha_{n+2} + \alpha_n). \end{aligned}$$

We proceed similarly with the function u'_2 . As $u'_2 = -\partial\psi'_\Phi/\partial x_1$, it follows that

$$\begin{aligned} u'_2 &= \sum_{n=0}^{+\infty} -\cos\theta j_{1,1}J'_n(j_{1,1})(a_n \cos(n\theta) + b_n \sin(n\theta)) - n \sin\theta(c_n \cos(n\theta) + d_n \sin(n\theta)) \\ &+ \sum_{n=0}^{+\infty} n \sin\theta J_n(j_{1,1})(b_n \cos(n\theta) - a_n \sin(n\theta)) + n \sin\theta(d_n \cos(n\theta) - c_n \sin(n\theta)) \quad \text{on } \partial\mathbb{B}_2. \end{aligned}$$

Identifying the cosine and sine terms in the relation above yields

- in front of the constant term:

$$-j_{1,1}J'_1(j_{1,1})a_1 - c_1 = \frac{j_{1,1}}{\sqrt{\pi}}\alpha_1,$$

- in front of the term $\cos\theta$:

$$-(j_{1,1}J'_2(j_{1,1}) + 2J_2(j_{1,1})) a_2 - 4c_2 = \frac{j_{1,1}}{\sqrt{\pi}}(2\alpha_0 + \alpha_2),$$

- in front of the term $\sin\theta$:

$$-(j_{1,1}J'_2(j_{1,1}) + 2J_2(j_{1,1})) b_2 - 4d_2 = \frac{j_{1,1}}{\sqrt{\pi}}\beta_2,$$

- in front of the term $\cos((n+1)\theta)$:

$$\begin{aligned} & (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1})) a_n - ((n+2)J_{n+2}(j_{1,1}) + j_{1,1}J'_{n+2}(j_{1,1})) a_{n+2} - 2(n+2)c_{n+2} \\ &= \frac{j_{1,1}}{\sqrt{\pi}}(\alpha_{n+2} + \alpha_n), \end{aligned}$$

- in front of the term $\sin((n+1)\theta)$:

$$\begin{aligned} & (nJ_n(j_{1,1}) - j_{1,1}J'_n(j_{1,1})) b_n - ((n+2)J_{n+2}(j_{1,1}) + j_{1,1}J'_{n+2}(j_{1,1})) b_{n+2} - 2(n+2)d_{n+2} \\ &= \frac{j_{1,1}}{\sqrt{\pi}}(\beta_{n+2} + \beta_n). \end{aligned}$$

We obtain the desired expression by combining together all the above relations. □

Proof of Lemma 5. Let us compute I : we have

$$\begin{aligned}
I[\mathbf{u}'_\Phi] &= \int_{\partial\mathbb{B}_2} \left(u'_1 \frac{\partial u'_1}{\partial \nu} + u'_2 \frac{\partial u'_2}{\partial \nu} \right) d\sigma \\
&= -\frac{j_{1,1}}{\sqrt{\pi}} \int_{\partial\mathbb{B}_2} \varphi \sin \theta \left[\sin \theta \frac{\partial^2 \psi'_\Phi}{\partial r^2} - \cos \theta \frac{\partial \psi'_\Phi}{\partial \theta} + \cos \theta \frac{\partial^2 \psi'_\Phi}{\partial r \partial \theta} \right] d\sigma \\
&\quad + \frac{j_{1,1}}{\sqrt{\pi}} \int_{\partial\mathbb{B}_2} \varphi \cos \theta \left[-\cos \theta \frac{\partial^2 \psi'_\Phi}{\partial r^2} - \sin \theta \frac{\partial \psi'_\Phi}{\partial \theta} + \sin \theta \frac{\partial^2 \psi'_\Phi}{\partial r \partial \theta} \right] d\sigma \\
&= \frac{j_{1,1}}{\sqrt{\pi}} \int_{\partial\mathbb{B}_2} \varphi \frac{\partial^2 \psi'_\Phi}{\partial r^2}.
\end{aligned}$$

Using the expansion (22) of ψ , we get

$$\left. \frac{\partial^2 \psi'_\Phi}{\partial r^2} \right|_{\partial\mathbb{B}_2} = \sum_{n=0}^{+\infty} j_{1,1}^2 (a_n \cos \theta + b_n \sin \theta) J_n''(j_{1,1}) + \sum_{n=1}^{+\infty} n(n-1) (c_n \cos(n\theta) + d_n \sin(n\theta)),$$

and we thus obtain

$$\begin{aligned}
I[\mathbf{u}'_\Phi] &= -\frac{j_{1,1}}{\sqrt{\pi}} \left[\pi j_{1,1}^2 \sum_{n=1}^{+\infty} (a_n \alpha_n + b_n \beta_n) J_n''(j_{1,1}) + \pi \sum_{n=1}^{+\infty} n(n-1) (c_n \alpha_n + d_n \beta_n) \right] \\
&= -j_{1,1}^2 \sum_{n=1}^{+\infty} \frac{j_{1,1}^2 J_n''(j_{1,1}) - n(n-1) J_n(j_{1,1})}{n J_n(j_{1,1}) - j_{1,1} J_n'(j_{1,1})} (\alpha_n^2 + \beta_n^2).
\end{aligned}$$

Now, by definition of Bessel functions of the first order, one has

$$J_n''(x) + \frac{1}{x} J_n'(x) + \left(1 - \frac{n^2}{x^2} \right) J_n(x) = 0$$

for every $x > 0$. We infer that $j_{1,1}^2 J_n''(j_{1,1}) - n(n-1) J_n(j_{1,1}) = (n - j_{1,1}^2) J_n(j_{1,1}) - j_{1,1} J_n'(j_{1,1})$, yielding finally

$$I = -j_{1,1}^2 \sum_{n=1}^{+\infty} \frac{(n - j_{1,1}^2) J_n(j_{1,1}) - j_{1,1} J_n'(j_{1,1})}{n J_n(j_{1,1}) - j_{1,1} J_n'(j_{1,1})} (\alpha_n^2 + \beta_n^2). \quad (27)$$

Recall that we assume $\varphi \in \langle 1 \rangle^\perp$ where 1 is the constant function and $\langle 1 \rangle$ is its L^2 orthogonal subspace.

According to (27), one has

$$\begin{aligned}
\langle d^2 \mathcal{F}(\mathbb{B}_2) \Phi, \Phi \rangle &= 2\pi I[\mathbf{u}'_\Phi] + 2\lambda_1(\mathbb{B}_2) \int_{\partial\mathbb{B}_2} \varphi^2 - \frac{2\lambda_1(\mathbb{B}_2)}{\pi} \left(\int_{\partial\mathbb{B}_2} \varphi \right)^2 \\
&= 2\pi j_{1,1}^2 \sum_{n=1}^{+\infty} \frac{n J_n(j_{1,1}) - j_{1,1} J_n'(j_{1,1}) - (n - j_{1,1}^2) J_n(j_{1,1}) - j_{1,1} J_n'(j_{1,1})}{n J_n(j_{1,1}) + j_{1,1} J_n'(j_{1,1})} (\alpha_n^2 + \beta_n^2).
\end{aligned}$$

Furthermore, recall that $x J_n'(x) = n J_n(x) - x J_{n+1}(x)$ for every $x \geq 0$, so that $j_{1,1} J_n'(j_{1,1}) - n J_n(j_{1,1}) = -j_{1,1} J_{n+1}(j_{1,1})$.

This leads to

$$\langle d^2 \mathcal{F}(\mathbb{B}_2) \Phi, \Phi \rangle = 2\pi j_{1,1}^4 \sum_{n=2}^{+\infty} \gamma_n (\alpha_n^2 + \beta_n^2) \quad \text{with } \gamma_n = \frac{J_n(j_{1,1})}{j_{1,1} J_{n+1}(j_{1,1})}. \quad (28)$$

(we recall that we have supposed φ orthogonal to $1, \cos \theta, \sin \theta$, so $a_0 = a_1 = b_1 = 0$. Since the first positive zero of J_n is greater than $j_{1,1}$ for all $n \geq 2$, one has $\gamma_n > 0$ for every $n \geq 2$. Furthermore, using that $J_n(x) \sim \frac{1}{n!} \left(\frac{x}{2}\right)^n$ as $n \rightarrow +\infty$, we infer that

$$\gamma_n \sim 2(n+1)/j_{1,1}^2 \quad \text{as } n \rightarrow +\infty.$$

Since the $H^{1/2}$ norm is given by

$$\|\varphi\|_{H^{1/2}} = \sum_{n=2}^{\infty} n(a_n^2 + b_n^2)$$

the conclusion follows. \square

4 Necessary optimality conditions: proof of Theorem 4

As alluded to in the introduction, Theorem 3 follows from Theorem 4, so that it is clearer to begin with the proof of the latter (nevertheless, we give a proof of 3 in section 5 that only relies on the computations of the semi-differential analysed in the present section). This theorem relies on several results on the semi-differentiability of eigenvalues. To write things down in a precise way, we consider a domain Ω whose boundary is of class \mathcal{C}^3 . Although $\lambda_1(\Omega)$ may not be simple, it has finite multiplicity, say N , and the map \mathcal{F} is semi-shape differentiable at Ω [20, Theorem 2.1, Chapter 1] (see also [13, Theorem 2.1 and Remark 2.2] and [15]), in the sense that, for any vector field $\Phi \in \mathcal{X}$, the limit

$$\langle \partial \mathcal{F}(\Omega), \Phi \rangle := \lim_{t \rightarrow 0^+} \frac{\mathcal{F}((\text{Id} + t\Phi)\Omega) - \mathcal{F}(\Omega)}{t}$$

exists. This fact relies on the semi-differentiability of λ_1 . By [20, Theorem 2.1, Chapter 1], if we let E_1 denote the first eigenspace associated with $\lambda_1(\Omega)$, there holds

$$\langle \partial \lambda_1(\Omega), \Phi \rangle = - \min_{\mathbf{u} \in E_1, \int_{\Omega} \|\mathbf{u}\|^2 = 1} \int_{\partial \mathbb{B}_3} \|(\nabla \mathbf{u})\nu\|^2 \langle \Phi, \nu \rangle.$$

Let $\{\mathbf{u}_1, \dots, \mathbf{u}_N\}$ be an orthonormal basis of E_1 . Using the differentiability of the volume, we deduce that

$$\begin{aligned} \langle \partial \mathcal{F}(\Omega), \Phi \rangle &= \frac{2}{3} \lambda_1(\Omega) |\Omega|^{\frac{1}{3}} \int_{\partial \Omega} \langle \Phi, \nu \rangle - |\Omega|^{\frac{2}{3}} \min_{\mathbf{u} \in E_1, \int_{\Omega} \|\mathbf{u}\|^2 = 1} \int_{\partial \Omega} \|(\nabla \mathbf{u})\nu\|^2 \langle \Phi, \nu \rangle \\ &= \min_{\mathbf{u} \in E_1, \int_{\Omega} \|\mathbf{u}\|^2 = 1} \int_{\partial \Omega} \left(\frac{2}{3} \lambda_1(\Omega) |\Omega|^{\frac{1}{3}} - |\Omega|^{\frac{2}{3}} \cdot \|(\nabla \mathbf{u})\nu\|^2 \right) \langle \Phi, \nu \rangle \\ &= \min_{(\alpha_i)_{i=1, \dots, N}, \sum_{i=1}^N \alpha_i^2 = 1} \sum_{i,j=1}^N \int_{\partial \Omega} \left(\frac{2}{3} \lambda_1(\Omega) |\Omega|^{\frac{1}{3}} \alpha_i^2 - |\Omega|^{\frac{2}{3}} \alpha_i \alpha_j \langle (\nabla \mathbf{u}_i)\nu, (\nabla \mathbf{u}_j)\nu \rangle \right) \langle \Phi, \nu \rangle. \end{aligned}$$

Let

$$M_{\Phi} := \frac{2}{3} \lambda_1(\Omega) |\Omega|^{\frac{1}{3}} \int_{\partial \Omega} \langle \Phi, \nu \rangle I_3 - |\Omega|^{\frac{2}{3}} \left(\int_{\partial \Omega} \langle (\nabla \mathbf{u}_i)\nu, (\nabla \mathbf{u}_j)\nu \rangle \langle \Phi, \nu \rangle \right)_{1 \leq i, j \leq N}.$$

I_3 is the 3×3 identity matrix. M_{Φ} is a real, symmetric matrix, and is thus diagonalisable. Let Σ_{Φ} be its spectrum. If Ω^* is optimal, then the optimality conditions read:

$$\forall \Phi \in \mathcal{X}, \Sigma_{\Phi} \subset [0; +\infty).$$

However, as $\Sigma_{-\Phi} = -\Sigma_{\Phi}$, we deduce that, if Ω is optimal then

$$\forall \Phi \in \mathcal{X}, \Sigma_{\Phi} = \{0\}.$$

As M_{Φ} is symmetric, we deduce that, for any Φ , there holds

$$M_{\Phi} = 0.$$

We thus obtain the following fact: if Ω^* is optimal then for any $(i, j) \in \{1, \dots, N\}^2$, for any $x \in \partial\Omega$,

$$\langle (\nabla \mathbf{u}_i)\nu, (\nabla \mathbf{u}_j)\nu \rangle = \frac{2}{3} \cdot \frac{\lambda_1(\Omega^*)}{|\Omega|^{\frac{1}{3}}} \delta_{i,j} = \frac{1}{\sigma_{\Omega^*}^2} \delta_{i,j}.$$

In this expression, $\delta_{i,j}$ is the usual Kronecker symbol and $\sigma_{\Omega^*}^2 = \frac{3}{2} \cdot \frac{|\Omega|^{\frac{1}{3}}}{\lambda_1(\Omega^*)}$.

However, since each \mathbf{u}_i is divergence free we have $\langle (\nabla \mathbf{u}_i)\nu, \nu \rangle = 0$ (see Remark 5). In particular, the multiplicity of the first eigenspace E_1 is at most 2 (we recall that we are in dimension 3 here).

Now, let us argue by contradiction and assume that $\lambda_1(\Omega^*)$ has multiplicity 2. Let $(\mathbf{u}_1, \mathbf{u}_2)$ be an orthonormal basis of E_1 . From the previous computation, we deduce that, for any $x \in \partial\Omega^*$, $\{\sigma_{\Omega^*}(\nabla \mathbf{u}_1)\nu, \sigma_{\Omega^*}(\nabla \mathbf{u}_2)\nu\}$ is an orthonormal basis of the tangent plane $T_x(\partial\Omega)$. Let us now argue using a scheme of proof inspired by [25, Proof of Theorem 2] to reach a contradiction.

For $i = 1, 2$, let \mathbf{v}_i be the solution of

$$\begin{cases} -\Delta \mathbf{v}_i + \nabla p_i = \lambda_1(\Omega^*)\mathbf{v}_i + \omega \mathbf{u}_i & \text{in } \Omega^*, \\ \nabla \cdot \mathbf{v}_i = 0 & \text{in } \Omega^*, \\ \mathbf{v}_i \in E_1^\perp, & \\ \mathbf{v}_i = -(\nabla \mathbf{u}_i)\nu & \text{on } \partial\Omega^*, \end{cases} \quad (29)$$

where E_1^\perp is the L^2 -orthogonal subspace to E_1 and

$$\omega = - \int_{\partial\Omega^*} \|(\nabla \mathbf{u}_i)\nu\|^2 = -\text{Per}(\partial\Omega^*) \cdot \frac{2}{3} \cdot \frac{\lambda_1(\Omega^*)}{|\Omega|^{\frac{1}{3}}}. \quad (30)$$

We have the following result:

Lemma 7. *For $i = 1, 2$, \mathbf{v}_i is uniquely defined.*

Proof of Lemma 7. The uniqueness of \mathbf{v}_i follows from the Fredholm alternative. Regarding the existence of \mathbf{v}_i , it is clear that the energy functional

$$\mathcal{E} : W_0^{1,2}(\Omega)^3 \ni \mathbf{v} \mapsto \frac{1}{2} \int_{\Omega^*} \|\nabla \mathbf{v}\|^2 - \frac{\lambda_1(\Omega^*)}{2} \int_{\Omega^*} \|\mathbf{v}\|^2$$

is coercive on

$$X_i := \{\mathbf{v} \in W_0^{1,2}(\Omega)^2, \nabla \cdot \mathbf{v} = 0, \mathbf{v} = -(\nabla \mathbf{u}_i)\nu \text{ on } \partial\Omega^*\} \cap E_1^\perp.$$

We let \mathbf{v}_i be a minimiser of \mathcal{E} on X_i . In particular, from the Euler-Lagrange equation we deduce that

$$-\Delta \mathbf{v}_i - \lambda_1(\Omega^*)\mathbf{v}_i \in (\{\mathbf{v} \in W_0^{1,2}(\Omega)^3, \nabla \cdot \mathbf{v} = 0\} \cap E_1^\perp).$$

Thus, there exist a function p_i and two coefficients $(\alpha_{i,j})_{j=1,2}$ such that

$$-\Delta \mathbf{v}_i - \lambda_1(\Omega^*) \mathbf{v}_i = -\nabla p_i + \sum_{j=1,2} \alpha_{i,j} \mathbf{u}_j.$$

Multiplying by \mathbf{u}_j , using the fact that $(\nabla \mathbf{u}_i) \nu$ and $(\nabla \mathbf{u}_j) \nu$ are orthogonal on $\partial\Omega^*$, we deduce that

$$\alpha_{i,j} = \omega \delta_{i,j}$$

where ω is defined by (30). □

Turning back to the proof of Theorem 4, observe that

$$\nabla(\langle \mathbf{v}_1, \mathbf{v}_2 \rangle) = (\nabla \mathbf{v}_1) \mathbf{v}_2 + (\nabla \mathbf{v}_2) \mathbf{v}_1.$$

Since \mathbf{v}_1 and \mathbf{v}_2 are orthogonal on $\partial\Omega^*$, we deduce that, letting X^τ denote the tangential part of a vector field X on $\partial\Omega^*$, there holds

$$((\nabla \mathbf{v}_1) \mathbf{v}_2)^\tau = -((\nabla \mathbf{v}_2) \mathbf{v}_1)^\tau.$$

Furthermore, since $\nabla \langle \mathbf{v}_1, \mathbf{v}_2 \rangle$ is colinear to ν , and since \mathbf{v}_1 is orthogonal to ν there holds

$$0 = \langle \mathbf{v}_1, \nabla \langle \mathbf{v}_1, \mathbf{v}_2 \rangle \rangle.$$

Developing the gradient yields

$$\begin{aligned} 0 &= \langle \mathbf{v}_1, \nabla \langle \mathbf{v}_1, \mathbf{v}_2 \rangle \rangle = \langle \mathbf{v}_1, (\nabla \mathbf{v}_1) \mathbf{v}_2 \rangle + \langle \mathbf{v}_1, (\nabla \mathbf{v}_2) \mathbf{v}_1 \rangle \\ &= \langle \mathbf{v}_1, (\nabla \mathbf{v}_2) \mathbf{v}_1 \rangle + \sum_{i,j} \mathbf{v}_{1,i} \mathbf{v}_{2,j} \frac{\partial \mathbf{v}_{1,i}}{\partial x_j} = \langle \mathbf{v}_1, (\nabla \mathbf{v}_2) \mathbf{v}_1 \rangle + \frac{1}{2} \sum_{i,j} \mathbf{v}_{2,j} \frac{\partial \mathbf{v}_{1,i}^2}{\partial x_j} \\ &= \langle \mathbf{v}_1, (\nabla \mathbf{v}_2) \mathbf{v}_1 \rangle \end{aligned}$$

a.e. on $\partial\Omega$, where we used once again the optimality condition to write that $\|\mathbf{v}_1\|^2$ is constant on $\partial\Omega^*$, and the fact that \mathbf{v}_2 is tangential.

Similarly,

$$0 = \langle \mathbf{v}_2, (\nabla \mathbf{v}_1) \mathbf{v}_2 \rangle = -\langle \mathbf{v}_2, (\nabla \mathbf{v}_2) \mathbf{v}_1 \rangle.$$

Here again the fact that \mathbf{v}_2 is tangential is instrumental. Thus $(\nabla \mathbf{v}_2) \mathbf{v}_1$ is orthogonal to both \mathbf{v}_1 and \mathbf{v}_2 , so that there exists a function f satisfying

$$(\nabla \mathbf{v}_2) \mathbf{v}_1 = f \nu.$$

For the same reason,

$$(\nabla \mathbf{v}_1) \mathbf{v}_2 = g \nu$$

for some function g . Finally, letting $[X, Y]$ be the Lie bracket of two vector fields, we deduce that

$$[(\nabla \mathbf{v}_1) \mathbf{v}_2, (\nabla \mathbf{v}_2) \mathbf{v}_1] = (f - g) \nu.$$

However, $(\nabla \mathbf{v}_2) \mathbf{v}_1$ and $(\nabla \mathbf{v}_1) \mathbf{v}_2$ are tangential, whence their Lie bracket also is, so that

$$[(\nabla \mathbf{v}_1) \mathbf{v}_2, (\nabla \mathbf{v}_2) \mathbf{v}_1] = 0.$$

In particular, this implies that $\partial\Omega^*$ admits a set of two pointwise orthogonal, commuting vector fields. This implies that the Riemann tensor associated with $\partial\Omega^*$ endowed with the induced metric is zero (we refer for instance to [34, Theorem 3.1.7]). In particular, the Gauß curvature of $\partial\Omega^*$ is zero everywhere, which is impossible.

5 The ball does not minimise F in dimension 3: proof of Theorem 3

As in the two-dimensional case, we need to describe precisely the eigenspace associated with $\lambda_1(\mathbb{B}_3)$. The description of this first eigenvalue is less standard and mostly due to Saks [44]. However, [44] is quite an involved article and, for the sake of convenience, we recall the main steps of [44] that we use to derive the results of this preliminary section in Appendix A.

Let $\omega > 0$ be the first positive root of the equation $\tan(x) = x$. The main result is the following:

Proposition 3. $\lambda_1(\mathbb{B}_3)$ has multiplicity 3. An orthogonal basis of the associated eigenspace E_1 is given by

$$\left\{ \mathbf{U}_1 = \psi(r) \begin{pmatrix} 0 \\ z \\ -y \end{pmatrix}, \mathbf{U}_2 = \psi(r) \begin{pmatrix} -z \\ 0 \\ x \end{pmatrix}, \mathbf{U}_3 = \psi(r) \begin{pmatrix} -y \\ x \\ 0 \end{pmatrix} \right\}$$

where

$$\psi(r) = \frac{\sin(\omega r)}{\omega^2 r^3} - \frac{\cos(\omega r)}{\omega r^2}.$$

Finally, for any $i \in \{1, 2, 3\}$, there holds

$$\int_{\mathbb{B}_3} \|\mathbf{U}_i\|^2 = \frac{2\pi}{3\omega^4} (\omega^2 - \sin^2 \omega) =: A.$$

As mentioned, we prove this multiplicity result in Appendix A. In particular, applying Theorem 4, the conclusion follows.

Let us now give another proof of Theorem 3 that only relies on the analysis of the semi-differential as given section 4. Recall that the semi-differential of λ_1 at B_3 in direction $\Phi \in \mathcal{X}$ (see Lemma 3 for the definition of \mathcal{X}) reads

$$\langle \partial\lambda_1(\mathbb{B}_3), \Phi \rangle = \min_{\substack{\mathbf{U} \in \text{Span}\{\mathbf{U}_i/A\}_{1 \leq i \leq 3} \\ \|\mathbf{U}\|_{L^2(\mathbb{B}_3)} = 1}} - \int_{\partial\mathbb{B}_3} \|\nabla\mathbf{U}\nu\|^2 \langle \Phi, \nu \rangle.$$

It follows that

$$\begin{aligned} \langle \partial\lambda_1(\mathbb{B}_3), \Phi \rangle &= \min_{\substack{(\alpha_i)_{1 \leq i \leq 3} \in \mathbb{R}^3 \\ \alpha_1^2 + \alpha_2^2 + \alpha_3^2 = 1}} - \frac{1}{A^2} \int_{\partial\mathbb{B}_3} \left\| \sum_{i=1}^3 \alpha_i \nabla\mathbf{U}_i \nu \right\|^2 \langle \Phi, \nu \rangle \\ &= \frac{1}{A^2} \mu_1(\mathcal{M}_{\partial\mathbb{B}_3}(\langle \Phi, \nu \rangle)), \end{aligned}$$

where $\mu_1(\mathcal{M}_{\partial\mathbb{B}_3}(\langle \Phi, \nu \rangle))$ denotes the lowest eigenvalue of the matrix

$$\mathcal{M}_{\partial\mathbb{B}_3}(\langle \Phi, \nu \rangle) = \left(- \int_{\partial\mathbb{B}_3} (\nabla\mathbf{U}_i \nu \cdot \nabla\mathbf{U}_j \nu) \langle \Phi, \nu \rangle \right)_{1 \leq i, j \leq 3}.$$

From the explicit expressions for the vector fields \mathbf{U}_i provided in Proposition 3, one computes successively

$$\nabla\mathbf{U}_1 \nu = \begin{pmatrix} 0 \\ -z\psi'(1) \\ y\psi'(1) \end{pmatrix}, \quad \nabla\mathbf{U}_2 \nu = \begin{pmatrix} z\psi'(1) \\ 0 \\ -x\psi'(1) \end{pmatrix}, \quad \nabla\mathbf{U}_3 \nu = \begin{pmatrix} -y\psi'(1) \\ x\psi'(1) \\ 0 \end{pmatrix}$$

and

$$\mathcal{M}_{\partial\mathbb{B}_3}(\langle\Phi, \nu\rangle) = \psi'(1)^2 \begin{pmatrix} -\int_{\partial\mathbb{B}_3} (y^2 + z^2)\langle\Phi, \nu\rangle & \int_{\partial\mathbb{B}_3} xy\langle\Phi, \nu\rangle & \int_{\partial\mathbb{B}_3} xz\langle\Phi, \nu\rangle \\ \int_{\partial\mathbb{B}_3} xy\langle\Phi, \nu\rangle & -\int_{\partial\mathbb{B}_3} (x^2 + z^2)\langle\Phi, \nu\rangle & \int_{\partial\mathbb{B}_3} yz\langle\Phi, \nu\rangle \\ \int_{\partial\mathbb{B}_3} xz\langle\Phi, \nu\rangle & \int_{\partial\mathbb{B}_3} yz\langle\Phi, \nu\rangle & -\int_{\partial\mathbb{B}_3} (x^2 + y^2)\langle\Phi, \nu\rangle \end{pmatrix}.$$

Furthermore, one easily computes $\psi'(1) = \sin \omega \neq 0$. Observe that $\mathcal{M}_{\partial\mathbb{B}_3}(\langle\Phi, \nu\rangle)$ is real symmetric. We infer from the previous computations that the semi-differential of \mathcal{F} , defined by (5), at \mathbb{B}_3 in direction Φ is given by

$$\begin{aligned} \langle\partial\mathcal{F}(\mathbb{B}_3), \Phi\rangle &= \frac{2}{3}\lambda_1(B_3)|\mathbb{B}_3|^{-1/3} \int_{\partial\mathbb{B}_3} \langle\Phi, \nu\rangle + \frac{|\mathbb{B}_3|^{2/3}}{A^2} \mu_1(\mathcal{M}_{\partial\mathbb{B}_3}(\langle\Phi, \nu\rangle)) \\ &= |\mathbb{B}_3|^{-1/3} \left(\frac{2}{3}\omega^2 \int_{\partial\mathbb{B}_3} \langle\Phi, \nu\rangle + \omega^2 \mu_1(\mathcal{M}_{\partial\mathbb{B}_3}(\langle\Phi, \nu\rangle)) \right) \\ &= |\mathbb{B}_3|^{-1/3} \mu_1 \left(\frac{2}{3}\omega^2 \int_{\partial\mathbb{B}_3} \langle\Phi, \nu\rangle I_3 + \omega^2 \mathcal{M}_{\partial\mathbb{B}_3}(\langle\Phi, \nu\rangle) \right) \end{aligned}$$

where I_3 denotes the identity matrix of size 3. Introduce the 3×3 matrix

$$\widehat{M}_\Phi := \frac{2}{3}\omega^2 \int_{\partial\mathbb{B}_3} \langle\Phi, \nu\rangle I_3 + \omega^2 \mathcal{M}_{\partial\mathbb{B}_3}(\langle\Phi, \nu\rangle)$$

Observe that the trace of the matrix \widehat{M}_Φ is

$$\text{Tr } \widehat{M}_\Phi = 2\omega^2 - 2\frac{\omega^4 \sin^2 \omega}{\omega^2 - \sin^2 \omega} = 0,$$

meaning, as expected, that the functional \mathcal{F} is dilation invariant. Using this observation, we claim that the non optimality of \mathbb{B}_3 will be proven whenever one shows that \widehat{M}_Φ is not the null matrix. Indeed, in that case, since \widehat{M}_Φ is real symmetric, we will get that \widehat{M}_Φ has two eigenvalues with opposite sign, yielding the existence of a perturbation which strictly decreases the functional \mathcal{F} . It suffices for instance to find Φ such that

$$\int_{\partial\mathbb{B}_3} yz\langle\Phi, \nu\rangle \neq 0.$$

Note that $\nu = x\vec{e}_x + y\vec{e}_y + z\vec{e}_z$ on $\partial\mathbb{B}_3$, in cartesian coordinates. We choose Φ given by $\Phi(x, y, z) = z\vec{e}_y$ and we are done.

6 Conclusion and open problems

There are several questions that we believe are interesting, but would require further work and, most likely, the use of different tools. Let us give some that we believe are the most ambitious.

Characterising the optimal domain Ω^* . Although we have proved that, in \mathbb{R}^2 , the ball \mathbb{B}_2 is likely to be an optimiser, it is not clear to us how one would approach such a result. Indeed, in the vectorial case, standard rearrangement and symmetrisation tools are bound to fail. One might then try to use an alternative approach to the Faber-Krahn inequality, typically using an overdetermined problem framework, but, to the best of our knowledge, no generalisation of the Serrin theorem can accommodate incompressibility constraints. In three dimensions, it seems useful to perform some numerical simulations to have at least an idea of the possible optimal sets. Following our Corollary 1, one can wonder whether it is a particular torus;

A priori regularity of the optimal domain. Following Theorem 1, we know that an optimal domain Ω^* exists in the class of quasi-open sets. Nevertheless, it seems extremely ambitious, at this stage, to develop an *a priori* regularity theory for this optimal set. Following the usual approach to such problems, see e.g. [28, chapter3], the first step in that direction would be to establish that Ω^* is, in fact, open. To derive further properties, the optimality conditions given in Theorem 4 would probably play a crucial role in the development of an appropriate blow-up theory.

Acknowledgments

The authors warmly thank Dorin Bucur, who provided the main argument in the proof of existence. I. M-F thanks Davide Buoso for helpful conversations. I. M-F was partially funded by a PSL Young Researcher Starting Grant 2023. The support of the CNRS through a 2023 PEPS project is also acknowledged.

Appendix

A The first Stokes eigenvalue in \mathbb{B}_3

In this appendix, we collect several facts about $\lambda_1(\mathbb{B}_3)$ and its associated eigenfunctions. The main goal is to prove the following proposition:

Proposition 4. $\lambda_1(\mathbb{B}_3)$ has multiplicity 3. An orthonormal basis of the associated eigenspace E_1 is given by

$$\left\{ \mathbf{U}_1 = \psi(r) \begin{pmatrix} 0 \\ z \\ -y \end{pmatrix}, \mathbf{U}_2 = \psi(r) \begin{pmatrix} -z \\ 0 \\ x \end{pmatrix}, \mathbf{U}_3 = \psi(r) \begin{pmatrix} -y \\ x \\ 0 \end{pmatrix} \right\}$$

where

$$\psi(r) = \frac{\sin(\omega r)}{\omega^2 r^3} - \frac{\cos(\omega r)}{\omega r^2}.$$

The proof of this proposition is essentially contained in [44], which actually gives more general results. For the reader's convenience however, we gather here the main facts necessary to obtain this result.

Eigenvalues of the curl operator. A typical approach to Stokes eigenvalue problems is to factorise the vector Laplacian Δ as $\Delta = \mathbf{curl}(\mathbf{curl})$ in the case of incompressible flows.

The spectral problem for the **curl** operator is the following: find (ξ, \mathbf{u}_ξ) , $\xi \in \mathbb{C}$, $\mathbf{u}_\xi \neq 0$ such that

$$\begin{cases} \mathbf{curl}(\mathbf{u}_\xi) = \xi \mathbf{u}_\xi & \text{in } \mathbb{B}_3, \\ \langle \mathbf{u}_\xi, \nu \rangle = 0 & \text{on } \partial \mathbb{B}_3. \end{cases} \quad (31)$$

In simply connected domains, it is possible to show the following spectral decomposition theorem [42]:

Theorem 5 (Spectral decomposition for the **curl**). *There exist two sequences $\{\mu_{k,\pm}\}_{k \in \mathbb{N}}$ of eigenfunctions ordered as follows*

$$-\infty \longleftarrow \mu_{k+1,-} \leq \mu_{k,-} \leq \dots \leq \mu_{1,-} < 0 < \mu_{1,+} \leq \dots \leq \mu_{k,+} \leq \mu_{k+1,+} \longrightarrow \infty,$$

and the associated eigenfunctions $\{\mathbf{u}_{k,\pm}\}_{k \in \mathbb{N}}$ form a Hilbert basis of the space

$$X := \{ \mathbf{u} \in L^2(\Omega)^3, \nabla \cdot \mathbf{u} = 0 \text{ in } \Omega, \langle \mathbf{u}, \nu \rangle = 0 \text{ on } \partial \Omega \}.$$

In this theorem, we use (implicitly) the fact that there is a natural notion of trace for zero-divergence vector fields.

Some explicit computations in the case of the ball. We now give a procedure to construct eigenfunctions. All the results stated are contained, in one way or another, in [44, 12]. We use the standard spherical coordinates (r, θ, ϕ) , and the associated orthonormal basis $(\vec{e}_r, \vec{e}_\theta, \vec{e}_\phi)$. Recall that in this coordinate frame we have

$$\begin{cases} x = r \sin \theta \cos(\phi), \\ y = r \sin \theta \sin(\phi), \\ z = r \cos(\phi) \end{cases} \quad \text{and} \quad \begin{cases} \vec{e}_r = \frac{x\vec{e}_x + y\vec{e}_y + z\vec{e}_z}{\sqrt{x^2 + y^2 + z^2}} \\ \vec{e}_\theta = \frac{(x\vec{e}_x + y\vec{e}_y)z - (x^2 + y^2)\vec{e}_z}{\sqrt{x^2 + y^2 + z^2}\sqrt{x^2 + y^2}} \\ \vec{e}_\phi = \frac{-y\vec{e}_x + x\vec{e}_y}{\sqrt{x^2 + y^2}} \end{cases}$$

Lemma 8. *Let (ξ, \mathbf{u}_ξ) be a solution of (31) in \mathbb{B}_3 and write, in spherical coordinates,*

$$\mathbf{u}_\xi = u_{\xi,r}\vec{e}_r + u_{\xi,\theta}\vec{e}_\theta + u_{\xi,\phi}\vec{e}_\phi.$$

Then:

1. *There holds $u_{\xi,r} \neq 0$.*
2. *$ru_{\xi,r}$ is an eigenfunction of the (scalar) Dirichlet-Laplace operator, associated with the eigenvalue ξ^2 .*

Proof of Lemma 8. 1. Recall that, in polar coordinates, we have (for a vector field $\mathbf{u} = u_r\vec{e}_r + u_\theta\vec{e}_\theta + u_\phi\vec{e}_\phi$),

$$\begin{aligned} \mathbf{curl}(\mathbf{u}) &= \frac{1}{r \sin \theta} \left(\frac{\partial(u_\phi \sin \theta)}{\partial \theta} - \frac{\partial u_\theta}{\partial \phi} \right) \vec{e}_r \\ &\quad + \frac{1}{r} \left(\frac{1}{\sin \theta} \frac{\partial u_r}{\partial \phi} - \frac{\partial(ru_\phi)}{\partial r} \right) \vec{e}_\theta \\ &\quad + \frac{1}{r} \left(\frac{\partial(ru_\theta)}{\partial r} - \frac{\partial u_r}{\partial \theta} \right) \vec{e}_\phi. \end{aligned}$$

Thus, if $u_{\xi,r} = 0$ we deduce that $(u_{\xi,\theta}, u_{\xi,\phi})$ solves (in particular) the system

$$\begin{cases} -\frac{\partial(ru_\phi)}{\partial r} = \xi r u_\theta, \\ \frac{\partial(ru_\theta)}{\partial r} = \xi r u_\phi. \end{cases}$$

Letting $w_{\theta/\phi} := ru_{\xi,\theta/\phi}$ we obtain

$$-\frac{\partial^2 w_{\theta/\phi}}{\partial r^2} = \xi^2 w_{\theta/\phi}, \quad w_{\theta/\phi}(0) = w'_{\theta/\phi}(0) = 0.$$

The Cauchy-Lipschitz theorem implies $w_{\theta/\phi} \equiv 0$, a contradiction.

2. Let (ξ, \mathbf{u}_ξ) be a solution of (31). Define $v := \langle x, \mathbf{u} \rangle = ru_{\xi,r}$. We have

$$-\Delta v = \langle x, \Delta \mathbf{u} \rangle - 2 \operatorname{div}(\mathbf{u}) = \xi^2 \langle x, \mathbf{u} \rangle.$$

Here, we used that $\operatorname{div}(\mathbf{u}) = 0$, whence $-\Delta \mathbf{u} = \mathbf{curl}(\mathbf{curl}(\mathbf{u}))$. Furthermore, $v = \langle x, \mathbf{u} \rangle = 0$ on $\partial \mathbb{B}_3$. As $u_{\xi,r} \neq 0$, the conclusion follows. □

Explicit expression for the lowest eigenvalue of the curl operator in \mathbb{B}_3 . Let $\omega > 0$ be defined as the first positive root of $x = \tan(x)$. We have the following proposition:

Proposition 5. ω is the lowest positive eigenvalue of the **curl** operator and $-\omega$ is the largest negative eigenvalue of the **curl** operator. Both have multiplicity 3.

In order to prove this result, recall the following description of the second eigenspace of the Laplacian:

Lemma 9. Let ψ be defined as

$$\psi(r) = \frac{\sin(\omega r)}{\omega^2 r^3} - \frac{\cos(\omega r)}{\omega r^2}.$$

The second eigenvalue μ_2 of the Dirichlet-Laplacian in \mathbb{B}_3 is ω^2 . Furthermore, $\mu_2(\mathbb{B}_3)$ has multiplicity 3. An orthogonal basis of the eigenspace $E_{\omega^2}^{-\Delta}$ is (v_1, v_2, v_3) , where

$$\begin{cases} v_1 : (r, \theta, \phi) \mapsto r\psi(r) \sin \theta \cos \phi, \\ v_2 : (r, \theta, \phi) \mapsto r\psi(r) \sin \theta \sin \phi, \\ v_3 : (r, \theta, \phi) \mapsto r\psi(r) \cos \theta. \end{cases}$$

To derive Proposition 5, let ξ_1 be the lowest positive eigenvalue of **curl**, and \mathbf{u}_1 be an associated eigenfunction. Then, by Lemma 8, we know that

$$\xi_1 \geq \sqrt{\mu_1(\mathbb{B}_3)},$$

where $\mu_1(\mathbb{B}_3)$ is the first eigenvalue of the Dirichlet Laplacian. In the case of the ball, in fact, we have

$$\xi_1 > \sqrt{\mu_1(\mathbb{B}_3)}. \quad (32)$$

Proof of (32). If we had $\xi_1 = \sqrt{\mu_1(\mathbb{B}_3)}$, then $\phi := ru_{\xi_1, r}$ is an eigenfunction of the Dirichlet-Laplacian with eigenvalue $\mu_1(\mathbb{B}_3)$. However, $\phi(0) = 0$, while, as $\mu_1(\mathbb{B}_3)$ is simple, any eigenfunction associated with μ_1 has a (strict) constant sign in \mathbb{B}_3 , a contradiction. \square

Proposition 5 is thus implied by the following lemma:

Lemma 10. For any $i \in \{1, 2, 3\}$, there exists a unique $\mathbf{u}_{i, \pm}$ such that:

1. $(\mathbf{u}_{i, \pm}, \pm\omega)$ is a solution of (31),
2. $\langle x, \mathbf{u}_{i, \pm} \rangle = v_i$.

The $\mathbf{u}_{i, \pm}$ have the following explicit expressions (in spherical coordinates):

$$\begin{cases} \mathbf{u}_{1, \pm} = \sin \theta \cos(\phi) r\psi(r) \vec{e}_r + \left(\frac{F}{r} \cos(\theta) \cos(\phi) \pm \frac{G}{r} \sin(\phi)\right) \vec{e}_\theta + \left(\pm \frac{G}{r} \cos(\theta) \cos(\phi) - \frac{F}{r} \sin(\phi)\right) \vec{e}_\phi, \\ \mathbf{u}_{2, \pm} = \sin \theta \sin(\phi) r\psi(r) \vec{e}_r + \left(\frac{F}{r} \cos(\theta) \sin(\phi) \mp \frac{G}{r} \cos(\phi)\right) \vec{e}_\theta + \left(\pm \frac{G}{r} \cos(\theta) \sin(\phi) + \frac{F}{r} \cos(\phi)\right) \vec{e}_\phi, \\ \mathbf{u}_{3, \pm} = \cos(\theta) r\psi(r) \vec{e}_r - \frac{F}{r} \sin \theta \vec{e}_\theta \mp \frac{G}{r} \sin \theta \vec{e}_\phi, \end{cases} \quad (33)$$

where $F = F(r)$, $G = G(r)$ satisfy

$$\begin{cases} F'' + \omega F = \frac{d}{dr} \left(\frac{1}{r^2} \psi(r) \right), \\ G'' + \omega G = -\omega \frac{\psi}{r^2}, \end{cases} \quad (34)$$

The expressions for F, G are explicit

$$\begin{cases} F(r) = \left(\frac{\cos(\omega r)}{\omega r} - \frac{\sin(\omega r)}{\omega^2 r^2} + \sin(\omega r) \right) \\ G(r) = - \left(\frac{\sin(\omega r)}{\omega r} - \cos(\omega r) \right), \end{cases}$$

and $F(1) \neq 0, G(1) = 0$.

Proof of Lemma 10. To study the existence of such functions, we can follow the approach of [44, Lemma 2]. Dropping the indices i in $\mathbf{u}_{i,\pm}$ and v_i , the function $\mathbf{u}_{\pm} = \pm \frac{v}{r} \vec{e}_r + w \vec{e}_\theta + z \vec{e}_\phi$ must satisfy

$$\begin{cases} \frac{1}{r \sin \theta} \left(\frac{\partial(z \sin \theta)}{\partial \theta} - \frac{\partial w}{\partial \phi} \right) = \sqrt{\mu} \frac{v}{r} \\ \frac{1}{r} \left(\frac{1}{\sin \theta} \frac{\partial v}{\partial \phi} - \frac{\partial(rz)}{\partial r} \right) = \pm \sqrt{\mu} w \\ \frac{1}{r} \left(\frac{\partial(rw)}{\partial r} - \frac{\partial v}{\partial \theta} \right) = \pm \sqrt{\mu} z. \end{cases} \quad (35)$$

We thus obtain

$$\frac{\partial^2(rw)}{\partial r^2} + \mu(rw) = \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial v}{\partial \theta} \right) \pm \frac{\sqrt{\mu}}{r \sin \theta} \frac{\partial v}{\partial \phi}.$$

We derive a similar equation for z , namely:

$$\frac{\partial^2(rz)}{\partial r^2} + \mu(rz) = \mp \frac{\sqrt{\mu}}{r} \frac{\partial v}{\partial \theta} + \frac{1}{\sin \theta} \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial v}{\partial \phi} \right).$$

Writing these two equations as a system in $\Phi_{w,\pm} := rw, \Phi_{z,\pm} := rz$, which satisfy $\Phi_{w/z,\pm}(0, \theta, \phi) = \partial_r \Phi_{w/z,\pm}(0, \theta, \phi) = 0$, we are thus tasked with solving

$$\frac{\partial^2 \Phi_{w,\pm}}{\partial r^2} + \omega \Phi_{w,\pm} = \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial v}{\partial \theta} \right) \pm \frac{\sqrt{\mu}}{r \sin \theta} \frac{\partial v}{\partial \phi}$$

and

$$\frac{\partial^2 \Phi_{z,\pm}}{\partial r^2} + \omega \Phi_{z,\pm} = \mp \frac{\sqrt{\mu}}{r} \frac{\partial v}{\partial \theta} + \frac{1}{\sin \theta} \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial v}{\partial \phi} \right).$$

However, using the solutions (F, G) given in the statement of the theorem, we can immediately check that the expressions given are the correct ones; indeed, the uniqueness of the pair $(\mathbf{u}_{i,\pm})$ follows from the first point of Lemma 8. \square

An important corollary of the explicit expression of the first eigenfunctions of the curl is the following:

Corollary 2. *Let $(\alpha_{i,\pm})_{1=1,\dots,3} \in \mathbb{R}^6$ and consider the field*

$$\mathbf{u} = \sum_{i,\pm} \alpha_{i,\pm} \mathbf{u}_{i,\pm}$$

where the $u_{i,\pm}$ are given by Lemma 10. Then

$$\mathbf{u} = 0 \text{ on } \partial \mathbb{B}_3$$

if, and only if,

$$\forall i \in \{1, 2, 3\}, \quad \alpha_{i,+} = -\alpha_{i,-}.$$

Proof of Proposition 4. We are now in a position to prove Proposition 4. Consider \mathbf{U} a first eigenfunction of the Dirichlet-Stokes operator, associated with the eigenvalue $\lambda_1(\mathbb{B}_3)$. We already observed that $\lambda_1(\mathbb{B}_3) > \mu_1(\mathbb{B}_3)$. As \mathbf{U} satisfies homogeneous Dirichlet boundary conditions, it follows in particular that

$$\langle \mathbf{U}, \nu \rangle = 0 \quad \text{on } \partial\mathbb{B}_3$$

and thus $\mathbf{U} \in X$, where X is defined in Theorem 5. We can thus decompose \mathbf{U} as

$$\mathbf{U} = \sum_{k \in \mathbb{N}} \alpha_{k,\pm} \mathbf{u}_{k,\pm}$$

in X . Consequently, we deduce that

$$\lambda_1(\mathbb{B}_3) \in \{\xi_{k,\pm}^2, \xi_{k,\pm} \text{ eigenvalue of the curl operator}\}.$$

In particular, from Lemma 10, we have

$$\lambda_1(\mathbb{B}_3) \geq \omega^2.$$

However, observe that according to Lemma 10, $\mathbf{u}_{1,+} - \mathbf{u}_{1,-}$ is an eigenfunction of the Stokes operator with eigenvalue ω^2 . Thus, we deduce that

$$\lambda_1(\mathbb{B}_3) = \omega^2$$

and, furthermore, that for any associated eigenfunction \mathbf{U} , we can decompose \mathbf{U} as

$$\mathbf{U} = \sum_{i=1}^3 \sum_{\pm} \alpha_{i,\pm} \mathbf{u}_{i,\pm}.$$

Since \mathbf{U} satisfies homogeneous Dirichlet boundary conditions, Corollary 2 implies that \mathbf{U} writes

$$\mathbf{U} = \sum_{i=1}^3 \alpha_i (\mathbf{u}_{i,+} - \mathbf{u}_{i,-}).$$

Thus, the eigenspace has dimension at most 3.

We now compute, for each $i \in \{1, 2, 3\}$, the function $\mathbf{V}_i := \mathbf{u}_{i,+} - \mathbf{u}_{i,-}$. In spherical coordinates, we have

$$\begin{cases} \mathbf{V}_1 = 2\frac{G}{r} \sin(\phi) \vec{e}_\theta + 2\frac{G}{r} \cos(\theta) \cos(\phi) \vec{e}_\phi, \\ \mathbf{V}_2 = -2\frac{G}{r} \cos(\phi) \vec{e}_\theta + 2\frac{G}{r} \cos(\theta) \sin(\phi) \vec{e}_\phi, \\ \mathbf{V}_3 = -2\frac{G}{r} \sin \theta \vec{e}_\phi. \end{cases} \quad (36)$$

Now, to get a nicer expression in cartesian coordinates $(\vec{e}_x, \vec{e}_y, \vec{e}_z)$, recall that

$$\vec{e}_r = \frac{1}{r} (x\vec{e}_x + y\vec{e}_y + z\vec{e}_z), \quad \vec{e}_\theta = \frac{xz\vec{e}_x + yz\vec{e}_y - (x^2 + y^2)\vec{e}_z}{r\sqrt{x^2 + y^2}}, \quad \vec{e}_\phi = \frac{-y\vec{e}_x + x\vec{e}_y}{\sqrt{x^2 + y^2}}.$$

Furthermore, observe that

$$\frac{G(r)}{r^2} = \psi(r).$$

We thus obtain, in cartesian coordinates:

$$\mathbf{V}_1 = 2\psi(r) \begin{pmatrix} 0 \\ z \\ -y \end{pmatrix}, \mathbf{V}_2 = 2\psi(r) \begin{pmatrix} -z \\ 0 \\ x \end{pmatrix}, \mathbf{V}_3 = 2\psi(r) \begin{pmatrix} -y \\ x \\ 0 \end{pmatrix}.$$

In particular, defining $\mathbf{U}_i := \frac{\mathbf{V}_i}{2}$, we observe that $\mathbf{U}_1, \mathbf{U}_2, \mathbf{U}_3$ are orthogonal, whence the conclusion: the eigenspace is three dimensional, and we have an orthogonal basis of it.

It remains to compute the normalization constant A , using a spherical change of coordinates and explicit computations. Let us provide some details hereafter. One has

$$\begin{aligned} \|\mathbf{U}_i\|_{L^2(\mathbb{B}_3)}^2 &= \int_{\mathbb{B}_3} (x^2 + y^2)\psi(r)^2 dx dy dz = \int_0^{2\pi} \int_0^\pi \int_0^1 r^4 \sin^3 \theta \psi(r)^2 dr d\theta d\varphi \\ &= 2\pi \int_0^\pi \sin^3 \theta d\theta \int_0^1 \left(\frac{\sin(\omega r)}{\omega^2 r} - \frac{\cos(\omega r)}{\omega} \right)^2 dr \\ &= \frac{8\pi}{3\omega^3} \left(\frac{2 \cos^2 \omega + \omega^2 + \omega \cos \omega \sin \omega - 2}{2\omega^2} \right). \end{aligned}$$

The desired expression follows by using that $\tan \omega = \omega$.

References

- [1] S. Ashbaugh and D. Bucur. On the isoperimetric inequality for the buckling of a clamped plate. *Zeitschrift für Angewandte Mathematik und Physik (ZAMP)*, 54(5):756–770, Sept. 2003.
- [2] F. Boyer and P. Fabrie. *Mathematical tools for the study of the incompressible Navier-Stokes equations and related models*, volume 183 of *Appl. Math. Sci.* New York, NY: Springer, 2013.
- [3] D. Bucur. Uniform concentration-compactness for Sobolev spaces on variable domains. *J. Differ. Equations*, 162(2):427–450, 2000.
- [4] D. Bucur. Minimization of the k -th eigenvalue of the Dirichlet Laplacian. *Arch. Ration. Mech. Anal.*, 206:1073–1083, 2012.
- [5] D. Bucur and P. Freitas. A free boundary approach to the Faber-Krahn inequality. In *Geometric and computational spectral theory. Lectures from the Séminaire de Mathématiques Supérieures, Centre de Recherches Mathématiques, Université de Montréal, Montréal, QC, Canada, June 15–26, 2015*, pages 73–86. Providence, RI: American Mathematical Society (AMS); Montreal: Centre de Recherches Mathématiques (CRM), 2017.
- [6] D. Bucur and N. Varchon. Global minimizing domains for the first eigenvalue of an elliptic operator with non-constant coefficients. *Electron. J. Diff. Eqns.*, 36:1–10, 2000.
- [7] D. Buoso. Analyticity and criticality results for the eigenvalues of the biharmonic operator. In *Geometric properties for parabolic and elliptic PDE’s*, volume 176 of *Springer Proc. Math. Stat.*, pages 65–85. Springer, [Cham], 2016.
- [8] D. Buoso and P. D. Lamberti. Eigenvalues of polyharmonic operators on variable domains. *ESAIM Control Optim. Calc. Var.*, 19(4):1225–1235, 2013.

- [9] D. Buoso and P. D. Lamberti. Shape deformation for vibrating hinged plates. *Math. Methods Appl. Sci.*, 37(2):237–244, 2014.
- [10] G. Buttazzo and G. Dal Maso. An existence result for a class of shape optimization problems. *Archive for Rational Mechanics and Analysis*, 122(2):183–195, June 1993.
- [11] J. Cantarella, D. DeTurck, H. Gluck, and M. Teytel. Isoperimetric problems for the helicity of vector fields and the Biot–Savart and curl operators. *Journal of Mathematical Physics*, 41(8):5615–5641, Aug. 2000.
- [12] J. Cantarella, D. DeTurck, H. Gluck, and M. Teytel. The spectrum of the curl operator on spherically symmetric domains. *Physics of Plasmas*, 7(7):2766–2775, July 2000.
- [13] F. Caubet, M. Dambrine, and R. Mahadevan. Shape derivative for some eigenvalue functionals in elasticity theory. *SIAM Journal on Control and Optimization*, 59(2):1218–1245, Jan. 2021.
- [14] S. Chandrasekhar and P. C. Kendall. On force-free magnetic fields. *The Astrophysical Journal*, 126:457, Sept. 1957.
- [15] T. Chatelain and M. Choulli. Clarke generalized gradient for eigenvalues. *Commun. Appl. Anal.*, 1(4):443–454, 1997.
- [16] W. G. Chinn and N. E. Steenrod. First concepts of topology. The geometry of mappings of segments, curves, circles and disks. New York: Random House; The L. W. Singer Company. VIII, 160 p. , 1966.
- [17] Y. Chitour, D. Kateb, and R. Long. Generic properties of the spectrum of the Stokes system with Dirichlet boundary condition in \mathbb{R}^3 . *Ann. Inst. Henri Poincaré, Anal. Non Linéaire*, 33(1):119–167, 2016.
- [18] S. J. Cox. The generalized gradient at a multiple eigenvalue. *J. Funct. Anal.*, 133(1):30–40, 1995.
- [19] M. Dambrine and D. Kateb. On the shape sensitivity of the first Dirichlet eigenvalue for two-phase problems. *Applied Mathematics & Optimization*, 63(1):45–74, jul 2010.
- [20] M. C. Delfour and J.-P. Zolésio. *Shapes and geometries. Analysis, differential calculus, and optimization*, volume 4 of *Adv. Des. Control*. Philadelphia, PA: SIAM, 2001.
- [21] A. Enciso, W. Gerner, and D. Peralta-Salas. Optimal convex domains for the first curl eigenvalue, 2022. preprint: <https://arxiv.org/abs/2202.09204>.
- [22] A. Enciso and D. Peralta-Salas. Non-existence of axisymmetric optimal domains with smooth boundary for the first curl eigenvalue. *Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5)*, 24(1):311–327, 2023.
- [23] G. Faber. Beweis, daß unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt. 1923.
- [24] A. Falocchi and F. Gazzola. Remarks on the 3d Stokes eigenvalue problem under Navier boundary conditions. *Annali di Matematica Pura ed Applicata (1923 -)*, 201(3):1481–1488, Oct. 2021.
- [25] W. Gerner. Isoperimetric problem for the first curl eigenvalue. *Journal of Mathematical Analysis and Applications*, 519(2):126808, Mar. 2023.

- [26] P. Grinfeld. Hadamard’s formula inside and out. *J. Optim. Theory Appl.*, 146(3):654–690, 2010.
- [27] A. Henrot. *Extremum Problems for Eigenvalues of Elliptic Operators*. Birkhäuser Basel, 2006.
- [28] A. Henrot, editor. *Shape optimization and spectral theory*. Berlin: De Gruyter, 2017.
- [29] A. Henrot. *Isoperimetric Inequalities for Eigenvalues of the Laplacian*, pages 47–88. Springer International Publishing, 2018. *Geometry of PDEs and Related Problems*.
- [30] A. Henrot and M. Pierre. *Shape variation and optimization. A geometrical analysis*, volume 28 of *EMS Tracts Math*. Zürich: European Mathematical Society (EMS), 2018.
- [31] J. P. Kelliher. On the vanishing viscosity limit in a disk. *Math. Ann.*, 343(3):701–726, 2009.
- [32] S. Kesavan. *Symmetrization and applications*, volume 3 of *Ser. Anal*. Hackensack, NJ: World Scientific, 2006.
- [33] E. Krahn. Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises. *Mathematische Annalen*, 94(1):97–100, Dec. 1925.
- [34] M. Kunzinger and R. Steinbauer. Riemannian geometry- lecture notes. <https://www.mat.univie.ac.at/stein/teaching/skripten/rg.pdf>.
- [35] P. D. Lamberti and M. Zaccaron. Shape sensitivity analysis for electromagnetic cavities. *Math. Methods Appl. Sci.*, 44(13):10477–10500, 2021.
- [36] D.-S. Lee and B. Rummeler. The eigenfunctions of the Stokes operator in special domains. III. *ZAMM, Z. Angew. Math. Mech.*, 82(6):399–407, 2002.
- [37] P.-L. Lions. The concentration-compactness principle in the calculus of variations. The locally compact case, part 1. *Annales de l’I.H.P. Analyse non linéaire*, 1(2):109–145, 1984.
- [38] D. Mazzoleni and A. Pratelli. Existence of minimizers for spectral problems. *J. Math. Pures Appl.*, 2013.
- [39] F. Mugnot, F. Murat, and J. P. Puel. Variation d’un point de retournement par rapport au domaine. *Commun. Partial Differ. Equations*, 4:1263–1297, 1979.
- [40] C. Nitsch and C. Trombetti. The classical overdetermined Serrin problem. *Complex Var. Elliptic Equ.*, 63(7-8):1107–1122, 2018.
- [41] J. H. Ortega and E. Zuazua. Generic simplicity of the eigenvalues of the Stokes system in two space dimensions. *Adv. Differ. Equ.*, 6(8):987–1023, 2001.
- [42] R. Rodríguez and P. Venegas. Numerical approximation of the spectrum of the curl operator. *Math. Comput.*, 83(286):553–577, 2014.
- [43] B. Rousselet and D. Chenaïs. Continuité et différentiabilité d’éléments propres: Application à l’optimisation de structures. *Applied Mathematics & Optimization*, 22(1):27–59, July 1990.
- [44] R. S. Saks. Solving of spectral problems for curl and Stokes operators. *Ufa Mathematical Journal*, 5(2):63–81, 2013.

- [45] J. Serrin. A symmetry problem in potential theory. *Arch. Ration. Mech. Anal.*, 43:304–318, 1971.
- [46] N. Steenrod. *The topology of fibre bundles*. Princeton, NJ: Princeton University Press, 1999.
- [47] K. Stollenwerk. Optimal shape of a domain which minimizes the first buckling eigenvalue. *Calc. Var. Partial Differ. Equ.*, 55(1):29, 2016.
- [48] K. Stollenwerk. On the optimal domain for minimizing the buckling load of a clamped plate. *Z. Angew. Math. Phys.*, 74(1):16, 2023.
- [49] B. Velichkov. *Existence and Regularity Results for Some Shape Optimization Problems*. Ed: Scuola Normale Superiore, 2015.