

# Lovász–Schrijver PSD-operator and the stable set polytope of claw-free graphs

Silvia M Bianchi, Mariana Escalante, Graciela Nasini, Annegret K Wagler

# ► To cite this version:

Silvia M Bianchi, Mariana Escalante, Graciela Nasini, Annegret K Wagler. Lovász–Schrijver PSD-operator and the stable set polytope of claw-free graphs. Discrete Applied Mathematics, 2023, 332, pp.70-86. 10.1016/j.dam.2023.01.012 . hal-04404041

# HAL Id: hal-04404041 https://hal.science/hal-04404041

Submitted on 18 Jan2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Lovász-Schrijver PSD-operator and the Stable Set Polytope of Claw-Free Graphs<sup>\*</sup>

Silvia M. Bianchi<sup>1</sup>, Mariana Escalante<sup>1</sup>, Graciela Nasini<sup>1</sup>, and Annegret K. Wagler<sup>2</sup>

<sup>1</sup>CONICET and FCEIA, Universidad Nacional de Rosario, Rosario, Argentina, {sbianchi,mariana,nasini}@fceia.unr.edu.ar <sup>2</sup>LIMOS (UMR 6158 CNRS), Université Clermont Auvergne, Clermont-Ferrand, France, annegret.wagler@uca.fr

August 6, 2019

#### Abstract

The subject of this work is the study of  $LS_+$ -perfect graphs defined as those graphs G for which the stable set polytope STAB(G) is achieved in one iteration of Lovász-Schrijver PSD-operator  $LS_+$ , applied to its edge relaxation ESTAB(G). In particular, we look for a polyhedral relaxation of STAB(G) that coincides with  $LS_+(ESTAB(G))$  and STAB(G) if and only if G is  $LS_+$ -perfect. An according conjecture has been recently formulated ( $LS_+$ -Perfect Graph Conjecture); here we verify it for the wellstudied class of claw-free graphs.

## 1 Introduction

The notion of a perfect graph was introduced by Berge in the early 1960s [1]. A graph is called *perfect* if each of its induced subgraphs has chromatic number equal to the size of a maximum cardinality clique in the subgraph.

Perfect graphs turned out to be an interesting and important class with a rich structure and a nice algorithmic behavior [24] and inspired numerous very interesting contributions to the literature for the past fifty years.

An early polyhedral characterization of perfect graphs was given by Chvátal in [10]: G is perfect if and only if STAB(G) coincides with its corresponding clique relaxation QSTAB(G). However, solving the Maximum Weight Stable Set Problem (MWSSP) for a perfect graph G by maximizing a linear objective

<sup>\*</sup>This work was supported by a MATH-AmSud cooperation (PACK-COVER), PID-CONICET 0277, and PICT-ANPCyT 0586.

function over QSTAB(G) does not work directly, but only via a detour involving a geometric representation of graphs [28] and the resulting semi-definite relaxation TH(G), the *theta body* of G. This is one of the main results in the seminal paper of Grötschel, Lovász and Schrijver [23]. Later on, the same authors proved a related result which connects a purely graph theoretic notion to polyhedrality of a typically nonlinear convex relaxation and to the integrality and equality of two fundamental polytopes:

**Theorem 1.** (Grötschel, Lovász and Schrijver [23, 24]) For every graph G, the following are equivalent:

- 1. G is perfect;
- 2. STAB(G) = QSTAB(G);
- 3.  $\operatorname{TH}(G) = \operatorname{STAB}(G);$
- 4.  $\operatorname{TH}(G) = \operatorname{QSTAB}(G);$
- 5. TH(G) is polyhedral.

Some years later, Lovász and Schrijver [29] introduced a semidefinite relaxation of STAB(G) which is stronger than TH(G) and it is obtained after applying once the LS<sub>+</sub> lift-and-project operator ( $N_+$  in [29]) to the edge relation of STAB(G), see Section 2.3 for details.

We denote this relaxation by  $LS_+(G)$ . Following the same line of reasoning used for perfect graphs, they proved that the MWSSP can be solved in polynomial-time for the class of graphs G for which  $LS_+(G) = STAB(G)$ . Those graphs have been called  $LS_+$ -perfect graphs in [5] (originally  $N_+$ -perfect in [3]). The fact that solving the MWSSP on  $LS_+$ -perfect graphs can be done in polynomial-time encourages the search of a complete description of their stable set polytope in terms of linear inequalities.

In this context, one of our goals is to obtain a characterization of  $LS_+$ -perfect graphs similar to the one given in Theorem 1 for perfect graphs. More precisely, we look for an appropriate polyhedral relaxation of STAB(G) playing the role of QSTAB(G) in Theorem 1, when we replace TH(G) by  $LS_+(G)$ .

Also, in [29] it is proved that every valid inequality for STAB(G) having as support a near-bipartite graph is valid for  $LS_+(G)$  (a graph is *near-bipartite* if the non-neighbors of every node induce a bipartite graph [37]).

This result enables the authors in [3] to establish the following:

**Conjecture 2** (LS<sub>+</sub>-Perfect Graph Conjecture). For every graph G, if LS<sub>+</sub>(G) = STAB(G) then every facet defining inequality of STAB(G) has near-bipartite support.

According to the findings in [37], if G is a near-bipatite graph then every facet defining inequality of STAB(G) has a complete join of antiwebs as a support graph, see Section 2 for details. Hence, Conjecture 2 can be restated as:

**Conjecture 3.** If G is an  $LS_+$ -perfect graph, the support graph of any facet defining inequality of STAB(G) is a complete join of antiwebs.

Later, in [42] Wagler defined for a given graph G the relaxation ASTAB<sup>\*</sup>(G) described by inequalities whose support are complete joins of antiweb subgraphs in G. In the same context, graphs for which STAB(G) coincides with ASTAB<sup>\*</sup>(G) are called *joined a-perfect* [11, 40]. Hence Conjecture 2 establishes that ASTAB<sup>\*</sup>(G) corresponds to the relaxation we are looking for to play the role of QSTAB(G) in Theorem 1, when we replace TH(G) by LS<sub>+</sub>(G).

Some progress has been made towards proving its validity since it was proposed in [3]. Conjecture 2 has been already verified for near-perfect graphs [3], for *fs-perfect graphs* [4, 5], for *webs* [16], for *line graphs* [17], and for several graph classes defined by clique cutsets [43, 44], see Section 2.3 for details and more precise definitions.

The aim of this contribution is to verify Conjecture 3 for a well studied graph class containing all webs, all line graphs and the complements of near-bipartite graphs: the class of claw-free graphs. Claw-free graphs attracted much attention due to their seemingly asymmetric behavior w.r.t. the stable set problem. On the one hand, the first combinatorial algorithms to solve the problem in polynomial time for claw-free graphs [30, 35] date back to 1980. Therefore, the polynomial equivalence of optimization and separation due to [24] implies that it is possible to optimize over the stable set polytope of a claw-free graph in polynomial time. On the other hand, the problem of characterizing the stable set polytope of claw-free graphs in terms of an explicit description by means of a facet-defining system, originally posed in [24], was open for several decades. This motivated the study of claw-free graphs and its different subclasses, that finally answered this long-standing problem only recently (see Section 2.4 for details).

To prove the validity of the conjecture on claw-free graphs we use two different approaches. On one hand, we take advantage of the known description of the stable set polytope for quasi-line graphs. On the other hand, we characterize the structure of the connected  $LS_+$ -perfect claw-free not quasi-line graphs without clique cutsets.

The paper is organized as follows: In Section 2.1 we present notation and basic results that will be used throughout the paper. In Section 2.2 we present the different relaxations we use of the well-known stable set polytope, and also families of graphs which are defined in terms of the facet defining inequalities in their stable set polytope. In Section 2.3, we present the State-of-the-Art on  $LS_+$ -perfect graphs (including families of  $LS_+$ -imperfect graphs needed for the subsequent proofs) and the results concerning the facet-description of their stable set polytopes from the literature. We devote a subsection to claw-free graphs (Section 2.4) and their relevant subclasses, presenting known valid inequalities describing the stable set polytope of certain claw-free graphs. In Section 3, we verify Conjecture 3 for the class of claw-free graphs that are claw-free. Finally, in Section 5 we present the conclusions and future lines of research on this topic, establishing connections with the previous results.

Parts of the here presented results appeared without proofs in [6, 7].

## 2 Definitions and preliminary results

#### 2.1 Graph theory

Throughout this work, G = (V, E) stands for a simple graph with node set V and edge set E. When we need to emphasize the relationship between the node set and the edge set with the graph, we may write V(G) and E(G). The complementary graph of G is denoted by  $\overline{G}$ .

Given  $U \subset V$ , G[U] is the subgraph of G induced by the nodes in U. i.e. having node set U and edge set  $\{uv : uv \in E, \{u, v\} \subset U\}$ . Given a graph G', if there is a subset of nodes of G inducing G' we say that G' is a node induced subgraph of G and write  $G' \subset G$ . Also, we denote by G - U the subgraph of G induced by the nodes in  $V \setminus U$ . For simplicity, we write G - u instead of  $G - \{u\}$ .

Given  $v \in V$ ,  $N_G(v)$  is the *neighbourhood* of v and  $N_G[v] = N_G(v) \cup \{v\}$  is the *closed neighbourhood* of v. When the graph is clear from the context, we simply write N(v) or N[v].

A stable set in G is a subset of mutually nonadjacent nodes in G and a *clique* is a subset of pairwise adjacent nodes in G. The cardinality of a stable set of maximum cardinality in G is denoted by  $\alpha(G)$ .

A *clique cutset* of a connected graph G is a clique whose removal disconnects G.

For  $n \in \mathbb{N}$ , [n] will denote the additive group defined on the set  $\{1, \ldots, n\}$ , with integer addition modulo n. Given  $a, b \in [n]$ , let b - a be the minimum non-negative integer t such that  $a + t = b \mod n$ . We denote by  $[a, b]_n$  the *circular interval* defined by the set  $\{a + s : 0 \leq s \leq b - a\}$ . Similarly,  $(a, b]_n$ ,  $[a, b)_n$ , and  $(a, b)_n$  correspond to  $[a, b]_n \setminus \{a\}$ ,  $[a, b]_n \setminus \{b\}$ , and  $[a, b]_n \setminus \{a, b\}$ , respectively.

For any positive integer number n,  $C_n$  denotes the cycle with set of nodes [n] and such that for all  $i \in [n]$ , i is adjacent to i + 1. Similarly,  $P_n$  denotes the path obtained by deleting the edge n1 in  $C_n$ .

A hole in the graph G is an induced cycle in the graph. Similarly, an *antihole* in G is an induced cycle in  $\overline{G}$ . The hole (antihole) is *odd* if it has an odd number of nodes.

Chudnosvsky et al. proved the well-known Strong Perfect Graph Theorem [8] (previously stated as a conjecture by Berge [1]) showing that perfect graphs are exactly those graphs having neither odd holes nor odd antiholes.

A wheel is a graph formed by a cycle and one extra node outside the cycle, connected to every node in it.

Given integer numbers k and n such that  $n \ge 2(k+1)$ ,  $W_n^k$  denotes the web graph with set of nodes [n] and edges i (i+t) for all  $i \in [n]$  and  $t \in \{1, \ldots, k\}$ . Observe that, for all  $i \in [n]$ ,  $[i, i+k]_n$  is a maximal clique of  $W_n^k$ . We say that a set of nodes  $U \subset [n]$  is a set of  $r \ge 1$  consecutive nodes in  $W_n^k$  if there exists  $i \in [n]$  such that  $U = [i, i+r)_n$ . The antiweb  $A_n^{k+1}$  is the complementary graph of a web  $W_n^k$ . It is easy to see that  $C_n = W_n^1 = A_n^k$  and  $\overline{C}_n = W_n^{k-1} = A_n^2$  with  $k = \lfloor \frac{n}{2} \rfloor$ . Also, complete graphs of n nodes are the antiwebs  $A_n^l$ . A graph is *near-bipartite* [37] if the non-neighbors of every node induce a bipartite graph. Observe that antiwebs are near-bipartite graphs.

The line graph of a graph G, denoted by L(G), is obtained by turning adjacent edges of G into adjacent nodes of the line graph. Then we say that a graph H is a *line graph* if there is a graph G such that H = L(G).

A graph is *claw-free* if it does not contain a claw as a node induced subgraph, where a *claw* is a four node graph with one node connected to a stable set of size three.

Let us now recall some operations on graphs that will be used in this paper.

Given two node disjoint graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  the complete join of them is the graph  $G_1 \vee G_2$  having node set  $V_1 \cup V_2$  and edge set  $E_1 \cup E_2 \cup \{uv : u \in V_1, v \in V_2\}$ . For a given graph G = (V, E) and  $U_1, U_2 \subset V$ such that  $U_1 \cap U_2 = \emptyset$ , we say that  $U_1$  is completely joined to  $U_2$  (in G) if  $G[U_1 \cup U_2] = G[U_1] \vee G[U_2]$ .

The graph G' is obtained from G after an *odd subdivision of the edge* e of G if it is obtained from G by replacing e by a path of odd length ([45]).

The stretching of a node is defined in [27]. Let v be a node of G with neighborhood N(v) and let  $A_1$  and  $A_2$  be nonempty subsets of N(v) such that  $A_1 \cup A_2 = N(v)$ , and  $A_1 \cap A_2 = \emptyset$ . A stretching of a node v is obtained as follows: remove v, introduce three nodes instead, called  $v_1, v_2$  and u, and add an edge between  $v_i$  and every node in  $\{u\} \cup A_i$  for  $i \in \{1, 2\}$ . Also, we say that H is a stretching of G if it is obtained from G after a finite number of node-stretching operations. It is not hard to see that an odd-subdivision of an edge is a sequence of node stretching operations.

#### 2.2 On the Stable Set Polytope

Given G = (V, E), the stable set polytope STAB(G) of G is defined as the convex hull of the incidence vectors of all stable sets of G.

The support graph of a valid inequality of STAB(G) is the subgraph of G induced by the nodes with nonzero coefficient in the inequality. A full-support inequality has G as its support graph. A graph G is said to be facet-defining if STAB(G) has a full-support facet-defining inequality.

The following is an immediate consequence of Theorem 4.1 in [10].

**Corollary 4.** If G is a facet-defining graph then G does not contain a clique cutset.

If  $G' \subset G$ , the inequality

$$\sum_{i \in V(G')} x_i \le \alpha(G')$$

is known as the *rank constraint* associated with G' and it is clearly valid for STAB(G).

A more general type of valid inequalities for STAB(G) is obtained from subgraphs of G which are complete join of graphs, called *joined constraints*: if  $G' \subset G$  and G' is the complete join of  $G_j$  with  $j = 1, \ldots, k$ , the constraint

$$\sum_{j=1}^{k} \frac{1}{\alpha(G_j)} \sum_{i \in V(G_j)} x_i \le 1,$$

is also valid for STAB(G).

Note that the inequalities are scaled to have right hand side 1 and they include rank constraints when k = 1. From complete joins of antiwebs, the *joined antiweb constraints* are obtained.

Two canonical relaxations of STAB(G) are defined in terms of rank constraints: the *edge relaxation* 

$$ESTAB(G) = \{ x \in [0, 1]^V : x_i + x_j \le 1, ij \in E \},\$$

and the *clique relaxation* 

$$QSTAB(G) = \{ x \in [0,1]^V : \sum_{i \in Q} x_i \le 1, \ Q \text{ maximal clique of } G \}.$$

We have  $STAB(G) \subset QSTAB(G) \subset ESTAB(G)$  for any graph. In addition, STAB(G) equals ESTAB(G) for bipartite graphs only, and QSTAB(G) for perfect graphs only [10].

A graph G is called *near-perfect graphs* (defined in [36]) if STAB(G) is obtained after adding to QSTAB(G) the rank inequality associated with the whole graph. Moreover, G is *rank-perfect* ([40]) if and only if STAB(G) is described by rank constraints only. By definition, rank-perfect graphs include all perfect and near-perfect graphs.

As subclasses of rank-perfect graphs, the class of *a-perfect graphs* is introduced in [41] as graphs G where STAB(G) is given by nonnegativity constraints and rank constraints associated with antiwebs. In particular, antiwebs are *a*perfect by [37, 41].

We denote by  $ASTAB^*(G)$  the linear relaxation of STAB(G) obtained by nonnegativity constraints and all joined antiweb constraints. By construction, we see that

$$STAB(G) \subset ASTAB^*(G) \subset QSTAB(G) \subset ESTAB(G).$$

Recall that graphs G with  $STAB(G) = ASTAB^*(G)$  are called joined a-perfect. Results from Shepherd [37] imply that all near-bipartite graphs are joined aperfect, and results from Wagler [43, 44] show that several graph classes defined by clique cutsets are further subclasses of joined a-perfect graphs.

#### **2.3** On the semidefinite relaxation $LS_+(G)$ of STAB(G)

In order to introduce the LS<sub>+</sub>-operator we denote by  $e_0, e_1, \ldots, e_n$  the vectors of the canonical basis of  $\mathbb{R}^{n+1}$  (where the first coordinate is indexed zero) and

 $S^n_+$  the convex cone of symmetric and positive semi-definite  $(n \times n)$ -matrices with real entries. Let  $K \subset [0,1]^n$  be a convex set and

$$\operatorname{cone}(K) = \left\{ \left( \begin{array}{c} x_0 \\ x \end{array} \right) \in \mathbb{R}^{n+1} : x = x_0 y; \ y \in K \right\}.$$

Then, the convex set  $M_+(K)$  is defined as:

$$M_{+}(K) = \{ Y \in S_{+}^{n+1} : \qquad Ye_{0} = \operatorname{diag}(Y),$$
$$Ye_{i} \in \operatorname{cone}(K),$$
$$Y(e_{0} - e_{i}) \in \operatorname{cone}(K), \ i = 1, \dots, n \}$$

where diag(Y) denotes the vector whose *i*-th entry is  $Y_{ii}$ , for every i = 0, ..., n. Projecting this lifting back to the space  $\mathbb{R}^n$  results in

$$\mathrm{LS}_+(K) = \left\{ x \in [0,1]^n : \begin{pmatrix} 1 \\ x \end{pmatrix} = Ye_0, \text{ for some } Y \in M_+(K) \right\}$$

In [29], Lovász and Schrijver proved that  $LS_+(K)$  is a relaxation of the convex hull of integer solutions in K and that

$$\mathrm{LS}^n_+(K) = \mathrm{conv}(K \cap \{0,1\}^n),$$

where  $LS^{0}_{+}(K) = K$  and  $LS^{k}_{+}(K) = LS_{+}(LS^{k-1}_{+}(K))$  for every  $k \ge 1$ .

In this work we focus on the behavior of a single application of the LS<sub>+</sub>-operator to the edge relaxation ESTAB(G) of the stable set polytope of a graph. Recall that  $\text{LS}_+(G) = \text{LS}_+(\text{ESTAB}(G))$ .

If  $LS_+(G) = STAB(G)$  then G is  $LS_+$ -perfect, and all other graphs are  $LS_+$ -imperfect.

According to Lemma 1.5 in [29], given a graph G, every valid inequality whose support graph is near-bipartite is also valid for  $LS_+(G)$ . Then, this result and the definition of  $ASTAB^*(G)$  imply that

$$LS_{+}(G) \subseteq ASTAB^{*}(G) \tag{1}$$

and Conjecture 3 establishes that the equality in equation (1) characterizes  $LS_+$ -perfect graphs or, in other words, that  $LS_+$ -perfect graphs coincide with joined a-perfect graphs.

It is known from [5] that every subgraph of an LS<sub>+</sub>-perfect graph is also LS<sub>+</sub>-perfect. Thus, exhibiting one LS<sub>+</sub>-imperfect subgraph G' in a graph G certifies the LS<sub>+</sub>-imperfection of G. This motivates the definition of *minimally* LS<sub>+</sub>-*imperfect graphs* as the LS<sub>+</sub>-imperfect graphs whose proper induced subgraphs are all LS<sub>+</sub>-perfect. The two smallest such graphs were found by [15] and [27] and are called  $G_{LT}$  and  $G_{EMN}$ , see Figure 1.

Observe that the graph  $G_{LT}$  is obtained after replicating a node in  $C_5$ . It is clear that odd cycles and their complements are LS<sub>+</sub>-perfect. Thus, node replication does not preserve LS<sub>+</sub>-perfection. Moreover, we easily obtain:



Figure 1: The graphs  $G_{LT}$  (on the left) and  $G_{EMN}$  (on the right).



Figure 2: Some node-stretchings  $(v_1, w, v_2 \text{ in black})$  of  $G_{LT}$  and  $G_{EMN}$ .

**Remark 5.** Given a graph G = (V, E) having  $C_5$  as a node induced subgraph, the graph obtained after replication of any node in  $C_5$  is  $LS_+$ -imperfect.

Further  $LS_+$ -imperfect graphs can be obtained by applying operations preserving  $LS_+$ -imperfection. In [27] it is shown:

**Theorem 6** ([27]). The stretching of a node preserves  $LS_+$ -imperfection.

Hence, all stretchings of  $G_{LT}$  and  $G_{EMN}$  are LS<sub>+</sub>-imperfect, see Figure 2 for some examples.

#### 2.4 On claw-free graphs

In several respects, claw-free graphs are generalizations of line graphs. An intermediate class between line graphs and claw-free graphs form *quasi-line graphs*, where the neighborhood of any node can be partitioned into two cliques (i.e., quasi-line graphs are the complements of near-bipartite graphs).

Quasi-line graphs can be divided into two subclasses: *fuzzy circular interval* graphs and semi-line graphs.

Let  $\mathcal{C}$  be a circle,  $\mathcal{I}$  a collection of intervals in  $\mathcal{C}$  without proper containments and common endpoints, and V a multiset of points in  $\mathcal{C}$ . A fuzzy circular interval graph  $G(V,\mathcal{I})$  has node set V and two nodes are adjacent if both belong to one interval  $I \in \mathcal{I}$ , where edges between different endpoints of the same interval may be omitted.

A graph is semi-line if it is either a line graph or a quasi-line graph without a representation as a fuzzy circular interval graph.

For fuzzy circular interval graphs it turned out ([14]) that so-called *clique* family inequalities suffice to describe their stable set polytope.

More specifically, Stauffer (in [38, 39]) verified a conjecture stated in [32] establishing that every facet-defining clique family inequality of a fuzzy circular

interval graph G is associated with a web in G. These inequalities have the form

$$(p-r)\sum_{i\in W} x_i + (p-r-1)\sum_{i\in W_o} x_i \le (p-r)\left\lfloor\frac{n}{p}\right\rfloor$$
(2)

where G[W] has a web subgraph  $W_n^{p-1}$  and  $r = n - \lfloor \frac{n}{p} \rfloor p \ge 1$ . Moreover,  $W \setminus V(W_n^{p-1})$  is the set of nodes in G having at least 2p-1 consecutive neighbors in  $W_n^{p-1}$ . Also, a node in  $W_o$  is adjacent to exactly 2p-2 consecutive nodes in  $W_n^{p-1}$ .

Chudnovsky and Seymour [9] extend results of Edmonds [12] and Edmonds and Pulleyblank [13] to semi-line graphs. Actually, they prove that if G is a semi-line graph then STAB(G) is given by clique constraints and rank constraints associated with the line graphs of 2-connected hypomatchable induced subgraphs (a graph is called *hypomatchable* if removing any of its nodes results in a graph having a perfect matching).

Then, semi-line graphs are rank-perfect with line graphs as only facet-defining subgraphs.

However, there are claw-free graphs which are not quasi-line. Although there are many important results on the facial structure of them [18, 19, 20, 21, 22, 26], in order to prove the conjecture on claw-free not quasi-line graphs we take into account their decomposition given in [9].

A node is simplicial if its neighborhood is a clique. A strip (G, a, b) is a (not necessarily connected) graph with two designated simplicial nodes a and b. Given two node-disjoint strips  $(G_1, a_1, b_1)$  and  $(G_2, a_2, b_2)$ , their composition is the union of  $G_1 \setminus \{a_1, b_1\}$  and  $G_2 \setminus \{a_2, b_2\}$  together with all edges between  $N_{G_1}(a_1)$  and  $N_{G_2}(a_2)$ , and between  $N_{G_1}(b_1)$  and  $N_{G_2}(b_2)$  A claw-free strip containing a 5-wheel as induced subgraph is a 5-wheel strip.

This composition operation can be generalized to more than two strips (see [9] for further details).

According to these definitions, in [31] it is proved that every claw-free not quasi-line graph G with  $\alpha(G) \geq 4$  admits a decomposition into strips, where at most one strip is quasi-line and all the remaining ones are 5-wheel strips having stability number at most 3.

There are only three "basic" types of 5-wheel strips (see Fig. 3) which can be extended by adding nodes belonging to the neighborhood of the 5-wheels (see [31] for details).

Note that a claw-free but not quasi-line graph G with  $\alpha(G) \ge 4$  containing a clique cutset may have a facet-defining subgraph G' with  $\alpha(G') = 3$  (inside a 5-wheel strip of type 3), see [34] for examples.

# 3 LS<sub>+</sub>-Perfect Graph Conjecture for claw free graphs

In this section we verify the validity of the  $LS_+$ -Perfect Graph Conjecture on claw-free graphs by analyzing the behavior of the  $LS_+$ -operator on all its rele-



Figure 3: The three types of basic 5-wheel strips.

vant subclasses. Additionally, we characterize minimally  $LS_+$ -imperfect graphs within these classes.

We start with those graphs having stability number two by recalling the following result:

**Theorem 7** ([5]). Let G be a graph with  $\alpha(G) = 2$  such that G - v is an odd antihole for some node v. G is LS<sub>+</sub>-perfect if and only if v is completely joined to  $V(G) \setminus \{v\}$ .

From this, we can prove:

**Theorem 8.** Let G be a connected imperfect graph with  $\alpha(G) = 2$ . If G is  $LS_+$ -perfect, then G is a complete join of odd antiholes and (possible empty) perfect graphs.

*Proof.* Since G is an imperfect graph and  $\alpha(G) = 2$ , G contains an odd antihole  $\overline{C}$ . If  $G = \overline{C}$ , we are done. Otherwise, if G is LS<sub>+</sub>-perfect then by Theorem 7, G is the complete join of  $\overline{C}$  and  $G' = G - V(\overline{C})$ .

If G' is a perfect graph then the result follows. Otherwise, G' is LS<sub>+</sub>-perfect and  $\alpha(G') = 2$  and we apply the same argument as for G.

Using Corollary 5.2 in [10] on the linear description of the stable set polytope of complete joins of graphs, we conclude:

**Corollary 9.** The  $LS_+$ -Perfect Graph Conjecture is true for graphs with stability number 2.

#### 3.1 Quasi-line graphs

Recall that quasi-line graphs divide into the two subclasses of semi-line graphs and fuzzy circular interval graphs.

As we have already mentioned, a way to attack the conjecture is from the polyhedral point of view. Using this approach, in Lemma 7 in [17], the authors characterized  $LS_+$ -perfect line graphs by showing:

**Theorem 10** ([17]). Let G be the line graph of a 2-connected hypomatchable graph. Then, G is  $LS_+$ -perfect if and only if either G is a complete graph or G is an odd cycle.

Additionally, Chudnovsky and Seymour [9] proved that the stable set polytope of a semi-line graph is given by rank constraints associated with cliques and the line graphs of 2-connected hypomatchable graphs. From these two results we directly conclude:

**Corollary 11.** The LS<sub>+</sub>-Perfect Graph Conjecture holds for semi-line graphs.

For fuzzy circular interval graphs, we make use of Theorem 5 in [16].

**Theorem 12** ([16]). If a web graph is  $LS_+$ -perfect then it is either a perfect or a minimally imperfect graph.

Using this result, we can prove:

**Theorem 13.** Let G be a facet-defining fuzzy circular interval graph. Then, G is  $LS_+$ -perfect if and only if it is an odd cycle or its complement.

*Proof.* If G is an odd cycle or its complement then clearly G is  $LS_+$ -perfect. Let us prove the converse.

Since G is facet-defining, from the results in [38, 39], we have that G is the support graph of a clique family inequality

$$(p-r)\sum_{i\in W} x_i + (p-r-1)\sum_{i\in W_o} x_i \le (p-r)\left\lfloor \frac{n}{p} \right\rfloor$$

associated with a web subgraph  $W_n^{p-1}$  of G[W]. Recall that every node in  $W \setminus V(W_n^{p-1})$  (resp. in  $W_o$ ) is adjacent to at least 2p - 1 (resp. to exactly 2p - 2) consecutive nodes of  $W_n^{p-1}$  and  $r \ge 1$  (see Section 2.4). By Theorem 12,  $W_n^{p-1}$  is LS<sub>+</sub>-perfect if and only if it is an odd cycle or its

By Theorem 12,  $W_n^{p-1}$  is LS<sub>+</sub>-perfect if and only if it is an odd cycle or its complement. Since G is LS<sub>+</sub>-perfect we restrict our attention to web subgraphs of G[W] where n = 2k + 1 and p = 2 or  $p = k \ge 3$ . In both cases, r = 1 follows and we have to prove that  $G = W_n^{p-1}$ .

Observe that if p = 2, then p - r - 1 = 0 and the inequality takes the form

$$\sum_{i \in W} x_i \le k.$$

Suppose that there exists  $v \in W \setminus V(W_{2k+1}^1)$ . Then, v is connected to  $t \geq 2p-1=3$  consecutive nodes in  $W_{2k+1}^1$ .

If  $t \ge 5$  and k = 2, G has a 5-wheel as induced subgraph, contradicting the fact that G is quasi-line. But, if  $t \ge 5$  and  $k \ge 3$ , G has a claw, again a contradiction. Then,  $t \le 4$ .

Now, if t = 3 (resp. t = 4) G contains an odd-subdivision of an edge of  $G_{LT}$  (resp.  $G_{EMN}$ ).

Since G is LS<sub>+</sub>-perfect then  $W \setminus V(W_{2k+1}^1) = \emptyset$ , or equivalently,  $G = W_{2k+1}^1 = C_{2k+1}$ .

Let us now analyze the case  $p = k \ge 3$  and  $G[W_o \cup W] = G$ .

Suppose that there exists  $v \in (W_o \cup W) \setminus V(W_{2k+1}^{k-1})$ . It holds that v is adjacent to at least 2k - 2 consecutive nodes in  $W_{2k+1}^{k-1}$ .



Figure 4: The gear and the 3-gear graphs.

It follows that the subgraph G' of G induced by  $V(W_{2k+1}^{k-1}) \cup \{v\}$  has stability number two. Since G' is LS<sub>+</sub>-perfect, from Theorem 8, v is completely joined to  $V(W_{2k+1}^{k-1})$ , contradicting that G is quasi-line.

Then, we conclude  $G[W_o \cup W] = W_{2k+1}^{k-1}$  or, equivalently, the complementary graph of  $C_{2k+1}$  and the proof is complete.

Clearly the theorem above implies that the  $LS_+$ -Perfect Graph Conjecture is true for fuzzy circular interval graphs. Since the class of quasi-line graphs divides into semi-line graphs and fuzzy circular interval graphs, we obtain, as direct consequence:

**Corollary 14.** The LS<sub>+</sub>-Perfect Graph Conjecture is true for quasi-line graphs.

#### 3.2 Claw-free graphs that are not quasi-line

It is left to treat the case of claw-free graphs that are not quasi-line. When  $\alpha(G) \geq 4$ , we study LS<sub>+</sub>-perfection by using the strip decomposition given in [31]. Actually, we prove that in this class there is no connected LS<sub>+</sub>-perfect graph G without clique cutset such that  $\alpha(G) \geq 4$ . Finally, when G has stability number 3, we obtain a decomposition of connected LS<sub>+</sub>-perfect graphs without clique cutset that allows us to conclude that the only graphs with these properties are the gear and the 3-gear graphs depicted in Figure 4.

In order to simplify the writing, we denote by  $\mathcal{G}$  the class of connected clawfree not quasi-line graphs without clique cutset.

We first prove the following:

#### **Theorem 15.** Let $G \in \mathcal{G}$ . If G is $LS_+$ -perfect then $\alpha(G) \leq 3$ .

*Proof.* Assume that G is LS<sub>+</sub>-perfect and  $\alpha(G) \geq 4$ .

As we have already pointed out in Section 2.4, according to [31], G has a decomposition into strips, where at most one strip is quasi-line and all the remaining ones have stability number at most 3 and contain a 5-wheel each. Since G is not quasi-line, it contains at least one 5-wheel strip G'. Also, recall that there are only three types of 5-wheel strips, Fig. 3 shows the "basic" types, which can be extended by adding nodes belonging to the neighborhood of the 5-wheels.



Figure 5:  $LS_+$ -imperfect subgraphs if G' is of type 1.



Figure 6:  $LS_+$ -imperfect subgraphs if G' is of type 2.

If G' is of type 3, then G' contains  $G_{LT}$ , induced by the squared nodes indicated in Fig. 3, contradicting the assumption that G is LS<sub>+</sub>-perfect. Let us analyze the cases where G' is of type 1 or 2.

Note further that G' is a proper subgraph of G since  $\alpha(G') \leq 3$  and  $\alpha(G) \geq 4$ . Then, G contains as subgraph the composition of G' and another strip graph, say G''. Then, there are two nodes in G'' playing the role of the two simplicial nodes of G' (the two black nodes in Fig. 3). Since G has no clique cutset, these two nodes are connected by a path P in G''.

If G' is of type 1 and P has even length, G has a node stretching of  $G_{EMN}$  as induced subgraph. Similarly, if P is of odd length it has a node stretching of  $G_{LT}$  (see the squared nodes in Fig. 5).

Similarly, if G' is of type 2 and P has even length, G has a node stretching of  $G_{LT}$  as induced subgraph. Finally, if P is odd G' has a node stretching of  $G_{EMN}$  (see the squared nodes in Fig. 6).

Hence, in all cases we contradict the LS<sub>+</sub>-perfection of G and we conclude that  $\alpha(G) \leq 3$ .

As an immediate consequence, we obtain:

**Corollary 16.** Every facet-defining claw-free not quasi-line graph G with  $\alpha(G) \geq 4$  is LS<sub>+</sub>-imperfect.

For claw-free graphs having stability number three, there is no decomposition known yet. Hence, we start with some technical results that will allow us to characterize  $LS_+$ -perfect graphs in  $\mathcal{G}$  with stability number three.

In what follows, given G = (V, E) and  $v \in V$ , we denote by  $G_v$  the subgraph of G induced by N[v]. It is clear that there is no stable set of size 3 in N(v) for every node v of a claw-free graph. The following remark directly follows: **Remark 17.** Let G = (V, E) be a claw-free graph. Then,  $\alpha(G_v) \leq 2$  for all  $v \in V$ . Moreover, given K, L, M three mutually disjoint nonempty subsets of V such that  $K \cup L$  is completely joined to M, it holds that  $\alpha(G[K \cup L]) \leq 2$ . In particular, if no node of K is adjacent to any node in L, then K and L are cliques.

In addition, if  $C \subset V(G)$  induces an odd antihole, we assume  $C = \{u_i : i \in [2k+1]\}$  with  $k \geq 2$  and, for all  $i \in [2k+1]$ ,  $u_i, u_{i+k}$  are not adjacent in G. We say that  $u_i$  and  $u_{i+1}$  are consecutive (in C), for all  $i \in [2k+1]$ .

We have the following result:

**Lemma 18.** Let G = (V, E) be a connected claw-free graph and let  $C \subset V$ induce an odd antihole in G. If there exists  $y \in V \setminus C$  such that  $N(y) \cap C \neq \emptyset$ and  $\alpha(G[C \cup \{y\}]) = 3$  then k = 2. Moreover, y has exactly two neighbors in Cwhich are consecutive.

*Proof.* Let  $C = \{u_i : i \in [2k+1]\}$  for some  $k \ge 2$  and  $y \in V$  such that  $N(y) \cap C \neq \emptyset$  and  $\alpha(G[C \cup \{y\}]) = 3$ . Then, there exists  $i \in [2k+1]$  such that  $S = \{y, u_i, u_{i+k}\}$  is a stable set of G.

If the nodes in S have a common neighbor z then  $S \cup \{z\}$  induces a claw. Therefore, y is not adjacent to any node in  $N(u_i) \cap N(u_{i+k})$ .

Since  $C \setminus N(u_i) = \{u_i, u_{i+k}, u_{i+k+1}\}, C \setminus N(u_{i+k}) = \{u_i, u_{i+k}, u_{i+2k}\}, \text{ and } y \text{ is not adjacent to } u_i, u_{i+k} \text{ it holds that } \emptyset \neq N(y) \cap C \subset \{u_{i+k+1}, u_{i+2k}\}.$ 

If  $N(y) \cap C = \{u_{i+2k}\}$  then  $\{u_{i+2k}, u_{i+1}, y, u_{i+k+1}\}$  induces a claw. Similarly, if  $N(y) \cap C = \{u_{i+k+1}\}$  then  $\{u_{i+k+1}, u_{i+1}, y, u_{i+2k}\}$  is a set inducing a claw. Therefore,  $N(y) \cap C = \{u_{i+k+1}, u_{i+2k}\}$ .

Finally, if  $k \ge 3$ ,  $u_{i+k+2} \ne u_{i+2k}$  then we have again a claw induced by the nodes  $\{u_{i+2k}, u_{i+1}, y, u_{i+k+2}\}$ . Then, k = 2 implying that the two neighbors of y in C are  $u_{i+3}$  and  $u_{i+4}$ , which are consecutive.

Let us focus on graphs  $G \in \mathcal{G}$  with stability number three. By Remark 17,  $\alpha(G_v) \leq 2$  for every node v. But, if there is  $C \subset N(v)$  inducing an odd antihole in G it holds that  $\alpha(G_v) = 2$ . If  $G_v$  is LS<sub>+</sub>-perfect, using Theorem 8 we obtain the next observation that will be useful in what remains of this section.

**Remark 19.** Let  $v \in V(G)$  such that  $G_v$  is  $LS_+$ -perfect. If there exists  $C \subset N(v)$  inducing an odd antihole in G, then C is completely joined to  $N[v] \setminus C$ .

The next result gives a decomposition of graphs in  $\mathcal{G}$  with stability number three such that every subgraph with stability number two is LS<sub>+</sub>-perfect.

**Lemma 20.** Let G = (V, E) be a graph such that  $G \in \mathcal{G}$ ,  $\alpha(G) = 3$ , and for all  $G' \subset G$  with  $\alpha(G') = 2$ , G' is  $LS_+$ -perfect. Then, V can be partitioned into the sets  $Q, C, Y_2, Y_3, Z$  such that C induces  $C_5$ ,  $|Y_3| \ge 2$ , while  $Q, Y_2$ , and Z are cliques. In addition,

(1) Q is completely joined to C and no node in Q is adjacent to any node in  $V \setminus (C \cup Q)$ ,

- (2)  $Y_2$  is completely joined to  $C \cup Y_3$ ,
- (3) Z is completely joined to  $Y_3$  and no node in Z is adjacent to any node in  $Q \cup C$ ,
- (4) every node in Y<sub>3</sub> has exactly two neighbors in C which are consecutive. Moreover, if two nodes in Y<sub>3</sub> share a neighbor in C then they are adjacent. Also, not all nodes in Y<sub>3</sub> have the same neighbors in C.

*Proof.* Since G is claw-free and not quasi-line, there exists a node v such that N(v) cannot be partitioned into two cliques. Then, the complementary graph of  $G_v \setminus \{v\}$  is not bipartite and it has a set of nodes C inducing an odd cycle. Since  $\alpha(G_v) \leq 2$ ,  $|C| \geq 5$ . Then C induces an antihole in G and we denote  $C = \{u_i : i \in [2k+1]\}$  for some  $k \geq 2$ .

Let  $W = N(v) \setminus C$  and  $Q = W \cup \{v\}$ . Clearly, C, Q is a partition of N[v]. By assumption  $G_v$  is LS<sub>+</sub>-perfect. Then, Remark 19 ensures that Q is completely joined to C. Thus, we have partially proved (1).

Since  $\alpha(G_v) = 2$  and  $\alpha(G) = 3$ ,  $V \setminus N[v] \neq \emptyset$ . Now, consider the partition of  $V \setminus N[v]$  into the sets X, Y, Z such that:

- $X = \{x \in V \setminus N[v] : N(x) \cap W \neq \emptyset\},\$
- $Y = \{y \in V \setminus N[v] : N(y) \cap W = \emptyset \text{ and } N(y) \cap C \neq \emptyset\}$ , and
- $Z = \{z \in V \setminus N[v] : N(z) \cap N[v] = \emptyset\}.$

Observe that, since  $\alpha(G) = 3$ ,  $\alpha(G_v) = 2$ , and no node in Z is connected with any node in N[v], we have that  $\alpha(G[Z]) \leq 1$  and then Z is a (possibly empty) clique.

In order to complete the proof of (1) we need to show that  $X = \emptyset$ . Let us assume that  $X \neq \emptyset$ .

Let  $x \in X$  and  $w \in N(x) \cap W$ . Since  $C \subset N(w)$  and we assume that  $G_w$  is  $LS_+$ -perfect, Remark 19 implies that C is completely joined to  $N[w] \setminus C$  and, in particular, C is completely joined to  $\{x\}$ . Then X is completely joined to C. Now, by Remark 17 considering  $K = \{v\}$ , L = X, and M = C it holds that X is a clique.

Since  $Q \cup X$  is completely joined to C, by Remark 17 with M = C, K = X, and L = Q we obtain that  $\alpha(G[Q \cup X]) \leq 2$ . Then  $\alpha(G[N[v] \cup X]) = 2$ . Since  $\alpha(G) = 3$ ,  $Y \cup Z \neq \emptyset$ . If  $Y = \emptyset$ , X is a clique cutset, contradicting the hypothesis. Then,  $Y \neq \emptyset$ .

Let  $y \in Y$  and  $u \in C$  such that  $y \in N(u)$ . Again, considering Remark 17 with  $K = \{v\}, L = X \cup \{y\}$ , and  $M = \{u\}$ , we have that L is a clique. Therefore Y is completely joined to X.

So we have that  $C \cup Y \subset N(x)$  for every  $x \in X$  and by Remark 19 (using that  $G_x \text{ LS}_+$ -perfect) we get that C is completely joined to Y. Applying again Remark 17 for  $K = \{v\}, L = X \cup Y$  and M = C we obtain that  $X \cup Y$  is a clique and then  $\alpha(N[v] \cup X \cup Y) = 2$ . This implies  $Z \neq \emptyset$  and  $X \cup Y$  is a clique cutset, a contradiction to the assumption on G.



Figure 7: A scheme of a graph with a partition as in Lemma 20 (sets with bold lines stand for complete graphs, the  $\bowtie$  symbol indicates a complete join between two sets).

Then,  $X = \emptyset$  and (1) holds.

Since G is connected, (1) implies that  $Y \neq \emptyset$ . Moreover,  $|Y| \ge 2$  otherwise Y is a clique cutset.

Let  $y \in Y$  and  $u \in C$  such that  $y \in N(u)$ . Since C is completely joined to Q, Remark 17 with K = Q,  $L = \{y\}$  and  $M = \{u\}$  implies that Q is a clique.

Let us show that any two nodes in Y sharing a node in U are adjacent. Indeed, consider  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$  such that  $\{y_1, y_2\} \subset N(u)$  for some  $u \in C$ . It follows from Remark 17 with K = Q,  $L = \{y_1, y_2\}$  and  $M = \{u\}$  that L is a clique.

Now, we split Y into the sets  $Y_i = \{y \in Y : \alpha(G[\{y\} \cup C]) = i\}$  for i = 2, 3. In this way, V is partitioned into C, Q, Y<sub>2</sub>, Y<sub>3</sub>, and Z. Recall that Q is a nonempty clique  $(v \in Q)$ .

In order to prove (2), observe that Theorem 8 ensures that  $Y_2$  is completely joined to C. Since two nodes in Y sharing a neighbor in C are adjacent, it follows that  $Y_2$  is a clique. Moreover,  $Y_2$  is completely joined to  $C \cup Y_3$ .

Suppose now that  $Y_3 = \emptyset$ . Then  $\alpha(G[N[v] \cup Y]) = 2$  and  $Z \neq \emptyset$  follows. Thus,  $Y_2$  is a clique cutset and we arrive to a contradiction. Then,  $Y_3 \neq \emptyset$ . If  $|Y_3| = 1$  and using (2) we have Y is a clique cutset, a contradiction. Thus,  $|Y_3| \geq 2$ . Moreover, by Lemma 18, C induces  $C_5$  and every node in  $Y_3$  is adjacent to two consecutive nodes in C.

To complete the proof of (3), suppose that there are  $y \in Y_3$  and  $z \in Z$  which are not adjacent. It holds that there exists  $i \in [5]$  such that  $\{y, u_i, u_{i+2}, z\}$  is a stable set of size 4, which is a contradiction on  $\alpha(G) = 3$ . Then, Z is completely joined to  $Y_3$ .

Finally, not all nodes in  $Y_3$  have the same two neighbors in C, say  $u_i, u_{i+1}$  for some  $i \in [5]$ . Otherwise,  $Y_2 \cup \{u_i, u_{i+1}\}$  is a clique cutset. This completes the proof of (4).

It is not hard to check that the support graphs of facet-defining inequalities



Figure 8: Subgraph induced by  $C_5 \cup \{v, y, y', y^*\}$ , the bold edges indicate the  $G_{EMN}$ . Removing  $y_*$  yields the gear.

of the gear and the 3-gear graphs are complete and 5-wheel graphs. Then, the gear and the 3-gear graphs are  $LS_+$ -perfect. Also, since any node of these graphs belongs to a 5-hole (see Figure 4), by Remark 5 we have the following observation:

**Remark 21.** The graph obtained after the replication of any node in the gear or the 3-gear graph is  $LS_+$ -imperfect.

Finally, we obtain:

**Theorem 22.** Let  $G \in \mathcal{G}$  an  $LS_+$ -perfect graph with  $\alpha(G) \geq 3$ . Then G is the gear or the 3-gear graph.

*Proof.* By Theorem 15,  $\alpha(G) = 3$ . Since G is LS<sub>+</sub>-perfect, by the previous lemma we can consider the partition of V(G) into the sets Q, C, Y<sub>2</sub>, Y<sub>3</sub>, and Z.

Recall that  $Q \neq \emptyset$  and let  $v \in Q$ . Moreover, we know that  $|Y_3| \ge 2$ , not all the nodes in  $Y_3$  have the same two consecutive neighbors in C, and two nodes in  $Y_3$  having a common neighbor in C are adjacent. Let us now prove that adjacent nodes in  $Y_3$  share a common neighbor in C.

Let  $y \neq y' \in Y_3$  such that y is adjacent to y' and assume that they do not share a common neighbor in C. W.l.o.g. we can assume that  $N(y) \cap C = \{u_1, u_2\}$ and  $N(y') \cap C = \{u_3, u_4\}$ . Then,  $\{v, u_1, y, y', u_4, u_5\}$  induces the graph  $G_{LT}$ contradicting the fact that G is LS<sub>+</sub>-perfect.

Let us prove that if there is a node in C completely joined to  $Y_3$  then G is the gear graph.

W.l.o.g. assume that  $u_2 \in N(y)$  for all  $y \in Y_3$ . In this case  $Y_3$  is a clique. Then  $Z = \emptyset$ , otherwise,  $Y_2 \cup Y_3$  is a clique cutset. Since not all the nodes in  $Y_3$  have the same neighbors in C, there exist  $y, y' \in Y_3$  such that  $N(y) \cap C = \{u_1, u_2\}$  and  $N(y') \cap C = \{u_2, u_3\}$ .

If there is  $y^* \in Y_2$  then,  $\{v, u_1, u_3, y, y', y^*\}$  induces  $G_{EMN}$  (see Figure 8), again a contradiction to the fact that G is LS<sub>+</sub>-perfect. Then,  $Y_2 = \emptyset$ .

It is easy to see that every node in Q is a replication of v in G. Due to Remark 21, the LS<sub>+</sub>-perfection of G forbids the replications of v and then G is the gear graph.

Let us consider the case where not all the nodes in  $Y_3$  have a common neighbor in C. W.l.o.g. we can assume that there are two nonadjacent nodes y, y' in  $Y_3$  such that  $N(y) \cap C = \{u_1, u_2\}$  and  $N(y') \cap C = \{u_3, u_4\}$ .



Figure 9: A node stretching of  $G_{EMN}$ 



Figure 10: The graph *H*. Black nodes induce a stretching of  $G_{EMN}$ .

Observe that  $Y_2 = Z = \emptyset$ . Indeed, if there is  $y^* \in Y_2$  the set  $\{y^*, y, y', u_5\}$  induces a claw. If  $z \in Z$ , from Lemma 20 (3), z is adjacent to every node in  $Y_3$  and then  $\{z, y, y'\} \cup C$  induces a node stretching of  $G_{EMN}$  (see Figure 9).

Then, V(G) is partitioned into Q, C and  $Y_3$ . Also,  $\{v\} \cup C$  induces a 5-wheel for all  $v \in Q$ ,  $N(y) \cap C = \{u_1, u_2\}$ , and  $N(y') \cap C = \{u_3, u_4\}$ . Since G has no clique cutset, there must be another node  $t \in Y_3$  which is not a replication either of y or y'.

Let us first analyze the case when no node of  $Y_3$  is adjacent to  $\{u_2, u_3\}$ . In this case, w.l.o.g. we can assume that  $N(t) \cap C = \{u_1, u_5\}$ . Again, since G has no clique cutset, there must exist another node y'' which is not a replication of any node in  $\{y, t, y'\}$ . Then,  $N(y'') \cup C = \{u_4, u_5\}$ . Figure 10 shows the graph  $\hat{H}$  induced by  $C \cup \{v, y, y', t, y''\}$ . Observe that the black nodes in  $\hat{H}$  induces a stretching of  $G_{EMN}$ , a contradiction to the fact that G is  $LS_+$ -perfect.

It only remains to analyze the case when  $N(t) \cap C = \{u_2, u_3\}$ .

If there is another node  $y'' \in Y_3$ , by Remark 21, it is not a replication of any node in  $\{y, t, y'\}$  and w.l.o.g. we can assume that  $N(y'') = \{u_1, u_5\}$ . It is easy to see that the subgraph induced by  $C \cup \{v, y, y', t, y''\}$  is the graph  $\hat{H}$  after relabeling its nodes (see Figure 11), a contradiction.

Then, there is no other node in  $Y_3$  and G is the graph induced by  $C \cup \{v, y, y', t\}$  which is the 3-gear graph (see Figure 12).

Since the gear and the 3-gear graphs are not facet-defining, from Theorem



Figure 11: The graph  $\hat{H}$  with a relabeling of its nodes.



Figure 12: Subgraph induced by  $C_5 \cup \{v, y, y', t\}$ , called a 3-gear.

22 we conclude:

**Corollary 23.** Every facet-defining claw-free not quasi-line graph G with  $\alpha(G) \geq 3$  is LS<sub>+</sub>-imperfect.

Combining corollaries 9, 14, and 23, we obtain our main contribution:

**Theorem 24.** The  $LS_+$ -Perfect Graph Conjecture is true for claw-free graphs.

# 4 On minimally LS<sub>+</sub>-imperfect claw-free graphs

In this section we address our attention to minimally  $LS_+$ -imperfect graphs that are claw-free. These graphs are facet-defining and therefore the results in Section 3 will be reused in order to describe all minimally  $LS_+$ -imperfect claw-free graphs.

There are some known results on this matter. In [2], we consider for each  $k \ge 2$ , a family of graphs with 2k+2 nodes, having a 2k+1-antihole as subgraph. Theorem 4.3 in the paper, states that the only minimally LS<sub>+</sub>-imperfect graph with stability number two in the family is the graph for which the node outside the antihole has degree 2k. This graph is called  $H_k$ .

**Lemma 25.** Let G = (V, E) be a minimally  $LS_+$ -imperfect graph with  $\alpha(G) = 2$ . Then, G is either  $G_{LT}$ ,  $G_{EMN}$ , or  $H_k$ , for some  $k \ge 3$ .

*Proof.* Since G is imperfect, minimally LS<sub>+</sub>-imperfect and  $\alpha(G) = 2$ , G has a (2k+1)-antihole induced by  $C \subset V$  and a node  $v \in V \setminus C$  outside C. By Theorem



Figure 13: The graphs  $H^1$  and  $H^2$ , for k = 2, t = 1, i = 1 and j = 2, 3, respectively.

7, v is not completely joined to C. Since G is minimally LS<sub>+</sub>-imperfect it holds that  $V = C \cup \{v\}$  and either k = 2 implying that  $G = G_{LT}$  or  $G = G_{EMN}$ , or  $k \geq 3$  and from the discussion above we get  $G = H_k$ .

Remind that the facet-defining semi-line graphs are line graphs (of 2-connected hypomatchable graphs). In addition, Corollary 10 in [17] establishes that the line graph of graph H is minimally LS<sub>+</sub>-imperfect if and only if H has a particular structure that can be described as follows: H is an odd cycle  $C_{2k+1}$ ,  $k \ge 2$ , with a path of odd length  $P_t$ ,  $t \ge 1$  attached to nodes i and j in  $C_{2k+1}$  (observe that the path can have length one and H can have a double edge). It is not hard to see that every such graph H is an odd-subdivision of one of the graphs depicted in Figure 13.

Observe that the line graphs of  $H^1$  and  $H^2$  are  $G_{LT}$  and  $G_{EMN}$ , respectively. From Lemma 2 in [17] we can easily see that if a graph G is an odd subdivision of an edge e in a graph G', L(G) can be obtained from L(G') by stretching of its node e.

This allows us to conclude:

**Lemma 26.** Let G be a minimally  $LS_+$ -imperfect semi-line graph. Then G is a stretching of  $G_{LT}$  or a stretching of  $G_{EMN}$ .

Let us now consider fuzzy circular interval graphs.

**Theorem 27.** Let G be a minimally  $LS_+$ -imperfect fuzzy circular interval graph. Then, G is either  $W_{10}^2$ ,  $G_{LT}$ ,  $G_{EMN}$ ,  $G = H_k$  for some  $k \ge 3$ , a stretching of  $G_{LT}$ , or a stretching of  $G_{EMN}$ .

*Proof.* Let assume that G is neither  $W_{10}^2$ ,  $G_{LT}$  nor  $G_{EMN}$ . Since G is facetdefining, we can follow the reasoning in the proof of Theorem 13.

Remind that G is the support graph of a clique family inequality (2) associated with a web subgraph  $W_n^{p-1}$  of G. Since G is minimally  $LS_+$ -imperfect then  $W_n^{p-1}$  is  $LS_+$ -perfect and there exists a node v of G outside the web. Again Theorem 12 implies that the web  $W_n^{p-1}$  in question is an odd cycle or its complement and we obtain that n = 2k + 1 with  $k \ge 3$  and p = 2 or  $p = k \ge 3$ . In both cases, r = 1.

As we have already observed in the proof of Theorem 13, when p = 2 it holds that p - r - 1 = 0 and v is connected to  $t \ge 2p - 1 = 3$  consecutive nodes in  $W_{2k+1}^1$ . As  $G_{LT}$  and  $G_{EMN}$  cannot be subgraphs of a minimally LS<sub>+</sub>-imperfect graph then  $k \geq 3$  and G has an stretching of  $G_{LT}$  or of  $G_{EMN}$  as a subgraph. Since the latter graphs are minimally LS<sub>+</sub>-imperfect then G is one of them.

In the case  $p = k \ge 3$ , we have also observed that v is adjacent to at least 2k - 2 consecutive nodes in  $W_{2k+1}^{k-1}$ . Thus, the subgraph of G induced by  $V(W_{2k+1}^{k-1}) \cup \{v\}$  has stability number two. Then, Lemma 25 implies that the only possible choice is that  $G = H_k$  for some  $k \ge 3$ .

Concerning claw-free not quasi-line graphs we have seen in Theorem 15 that if the graph has stability number at least 4, it has a  $G_{LT}$  or a  $G_{EMN}$  as node induced subgraph.

**Remark 28.** There is no minimally  $LS_+$ -imperfect graph in the class of clawfree not quasi-line graphs with stability number at least four.

A similar result holds when the stability number is equal to three. In this case we make use of Theorem 22.

**Theorem 29.** There is no minimally  $LS_+$ -imperfect graph in the class of clawfree not quasi-line graphs with stability number three.

*Proof.* Let  $G \in \mathcal{G}$  be a minimally LS<sub>+</sub>-imperfect graph with stability number three. It is clear that G cannot have a minimally LS<sub>+</sub>-imperfect graph with stability number two as induced subgraph, then we can consider the partition  $Q, C, Y_2, Y_3$ , and Z of V(G) given by Lemma 20. By doing so, we can follow the reasoning in the proof of Theorem 22 and conclude that if G is LS<sub>+</sub>-imperfect and has stability number three, then it has a node stretching of  $G_{LT}$  or  $G_{EMN}$ as a subgraph. But, since G is minimally LS<sub>+</sub>-imperfect then G is one of these graphs itself. Then G is a line graph and we arrive to a contradiction.

From the previous results we have that minimally  $LS_+$ -imperfect claw free graphs are fuzzy circular interval graphs, that is:

**Corollary 30.** Let G be a minimally  $LS_+$ -imperfect claw-free graph. Then, G is  $W_{10}^2$ ,  $G_{LT}$ ,  $G_{EMN}$ ,  $G = H_k$  for some  $k \ge 3$ , a stretching of  $G_{LT}$ , or a stretching of  $G_{EMN}$ .

## 5 Conclusion and future research

The context of this work was the study of  $LS_+$ -perfect graphs, i.e., graphs where a single application of the Lovász-Schrijver PSD-operator  $LS_+$  to the edge relaxation yields the stable set polytope. Hereby, we are particularly interested in finding an appropriate polyhedral relaxation P(G) of STAB(G) that coincides with  $LS_+(G)$  and STAB(G) if and only if G is  $LS_+$ -perfect. An according conjecture has been recently formulated ( $LS_+$ -Perfect Graph Conjecture); here we verified it for the well-studied class of claw-free graphs (Theorem 24). Note further that, besides verifying the  $LS_+$ -Perfect Graph Conjecture for claw-free graphs, we obtained a description of all minimally  $LS_+$ -imperfect claw-free graphs (Corollary 30).

For that, it surprisingly turned out that it was not necessary to make use of the description of STAB(G) for claw-free not quasi-line graphs G

- with  $\alpha(G) = 2$  (by Cook, see [36]),
- with  $\alpha(G) = 3$  (by Pêcher, Wagler [33]),
- with  $\alpha(G) \ge 4$  (by Galluccio, Gentile, Ventura [18, 19, 20]).

From the presented results and proofs, we can draw some further conclusions. First of all, we can determine the subclass of joined *a*-perfect graphs to which all  $LS_+$ -perfect claw-free graphs belong to. In [25], it is suggested to call a graph *G m*-perfect if the only facets of STAB(G) are associated with cliques and minimally imperfect graphs. According to [11], *G* is *joined m*-perfect if its facet defined graphs are complete graphs, minimally imperfect graphs and their complete joins. The results from Section 3 provide the complete list of all facet-defining  $LS_+$ -perfect claw-free graphs:

- complete graphs,
- odd holes and odd antiholes,
- complete joins of odd antihole(s) and a (possibly empty) complete graph.

Hence, we conclude:

**Corollary 31.** All  $LS_+$ -perfect claw-free graphs are joined m-perfect.

Finally, the subject of the present work has parallels to the well-developed research area of perfect graph theory also in terms of polynomial time computability. In fact, it has the potential of reaching even stronger results due the following reasons. Actually, in [29], the authors prove that calculating the value

$$\eta_+(G) = \max \mathbf{1}x, x \in \mathrm{LS}_+(G)$$

can be obtained with arbitrary precision in polynomial time for every graph G, even in the weighted case. Thus, the stable set problem can be solved in polynomial time for a strict superset of perfect graphs, the LS<sub>+</sub>-perfect graphs, by  $\alpha(G) = \eta_+(G)$ . Hence, our future lines of research include to find

- new families of graphs where the conjecture holds (e.g., by characterizing the minimally LS<sub>+</sub>-imperfect graphs within the class),
- new subclasses of LS<sub>+</sub>-perfect or joined a-perfect graphs,
- classes of graphs G where STAB(G) and  $LS_+(G)$  are "close enough" to have  $\alpha(G) = \lfloor \eta_+(G) \rfloor$ .

In particular, the class of graphs G with  $\alpha(G) = \lfloor \eta_+(G) \rfloor$  can be expected to be large since  $\mathrm{LS}_+(G)$  is a much stronger relaxation of  $\mathrm{STAB}(G)$  than  $\mathrm{TH}(G)$ . In all cases, the stable set problem could be approximated with arbitrary precision in polynomial time in these graph classes by optimizing over  $\mathrm{LS}_+(G)$ . Finally, note that  $\mathrm{LS}_+(P(G))$  with

$$STAB(G) \subseteq P(G) \subseteq ESTAB(G)$$

clearly gives an even stronger relaxation of STAB(G) than  $LS_+(G)$ . However, already approximating with arbitrary precision over  $LS_+(QSTAB(G))$  cannot be done in polynomial time anymore for all graphs G by [29]. Hence,  $LS_+$ perfect graphs or their generalizations satisfying  $\alpha(G) = \lfloor \eta_+(G) \rfloor$  are the most promising cases in this context.

## References

- C. Berge, "Perfect graphs", Six Papers on Graph Theory, Calcutta: Indian Statistical Institute, pp. 1–21, 1963.
- [2] S. Bianchi, M. Escalante, G. Nasini, "Some forbidden subgraphs for LS<sub>+</sub>perfection", Annals of the Latin-Iberoamerican Conference on Operations Research (CLAIO), 2016.
- [3] S. Bianchi, M. Escalante, G. Nasini, and L. Tunçel, "Near-perfect graphs with polyhedral N<sub>+</sub>(G)," *Electronic Notes in Discrete Mathematics*, vol. 37, pp. 393–398, 2011.
- [4] S. Bianchi, M. Escalante, G. Nasini, and L. Tunçel, "Lovász-Schrijver PSDoperator and a superclass of near-perfect graphs", *Electronic Notes in Discrete Mathematics*, vol. 44, pp. 339–344, 2013.
- [5] S. Bianchi, M. Escalante, G. Nasini, and L. Tunçel, "Lovász-Schrijver PSDoperator, near-perfect and near-bipartite graphs", *Mathematical Program*ming., vol. 162 (1-2), pp. 201–223, 2017.
- [6] S. Bianchi, M. Escalante, G. Nasini, and A. Wagler, "Lovász-Schrijver PSD-Operator on Claw-Free Graphs" In: Cerulli R., Fujishige S., Mahjoub A. (eds) Combinatorial Optimization. ISCO 2016. Lecture Notes in Computer Science, vol. 9849, pp. 59–70, 2016.
- [7] S. Bianchi, M. Escalante, G. Nasini, and A. Wagler, "Minimally LS<sub>+</sub>imperfect claw-free graphs" In: Bordeaux Graph Workshop 2016 Booklet, pp. 29–32, 2016.
- [8] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas, "The Strong Perfect Graph Theorem," Annals of Mathematics, vol. 164, pp. 51–229, 2006.

- [9] M. Chudnovsky and P. Seymour, *The structure of claw-free graphs*, manuscript (2004).
- [10] V. Chvátal, "On Certain Polytopes Associated with Graphs," J. Combin. Theory (B), vol. 18, pp. 138–154, 1975.
- [11] S. Coulonges, A. Pêcher, and A. Wagler, "Characterizing and bounding the imperfection ratio for some classes of graphs," *Math. Programming A*, vol. 118, pp. 37–46, 2009.
- [12] J.R. Edmonds, "Maximum Matching and a Polyhedron with (0,1) Vertices," J. Res. Nat. Bur. Standards, vol. 69B, pp. 125–130, 1965.
- [13] J.R. Edmonds and W.R. Pulleyblank, "Facets of 1-Matching Polyhedra," In: C. Berge and D.R. Chuadhuri (eds.) Hypergraph Seminar. Springer, pp. 214–242, 1974.
- [14] F. Eisenbrand, G. Oriolo, G. Stauffer, and P. Ventura, The stable set polytope of quasi-line graphs. Combinatorica, 28:45–67, 2008.
- [15] M. Escalante, M.S. Montelar, and G. Nasini, "Minimal N<sub>+</sub>-rank graphs: Progress on Lipták and Tunçel's conjecture," *Operations Research Letters*, vol. 34, pp. 639–646, 2006.
- [16] M. Escalante and G. Nasini, "Lovász and Schrijver N<sub>+</sub>-relaxation on web graphs," *Lecture Notes in Computer Sciences*, vol. 8596 (ISCO 2014), pp. 221–229, 2014.
- [17] M. Escalante, G. Nasini, and A. Wagler, "Characterizing  $N_+$ -perfect line graphs", International Transactions in Operational Research, 2015, DOI: 10.1111/itor.12275.
- [18] A. Galluccio, C. Gentile, and P. Ventura. Gear composition and the stable set polytope. Operations Research Letters, 36:419–423, 2008.
- [19] A. Galluccio, C. Gentile, and P. Ventura. The stable set polytope of clawfree graphs with stability number at least four. I. Fuzzy antihat graphs are W-perfect, Journal of Combinatorial Theory Series B 107, 92-122, 2014
- [20] A. Galluccio, C. Gentile, and P. Ventura. The stable set polytope of clawfree graphs with stability number at least four. II. Striped graphs are Gperfect, Journal of Combinatorial Theory Series B 108, 1-28, 2014
- [21] A. Galluccio and A. Sassano, The Rank Facets of the Stable Set Polytope for Claw-Free Graphs. J. Comb. Theory B 69 (1997) 1–38.
- [22] R. Giles and L.E. Trotter. On stable set polyhedra for K1,3 -free graphs. J. of Combinatorial Theory, 31:313–326, 1981.

- [23] M. Grötschel, L. Lovász, and A. Schrijver, "The Ellipsoid Method and its Consequences in Combinatorial Optimization," *Combinatorica*, vol. 1, pp. 169–197, 1981.
- [24] M. Grötschel, L. Lovász, and A. Schrijver, Geometric Algorithms and Combinatorial Optimization. Springer-Verlag (1988)
- [25] A. Koster and A. Wagler, Comparing imperfection ratio and imperfection index for graph classes, RAIRO Operations Research, vol. 42, pp. 485-500, 2008.
- [26] T.M. Liebling, G. Oriolo, B. Spille, and G. Stauffer. On non-rank facets of the stable set polytope of claw-free graphs and circulant graphs. Mathematical Methods of Operations Research, 59(1):25–35, 2004.
- [27] L. Lipták and L. Tunçel, "Stable set problem and the lift-and-project ranks of graphs," *Math. Programming A*, vol. 98, pp. 319–353, 2003.
- [28] L. Lovász, "On the Shannon capacity of a graph," *IEEE Trans. Inf. Theory*, vol. 25, pp. 1–7, 1979.
- [29] L. Lovász and A. Schrijver, "Cones of matrices and set-functions and 0-1 optimization," SIAM J. on Optimization, vol. 1, pp. 166–190, 1991.
- [30] G. Minty, "On maximal independent sets of vertices in claw-free graphs", J. of Combinatorial Theory, vol. 28, pp. 284–304, 1980.
- [31] G. Oriolo, U. Pietropaoli, and G. Stauffer, "A new algorithm for the maximum weighted stable set problem in claw-free graphs", in A. Lodi, A. Panconesi and G. Rinaldi (eds), IPCO 2008, Vol. 5035 of Lecture Notes in Computer Science, Springer, Bertinoro, Italy, May 26-28, pp. 77–96, 2008.
- [32] A. Pêcher, A. Wagler, "Almost all webs are not rank-perfect", Math. Programming B 105 (2006) 311–328
- [33] A. Pêcher, A. Wagler, "On facets of stable set polytopes of claw-free graphs with stability number three", Discrete Mathematics 310 (2010) 493–498
- [34] U. Pietropaoli, A. Wagler, "Some results towards the description of the stable set polytope of claw-free graphs", Proc. of ALIO/EURO Workshop on Applied Combinatorial Optimization, Buenos Aires, 2008
- [35] N. Sbihi, "Algorithme de recherche d'un stable de cardinalité maximum dans un graphe sans étoile", Discrete Mathematics 29, 53–76, 1980.
- [36] F.B. Shepherd, "Near-Perfect Matrices," Math. Programming, vol. 64, pp. 295–323, 1994.
- [37] F.B. Shepherd, "Applying Lehman's Theorem to Packing Problems," Math. Programming, vol. 71, pp. 353–367, 1995.

- [38] G. Stauffer, "On the Stable set polytope of claw-free graphs". Phd thesis, 2005, Swiss Institute of Technology in Lausanne.
- [39] G. Stauffer, personal communication (2011).
- [40] A. Wagler, *Critical Edges in Perfect Graphs.* PhD thesis, TU Berlin and Cuvillier Verlag Göttingen (2000).
- [41] A. Wagler, "Antiwebs are rank-perfect," 4OR, vol. 2, pp. 149-152, 2004.
- [42] A. Wagler, "On rank-perfect subclasses of near-bipartite graphs," 4OR, vol. 3, pp. 329–336, 2005.
- [43] A. Wagler, "Lovász-Schrijver PSD-operator on some graph classes defined by clique cutsets," In: Lee J., Rinaldi G., Mahjoub A. (eds) Combinatorial Optimization: ISCO 2018. Lecture Notes in Computer Science, vol 10856. Springer, Cham, 2018.
- [44] A. Wagler, "On the Lovász-Schrijver PSD-operator on graph classes defined by clique cutsets," to appear in *Discrete Applied Mathematics*, DOI: 10.1016/j.dam.2019.07.017
- [45] L.A. Wolsey, "Further facet generating procedures for vertex packing polytopes". Math. Program., vol. 11, pp. 158–163, 1976.