



HAL
open science

Celullar Economics in Cell-Free

Olivier Borkowski

► **To cite this version:**

Olivier Borkowski. Celullar Economics in Cell-Free. Master. Université paris-saclay, France. 2024.
hal-04405950

HAL Id: hal-04405950

<https://hal.science/hal-04405950>

Submitted on 19 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Celular Economics in Cell-Free

Olivier Borkowski, PhD

Permanent researcher in Synthetic Biology
Institut Micalis, INRAE - AgroParisTech - Univ. Paris-Saclay, France
SyBER team
Email: Olivier.Borkowski@inrae.fr



Micalis Institute




21 Research Teams

- 340 people
- 130 scientists and engineers
- 140 PhD students and post-docs
- 70 scientific and administrative personnel

Themes

- (1) Emergence, control, and adaptation of pathogenic microorganisms
- (2) Microbial food and intestinal ecosystems
- (3) Synthetic and systems biology

Understand and use the relationship between protein production and bacterial physiology



Bacterial Physiology

Bacterial physiology is a branch of biology that aims to understand overarching principles of cellular reproduction:

What are the fundamental principles underlying the regulation and coordination of biosynthesis in a given growth environment?

-> Bacterial growth law

Bacterial Physiology

Bacterial physiology is a branch of biology that aims to understand overarching principles of cellular reproduction:

What are the fundamental principles underlying the regulation and coordination of biosynthesis in a given growth environment?

-> Bacterial growth law

Some major principles studied in bacterial physiology:


- Link between bacterial growth and nutrient concentrations in culture media (Monod, 1942)
- Link between the macromolecular composition and the cell doubling time (Schaechter et al., 1958)
- Linear relationships between RNA/DNA, RNA/protein, protein/DNA and growth rate (Bremer & Dennis, 1974)
- Increasing allocation of cellular resources to ribosome synthesis with increasing growth rate (Hwa & Scott, 2010)

-> New technology allows novel experiments


Bacterial Physiology

Experimental setup

Genomic regulations




E. coli living cells




Bacterial Physiology

Experimental setup

Genomic regulations



E. coli living cells




Intertwined Effects

-> New technology to decouple the many effects observed in living cells

Cell-free reaction:

Protein expression in an open system

A direct manipulation of the environment of protein production because it is an open system:




Carlson et al. Biotechnology advances (2012)

Cell-free reaction:


Protein expression in an open system

A direct manipulation of the environment of protein production because it is an open system:

- An alternative technology to study central process observed in living cells (replication, transcription, translation, degradation, metabolism)
- A versatile technology for bioengineering (Use in Synthetic Biology)



Carlson et al. *Biotechnology advances* (2012)



Nadanai, L. *Frontiers in Bioengineering and Biotechnology* (2020)

1. Description of Cell-Free Systems

History: First use of cell-free

What is the composition of a cell-free reaction?

Advantages of producing protein outside of the cell

Measurements in 384-well plate

Current existing protocol to produce bacterial cell extract

Alternative cell-free approach, the pure system

2. Current Applications of Cell-Free Systems in Academia and Industry

Cell-free reaction:

First major use of cell-free

Cell-free was first used in the 1960's by Nirenberg and Matthaei to understand the genetic code

American & German biochemist

Johannes Heinrich
Matthaei


*THE DEPENDENCE OF CELL-FREE PROTEIN SYNTHESIS IN E. COLI
UPON NATURALLY OCCURRING OR SYNTHETIC
POLYRIBONUCLEOTIDES*

BY MARSHALL W. NIRENBERG AND J. HEINRICH MATTHAEI*

NATIONAL INSTITUTES OF HEALTH, BETHESDA, MARYLAND

Communicated by Joseph E. Smadel, August 3, 1961

A stable cell-free system has been obtained from *E. coli* which incorporates C¹⁴-valine into protein at a rapid rate. It was shown that this apparent protein synthesis was energy-dependent, was stimulated by a mixture of L-amino acids, and was markedly inhibited by RNAase, puromycin, and chloramphenicol.¹ The present communication describes a novel characteristic of the system, that is, a requirement for template RNA, needed for amino acid incorporation even in the




Marshall Warren
Nirenberg

Cell-free reaction:


First major use of cell-free

Nirenberg's laboratory notebooks




Cell-free reaction:

History



Cell-free reaction:


What is the composition of a cell-free reaction?





Melinek B, et al. BioProcess International (2020)

Cell-free reaction:

What is the composition of a cell-free reaction?




Melinek B, et al. *BioProcess International* (2020)



Seok Hoon Hong et al. *Frontiers in Chemistry* 2014

Cell-free reaction: Cell-extract preparation



Melinek B, et al. *BioProcess International* (2020)

Physical disruption

Sonication




French press




Chemical disruption

Autolysis





Lysozyme




Cell-free reaction:

Protocol for home-made cell-extract preparation



Cell-free reaction: Buffer composition




Melinek B, et al. *BioProcess International* (2020)

Amino Acid
tRNA
Mg-glutamate
K-glutamate
Folnic Acid
Spermidine
Coenzyme A
NAD
3-PGA
AMPC
Nucleoside triphosphate (NTP)

Translation
Translation
Ribosome maintenance
Ribosome stability
A precursor for formylmethionine (fMet)
Ribosome activity /RNA stability
Regenerate ATP via glycolysis
Regenerate ATP via glycolysis
Regenerate ATP
regulating glycolysis
Transcription



Cell-free reaction:

Measurements in 384-well plate




Melinek B, et al. *BioProcess International* (2020)

1 to 20 μ l reactions
(in 384-well)





Cell-free reaction:

Measurements in 384-well plate




Melinek B, et al. *BioProcess International* (2020)

Translation




Transcription



Cell-free reaction:

Current existing protocol to produce bacterial cell extract




Melinek B, et al. *BioProcess International* (2020)

Table 1. Current existing protocols to produce cell-free lysates from bacteria cells

Organism	Bacteria Phylum
<i>E. coli</i>	Proteobacteria
<i>B. subtilis</i>	Firmicutes
<i>L. lactis</i>	Firmicutes
<i>B. megaterium</i>	Firmicutes
<i>C. autoethanogenum</i>	Firmicutes
<i>V. natriegens</i>	Proteobacteria
<i>P. putida</i>	Proteobacteria
<i>E. fergusonii</i>	Proteobacteria
<i>S. enterica</i>	Proteobacteria
<i>K. oxytoca</i>	Proteobacteria
<i>C. glutamicum</i>	Actinobacteria
<i>S. venezuelae</i>	Actinobacteria

Cell-free reaction:

Types of cell-free extract




Melinek B, et al. *BioProcess International* (2020)

	Soluble proteins	Membrane proteins	Glycoproteins
Archae / Protozoa	✓		
<i>E.coli</i>	✓		
Wheat Germ	✓		
Yeast	✓		✓ Addition of microsomes
Rabbit Reticulocyte	✓		✓ Addition of exogenous microsomes necessary
Tobacco	✓	✓	✓
Insect	✓	✓	✓
CHO / Human	✓	✓	✓

Anne Zemella et al. *ChemBioChem* 2015


Cell-free reaction:

Alternative cell-free approach, the pure system



Melinek B, et al. *BioProcess International* (2020)

36 PURE protein components produced using the OnePot method + tetra-(His)₆-tagged 70S ribosomes purified using affinity chromatography




Lavickova, Barbora, et al. *ACS synthetic biology* (2019)

Cell-free reaction:

Advantages of producing protein outside of the cell

- Fast reaction to produce protein
- No need for transformation
- No GMO
- No interaction with a living host
(No mutation, No burden)
- Decoupling effect
(e.g. Metabolic burden vs production burden)
- Low volume of reaction
- Higher level of control: Amount of DNA, Composition of the reaction.
- Work with Linear DNA



Cell-free Quiz



Current Applications of Cell-Free Systems in Academia and Industry

Fundamental science: cell-free to understand living cells

Example 1: Exploring Resource Competition through Cell-Free Systems

Applied Science: Bridging Theory to Practice



Example 2: Cell-Free Biosensor for food industry and medicine

Industrial Applications: Transforming Knowledge into Innovation

Example 1 :

Exploring Resource Competition through Cell-Free Systems


- Resource allocation *in vivo*
Production burden



Example 1 :

Exploring Resource Competition through Cell-Free Systems




- Resource allocation *in vitro*
Production burden



Example 1 :

Exploring Resource Competition through Cell-Free Systems




- Resource allocation *in vitro*
Tuning the cell-free reaction



Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource allocation *in vitro*
Different levels of protein production




Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource allocation *in vitro*

Different types of protein production







Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource allocation *in vitro*

Burden in different growth conditions





Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource competition *in vitro*


Measure with a metabolic pathway



Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource competition *in vitro*
Measure with a metabolic pathway




Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource competition *in vitro*


Measure with an inactivated metabolic pathway



Example 1 :

Exploring Resource Competition through Cell-Free Systems


- Resource competition *in vitro*
Measure with a metabolic pathway



Example 1 :

Exploring Resource Competition through Cell-Free Systems

- Resource allocation
 - Decouple production and metabolic burden



Fundamental science: cell-free to understand living cells

Example 1: Exploring Resource Competition through Cell-Free Systems


Applied Science: Bridging Theory to Practice

Example 2: Cell-Free Biosensor for food industry and medicine

Industrial Applications: Transforming Knowledge into Innovation


Example 1:

Cell-Free Biosensor for food industry and medicine



Example 1:


Cell-Free Biosensor for food industry and medicine



Example 1:

Cell-Free Biosensor for food industry and medicine


- Properties of a biosensor



Example 1:

Cell-Free Biosensor for food industry and medicine

- Properties of the transducer





Cell-free reaction:



Applied Science: Bridging Theory to Practice

- Efficiency of the biosensor

a




b



Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Efficiency of the biosensor



Do you have an idea to use
cell-free?

Fundamental science: cell-free to understand living cells

Applied Science: Bridging Theory to Practice

Industrial Applications: Transforming Knowledge into Innovation

Cell-free reaction:

Current uses of cell-free system

Fundamental science: cell-free to understand living cells

- Central functions
 - DNA replication
 - Transcription and translation process
 - Metabolism and energy production
- Resource allocation *in vitro*
- Construction hybrid systems
 - Multi-species systems
 - Nonstandard (unnatural) amino acids
 - Eukaryote properties (Glycolysation)
- Constructing artificial cells



Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

DNA replication

DNA replication system reconstructed by expressing 13 central genes encoding initiator (*dnaA*), DNA helicase (*dnaB*), helicase loader (*dnaC*), RNA primase (*dnaG*) and DNA polymerase III.



Omitting *dnaC* (one of the 13 genes) is the most important effect.


Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

DNA replication

DNA replication system reconstructed by expressing 1 gene of the bacteriophage phi 29 DNA replication .



Use for replication of large DNA genomes




Libicher, K, nat comms (2020)

Cell-free reaction:



Fundamental science: cell-free to understand living cells

- Central functions

Transcription process: T7 polymerase



Gene circuit performance characterization



Cell-free reaction:

Fundamental science: cell-free to understand living cells



- Central functions

Transcription process: modified promoter to recruits more T7 polymerase

Control environment: Fixed purified T7 RNA polymerase concentration, Fixed plasmid concentration, Fixed lysate and buffer concentration (ribosome, NTPs, Amino Acides etc..)

One changing elements = promoter sequence

Specific effects measured = Protein production, RNA production and DNA template degradation




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions


Transcription



Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions
- Transcription



σ -factor	Function	Group
σ^D/σ^{70}	Housekeeping genes	1
σ^S/σ^{38}	Stationary phase	2
σ^H/σ^{32}	Heat shock response	3
σ^F/σ^{28}	Motility genes	3
σ^E/σ^{24}	Extracytoplasmic function	4
$\sigma^{fecI}/\sigma^{19}$	Extracytoplasmic function	4
σ^N/σ^{54}	Nitrogen metabolism	

Cell-free reaction:


Fundamental science: cell-free to understand living cells

- Central functions

Translation: Optimizing translation process

Component	Concentration			
Mg-glutamate (mM)	0.4	1.2	2	4
K-glutamate (mM)	8	24	40	80
Amino Acid (mM)	0.15	0.45	0.75	1.5
tRNA (mg/ml)	0.02	0.06	0.1	0.2
CoA (mM)	0.026	0.078	0.13	0.26
NAD (mM)	0.033	0.099	0.165	0.33
cAMP (mM)	0.075	0.225	0.375	0.75
Folinic Acid (mM)	0.0068	0.0204	0.034	0.068
Spermidine (mM)	0.1	0.3	0.5	1
3-PGA (mM)	3	9	15	30
NTP (mM)	0.15	0.45	0.75	1.5

Combinatorial space = $4^{11} = 4\,194\,304$ compositions




- Can we improve protein production for a specific lysate without increasing the price of cell-free reaction?
- Can we provide efficient predictions of protein production?
- Can we highlight the critical parameters involved in protein production?

Cell-free reaction:


Fundamental science: cell-free to understand living cells

- Central functions


Translation: Optimizing translation process



Liquid Handling Robot



Screening




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

Translation: Optimizing translation process




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

Translation: Optimizing translation process



⇒ Components that improve yield at high concentration: Mg-glutamate, K-glutamate, Amino Acids and NTPs



Components that improve yield at low concentration: Spermidine, 3-PGA

Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

Translation: Optimizing translation process



Transcription vs translation:


Our data suggests that we mainly improved translation via composition optimization

Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

Translation: engineering translation machinery





Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

Metabolism and energy production




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Central functions

Metabolism and energy production

Study of composition in metabolites in different cell-free process





-> improve cell-free efficiency, supplemented with proteins to observe the impact on each part of the metabolism

Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Construction hybrid systems
- Multi-species systems




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Construction hybrid systems


Nonstandard (unnatural) amino acids, more orthogonality for a better predictability



Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Construction hybrid systems
- Nonstandard (unnatural) amino acids




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Construction hybrid systems

Multi-species systems




Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Construction hybrid systems


Eukaryote properties (Glycolysation)



Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Constructing artificial cells





Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Constructing artificial cells


Liposomes



Cell-free reaction:

Fundamental science: cell-free to understand living cells


- Constructing artificial cells: Communication / exchange with the environment



Cell-free reaction:

Fundamental science: cell-free to understand living cells



- Constructing artificial cells : compartmentation



Cell-free reaction:

Fundamental science: cell-free to understand living cells

- Constructing artificial cells : compartmentation



Cell-free reaction:

Current uses of cell-free system

Applied Science: Bridging Theory to Practice


- Prototyping genetic circuits and pathways
 - Genetic parts
 - Metabolic engineering
- Developing biosensors
 - Optimize cell-free biosensor
 - Biocomputing
 - Materials and cell-free
 - Medical diagnostics
 - Pollution diagnostics
- Bio-machine assembly
 - Phages selection
 - Antibodies

Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Prototyping genetic circuits and pathways



Genetic parts



Cell-free reaction:

Applied Science: Bridging Theory to Practice


- Prototyping genetic circuits and pathways
- Genetic parts



Cell-free reaction:

Applied Science: Bridging Theory to Practice


- Prototyping genetic circuits and pathways
Genetic parts: regulated systems



Cell-free reaction:

Applied Science: Bridging Theory to Practice



- Prototyping genetic circuits and pathways
Genetic parts: Linear fragment vs plasmid



Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Prototyping genetic circuits and pathways
- Genetic parts: Linear fragment vs plasmid




Cell-free reaction:

Applied Science: Bridging Theory to Practice

Prototyping genetic circuits and pathways

Metabolic engineering: combinatorial exploration




Example of : 6⁴ combinations

Cell-free reaction:

Applied Science: Bridging Theory to Practice

Metabolic engineering: combinatorial exploration

6⁴ plasmids






Cell-free reaction:

Applied Science: Bridging Theory to Practice

Metabolic engineering: combinatorial exploration

$6 \times 4 = 24$ plasmids





Cell-free reaction:

Applied Science: Bridging Theory to Practice

Prototyping genetic circuits and pathways

Metabolic engineering: combinatorial exploration




Cell-free reaction:

Applied Science: Bridging Theory to Practice


Developing biosensors
Biocomputing



Perceptron weights (w_i) are learned *in silico* to increase classifier accuracy



Binary classification:




Cell-free reaction:

Applied Science: Bridging Theory to Practice


Developing biosensors

Biocomputing: logic gates

AND		
I_1	I_2	out
0	0	0
0	1	0
1	0	0
1	1	1



OR		
I_1	I_2	out
0	0	0
0	1	1
1	0	1
1	1	1










Cell-free reaction:

Applied Science: Bridging Theory to Practice

Developing biosensors

Biocomputing: 4 biosensors






Cell-free reaction:

Applied Science: Bridging Theory to Practice

Developing biosensors

Biocomputing: tuning the weights






Cell-free reaction:

Applied Science: Bridging Theory to Practice

Developing biosensors


Biocomputing: perceptron



Cell-free reaction:

Applied Science: Bridging Theory to Practice


- Developing biosensors
Medical diagnostics



Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Developing biosensors
- Medical diagnostics



Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Developing biosensors
Materials and cell-free

Johnson & Johnson's QuickFire Challenge
Johnson & Johnson was Live • Follow

BioFabrics
Synthetic Biology + Textiles

Biosensor for Zika Virus on Thread

Thread Alone - Zika + Zika

Biosensor Design
A U G C

Cell-Free Reaction
Freeze Drying

Synthetic Biology Biosensor
INPUT → Environment-responsive promoter → Gene → Gene → Gene → OUTPUT
• Engineered gene circuits
• Transcriptional changes

Response to Target

Clean
Dangerous Splash Detected

Wearable Diagnostics?
Animal Sentinels
Engineered Biosensors
Paper-based Diagnostics

Integration with Internet of Things
#ChampionsOfScience


WYSS 3607 HD BROAD INSTITUTE

Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Developing biosensors



Materials and cell-free



Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Developing biosensors
- Pollution detection









Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Developing biosensors


Pollution detection



Cell-free reaction:

Applied Science: Bridging Theory to Practice


- Bio-machine assembly
- Antibodies (eukaryotic cell-free)



Cell-free reaction:

Applied Science: Bridging Theory to Practice


- Bio-machine assembly
- Phages selection



Cell-free reaction:

Applied Science: Bridging Theory to Practice

- Bio-machine assembly
- Phages selection



Encoded by a 40 kbp DNA program composed of about 60 genes

Fundamental science: cell-free to understand living cells

Applied Science: Bridging Theory to Practice

Industrial Applications: Transforming Knowledge into Innovation

Cell-free reaction:

Industrial Applications: Transforming Knowledge into Innovation

Pharmaceuticals

- Monoclonal Antibodies & Antimicrobial Peptides
 - LenioBio (Germany)
 - Cube Biotech (Germany)
 - BioCat (Germany)
- Vaccines
 - Ntxpress (New Mexico, USA)
 - CPI (UK)
 - Sutro Biopharma (California, USA)
- Small Molecules
 - Creativebiomart (USA, Europe, Canada)
- Membrane Proteins
 - Creativebiomart (USA, Europe, Canada)
 - Desgin Pharma (Massachussets, USA)
 - Synthelis (France)

Pollution

- CO2 capture
 - Ensovi (Massachussets, USA)
- Biosensor
 - Stemloop (Illinois, USA)

On-Demand


- Purified proteins
 - Tierra bioscience (California, USA)
 - CFS (Japan)
 - Cell-free (Ontario, Canada)

Cell-free kit providers


- New England Biolabs (Massachussets, USA)
- Thermo Fisher (Massachussets, USA)
- Biotechrabbit (Germany)
- arbor bioscience (Michigan, USA)

Scale-up

a




Approximate batch size	V = 1 µl	V = 1 ml	V = 100 l	V = 1,000 l (continuous operation)
Example product	Laboratory use T7 RNA polymerase	Point-of-use vaccine dose	Pharmaceutical antibody-drug conjugate	Commodity grade subtilisin
Commercial price (\$/mg)	\$13,250	\$200	\$5,500	\$0.022
Estimated cell-free yield (mg/ml reaction)	1.0	0.02	0.250	2.0
Estimated purification cost	0x	0x	10x	1x
Estimated breakeven cost (\$/ml reaction)	\$13,250	\$4	\$120	\$0.022



Cell-free reaction:

Additional use / challenges with cell-free


Cell-free for education



myTXTL® Cell-Free Protein Expression kits for Igem

National Institute of Standards and Technology

Towards reproducible cell-free systems



Acknowledgements



Matthieu Jules

Léa Wagner

Olivier Delumeau

Vincent Sauveplane

TOM ELLIS LAB

Tom Ellis

Guy-Bart Stan Lab

Imperial College
London

Brooke Rothschild-Mancinelli

Carlos Bricio

Francesca Ceroni

Michela Murgiano



Bio-RetroSynth

Jean-Loup Faulon

Institut Pasteur

Fabrice Agou

Mathilde Koch

Amir Pandi

Agnès Zettor

