



HAL
open science

Ramsey theory without pigeonhole principle and the adversarial Ramsey principle

Noé de Rancourt

► **To cite this version:**

Noé de Rancourt. Ramsey theory without pigeonhole principle and the adversarial Ramsey principle. Transactions of the American Mathematical Society, 2020, 373 (7), pp.5025-5056. 10.1090/tran/8063 . hal-04406796

HAL Id: hal-04406796

<https://hal.science/hal-04406796>

Submitted on 19 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Ramsey theory without pigeonhole principle and the adversarial Ramsey principle

N. de Rancourt

Abstract

We develop a general framework for infinite-dimensional Ramsey theory with and without pigeonhole principle, inspired by Gowers' Ramsey-type theorem for block sequences in Banach spaces and by its exact version proved by Rosendal. In this framework, we prove the adversarial Ramsey principle for Borel sets, a result conjectured by Rosendal that generalizes at the same time his version of Gowers' theorem and Borel determinacy of games on integers.

1 Introduction

This paper has two main goals. The first is to develop an abstract formalism for infinite-dimensional Ramsey theory, enabling to prove both Ramsey results with a pigeonhole principle (like Mathias–Silver's theorem [14, 20]) and Ramsey results without a pigeonhole principle, like Gowers' Ramsey-type theorem for block sequences in Banach spaces [7] and its exact version given by Rosendal [18]. The second goal is to prove the *adversarial Ramsey principle* for Borel sets, a result conjectured by Rosendal [19], unifying his exact version of Gowers' theorem with Borel determinacy of games on integers. In order to motivate the rest of the paper and to recall the statement of the latter results and conjectures, let us begin with some history.

Infinite-dimensional Ramsey theory is a branch of Ramsey theory where we color *infinite-dimensional* objects, that are, *sequences of points* of some space, and where we want to find homogeneous subspaces. The fundamental result in infinite-dimensional Ramsey theory is Mathias–Silver's theorem, proved independently in 1968 by Mathias [14] and in 1970 by Silver [20], saying that if \mathcal{X} is an analytic subset of $[\omega]^\omega$ (the set of all infinite subsets of ω , endowed with the topology inherited from the Cantor space $\mathcal{P}(\omega) = 2^\omega$ with the product topology) then for every infinite $M \subseteq \omega$, there exists an infinite $N \subseteq M$ (called a *homogeneous* set) such that either for every infinite $S \subseteq N$, we have $S \in \mathcal{X}$, or for every infinite $S \subseteq N$, we have $S \in \mathcal{X}^c$. (A set $\mathcal{X} \subseteq [\omega]^\omega$ satisfying

Mathematics Subject Classification (2010): 05D10, 03E60

Keywords: Infinite-dimensional Ramsey theory, Gowers' games, Determinacy

the conclusion of this theorem will be called a *Ramsey set*.) An important remark is that in this theorem (and in particular, in all of its proofs using the classical technique of combinatorial forcing, see for example [21], Chapter 1, for a presentation of this technique), the sets M and N are often seen as *subspaces* (elements of a poset) while the set S is rather seen as an *infinite sequence*, the increasing sequence of its elements; this distinction between subspaces and sequences of points will appear in all results presented in this paper.

The proof of Mathias–Silver’s theorem uses in an essential way the *pigeonhole principle*, i.e. the trivial fact that for every infinite $M \subseteq \omega$ and every $A \subseteq \omega$, there exists an infinite $N \subseteq M$ such that either $N \subseteq A$, or $N \subseteq A^c$. In the decades that followed the proof of this theorem, several similar results arose in different contexts (words, trees, etc.). All of these results have the same form: we color infinite sequences of points satisfying some structural condition (being increasing, being block sequences, etc.) and the theorem ensures that we can find a monochromatic subspace. The proof of each of these results relies on an analogue of the pigeonhole principle, whose proof is in most cases way less trivial as in ω . In general, a *pigeonhole principle* is a one-dimensional Ramsey result, i.e. a result where we color points and we want to find a monochromatic subspace. A lot of these pigeonhole principles, and the infinite-dimensional Ramsey results they imply, can be found in Todorćević’s book [21], where a general framework to deduce an infinite dimensional Ramsey result from its associated pigeonhole principle is also developed.

The first infinite-dimensional Ramsey-type result that was not relying on a pigeonhole principle was proved by Gowers, in the 90’s. The aim of Gowers was to solve a celebrated problem asked by Banach, the homogeneous space problem, asking whether ℓ_2 was the only infinite-dimensional Banach space, up to isomorphism, that was isomorphic to all of its closed, infinite-dimensional subspaces. Gowers proved a dichotomy [7] that, combined with a result by Komorowski and Tomczak-Jaegermann [11], provided a positive answer to Banach’s question. The proof of this dichotomy relies on a Ramsey-type theorem in separable Banach spaces, that we will state now. The reader who is not familiar with Banach space geometry can skip this part, since it will only be relevant to understanding sections 4 and 5 of this paper.

In this paper, to save writing, we will only consider real Banach spaces, but the results we present here adapt to the complex case. Let E be a Banach space. Recall that a (*Schauder*) *basis* of E is a sequence $(e_i)_{i \in \omega}$ such that every $x \in E$ can be written in a unique way as an infinite sum $\sum_{i=0}^{\infty} x^i e_i$, where $x^i \in \mathbb{R}$. In this case, the *support* of the vector x , denoted by $\text{supp}(x)$, is defined as the set $\{i \in \omega \mid x^i \neq 0\}$. A *block sequence* of (e_i) is an infinite sequence $(x_n)_{n \in \omega}$ of nonzero vectors of E with $\text{supp}(x_0) < \text{supp}(x_1) < \dots$ (here, for two nonempty sets of integers A and B , the notation $A < B$ means that $\forall i \in A \forall j \in B \ i < j$). It can be shown that a block sequence is a basis of the closed subspace it spans, such a space being called a *block subspace*. A basis, or a block sequence, is said to be *normalized* if all of its terms have norm 1. In the rest of this article, unless otherwise specified, every basis and every block sequence in a Banach space will be normalized.

For X a block subspace of E , let $[X]$ denote the set of block sequences all of whose terms are in X . We can equip $[E]$ with a natural topology by seeing it as a subspace of $(S_E)^\omega$ with the product topology (where S_E denotes the unit sphere of E with the norm topology), which makes it a Polish space. For $\mathcal{X} \subseteq [E]$ and $\Delta = (\Delta_n)_{n \in \omega}$ a sequence of positive real numbers, let $(\mathcal{X})_\Delta$ denote the Δ -expansion of \mathcal{X} , that is, the set $\{(x_n)_{n \in \omega} \in [E] \mid \exists (y_n)_{n \in \omega} \in \mathcal{X} \forall n \in \omega \|x_n - y_n\| \leq \Delta_n\}$. In order to state Gowers' theorem, we need a last definition.

Definition 1.1. Let X be a block subspace of E . *Gowers' game* below X , denoted by G_X , is the following infinite two-players game (whose players will be denoted by **I** and **II**):

$$\begin{array}{ccccccc} \mathbf{I} & Y_0 & & Y_1 & & \dots & \\ \mathbf{II} & & y_0 & & y_1 & & \dots \end{array}$$

where the Y_i 's are block subspaces of X , and the y_i 's are normalized vectors of E with finite support, with the constraints for **II** that for all $i \in \omega$, $y_i \in Y_i$ and $\text{supp}(y_i) < \text{supp}(y_{i+1})$. The outcome of the game is the sequence $(y_i)_{i \in \omega} \in [E]$.

In this paper, when dealing with games, we shall use a convention introduced by Rosendal: we omit to set a winning condition when defining a game, but rather associate an *outcome* to the game and say that a player has a strategy to force this outcome to belong to some fixed set. For example, saying that player **II** has a strategy to reach a set $\mathcal{X} \subseteq [E]$ in the game G_X means that she has a winning strategy in the game whose rules are those of G_X and whose winning condition is the fact that the outcome belongs to \mathcal{X} .

We can now state Gowers' theorem:

Theorem 1.2 (Gowers' Ramsey-type theorem). *Let $\mathcal{X} \subseteq [E]$ be an analytic set, $X \subseteq E$ be a block subspace, and Δ be an infinite sequence of positive real numbers. Then there exists a block subspace Y of X such that either $[Y] \subseteq \mathcal{X}^c$, or player **II** has a strategy in G_Y to reach $(\mathcal{X})_\Delta$.*

While one of the possible conclusions of this theorem, $[X] \subseteq \mathcal{X}^c$, is very similar to "For every infinite $S \subseteq M$, we have $S \in \mathcal{X}^c$ " in Mathias–Silver's theorem, the other one is much weaker, for two reasons: the use of metrical approximation and the use of a game. As we will see later, the necessity of the approximation is due to a lack of finiteness, while the necessity for one of the possible conclusions to involve a game matters much more and is due to the lack of a pigeonhole principle in this context. In some Banach spaces, a pigeonhole principle holds, and in these spaces, Gowers gave a strengthening of his theorem, involving no game, that we will introduce now. We start by stating the general form of the relevant pigeonhole principle in the context of Banach spaces; since an exact pigeonhole principle is never satisfied in this context, and would anyways be useless since approximation is needed for other reasons, we will only state an approximate pigeonhole principle. For a Banach space E , a set $A \subseteq S_E$, and $\delta > 0$, denote by $(A)_\delta$ the δ -expansion of A , that is, the set $\{x \in S_E \mid \exists y \in A \|x - y\| \leq \delta\}$.

Definition 1.3. Say that a Banach space E with a Schauder basis satisfies the *approximate pigeonhole principle* if for every $A \subseteq S_E$, for every block subspace $X \subseteq E$, and for every $\delta > 0$, there exists a block subspace $Y \subseteq X$ such that either $S_Y \subseteq A^c$, or $S_Y \subseteq (A)_\delta$.

Recall that an infinite-dimensional Banach space E is said to be c_0 -saturated if c_0 can be embedded in all of its infinite-dimensional, closed subspaces. A combination of results by Milman [15], Gowers [6], and Odell and Schlumprecht [16], shows the following:

Theorem 1.4. *A space E with a Schauder basis satisfies the approximate pigeonhole principle if and only if it is c_0 -saturated.*

Thus, in c_0 -saturated spaces, we have a strengthening of Gowers' theorem:

Theorem 1.5 (Gowers' Ramsey-type theorem for c_0). *Suppose that E is c_0 -saturated. Let $\mathcal{X} \subseteq [E]$ be an analytic set, $X \subseteq E$ be a block subspace, and Δ be an infinite sequence of positive real numbers. Then there exists a block subspace Y of X such that either $[Y] \subseteq \mathcal{X}^c$, or $[Y] \subseteq (\mathcal{X})_\Delta$.*

For a complete survey of Gowers' Ramsey-type theory in Banach spaces, see [1], Part B, Chapter IV.

In 2010, in [18], Rosendal proved an exact version (without approximation) of Gowers' theorem, in countable vector spaces, which easily implies Gowers' theorem in Banach spaces. In this theorem, in order to remove the approximation, the non-game-theoretical conclusion has to be weakened by introducing a new game, the *asymptotic game*. We present here Rosendal's theorem in more details. Let E be a countably infinite-dimensional vector space over an at most countable field K and $(e_i)_{i \in \omega}$ be a basis (in the algebraic sense) of E . The notions of *support* of a vector and of a *block sequence* relative to this basis are defined exactly in the same way as in a Banach space with a Schauder basis (this time, since there is no norm, there is obviously no notion of normalized bases or block sequences). A *block subspace* is a subspace of E spanned by a block sequence. Observe that in this setting, since every vector has finite support, every infinite-dimensional subspace of E has a further subspace that is a block subspace. Given a block subspace $X \subseteq E$, define the two following games:

Definition 1.6.

1. *Gowers' game* below X , denoted by G_X , is defined in the following way:

$$\begin{array}{llll} \mathbf{I} & Y_0 & Y_1 & \dots \\ \mathbf{II} & y_0 & y_1 & \dots \end{array}$$

where the Y_i 's are block subspaces of X , and the y_i 's are nonzero elements of E , with the constraint for **II** that for all $i \in \omega$, $y_i \in Y_i$. The outcome of the game is the sequence $(y_i)_{i \in \omega} \in E^\omega$.

2. The *asymptotic game* below X , denoted by F_X , is defined in the same way as G_X , except that this time, the Y_i 's are moreover required to have finite codimension in X .

We endow E with the discrete topology and E^ω with the product topology; since E is countable, E^ω is a Polish space. Rosendal's theorem is the following:

Theorem 1.7 (Rosendal). *Let \mathcal{X} be an analytic subset of E^ω . Then for every block subspace $X \subseteq E$, there exists a block subspace $Y \subseteq X$ such that either **I** has a strategy in F_Y to reach \mathcal{X}^c , or **II** has a strategy in G_Y to reach \mathcal{X} .*

Say that a set $\mathcal{X} \subseteq E^\omega$ is *strategically Ramsey* if it satisfies the conclusion of this theorem. Here, the use of an asymptotic game in one side of the alternative is (as we will see in the rest of this paper) not much weaker than a non-game-theoretical conclusion as in Mathias–Silver's theorem.

In the same paper as the last theorem, Rosendal, inspired by the work of Pelczar [17], and by a common work with Ferenczi [4], introduced a new Ramsey principle which is, unlike Theorem 1.7, symmetrical. His result was then refined in [19]. It involves two games, known as the *adversarial Gowers' games*, obtained by mixing the games G_X and F_X .

Definition 1.8.

1. For a block subspace $X \subseteq E$, the game A_X is defined in the following way:

$$\begin{array}{ccccccc} \mathbf{I} & & x_0, Y_0 & & x_1, Y_1 & & \dots \\ \mathbf{II} & X_0 & & y_0, X_1 & & y_1, X_2 & \dots \end{array}$$

where the x_i 's and the y_i 's are nonzero vectors of X , the X_i 's are block subspaces of X , and the Y_i 's are block subspaces of X with finite codimension. The rules are the following:

- for **I**: for all $i \in \omega$, $x_i \in X_i$;
- for **II**: for all $i \in \omega$, $y_i \in Y_i$;

and the outcome of the game is the sequence $(x_0, y_0, x_1, y_1, \dots) \in E^\omega$.

2. The game B_X is defined in the same way as A_X , except that this time the X_i 's are required to have finite codimension in X , whereas the Y_i 's can be arbitrary block subspaces of X .

The result Rosendal proves in [19] is the following:

Theorem 1.9 (Rosendal). *Let $\mathcal{X} \subseteq E^\omega$ be Σ_3^0 or Π_3^0 . Then for every block subspace $X \subseteq E$, there exists a block subspace $Y \subseteq X$ such that either **I** has a strategy in A_Y to reach \mathcal{X} , or **II** has a strategy in B_Y to reach \mathcal{X}^c .*

Say that a set $\mathcal{X} \subseteq E^\omega$ is *adversarially Ramsey* if it satisfies the conclusion of this theorem. Then, a natural question to ask is for which complexity of the set \mathcal{X} one can ensure that it is adversarially Ramsey.

There are two things to observe. First, let $\mathcal{X} \subseteq E^\omega$ and define $\mathcal{X}' = \{(x_i)_{i \in \omega} \in E^\omega \mid (x_{2i})_{i \in \omega} \in \mathcal{X}\}$. Then by forgetting the contribution of player **II** to the outcome of the adversarial Gowers' games and switching the roles of players **I** and **II**, we see that \mathcal{X} is strategically Ramsey if and only if \mathcal{X}' is adversarially Ramsey. So, for a suitable class Γ of subsets of Polish spaces, saying that all Γ -subsets of E^ω are adversarially Ramsey is stronger than saying that all Γ -subsets of E^ω are strategically Ramsey. The second remark is that, if the field K is infinite, then the adversarial Ramsey property for Γ -subsets of E^ω also implies that all Γ -subsets of ω^ω are determined. To see this, observe that when playing vectors in A_X or B_X , no matter the constraint imposed by the other player, players **I** and **II** have total liberty for choosing the first non-zero coordinate of the vectors they play. Therefore, by making \mathcal{X} only depend on the first nonzero coordinate of each vector played, we recover a classical Gale-Stewart game in $(K^*)^\omega$. For this reason, there is no hope, in *ZFC*, to prove the adversarial Ramsey property for a class larger than Borel sets. Then, Rosendal asks the following questions in [19]:

Question 1.10 (Rosendal). *Is every Borel set adversarially Ramsey?*

Question 1.11 (Rosendal). *In the presence of large cardinals, is every analytic set adversarially Ramsey?*

This paper is organised as follows. In section 2, we define an abstract setting for infinite-dimensional Ramsey theory, called the setting of *Gowers spaces*, where the asymptotic game, Gowers' game, and the adversarial Gowers' games can be defined. In this general setting, we prove that the answer to Question 1.10 is yes, and that the answer to Question 1.11 is yes in the presence of a measurable cardinal. In section 3, we prove a general version of Rosendal's Theorem 1.7 in Gowers spaces; then, we introduce the pigeonhole principle in these spaces and we show that if this principle holds, then we can refine the abstract Rosendal's theorem to a result with an asymptotic game in both sides, which Mathias-Silver's theorem is an easy consequence. In section 4, we give approximate versions of the results of sections 2 and 3 enabling to work in Polish metric spaces instead of countable spaces. Finally, in section 5, we develop a general framework to deduce a genuine, non game-theoretical Ramsey conclusion (of the form "in some subspace, every sequence satisfying some structural property is in \mathcal{X} ") from a conclusion involving a strategy of player **I** in the asymptotic game. The results of this section, combined with these of section 4, will for instance have Gowers' Theorems 1.2 and 1.5 as an immediate consequence.

2 Gowers spaces and the adversarial Ramsey property

In this section, we will introduce the notion of a *Gowers space*, which will be our abstract setting for infinite-dimensional Ramsey theory; then, we will prove in this setting

the adversarial Ramsey principle, our most general Ramsey result without pigeonhole principle, which will give positive answers to Questions 1.10 and 1.11.

Inspired by the examples given in the introduction, we define a formalism with two notions, a notion of *subspaces* and a notion of *points*. The idea is that we will color infinite sequences of points and try to find subspaces such that lots of sequences of points in this subspace share the same color.

For X a set, denote by $\text{Seq}(X) = X^{<\omega} \setminus \{\emptyset\}$ the set of all nonempty finite sequences of elements of X . For $s, t \in X^{<\omega}$, denote by $|s|$ the length of s (i.e. the integer n such that $s \in X^n$), by $s \subseteq t$ the fact that s is an initial segment of t , and by $s \hat{\ } t$ the concatenation of s and t . If $x \in X$, the concatenation of s and of the sequence (x) whose unique term is x will be abusively denoted by $s \hat{\ } x$.

Definition 2.1. A *Gowers space* is a quintuple $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$, where P is a nonempty set (the set of *subspaces*), X is an at most countable nonempty set (the set of *points*), \leq and \leq^* are two quasiorders on P (i.e. reflexive and transitive binary relations), and $\triangleleft \subseteq \text{Seq}(X) \times P$ is a binary relation, satisfying the following properties:

1. for every $p, q \in P$, if $p \leq q$, then $p \leq^* q$;
2. for every $p, q \in P$, if $p \leq^* q$, then there exists $r \in P$ such that $r \leq p$, $r \leq q$ and $p \leq^* r$;
3. for every \leq -decreasing sequence $(p_i)_{i \in \omega}$ of elements of P , there exists $p^* \in P$ such that for all $i \in \omega$, we have $p^* \leq^* p_i$;
4. for every $p \in P$ and $s \in X^{<\omega}$, there exists $x \in X$ such that $s \hat{\ } x \triangleleft p$;
5. for every $s \in \text{Seq}(X)$ and every $p, q \in P$, if $s \triangleleft p$ and $p \leq q$, then $s \triangleleft q$.

Say that $p, q \in P$ are *compatible* if there exists $r \in P$ such that $r \leq p$ and $r \leq q$. To save writing, we will often write $p \lesssim q$ when $p \leq q$ and $q \leq^* p$. Observe that by 2., the p^* in 3. can be chosen in such a way that $p^* \leq p_0$.

In most usual cases, the fact that $s \triangleleft p$ will only depend on p and on the last term of s ; the spaces satisfying this property will be called *forgetful Gowers spaces*. In these spaces, we will allow ourselves to view \triangleleft as a binary relation on $X \times P$. However, for some applications (see, for example, the proof of Theorem 4.6), it is sometimes useful to make the fact that $s \triangleleft p$ also depend on the length of the sequence s , and we could imagine that even all the terms of s would be relevant in some contexts.

When thinking about a Gowers space, we should have the two following examples in mind:

- The *Mathias–Silver space* $\mathcal{N} = ([\omega]^\omega, \omega, \subseteq, \subseteq^*, \triangleleft)$, where $[\omega]^\omega$ is the set of all infinite sets of integers, $M \subseteq^* N$ iff $M \setminus N$ is finite and $(x_0, \dots, x_n) \triangleleft M$ iff $x_n \in M$. Here, we have that $M \lesssim N$ iff M is a cofinite subset of N , and M and N are compatible iff $M \cap N$ is infinite.

- The *Rosendal space* over an at most countable field K , $\mathcal{R}_K = (P, E \setminus \{0\}, \subseteq, \subseteq^*, \triangleleft)$, where E is a countably infinite-dimensional K -vector space with a basis $(e_i)_{i \in \omega}$, P is the set of all block subspaces of E relative to this basis, $X \subseteq^* Y$ iff Y contains some finite-codimensional block subspace of X , and $(x_0, \dots, x_n) \triangleleft X$ iff $x_n \in X$. Here, we have that $X \lesssim Y$ iff X is a finite-codimensional subspace of Y , and X and Y are compatible iff $X \cap Y$ is infinite-dimensional.

Observe that both of these spaces are forgetful, so we could have defined \triangleleft as a relation between points and subspaces (and that is what we will do, in such cases, in the rest of this paper); in this way, in both cases, \triangleleft is the membership relation. It is easy to verify that, for these examples, the axioms 1., 2., 4., and 5. are satisfied; we briefly explain how to prove 3.. For the Mathias–Silver space, if $(M_i)_{i \in \omega}$ is a \subseteq -decreasing sequence of infinite subsets of ω , then we can, for each $i \in \omega$, choose $n_i \in M_i$ in such a way that the sequence $(n_i)_{i \in \omega}$ is increasing, and let $M^* = \{n_i \mid i \in \omega\}$. Then the set M^* is as wanted. For the Rosendal space, the idea is the same: given $(F_i)_{i \in \omega}$ a decreasing sequence of block subspaces of E , we can pick, for each i , a nonzero vector $x_i \in F_i$, in such a way that for $i \geq 1$, we have $\text{supp}(x_{i-1}) < \text{supp}(x_i)$. In this way, $(x_i)_{i \in \omega}$ is a block sequence, and the block subspace F^* spanned by this sequence is as wanted.

Also observe that in the definition of the Rosendal space, choosing $E \setminus \{0\}$ and not E for the set of points is totally arbitrary, and here, we only made this choice in order to use the same convention as Rosendal in his papers [18, 19]; but the results we will show apply as well when the set of points is E . Also, we could have taken for P the set of all infinite-dimensional subspaces of E (where, here, the relation \subseteq^* is defined by $X \subseteq^* Y$ iff $X \cap Y$ has finite codimension in X) instead of only block subspaces. However, the abstract results we will prove are slightly stronger in the case when we consider only block subspaces; this is due to the fact that, while every infinite-dimensional subspace of E contains a block subspace, there are finite-codimensional subspaces that do not contain any finite-codimensional block subspace.

In the rest of this section, we fix a Gowers space $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$. For $p \in P$, define the *adversarial Gowers' games below* p as follows:

Definition 2.2.

1. The game A_p is defined in the following way:

I	x_0, q_0	x_1, q_1	\dots
II	p_0	y_0, p_1	$y_1, p_2 \quad \dots$

where the x_i 's and the y_i 's are elements of X , and the p_i 's and the q_i 's are elements of P . The rules are the following:

- for **I**: for all $i \in \omega$, $(x_0, y_0, \dots, x_{i-1}, y_{i-1}, x_i) \triangleleft p_i$ and $q_i \lesssim p_i$;
- for **II**: for all $i \in \omega$, $(x_0, y_0, \dots, x_i, y_i) \triangleleft q_i$ and $p_i \leq p$.

The outcome of the game is the sequence $(x_0, y_0, x_1, y_1, \dots) \in X^\omega$.

2. The game B_p is defined in the same way as A_p , except that this time we require $p_i \lesssim p$, whereas we only require $q_i \leq p$.

As in the particular case of vector spaces, we can define the *adversarial Ramsey property* for subsets of X^ω :

Definition 2.3. A set $\mathcal{X} \subseteq X^\omega$ is said to be *adversarially Ramsey* if for every $p \in P$, there exists $q \leq p$ such that either player **I** has a strategy to reach \mathcal{X} in A_q , or player **II** has a strategy to reach \mathcal{X}^c in B_q .

Informally, the adversarial Ramsey property for \mathcal{X} means that up to taking a subspace, one of the players has a winning strategy in the game that is the most difficult for him. Observe that the property that **I** has a strategy in A_p to reach some set \mathcal{X} (resp. the property that **II** has a strategy in B_p to reach \mathcal{X}^c) is strongly hereditary in the sense that if **I** has a strategy to reach \mathcal{X} in A_p , then he also has one in $A_{p'}$ for every $p' \leq^* p$ (and the same holds for **II** in B_p). Indeed, we can simulate a play of $A_{p'}$ with a play of A_p : when, in $A_{p'}$, player **I**'s strategy tells him to play x_i and q_i , then in A_p he can play the same x_i and a q'_i such that $q'_i \lesssim p'$ and $q'_i \leq q_i$, in such a way that the next y_i played by **II** in $A_{p'}$ will be also playable in A_p (the existence of such a q'_i is guaranteed by condition 2. in the definition of a Gowers space). And when, in $A_{p'}$, player **II** plays y_i and p'_{i+1} , then in A_p , **I** can make her play the same y_i and a p_{i+1} such that $p_{i+1} \leq p$ and $p_{i+1} \leq p'_{i+1}$, in such a way that the next x_{i+1} played by **I** in A_p according to his strategy will also be playable in $A_{p'}$. In this way, the outcomes of both games are the same, and since **I** reaches \mathcal{X} in A_p , then he also does in $A_{p'}$.

On the other hand, it is clear that if **I** has a strategy to reach some set \mathcal{X} in A_p , then he also has one in B_p , so **II** cannot have a strategy to reach \mathcal{X}^c in B_p . Thus, the fact that \mathcal{X} has the adversarial Ramsey property gives a genuine dichotomy between two disjoint and strongly hereditary classes of subspaces.

We endow the set X with the discrete topology and the set X^ω with the product topology. The main result of this section is the following:

Theorem 2.4 (Adversarial Ramsey principle, abstract version). *Every Borel subset of X^ω is adversarially Ramsey.*

In the case of the Rosendal space, the adversarial Gowers games defined here are exactly the same as those defined in the introduction. Thus, Theorem 2.4 applied to this space provides a positive answer to Question 1.10.

Also observe that if $P = \{\mathbb{1}\}$ and if we have $s \triangleleft \mathbb{1}$ for every $s \in \text{Seq}(X)$, then both $A_{\mathbb{1}}$ and $B_{\mathbb{1}}$ are the classical Gale-Stewart game in X , so the adversarially Ramsey subsets of X^ω are exactly the determined ones. So in this space, Theorem 2.4 is nothing more than Borel determinacy for games on integers; hence, we get that Theorem 2.4 has at least the metamathematical strength of Borel determinacy for games on integers. Therefore, by the work of Friedman [5], any proof of Theorem 2.4 should make use of the powerset axiom and of the replacement scheme. We also get that it is not provable in *ZFC* that every analytic (or coanalytic) set in every Gowers space is adversarially

Ramsey. Actually, it turns out that there is a large class of Gowers spaces for which Borel determinacy can be recovered from the version of Theorem 2.4 in these spaces; this will be shown in a forthcoming paper [3].

We will deduce Theorem 2.4 from Borel determinacy for games on real numbers. For this purpose, we follow an approach used first by Kastanas in [9]: in this paper Kastanas deduced the Ramsey property for subsets of $[\omega]^\omega$ from the determinacy of a game (a similar game was already introduced by Mansfield in [12], although it was only used in the open case, in order to estimate the complexity of a homogeneous set). In what follows, we adapt Kastanas' game in order to get the adversarial Ramsey property.

Definition 2.5. For $p \in P$, *Kastanas' game* K_p below p is defined as follows:

$$\begin{array}{llll} \mathbf{I} & x_0, q_0 & x_1, q_1 & \dots \\ \mathbf{II} & p_0 & y_0, p_1 & y_1, p_2 \dots \end{array}$$

where the x_i 's and the y_i 's are elements of X , and the p_i 's and the q_i 's are elements of P . The rules are the following:

- for **I**: for all $i \in \omega$, $(x_0, y_0, \dots, x_{i-1}, y_{i-1}, x_i) \triangleleft p_i$ and $q_i \leq p_i$;
- for **II**: $p_0 \leq p$, and for all $i \in \omega$, $(x_0, y_0, \dots, x_i, y_i) \triangleleft q_i$ and $p_{i+1} \leq q_i$.

The outcome of the game is the sequence $(x_0, y_0, x_1, y_1, \dots) \in X^\omega$.

The exact result we will show is the following:

Proposition 2.6. *Let $p \in P$ and $\mathcal{X} \subseteq X^\omega$.*

1. *If **I** has a strategy to reach \mathcal{X} in K_p , then there exists $q \leq p$ such that **I** has a strategy to reach \mathcal{X} in A_q ;*
2. *If **II** has a strategy to reach \mathcal{X}^c in K_p , then there exists $q \leq p$ such that **II** has a strategy to reach \mathcal{X}^c in B_q .*

Once this proposition is proved, Theorem 2.4 will immediately follow from the Borel determinacy of Kastanas' game.

Since the proofs of 1. and 2. of Proposition 2.6 are exactly similar, we only prove 2.. In order to do this, let us introduce some notation. During the whole proof, we fix a strategy τ for **II** in K_p to reach \mathcal{X}^c . Following the terminology introduced by Ferenczi and Rosendal in [4], a partial play of K_p ending with a move of **II** and during which **II** always plays according to her strategy will be called a *state*. Say that a state \mathcal{J} *realises* a finite sequence $(x_0, y_0, x_1, y_1, \dots, x_{n-1}, y_{n-1})$ if \mathcal{J} has the form $(p_0, x_0, \dots, q_{n-1}, y_{n-1}, p_n)$; say that a state realising a sequence of length $2n$ *has rank* n . Define in the same way the notions of a *total state* (which is a total play of K_p) and of realisation for a total state; the restriction of a total state $\mathcal{J} = (p_0, x_0, q_0, y_0, p_1, \dots)$ to a state of rank n , denoted by $\mathcal{J} \upharpoonright_n$, is the state $(p_0, x_0, \dots, q_{n-1}, y_{n-1}, p_n)$. If an infinite sequence $(x_0, y_0, x_1, y_1, \dots)$ is realised by a total state, then this sequence belongs to \mathcal{X}^c .

We will use the following lemma:

Lemma 2.7. *Let \mathcal{S} be an at most countable set of states, and $r \in P$. Then there exists $r^* \leq r$ satisfying the following property: for all $\mathcal{J} \in \mathcal{S}$ and $x, y \in X$ if there exists $u, v \in P$ such that:*

1. **I** can legally continue the play \mathcal{J} by the move (x, u) ;
2. $\tau(\mathcal{J} \frown (x, u)) = (y, v)$;
3. v and r^* are compatible;

then there exists $u', v' \in P$ satisfying 1., 2., and 3. and such that, moreover, we have $r^* \leq^* v'$.

Proof. Let $(\mathcal{J}_n, x_n, y_n)_{n \in \omega}$ be a (non-necessarily injective) enumeration of $\mathcal{S} \times X^2$. Define $(r_n)_{n \in \omega}$ a decreasing sequence of elements of P in the following way. Let $r_0 = r$. For $n \in \omega$, suppose r_n defined. If there exists a pair $(u, v) \in P^2$ such that:

- **I** can legally continue the play \mathcal{J}_n by the move (x_n, u) ;
- $\tau(\mathcal{J}_n \frown (x_n, u)) = (y_n, v)$;
- v and r_n are compatible;

then choose (u_n, v_n) such a pair and let r_{n+1} be a common lower bound to r_n and v_n . Otherwise, let $r_{n+1} = r_n$. This achieves the construction.

By the definition of a Gowers space, there exists $r^* \in P$ such that $r^* \leq r$ and for all $n \in \omega$, $r^* \leq^* r_n$. We show that r^* is as required. Let $n \in \omega$, and suppose that there exists $(u, v) \in P^2$ satisfying properties 1., 2., and 3. as in the statement of the lemma for the triple $(\mathcal{J}_n, x_n, y_n)$. Since $r^* \leq^* r_n$ and since v and r^* are compatible, then v and r_n are also compatible. This show that the pair (u_n, v_n) has been defined; by construction, this pair satisfies properties 1. and 2. for $(\mathcal{J}_n, x_n, y_n)$, and we have $r_{n+1} \leq v_n$, so $r^* \leq^* v_n$, which shows that $(u', v') = (u_n, v_n)$ is as required. □

Proof of Proposition 2.6. Define $(q_n)_{n \in \omega}$ a decreasing sequence of elements of P and $(\mathcal{S}_n)_{n \in \omega}$ a sequence where, for every $n \in \omega$, \mathcal{S}_n is an at most countable set of states of rank n , in the following way. Let $q_0 = \tau(\emptyset)$ and $\mathcal{S}_0 = \{(\tau(\emptyset))\}$. For $n \in \omega$, suppose q_n and \mathcal{S}_n being defined. Let q_{n+1} be the result of the application of Lemma 2.7 to q_n and the set of states \mathcal{S}_n . For $\mathcal{J} \in \mathcal{S}_n$, let $A_{\mathcal{J}}$ be the set of all pairs (x, y) such that there exists $(u, v) \in P^2$ satisfying:

1. **I** can legally continue the play \mathcal{J} by the move (x, u) ;
2. $\tau(\mathcal{J} \frown (x, u)) = (y, v)$;
3. v and q_{n+1} are compatible.

Then by construction of q_{n+1} , for all $(x, y) \in A_{\mathcal{J}}$, there exists a pair $(u, v) \in P^2$ satisfying 1., 2., and 3., and such that moreover $q_{n+1} \leq^* v$. For each $(x, y) \in A_{\mathcal{J}}$, choose $(u_{\mathcal{J},x,y}, v_{\mathcal{J},x,y})$ such a pair. Let $\mathcal{S}_{n+1} = \{\mathcal{J} \cap (x, u_{\mathcal{J},x,y}, y, v_{\mathcal{J},x,y}) \mid \mathcal{J} \in \mathcal{S}_n, (x, y) \in A_{\mathcal{J}}\}$; this is clearly a countable set of states of rank $n + 1$. This achieves the construction.

Now let $q \in P$ be such that $q \leq q_0$ and for all $n \in \omega$, we have $q \leq^* q_n$. Observe that since $q_0 \leq p$, we have $q \leq p$. We show that q is as required, by describing a strategy for **II** in B_q to reach \mathcal{X}^c . In order to do this, we simulate the play $\mathcal{J} = (v_0, x_0, u_0, y_0, v_1, \dots)$ of B_q that **I** and **II** are playing by a play $\mathcal{J}' = (v'_0, x_0, u'_0, y_0, v'_1, \dots)$ of K_p having the same outcome and during which **II** always plays according to her strategy τ . This will ensure that the outcome $(x_0, y_0, x_1, y_1, \dots)$ of both games lies in \mathcal{X}^c and so that the strategy for **II** in B_q that we described enables her to reach her goal. We do this construction in such a way that at each turn n , the following conditions are kept satisfied:

- (a) $\mathcal{J}'_{\uparrow n} \in \mathcal{S}_n$;
- (b) $v_n \leq v'_n$.

The moves of the players at the $(n + 1)^{\text{th}}$ turn in both games that are described in the following proof are represented in the diagrams below. The third diagram, called “Fictive K_p ”, represents a fictive situation that will be studied for technical reasons in the proof, and in which the moves of both players are the same as in K_p until the n^{th} turn but differ from the $(n + 1)^{\text{th}}$ turn.

B_q	I	...	x_n, u_n	
	II	...	\dots, v_n	y_n, v_{n+1}
K_p	I	...	x_n, u'_n	
	II	...	\dots, v'_n	y_n, v'_{n+1}
Fictive K_p	I	...	x_n, u''_n	
	II	...	\dots, v'_n	y_n, v''_{n+1}

Let us describe the strategy of **II** in B_q . At the first turn, this strategy will consist in playing $v_0 = q$; and, according to her strategy τ , **II** will play $v'_0 = \tau(\emptyset)$ in K_p . Now, suppose that both games have been played until the n^{th} turn, that is, the last moves of player **II** in the games B_q and K_p are respectively v_n and v'_n . Player **I** plays (x_n, u_n) in B_q . By the rules of the game B_q and the induction hypothesis, we have that

$u_n \leq q \leq^* v_n \leq v'_n$; so there exists $u''_n \in P$ such that $u''_n \lesssim u_n$ and $u''_n \leq v'_n$. We also have that $(x_0, y_0, \dots, x_n) \triangleleft v_n \leq v'_n$, so it is legal for **I** to pursue the game K_p by playing (x_n, u''_n) ; this fictive situation is represented in the third diagram above, called “Fictive K_p ”. In this fictive situation, the strategy τ of **II** would lead her to answer with a move (y_n, v''_{n+1}) satisfying $(x_0, y_0, \dots, x_n, y_n) \triangleleft u''_n$ and $v''_{n+1} \leq u''_n$. We have, by construction of q , that $v''_{n+1} \leq u''_n \leq u_n \leq q \leq^* q_{n+1}$; so in particular, v''_{n+1} and q_{n+1} are compatible. Recalling that $\mathcal{J}_{\uparrow n} \in \mathcal{S}_n$, we see that the pair (u''_n, v''_n) witnesses that $(x_n, y_n) \in A_{\mathcal{J}_{\uparrow n}}$.

Now let us leave the fictive situation and come back to the “real” K_p . Since $(x_n, y_n) \in A_{\mathcal{J}_{\uparrow n}}$, we know that the pair $(u_{\mathcal{J}_{\uparrow n}, x_n, y_n}, v_{\mathcal{J}_{\uparrow n}, x_n, y_n})$ has been defined; we denote this pair by (u'_n, v'_{n+1}) . In the “real” K_p , we make **I** play (x_n, u'_n) . By definition of (u'_n, v'_{n+1}) , this move is legal, and **II** will answer, according to her strategy, with (y_n, v'_{n+1}) . Observe that the required condition (a) in the induction hypothesis is satisfied by these moves since, by the definition of \mathcal{S}_{n+1} , we have $\mathcal{J}_{\uparrow n} \widehat{\ } (x_n, u'_n, y_n, v'_{n+1}) \in \mathcal{S}_{n+1}$. We also have that $q \leq^* q_{n+1} \leq^* v'_{n+1}$, so there exists $v_{n+1} \in P$ such that $v_{n+1} \leq v'_{n+1}$ and $v_{n+1} \lesssim q$. For this reason, and since we also have (as we already saw) $(x_0, y_0, \dots, x_n, y_n) \triangleleft u''_n \leq u_n$, we get that (y_n, v_{n+1}) is a legal move for **II** in B_q , that satisfies the condition (b) in the induction hypothesis. So we just have to define her strategy as leading her to play this move, and this achieves the proof. \square

We actually proved a little more than Theorem 2.4. Say that the Gowers space \mathcal{G} is *analytic* if P is an analytic subset of a Polish space and if the relations \leq and \triangleleft are Borel subsets of P^2 and of $\text{Seq}(X) \times P$ respectively. For most of the spaces we actually use, P can be identified to an analytic subset of $\mathcal{P}(X)$, the relation \leq to the inclusion, and the relation $(x_0, \dots, x_n) \triangleleft p$ to the membership relation $x_n \in p$; thus, these spaces are analytic. This is, for instance, the case for the Mathias–Silver space and the Rosendal space introduced at the beginning of this section.

Now, say that a class Γ of subsets of Polish spaces is *suitable* if it contains the class of Borel sets and is stable under finite unions, finite intersections and Borel inverse images. Equip \mathbb{R} with its usual Polish topology, and \mathbb{R}^ω with the product topology. Then an easy consequence of Proposition 2.6 is the following:

Corollary 2.8. *Let Γ be a suitable class of subsets of Polish spaces. If every Γ -subset of \mathbb{R}^ω is determined, then for an analytic Gowers space $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$, every Γ -subset of X^ω is adversarially Ramsey.*

Proof. Fix $\mathcal{X} \subseteq X^\omega$ a Γ -subset, and $p \in P$. By Proposition 2.6, it is enough to show that in the game K_p , either player **I** has a strategy to reach \mathcal{X} , or player **II** has a strategy to reach \mathcal{X}^c . Let $\varphi : \mathbb{R} \rightarrow P$ be a surjective Borel mapping, and consider the following game K'_p :

$$\begin{array}{ccccccc} \mathbf{I} & & x_0, \tilde{q}_0 & & x_1, \tilde{q}_1 & & \dots \\ \mathbf{II} & \tilde{p}_0 & & y_0, \tilde{p}_1 & & y_1, \tilde{p}_2 & \dots \end{array}$$

where the x_i 's and the y_i 's are elements of X and the \tilde{p}_i 's and the \tilde{q}_i 's are real numbers, with the constraint that $\varphi(\tilde{p}_0) \leq p$, for all $i \in \omega$, $\varphi(\tilde{q}_i) \leq \varphi(\tilde{p}_i)$, $\varphi(\tilde{p}_{i+1}) \leq \varphi(\tilde{q}_i)$,

$(x_0, y_0, \dots, x_i) \triangleleft \varphi(\tilde{p}_i)$, and $(x_0, y_0, \dots, x_i, y_i) \triangleleft \varphi(\tilde{q}_i)$, and whose outcome is the sequence $(x_0, y_0, x_1, y_1, \dots) \in X^\omega$. This game is clearly equivalent to K_p : **I** has a strategy to reach \mathcal{X} in K_p if and only if he has one in K'_p , and **II** has a strategy to reach \mathcal{X}^c in K_p if and only if she has one in K'_p . Since K'_p is a game on real numbers with Borel rules and since \mathcal{X} is in Γ , we deduce that in this game, either **I** has a strategy to reach \mathcal{X} , or **II** has a strategy to reach \mathcal{X}^c , what concludes the proof. \square

Martin proved in [13] that if there exists a measurable cardinal, then every analytic subset of ω^ω is determined. He observed in this paper that his proof actually shows a little more: if κ is a measurable cardinal, if S is a set with the discrete topology such that $|S| < \kappa$ and if S^ω is endowed with the product topology, then every Σ_1^1 -subset of S^ω is determined. (Here, a Σ_1^1 -subset of a topological space X is defined as a set $A \subseteq X$ which is the first projection of some closed subset of $X \times \omega^\omega$.) In particular, if there exists a measurable cardinal, then every analytic subset of \mathbb{R}^ω is determined, and by the last corollary, every analytic set in an analytic Gowers space is adversarially Ramsey. This gives an answer to Question 1.11.

Recall that $PD_{\mathbb{R}}$ is the statement “every projective subset of \mathbb{R}^ω (with \mathbb{R} endowed with its usual Polish topology) is determined” and that $AD_{\mathbb{R}}$ is the statement “every subset of \mathbb{R}^ω is determined”. Corollary 2.8 shows in particular that, in an analytic Gowers space, under $PD_{\mathbb{R}}$, every projective set is adversarially Ramsey. Recall that Harrington and Kechris [8], and independently Woodin [22] proved that under PD , every projective subset of $[\omega]^\omega$ is Ramsey. Using ideas from Woodin’s proof, Bagaria and López-Abad [2] showed that under PD , every projective set of block sequences of a basis of a Banach space is strategically Ramsey (i.e. satisfies the conclusion of Gowers’ Theorem 1.2). Basing ourselves on these facts, we can formulate the following conjecture:

Conjecture 2.9. *Under PD , if the Gowers space $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$ is analytic, then every projective subset of X^ω is adversarially Ramsey.*

Clearly, the method presented in the present paper does not enable to prove this.

Also observe that the proof of Proposition 2.6 can almost entirely be done in $ZF + DC$; the only use of the full axiom of choice is made to choose $u''_n \in P$ such that $u''_n \lesssim u_n$ and $u''_n \leq v'_n$, and $v_{n+1} \in P$ such that $v_{n+1} \leq v'_{n+1}$ and $v_{n+1} \lesssim q$, so actually to apply axiom 2. in the definition of a Gowers space. For this reason, say that the Gowers space \mathcal{G} is *effective* if in this axiom 2., the subspace r can be chosen in an effective way, that is, if there exist a function $f : P^2 \rightarrow P$ such that for every $p, q \in P$, if $p \leq^* q$, then we have $f(p, q) \lesssim p$ and $f(p, q) \leq q$. For instance:

- The Mathias–Silver space is effective: indeed, if $M \subseteq^* N$, then we can take $f(M, N) = M \cap N$.
- The Rosendal space is effective. Indeed, if X and Y are block subspaces such that $X \subseteq^* Y$, let $(x_n)_{n \in \omega}$ be a block sequence spanning X . Then we can let $f(X, Y)$ be

the subspace spanned by the largest final segment of (x_n) all of whose terms are in Y (this subspace does not depend on the choice of (x_n)).

To prove Proposition 2.6 for an effective Gowers space, we only need dependant choices. Thus, we have the following result:

Corollary 2.10 ($ZF + DC + AD_{\mathbb{R}}$). *Let $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$ be an effective Gowers space such that P is a subset of a Polish space. Then every subset of X^ω is adversarially Ramsey.*

Proof. Recall that in $ZF + DC + AD$, every subset of a Polish space is either at most countable, or contains a Cantor set, and is thus in bijection with \mathbb{R} (this is a consequence of Theorem 21.1 in [10], that can be proved in $ZF + DC$). So if P is countable, then Kastanas' game can be viewed as a game on integers and is thus determined, and if P is uncountable, then Kastanas' game can be viewed as a game on real numbers, that is also determined. The conclusion follows from Proposition 2.6. □

As above, we can't prove in this way that the same result holds under AD instead of $AD_{\mathbb{R}}$, but we conjecture that it does so. As we will see in the next section, if this is true, this would imply that under AD , every subset of $[\omega]^\omega$ is Ramsey, which is still conjectural today. More generally, it would be interesting to know whether, in general, the determinacy of Γ -subsets of ω^ω is enough to prove the adversarial Ramsey property for Γ -sets in sufficiently regular Gowers spaces, for any suitable class Γ .

3 Strategically Ramsey sets and the pigeonhole principle

The aim of this section is to prove a version of Rosendal's Theorem 1.7 in the general setting of Gowers spaces. We also introduce the notion of the *pigeonhole principle* for a Gowers space and see that the latter result can be strengthened in the case where this principle holds. This will enable us to see the fundamental difference between the Mathias–Silver space and the Rosendal space over a field with at least three elements. We start by introducing Gowers' game and the asymptotic game in the setting of Gowers spaces, and the notion of a strategically Ramsey set. In this whole section, we fix a Gowers space $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$.

Definition 3.1. Let $p \in P$.

1. *Gowers' game below p* , denoted by G_p , is defined in the following way:

$$\begin{array}{llll} \text{I} & p_0 & p_1 & \dots \\ \text{II} & x_0 & x_1 & \dots \end{array}$$

where the x_i 's are elements of X , and the p_i 's are elements of P . The rules are the following:

- for **I**: for all $i \in \omega$, $p_i \leq p$;
- for **II**: for all $i \in \omega$, $(x_0, \dots, x_i) \triangleleft p_i$.

The outcome of the game is the sequence $(x_i)_{i \in \omega} \in X^\omega$.

2. The *asymptotic game below p* , denoted by F_p , is defined in the same way as G_p , except that this time we moreover require that $p_i \lesssim p$.

Definition 3.2. A set $\mathcal{X} \subseteq X^\omega$ is said to be *strategically Ramsey* if for every $p \in P$, there exists $q \leq p$ such that either player **I** has a strategy to reach \mathcal{X}^c in F_q , or player **II** has a strategy to reach \mathcal{X} in G_q .

The general version of Rosendal's Theorem 1.7 is then the following:

Theorem 3.3 (Abstract Rosendal's theorem). *Every analytic subset of X^ω is strategically Ramsey.*

Observe that Theorem 1.7 is exactly the result of the application of Theorem 3.3 to the Rosendal space.

Proof. We prove the result for Borel sets first. In order to do this, consider another space $\tilde{\mathcal{G}} = (P, X, \leq, \leq^*, \tilde{\triangleleft})$, where P , X , \leq , and \leq^* are the same as in \mathcal{G} , but we replace \triangleleft by the relation $\tilde{\triangleleft}$ defined by $(x_0, y_0, x_1, y_1, \dots, x_n, y_n) \tilde{\triangleleft} p$ iff $(y_0, y_1, \dots, y_n) \triangleleft p$, and $(x_0, y_0, x_1, y_1, \dots, x_n) \tilde{\triangleleft} p$ iff $(x_0, x_1, \dots, x_n) \triangleleft p$. Now, to each set $\mathcal{X} \subseteq X^\omega$, associate a set $\tilde{\mathcal{X}} \subseteq X^\omega$ defined by $(x_0, y_0, x_1, y_1, \dots) \in \tilde{\mathcal{X}} \Leftrightarrow (y_0, y_1, \dots) \in \mathcal{X}$. Then, when players try to reach $\tilde{\mathcal{X}}$ or $\tilde{\mathcal{X}}^c$ in the games A_p and B_p of $\tilde{\mathcal{G}}$, the p_i 's played by **II** and the x_i 's played by **I** don't matter at all; so a strategy for **I** in the game A_p of $\tilde{\mathcal{G}}$ to reach $\tilde{\mathcal{X}}^c$ becomes a strategy for **I** in the game F_p of \mathcal{G} to reach \mathcal{X}^c , and a strategy for **II** in the game B_p of $\tilde{\mathcal{G}}$ to reach $\tilde{\mathcal{X}}$ becomes a strategy for **II** in the game G_p of \mathcal{G} to reach \mathcal{X} . Thus, the strategical Ramsey property for \mathcal{X} in \mathcal{G} is equivalent to the adversarial Ramsey property for $\tilde{\mathcal{X}}^c$ in $\tilde{\mathcal{G}}$, so the strategical Ramsey property for Borel sets in \mathcal{G} follows from Theorem 2.4.

From the result for Borel sets, we now deduce the result for arbitrary analytic sets using an unfolding argument (see [10], section 21, for a general presentation of the method of unfolding). Let $\mathcal{X} \subseteq X^\omega$ be analytic, and $p \in P$. Let $X' = X \times \{0, 1\}$, whose elements will be denoted by the letters (x, ε) . Define the binary relation $\triangleleft' \subseteq \text{Seq}(X') \times P$ by $(x_0, \varepsilon_0, \dots, x_n, \varepsilon_n) \triangleleft' p$ if $(x_0, \dots, x_n) \triangleleft p$, and consider the Gowers space $\mathcal{G}' = (P, X', \leq, \leq^*, \triangleleft')$. In this proof, we will use the notations F_q and G_q to denote respectively the asymptotic game and Gowers' game in the space \mathcal{G} , whereas the notations F'_q and G'_q will be used for these games in the space \mathcal{G}' . We denote by π the projection $X'^\omega \rightarrow X^\omega$. Let $\mathcal{X}' \subseteq X'^\omega$ be a G_δ set such that $\mathcal{X} = \pi(\mathcal{X}')$. Since \mathcal{X}' is \mathcal{G}_δ , it is strategically Ramsey by the first part of this proof; let $q \leq p$ witnessing so. If player **II** has a strategy in G'_q to reach \mathcal{X}' , then a run of the game G_q where **II** uses this strategy but omits to display the ε_i 's produces an outcome lying in \mathcal{X} ; hence, **II** has a strategy to reach \mathcal{X} in G_q . Then, our result will follow from the following fact:

Fact 3.4. *If \mathbf{I} has a strategy to reach \mathcal{X}'^c in F'_q , then he has a strategy to reach \mathcal{X}^c in F_q .*

Proof. Let τ' be a strategy enabling \mathbf{I} to reach \mathcal{X}'^c in F'_q . In order to save notation, in this proof, we consider that in the games F'_q and F_q , player \mathbf{II} is allowed not to respect the rules (i.e. to play x_i 's such that $(x_0, \dots, x_i) \not\triangleleft p_i$), but loses the game if she does. Then, the strategy τ' can be viewed as a mapping $X'^{<\omega} \rightarrow P$ such that for every $(x_0, \varepsilon_0, \dots, x_{n-1}, \varepsilon_{n-1}) \in X'^{<\omega}$, we have $\tau'(x_0, \varepsilon_0, \dots, x_{n-1}, \varepsilon_{n-1}) \lesssim q$. Observe that if $(p_j)_{j \in J}$ is a finite family of elements of P such that $\forall j \in J, p_j \lesssim q$, then by applying iteratively the property 2. in the definition of a Gowers space, we can get $p^* \in P$ such that $p^* \lesssim q$ and $\forall j \in J, p^* \leq p_j$. Thus, for every $(x_0, \dots, x_{n-1}) \in X^{<\omega}$, we can choose $\tau(x_0, \dots, x_{n-1}) \in P$ such that $\tau(x_0, \dots, x_{n-1}) \lesssim q$ and such that for every $(\varepsilon_0, \dots, \varepsilon_{n-1}) \in \{0, 1\}^n$, we have $\tau(x_0, \dots, x_{n-1}) \leq \tau'(x_0, \varepsilon_0, \dots, x_{n-1}, \varepsilon_{n-1})$. We have hence defined a mapping $\tau : X^{<\omega} \rightarrow P$; we show that this is a strategy for \mathbf{I} in F_q enabling him to reach \mathcal{X}^c .

Consider a run of the game F_q during which \mathbf{II} respects the rules and \mathbf{I} plays according to his strategy τ :

$$\begin{array}{cccc} \mathbf{I} & p_0 & p_1 & \dots \\ \mathbf{II} & x_0 & x_1 & \dots \end{array}$$

We have to show that $(x_i)_{i \in \omega} \notin \mathcal{X}$, that is, for every $(\varepsilon_i)_{i \in \omega} \in \{0, 1\}^\omega$, $(x_i, \varepsilon_i)_{i \in \omega} \notin \mathcal{X}'$. Let $(\varepsilon_i)_{i \in \omega} \in \{0, 1\}^\omega$; it is enough to show that $(x_i, \varepsilon_i)_{i \in \omega}$ is the outcome of a run of the game F'_q during which \mathbf{I} always follows his strategy τ' and \mathbf{II} always respects the rules. Letting $p'_i = \tau'(x_0, \varepsilon_0, \dots, x_{n-1}, \varepsilon_{n-1})$, this means that during the following run of the game F'_q , player \mathbf{II} always respects the rules:

$$\begin{array}{cccc} \mathbf{I} & p'_0 & p'_1 & \dots \\ \mathbf{II} & x_0, \varepsilon_0 & x_1, \varepsilon_1 & \dots \end{array}$$

But for every $i \in \omega$, we have that $p_i = \tau(x_0, \dots, x_{n-1})$ and $p'_i = \tau'(x_0, \varepsilon_0, \dots, x_{n-1}, \varepsilon_{n-1})$, so by definition of τ , we have $p_i \leq p'_i$. Since player \mathbf{II} respects the rules in F_q , we have that $(x_0, \dots, x_i) \triangleleft p_i$, so $(x_0, \varepsilon_0, \dots, x_i, \varepsilon_i) \triangleleft p'_i$, and \mathbf{II} also respects the rules in F'_q . This concludes the proof. □

□

Observe that in the proof of Theorem 3.3, we only need Theorem 2.4 for G_δ sets, and hence determinacy for G_δ games. Hence, unlike Theorem 2.4 in its generality, the last result is provable in ZC . Actually, as previously, for effective Gowers spaces, it is even provable in $Z + DC$.

Again, we actually proved a little more. For a suitable class Γ of subsets of Polish spaces, let $\exists\Gamma$ be the class of projections of Γ -sets; in other words, for A a subset of a Polish space Y , we have $A \in \exists\Gamma$ if and only if there exist $B \in Y \times 2^\omega$ such that $B \in \Gamma$ and A is the first projection of B (we could have taken any uncountable Polish space instead

of 2^ω in this definition, since Γ is closed under Borel inverse images). Then the proof of Theorem 3.3, combined with Corollaries 2.8 and 2.10, actually shows the following:

Corollary 3.5.

1. Let Γ be a suitable class of subsets of Polish spaces. If every Γ -subset of \mathbb{R}^ω is determined, then for an analytic Gowers space $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$, every $\exists\Gamma$ -subset of X^ω is strategically Ramsey.
2. ($ZF + DC + AD_{\mathbb{R}}$) Let $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$ be an effective Gowers space such that P is a subset of a Polish space. Then every subset of X^ω is strategically Ramsey.

In particular, if there exists a measurable cardinal, then in an analytic Gowers space, every Σ_2^1 -set is strategically Ramsey.

The rest of this section aims to explain how we can, in certain cases, get symmetrical Ramsey results like Mathias–Silver’s theorem from Theorem 3.3, which is asymmetrical. By *asymmetrical*, we mean here that unlike Mathias–Silver’s theorem, in Theorem 3.3, both possible conclusion don’t have the same form. Actually, one of these conclusions is stronger than the other (and, as it will turn out later, *strictly* stronger in general), as is shown by the following lemma.

Lemma 3.6. *Let $\mathcal{X} \subseteq X^\omega$ and $p \in P$. Suppose that **I** has a strategy in F_p to reach \mathcal{X} . Then **II** has a strategy in G_p to reach \mathcal{X} .*

Proof. Fix τ a strategy enabling **I** to reach \mathcal{X} in F_p . We describe a strategy for **II** in G_p by simulating a play $(q_0, x_0, q_1, x_1, \dots)$ of G_p by a play $(p_0, x_0, p_1, x_1, \dots)$ of F_p having the same outcome and during which **I** always plays according to τ ; this will ensure that $(x_0, x_1, \dots) \in \mathcal{X}$ and that this play of G_p will be winning for **II**.

Suppose that the first n turns of both games have been played, which means that the p_i ’s, the q_i ’s and the x_i ’s have been chosen for every $i < n$. For the next turn, in G_p , player **I** plays $q_n \leq p$, and in F_p , the strategy τ tells **I** to play $p_n \lesssim p$. Then $q_n \leq^* p_n$, so by axiom 2. in the definition of a Gowers space, there exists $r_n \in P$ such that $r_n \leq p_n$ and $r_n \leq q_n$. Let $x_n \in X$ such that $(x_0, \dots, x_n) \triangleleft r_n$ (existing by axiom 4.). Then x_n can be legally played by **II** in both F_p and G_p , what concludes the proof. □

Actually, the fact that **I** has a strategy in F_p to reach some set \mathcal{X} is in general much stronger than the fact, for **II**, to have a strategy in G_p to reach the same set, and the first statement is in fact very close to a “genuine” Ramsey statement. By a “genuine” Ramsey statement, we mean a non-game-theoretical statement of the form “every sequence $(x_n)_{n \in \omega}$ such that $\forall n \in \omega (x_0, \dots, x_n) \triangleleft p$, and moreover satisfying some structural condition, belongs to \mathcal{X} ”, like in the conclusions of Mathias–Silver’s theorem. In the case of the Mathias–Silver space, the link between the existence of a strategy for **I** in the asymptotic game and a genuine Ramsey statement is given by the following lemma:

Lemma 3.7. *Work in the Mathias–Silver space, and let $\mathcal{X} \subseteq \omega^\omega$. Suppose that, for some $M \in [\omega]^\omega$, player **I** has a strategy in F_M to reach \mathcal{X} . Then there exists an infinite $N \subseteq M$ such that every infinite $S \subseteq N$ belongs to \mathcal{X} (here, we identify infinite subsets of ω with increasing sequences of integers).*

Obviously, a weak converse of this lemma holds: if every infinite $S \subseteq M$ belongs to \mathcal{X} , then **I** has a strategy in F_M to reach \mathcal{X} . Indeed, he can always ensure that the outcome of this game is an increasing sequence.

Proof of Lemma 3.7. Without loss of generality, assume $M = \omega$. As in the proof of Fact 3.4, consider that in F_ω , player **II** is allowed to play against the rules, but loses if she does. Let τ be a strategy for player **I** in F_ω , enabling him to reach \mathcal{X} ; in this context, this strategy can be viewed as a mapping associating to each finite sequence of integers a cofinite subset of ω . Without loss of generality, we can assume that these cofinite subsets are final segments of ω ; for $s \in \omega^{<\omega}$, let $\tau_0(s) = \min \tau(s)$. Now define, by induction, a strictly increasing sequence $(n_i)_{i \in \omega}$ of integers in the following way: let $n_0 = \tau_0(\emptyset)$, and for $i \in \omega$, let n_{i+1} be the maximum of $n_i + 1$ and of the $\tau_0(n_{i_0}, \dots, n_{i_{k-1}})$'s for $k \in \omega$ and $0 \leq i_0 < \dots < i_{k-1} = i$. Let $N = \{n_i \mid i \in \omega\}$; then N is as required. Indeed, an infinite subset of N has the form $\{n_{i_k} \mid k \in \omega\}$ for a strictly increasing sequence of integers $(i_k)_{k \in \omega}$. To prove that $(n_{i_k})_{k \in \omega} \in \mathcal{X}$, it is enough to prove this sequence is the outcome of some legal run of the game F_ω during which player **I** always plays according to the strategy τ . In other words, letting, for all $k \in \omega$, $P_k = \tau(n_{i_0}, \dots, n_{i_{k-1}})$, we have to show that during the following run of the game F_ω , player **II** always respects the rules:

$$\begin{array}{ccccccc} \mathbf{I} & P_0 & & P_1 & & \dots & \\ \mathbf{II} & & n_{i_0} & & n_{i_1} & & \dots \end{array}$$

But by construction, we have that $n_{i_0} \geq n_0 = \tau_0(\emptyset) = \min P_0$, and for $k \geq 1$, $n_{i_k} \geq n_{i_{k-1}+1} \geq \tau_0(n_{i_0}, \dots, n_{i_{k-1}}) = \min P_k$, which concludes the proof. \square

The setting of Gowers spaces does not give enough structure to get such a result in general. A general version of this result will be given in section 5, in the setting of *approximate asymptotic spaces* with some additional structure; and, in a very different way, the setting of *Ramsey spaces* presented in [21] is also convenient to get non game-theoretical infinite-dimensional Ramsey results.

We now introduce the *pigeonhole principle*, a property of Gowers spaces under which we can get a Ramsey result involving a strategy for player **I** in the asymptotic game in both sides. This result is the best possible we can get in this general setting, and Lemma 3.7 shows that this is the right analogue of Mathias–Silver’s theorem.

In the rest of this paper, denote by $q \subseteq_s A$, for $q \in P$, $s \in X^{<\omega}$ and $A \subseteq X$, the fact that for every $x \in X$ such that $s \hat{\ } x \triangleleft q$, we have $x \in A$. This notation could sound strange, however, in spaces where $P \subseteq \mathcal{P}(X)$ and $(x_0, \dots, x_n) \triangleleft q \Leftrightarrow x_n \in q$, we have that $q \subseteq_s A$ iff $q \subseteq A$. Let us introduce the pigeonhole principle.

Definition 3.8. The Gowers space \mathcal{G} is said to satisfy the *pigeonhole principle* if for every $p \in P$, $s \in X^{<\omega}$ and $A \subseteq X$, there exists $q \leq p$ such that either $q \subseteq_s A$, or $q \subseteq_s A^c$.

The pigeonhole principle trivially holds in the Mathias–Silver space. It also holds in the Rosendal space over the field \mathbb{F}_2 : this is actually a rephrasing of Hindman’s theorem for FIN (see for example [21], Theorem 2.25). However, it does not hold in the Rosendal space over K , for $K \neq \mathbb{F}_2$: to see this, take for example for A the set of all vectors whose first nonzero coordinate is 1. Note that apart from this trivial obstruction, the pigeonhole principle does not hold in the Rosendal space for much more intrinsic reasons. Indeed, consider the *projective Rosendal space*, i.e. the forgetful Gowers space $\mathcal{PR}_K = (P, \mathbb{P}(E), \subseteq, \subseteq^*, \subseteq)$, where $\mathbb{P}(E)$ is a countably infinite-dimensional projective space over the field K (that is, the set of vector lines of some countably infinite-dimensional K -vector space E), P is the set of block subspaces of E relative to a fixed basis $(e_i)_{i \in \omega}$ of E , \subseteq^* is the inclusion up to finite codimension as in the definition of the Rosendal space, and where since the space is forgetful, the relation usually denoted by \triangleleft is viewed as a relation between points and subspaces, here the inclusion. Then for $K \neq \mathbb{F}_2$, the pigeonhole principle still does not hold in \mathcal{PR}_K : take for example for A the set of all vector lines Kx , where the first and the last non-zero coordinates of x are equal.

Under the pigeonhole principle, we will show a form of converse assertion to Lemma 3.6:

Proposition 3.9. *Suppose that the Gowers space \mathcal{G} satisfies the pigeonhole principle. Let $\mathcal{X} \subseteq X^\omega$ and $p \in P$. If player **II** has a strategy in G_p to reach \mathcal{X} , then there exists $q \leq p$ such that **I** has a strategy in F_q to reach \mathcal{X} .*

Before proving this proposition, let us make some remarks. First, Proposition 3.9 immediately implies the following corollary:

Corollary 3.10. *Suppose that the Gowers space \mathcal{G} satisfies the pigeonhole principle. Let $\mathcal{X} \subseteq X^\omega$ be a strategically Ramsey set. Then for all $p \in P$, there exists $q \leq p$ such that in F_q , player **I** has a strategy either to reach \mathcal{X} , or to reach \mathcal{X}^c .*

This corollary has some kind of converse. Indeed, for every $s \in X^{<\omega}$, consider the Gowers space $\mathcal{G}^s = (P, X, \leq, \leq^*, \triangleleft^s)$, where P , X , \leq and \leq^* are the same as in \mathcal{G} and where $t \triangleleft^s p \Leftrightarrow s \hat{\ } t \triangleleft p$. Then it is not hard to see that \mathcal{G} satisfies the pigeonhole principle if and only if the conclusion of Corollary 3.10 holds for all sets of the form $\{(x_n)_{n \in \omega} \mid x_0 \in A\}$ (where $A \subseteq X$), in the space \mathcal{G}^s for every s . Indeed, in \mathcal{G}^s , player **I** has a strategy in F_q to reach the set $\{(x_n)_{n \in \omega} \mid x_0 \in A\}$ if and only if there exists $q_0 \lesssim q$ such that $q_0 \subseteq_s A$. In the particular case of a forgetful space, the satisfaction of the conclusion of the last corollary for clopen sets is thus equivalent to the pigeonhole principle.

Also observe that Corollary 3.10 applied to the Mathias–Silver space, combined with Lemma 3.7, gives that a set $\mathcal{X} \subseteq [\omega]^\omega$ is Ramsey (in the sense of Mathias–Silver’s theorem) if and only if it is strategically Ramsey in the Mathias–Silver space (when seen as a subset of ω^ω). In particular, Mathias–Silver’s theorem is a consequence of the abstract Rosendal’s Theorem 3.3.

We now prove Proposition 3.9.

Proof of Proposition 3.9. Fix τ a strategy for **II** in G_p to reach \mathcal{X} . As in the proof of Proposition 2.6, call a *state* a partial play of G_p either empty or ending with a move of **II**, during which **II** always plays according to her strategy. Say that a state *realises* a sequence $(x_0, \dots, x_{n-1}) \in X^{<\omega}$ if it has the form $(p_0, x_0, \dots, p_{n-1}, x_{n-1})$. Define in the same way the notion of a *total state* (which is a total play of G_p) and of realisation for a total state; if an infinite sequence is realised by some total state, then it belongs to \mathcal{X} . Say that a point $x \in X$ is *reachable* from a state \mathcal{J} if there exists $r \leq p$ such that $\tau(\mathcal{J} \hat{\ } r) = x$. Denote by $A_{\mathcal{J}}$ the set of all points that are reachable from the state \mathcal{J} . We will use the following fact.

Fact 3.11. *For every state \mathcal{J} realising a finite sequence s , and for every $q \leq p$, there exists $r \leq q$ such that $r \subseteq_s A_{\mathcal{J}}$.*

Proof. Otherwise, by the pigeonhole principle, there would exist $r \leq q$ such that $r \subseteq_s (A_{\mathcal{J}})^c$. But then **I** could play r after the partial play \mathcal{J} , and **II** would answer, according to her strategy, by $x = \tau(\mathcal{J} \hat{\ } r)$ that should satisfy $s \hat{\ } x \triangleleft r$. Since $r \subseteq_s (A_{\mathcal{J}})^c$, this would imply that $x \in (A_{\mathcal{J}})^c$. But we also have, by definition of $A_{\mathcal{J}}$, that $x \in A_{\mathcal{J}}$, a contradiction. □

Now let $(s_n)_{n \in \omega}$ be an enumeration of $X^{<\omega}$ such that if $s_m \subseteq s_n$, then $m \leq n$. We define, for some $n \in \omega$, a state \mathcal{J}_n realising s_n , by induction in the following way: $\mathcal{J}_0 = \emptyset$ and for $n \geq 1$, letting $s_n = s_m \hat{\ } x$ for some $m < n$ and some $x \in X$,

- if \mathcal{J}_m has been defined and if x is reachable from \mathcal{J}_m , then choose a $r \leq p$ such that $x = \tau(\mathcal{J}_m \hat{\ } r)$ and put $\mathcal{J}_n = \mathcal{J}_m \hat{\ } (r, x)$,
- otherwise, \mathcal{J}_n is not defined.

Observe that if \mathcal{J}_n is defined and if $s_m \subseteq s_n$, then \mathcal{J}_m is defined and $\mathcal{J}_m \subseteq \mathcal{J}_n$.

We now define a \leq -decreasing sequence $(q_n)_{n \in \omega}$ of elements of P in the following way: $q_0 = p$ and

- if \mathcal{J}_n is defined, then q_{n+1} is the result of the application of Fact 3.11 to \mathcal{J}_n and q_n ;
- $q_{n+1} = q_n$ otherwise.

Finally, let $q \leq p$ be such that for every $n \in \omega$, $q \leq^* q_n$. We will show that **I** has a strategy in F_q to reach \mathcal{X} . We describe this strategy on the following play of F_q :

I	u_0	u_1	\dots
II	x_0	x_1	\dots

We actually show that **I** can always play preserving the fact that, if $n_i \in \omega$ is such that $s_{n_i} = (x_0, \dots, x_{i-1})$, then \mathcal{J}_{n_i} is defined. This will be enough to conclude: indeed, $\bigcup_{i \in \omega} \mathcal{J}_{n_i}$ will be a total state realising the sequence $(x_i)_{i \in \omega}$, showing that this sequence belongs to \mathcal{X} .

Suppose that the i^{th} turn of the play has just been played, so the sequence $s_{n_i} = (x_0, \dots, x_{i-1})$ has been defined, in such a way that \mathcal{J}_{n_i} is defined. Then by construction of q_{n_i+1} , we have that $q_{n_i+1} \subseteq_{s_{n_i}} A_{\mathcal{J}_{n_i}}$. We let **I** play some u_i such that $u_i \lesssim q$ and $u_i \leq q_{n_i+1}$. Then $u_i \subseteq_{s_{n_i}} A_{\mathcal{J}_{n_i}}$, so whatever is the x_i that **II** answers with, this x_i is reachable from \mathcal{J}_{n_i} . So if $s_{n_{i+1}} = s_{n_i} \hat{\ } x_i$, then $\mathcal{J}_{n_{i+1}}$ has been defined, and the wanted property is preserved. □

Observe that this proof can be done in $ZF + DC$, even if the space \mathcal{G} is not supposed effective.

4 Approximate Gowers spaces

For some applications, for instance in Banach-space theory, it would be useful to have results similar to those presented in the previous sections, but for spaces with an uncountable set of points. However, it turns out that in the definition of a Gowers space, this hypothesis is necessary: without it, Theorems 2.4 and 3.3 are not true in general. We begin this section with presenting a counterexample for this. Then, following an idea introduced by Gowers for his Ramsey-type Theorem 1.2, we introduce a metric version of Gowers spaces, allowing us to get approximate Ramsey-type results in situations where the set of points is uncountable. The results of this section, along with these of the next section, will allow us to directly recover results like Gowers' Theorems 1.2 and 1.5. The interest of the spaces we introduce here is more practical than theoretical: their main aim is to allow applications, for instance in Banach-space geometry.

Proposition 4.1. *There exists a space $\mathcal{G} = (P, X, \leq, \leq^*, \triangleleft)$ satisfying all the axioms of forgetful Gowers spaces, apart from the fact that here, X is not a countable set but an uncountable Polish space, in which not every Borel set is strategically Ramsey, neither every Borel set is adversarially Ramsey.*

Sketch of proof. Work in the \mathbb{R} -vector space \mathbb{R}^ω , endowed with the product topology. For $x = (x^i)_{i \in \omega} \in \mathbb{R}^\omega$, let $\text{supp}(x) = \{i \in \omega \mid x^i \neq 0\}$. Relative to this notion of support, define as usual the notion of a block-sequence, and define a block subspace as a closed subspace spanned by a block-sequence. Let P be the set of block-subspaces, and $X = \mathbb{R}^\omega \setminus \{0\}$. The space \mathcal{G} is defined similarly as the Rosendal space: $\mathcal{G} = (P, X, \subseteq, \subseteq^*, \in)$.

We sketch the construction of a Borel set $\mathcal{X} \subseteq X^\omega$ that is not strategically Ramsey (here, X^ω is endowed with the product of the Polish topologies on X). Observe that the set $\{(x_n)_{n \in \omega} \in X^\omega \mid (x_0, x_2, \dots) \in \mathcal{X}\}$ is also Borel and is not adversarially Ramsey.

For $x = (x^i)_{i \in \omega} \in X$, let $N(x) = x^{\min \text{supp}(x)}$. Let \tilde{P} denote the set of block-sequences; this is a closed subset of X^ω , so there is a Borel isomorphism $\varphi : \mathbb{R}^* \rightarrow \tilde{P}$. We define the set \mathcal{X} in the following way: $(x_n)_{n \in \omega}$ is in \mathcal{X} if and only if x_1 is equal to a term of the block sequence $\varphi(N(x_0))$. Then \mathcal{X} is not strategically Ramsey. Indeed, given

$p \in P$, player **II** cannot have a strategy in G_p to reach \mathcal{X} , because if she played x_0 at the first turn, then it is not hard to see that at the second turn, player **I** can always play a block-subspace $p_1 \subseteq p$ containing no term of the block-sequence $\varphi(N(x_0))$. And player **I** cannot either have a strategy in F_p to reach \mathcal{X}^c , because at the first turn, player **II** can always play a vector x_0 such that $\varphi(N(x_0))$ is a block-sequence spanning the block-subspace p . So the subspace $p_1 \approx p$ played by **I** will necessarily contain a term of the sequence $\varphi(N(x_0))$, and **II** will be able to play this term to reach her goal. \square

Remark 4.2. In the last proof, if we endow X^ω with the product of the discrete topologies on X instead of the product of the Polish topologies, then the counterexample \mathcal{X} we built is actually clopen.

We now define the metric version of Gowers spaces we will use for the rest of this paper.

Definition 4.3. An *approximate Gowers space* is a sextuple $\mathcal{G} = (P, X, d, \leq, \leq^*, \triangleleft)$, where P is a nonempty set, X is a nonempty Polish space, d is a compatible distance on X , \leq and \leq^* are two quasiorders on P , and $\triangleleft \subseteq X \times P$ is a binary relation, satisfying the same axioms 1. – 3. as in the definition of a Gowers' space and satisfying moreover the two following axioms:

4. for every $p \in P$, there exists $x \in X$ such that $x \triangleleft p$;
5. for every $x \in X$ and every $p, q \in P$, if $x \triangleleft p$ and $p \leq q$, then $x \triangleleft q$.

The relation \approx and the compatibility relation on P are defined in the same way as for a Gowers space.

For $p \in P$, define the games A_p, B_p, F_p , and G_p exactly in the same way as for Gowers spaces (see Definitions 2.2 and 3.1), except that the rules $(x_0, y_0, \dots, x_{i-1}, y_{i-1}, x_i) \triangleleft p_i$ and $(x_0, y_0, \dots, x_i, y_i) \triangleleft q_i$ in the definition of A_p and B_p and the rule $(x_0, \dots, x_i) \triangleleft p_i$ in the definition of F_p and G_p , will be naturally replaced with respectively $x_i \triangleleft p_i, y_i \triangleleft q_i$ and $x_i \triangleleft p_i$. The outcome is, there, an element of X^ω .

Observe that, with this definition, approximate Gowers spaces are always forgetful, that is, the relation \triangleleft is defined as a subset of $X \times P$ and not as a subset of $\text{Seq}(X) \times P$. This technical restriction is needed for our results (in particular in the proof of Theorem 4.6).

In the rest of this section, we fix an approximate Gowers space $\mathcal{G} = (P, X, d, \leq, \leq^*, \triangleleft)$. An important notion in the setting of approximate Gowers spaces is that of *expansion*.

Definition 4.4.

1. Let $A \subseteq X$ and $\delta > 0$. The δ -*expansion* of A is the set $(A)_\delta = \{x \in X \mid \exists y \in A \ d(x, y) \leq \delta\}$;

2. Let $\mathcal{X} \subseteq X^\omega$ and $\Delta = (\Delta_n)_{n \in \omega}$ be a sequence of positive real numbers. The Δ -expansion of \mathcal{X} is the set $(\mathcal{X})_\Delta = \{(x_n)_{n \in \omega} \in X^\omega \mid \exists (y_n)_{n \in \omega} \in \mathcal{X} \forall n \in \omega d(x_n, y_n) \leq \Delta_n\}$.

We can now define the notions of adversarially Ramsey sets and of strategically Ramsey sets in an approximate Gowers space:

Definition 4.5. Let $\mathcal{X} \subseteq X^\omega$.

1. We say that \mathcal{X} is *adversarially Ramsey* if for every sequence Δ of positive real numbers and for every $p \in P$, there exists $q \leq p$ such that either player **I** has a strategy in A_q to reach $(\mathcal{X})_\Delta$, or player **II** has a strategy in B_q to reach $(\mathcal{X}^c)_\Delta$.
2. We say that \mathcal{X} is *strategically Ramsey* if for every sequence Δ of positive real numbers and for every $p \in P$, there exists $q \leq p$ such that either player **I** has a strategy in F_q to reach \mathcal{X}^c , or player **II** has a strategy in G_q to reach $(\mathcal{X})_\Delta$.

Observe that if $\mathcal{G}_0 = (P, X, \leq, \leq^*, \triangleleft)$ is a forgetful Gowers space (where we consider \triangleleft as a subset of $X \times P$), then we can turn it into an approximate Gowers space $\mathcal{G}'_0 = (P, X, d, \leq, \leq^*, \triangleleft)$ by taking for d the discrete distance on X ($d(x, y) = 1$ for $x \neq y$). In this way, for $0 < \delta < 1$ and $A \subseteq X$ we have $(A)_\delta = A$, and for Δ a sequence of positive real numbers strictly lower than 1 and for $\mathcal{X} \subseteq X^\omega$, we have $(\mathcal{X})_\Delta = \mathcal{X}$. So for a set $\mathcal{X} \subseteq X^\omega$, the definition of being adversarially or strategically Ramsey in \mathcal{G}_0 and in \mathcal{G}'_0 coincide. Therefore, we will consider forgetful Gowers spaces as particular cases of approximate Gowers spaces.

Another interesting family of examples of approximate Gowers spaces is the following. Given a Banach space E with a Schauder basis $(e_i)_{i \in \omega}$, we can consider the *canonical approximate Gowers space over E* , $\mathcal{G}_E = (P, S_E, d, \subseteq, \subseteq^*, \epsilon)$, where P is the set of all block subspaces of E , S_E is the unit sphere of E , d the distance given by the norm, and $X \subseteq^* Y$ if and only if Y contains some finite-codimensional block subspace of X . We will see in the next section how to get Gowers' Theorems 1.2 and 1.5 from the study of this space.

The results that generalize Theorems 2.4 and 3.3 to adversarial Gowers spaces are the following:

Theorem 4.6.

1. Every Borel subset of X^ω is adversarially Ramsey;
2. Every analytic subset of X^ω is strategically Ramsey.

Proof. Observe that to prove 2., it is actually sufficient to prove the following apparently weaker result: for every $\mathcal{X} \subseteq X^\omega$ analytic, for every sequence Δ of positive real numbers and for every $p \in P$, there exists $q \leq p$ such that either player **I** has a strategy in F_q to reach $(\mathcal{X}^c)_\Delta$, or player **II** has a strategy in G_q to reach $(\mathcal{X})_\Delta$. Indeed, if \mathcal{X} is analytic,

then $(\mathcal{X})_{\frac{\Delta}{2}}$ is analytic too; so applying the last result to $(\mathcal{X})_{\frac{\Delta}{2}}$ and to the sequence $\frac{\Delta}{2}$, and using the fact that $\left(\left((\mathcal{X})_{\frac{\Delta}{2}}\right)^c\right)_{\frac{\Delta}{2}} \subseteq \mathcal{X}^c$ and $\left((\mathcal{X})_{\frac{\Delta}{2}}\right)_{\frac{\Delta}{2}} \subseteq (\mathcal{X})_{\Delta}$, we get that \mathcal{X} is strategically Ramsey.

Now let $D \subseteq X$ be a countable dense subset, and Δ be a sequence of positive real numbers. Consider the Gowers space $\mathcal{G}_{\Delta} = (P, D, \leq, \leq^*, \triangleleft_{\Delta})$, where \triangleleft_{Δ} is defined by $(y_0, \dots, y_n) \triangleleft_{\Delta} p$ if there exists $x_n \in X$ with $x_n \triangleleft p$ and $d(x_n, y_n) < \Delta_n$. To avoid confusion, denote by A_p, B_p, F_p and G_p the games in the space \mathcal{G} , and by $A_p^{\Delta}, B_p^{\Delta}, F_p^{\Delta}$ and G_p^{Δ} the games in the space \mathcal{G}_{Δ} .

If \mathcal{X} is Borel (resp. analytic) then the set $\mathcal{X} \cap D^{\omega}$ is Borel (resp. analytic) too (when D is endowed by the discrete topology), so it is adversarially (resp. strategically) Ramsey in \mathcal{G}_{Δ} . So to prove the theorem, it is enough to show that for every $p \in P$, we have that:

- (i) if player **I** has a strategy in F_p^{Δ} to reach \mathcal{X}^c , then he has a strategy in F_p to reach $(\mathcal{X}^c)_{\Delta}$;
- (ii) if player **II** has a strategy in G_p^{Δ} to reach \mathcal{X} , then she has a strategy in G_p to reach $(\mathcal{X})_{\Delta}$;
- (iii) if player **I** has a strategy in A_p^{Δ} to reach \mathcal{X} , then he has a strategy in A_p to reach $(\mathcal{X})_{\Delta}$;
- (iv) if player **II** has a strategy in B_p^{Δ} to reach \mathcal{X}^c , then she has a strategy in B_p to reach $(\mathcal{X}^c)_{\Delta}$.

We only prove (i) and (ii); the proofs of (iii) and (iv) are naturally obtained by combining the proofs of (i) and (ii).

- (i) As usual, we fix a strategy for **I** in F_p^{Δ} , enabling him to reach \mathcal{X}^c , and we describe a strategy for **I** in F_p to reach $(\mathcal{X}^c)_{\Delta}$ by simulating a play $(p_0, x_0, p_1, x_1, \dots)$ of F_p by a play $(p_0, y_0, p_1, y_1, \dots)$ of F_p^{Δ} in which **I** always plays using his strategy; we moreover suppose that the same subspaces are played by **I** in both games.

Suppose that in both games, the first n turns have been played, so the p_i 's, the x_i 's and the y_i 's are defined for $i < n$. According to his strategy, in F_p^{Δ} , **I** plays some $p_n \lesssim p$. Then we let **I** play the same p_n in F_p , and in this game, **II** answers with $x_n \in X$ such that $x_n \triangleleft p_n$. Then we choose $y_n \in D$ such that $d(x_n, y_n) < \Delta_n$; by the definition of \triangleleft_{Δ} , we have that $(y_0, \dots, y_n) \triangleleft_{\Delta} p_n$, so we can let **II** play y_n in F_p^{Δ} , and the games can continue!

Due to the choice of the strategy of **I** in F_p^{Δ} , we get that $(y_n)_{n \in \omega} \in \mathcal{X}^c$, so $(x_n)_{n \in \omega} \in (\mathcal{X}^c)_{\Delta}$ as wanted.

- (ii) We simulate a play $(p_0, x_0, p_1, x_1, \dots)$ of G_p by a play $(p_0, y_0, p_1, y_1, \dots)$ of G_p^{Δ} where **II** uses a strategy to reach \mathcal{X} , and we moreover suppose that **I** plays the

same subspaces in both games. Suppose that the first n turns of boths games have been played. In G_p , **I** plays p_n . We make **I** copy this move in G_p^Δ , and according to her strategy, **II** answers, in this game, by a $y_n \in D$ such that $(y_0, \dots, y_n) \triangleleft_\Delta p_n$. We can find $x_n \in X$ such that $x_n \triangleleft p_n$ and $d(x_n, y_n) < \Delta_n$; we let **II** play this x_n in G_p and the games continue. At the end, we have that $(y_n)_{n \in \omega} \in \mathcal{X}$, so $(x_n)_{n \in \omega} \in (\mathcal{X})_\Delta$ as wanted.

□

Say that the approximate Gowers space \mathcal{G} is *analytic* if P is an analytic subset of a Polish space, if the relation \leq is a Borel subset of P^2 , and if for every open set $U \subseteq X$, the set $\{p \in P \mid \exists x \in U \ x \triangleleft p\}$ is a Borel subset of P . Also recall that if Y is a Polish space, and if $\mathcal{F}(Y)$ is the set of all closed subsets of Y , the *Effros Borel structure* on $\mathcal{F}(Y)$ is the σ -algebra generated by the sets $\{F \in \mathcal{F}(Y) \mid F \cap U \neq \emptyset\}$ where U varies over open subsets of Y ; with this σ -algebra, $\mathcal{F}(Y)$ is a standard Borel space (see for example [10], Theorem 12.6). If P is an analytic subset of $\mathcal{F}(X)$ endowed with the Effros Borel structure, and if \subseteq and \triangleleft are respectively the inclusion and the membership relation, then \mathcal{G} is an analytic approximate Gowers space. This is, for instance, the case of the canonical approximate Gowers space \mathcal{G}_E over a Banach space E with a basis: indeed, the fact that $F \in \mathcal{F}(S_E)$ is the unit sphere of a block subspace of E can be written “there exists a block sequence $(x_i)_{i \in \omega}$ such that for every U in a countable basis of open subsets of S_E , $F \cap U \neq \emptyset$ if and only if there exists $n \in \omega$ and $(a_i)_{i < n} \in \mathbb{Q}^n \setminus \{0\}$ with $\frac{\sum_{i < n} a_i x_i}{\|\sum_{i < n} a_i x_i\|} \in U$ ”.

Observe that if \mathcal{G} is an analytic approximate Gowers space and Δ a sequence of positive real numbers, then the Gowers space \mathcal{G}_Δ defined in the proof of Theorem 4.6 is analytic. So this proof, combined with Corollaries 2.8 and 3.5, gives us the following:

Corollary 4.7. *Let Γ be a suitable class of subsets of Polish spaces. Suppose that every Γ -subset of \mathbb{R}^ω is determined. Then for every analytic approximate Gowers space $\mathcal{G} = (P, X, d, \leq, \leq^*, \triangleleft)$, we have that:*

1. every Γ -subset of X^ω is adversarially Ramsey;
2. every $\exists \Gamma$ -subset of X^ω is strategically Ramsey.

However, it is not straightforward, in the setting of approximate Gowers spaces, to get results in $ZF + DC + AD_{\mathbb{R}}$, because the proof of Theorem 4.6 uses the full axiom of choice. Indeed, since there is, in general, an uncountable number of subspaces, in the proof of (ii) (and the same will happen in the proofs of (iii) and (iv)), player **II** needs *AC* to choose x_n such that $d(x_n, y_n) < \Delta_n$ and $x_n \triangleleft p_n$. However, under a slight restriction, we can get a positive result. Define the notion of an *effective approximate Gowers space* exactly in the same way as for effective Gowers spaces. Effective forgetful Gowers spaces are obviously effective when seen as approximate Gowers spaces, but also, the canonical approximate Gowers space \mathcal{G}_E is effective (this can be shown in the same way as for the

Rosendal space). If \mathcal{G} is an effective approximate Gowers space and Δ a sequence of positive real numbers, then the Gowers space \mathcal{G}_Δ defined in the proof of Theorem 4.6 is also effective. And we have:

Corollary 4.8 ($ZF + DC + AD_{\mathbb{R}}$). *Let $\mathcal{G} = (P, X, d, \leq, \leq^*, \triangleleft)$ be an effective approximate Gowers space such that P is a subset of a Polish space, and such that for every $p \in P$, the set $\{x \in X \mid x \triangleleft p\}$ is closed in X . Then every subset of X^ω is adversarially Ramsey and strategically Ramsey.*

Proof. We follow the proof of Theorem 4.6, using Corollaries 2.10 and 3.5 to get that the set $\mathcal{X} \cap D^\omega$ is adversarially Ramsey and strategically Ramsey in \mathcal{G}_Δ . The only thing to do is to verify that the proofs of (i)–(iv) can be carried out with only DC instead of AC ; as previously, we only do it for (i) and (ii). In the proof of (i), we have to be able to choose $y_n \in D$ such that $d(x_n, y_n) < \Delta_n$; this can be done by fixing, at the beginning of the proof, a well-ordering of D , and by choosing, each time, the least such y_n . In the proof of (ii), the difficulty is to choose x_n ; so we have to prove that given $p \in P$, $n \in \omega$, and $y \in D$, if there exists $x \in X$ with $x \triangleleft p$ and $d(x, y) < \Delta_n$, then we are able to choose such an x without using AC .

Using countable choices, for every $y \in D$ and $n \in \omega$, we choose $f_{y,n} : \omega^\omega \rightarrow B(y, \Delta_n)$ a continuous surjection. Given p, n and y as in the previous paragraph, we can let $F = \{u \in \omega^\omega \mid f_{y,n}(u) \triangleleft p\}$, a closed subset of ω^ω . Consider $T \subseteq \omega^{<\omega}$ the unique pruned tree such that $F = [T]$. Then we can let u be the leftmost branch of T and let $x = f_{y,n}(u)$. □

We now introduce the pigeonhole principle in an approximate Gowers space and its consequences. We actually only need an approximate pigeonhole principle in this setting. For $q \in P$ and $A \subseteq X$, we abusively write $q \subseteq A$ to say that $\forall x \in X (x \triangleleft q \Rightarrow x \in A)$.

Definition 4.9. The approximate Gowers space \mathcal{G} is said to satisfy the *pigeonhole principle* if for every $p \in P$, $A \subseteq X$, and $\delta > 0$ there exists $q \leq p$ such that either $q \subseteq A^c$, or $q \subseteq (A)_\delta$.

For example, by Theorem 1.4, the canonical approximate Gowers space \mathcal{G}_E satisfies the pigeonhole principle if and only if E is c_0 -saturated.

As for Gowers spaces, we have the following proposition:

Proposition 4.10. *Suppose that the approximate Gowers space \mathcal{G} satisfies the pigeonhole principle. Let $\mathcal{X} \subseteq X^\omega$, $p \in P$ and Δ be a sequence of positive real numbers. If player **II** has a strategy in G_p to reach \mathcal{X} , then there exists $q \leq p$ such that player **I** has a strategy in F_q to reach $(\mathcal{X})_\Delta$.*

Before proving this proposition, let us make some remarks. Using again the fact that $\left((\mathcal{X})_{\frac{\Delta}{2}}\right)_{\frac{\Delta}{2}} \subseteq (\mathcal{X})_\Delta$, we deduce from Proposition 4.10 the following corollary:

Corollary 4.11. *Suppose that the approximate Gowers space \mathcal{G} satisfies the pigeonhole principle. Let $\mathcal{X} \subseteq X^\omega$ be a strategically Ramsey set. Then for every $p \in P$ and every sequence Δ of positive real numbers, there exists $q \leq p$ such that in F_q , player **I** either has a strategy to reach \mathcal{X}^c , or has a strategy to reach $(\mathcal{X})_\Delta$.*

Conversely, it is not hard to see that if the conclusion of Corollary 4.11 holds for all sets of the form $\{(x_n)_{n \in \omega} \in X^\omega \mid x_0 \in F\}$, where $F \subseteq X$ is closed, then the space \mathcal{G} satisfies the pigeonhole principle.

Also observe that if \mathcal{G}_0 is a forgetful Gowers space, and if \mathcal{G}'_0 is the associated approximate Gowers space, then the pigeonhole principle in \mathcal{G}_0 is equivalent to the pigeonhole principle in \mathcal{G}'_0 , and Proposition 4.10 and Corollary 4.11 are respectively the same as Proposition 3.9 and Corollary 3.10.

We now prove Proposition 4.10.

Proof of Proposition 4.10. Unlike the previous results about approximate Gowers spaces, here we cannot deduce this result from its exact version; thus, we adapt the proof of Proposition 3.9. To save notation, we show that there exists $q \leq p$ such that **I** has a strategy in F_q to reach $(\mathcal{X})_{3\Delta}$.

We fix τ a strategy for **II** in G_p to reach \mathcal{X} . As usual, call a *state* a partial play of G_p either empty or ending with a move of **II**, during which **II** always plays according to her strategy. Say that a state *realises* a sequence $(x_0, \dots, x_{n-1}) \in X^{<\omega}$ if it has the form $(p_0, x_0, \dots, p_{n-1}, x_{n-1})$. The *length* of the state \mathcal{J} , denoted by $|\mathcal{J}|$, is the length of the sequence it realises. Define in the same way the notion of a *total state* (which is a total play of G_p) and of realisation for a total state; if an infinite sequence is realised by a total state, then it belongs to \mathcal{X} . Say that a point $x \in X$ is *reachable* from a state \mathcal{J} if there exists $r \leq p$ such that $\tau(\mathcal{J} \hat{\ } r) = x$. Denote by $A_{\mathcal{J}}$ the set of all points that are reachable from the state \mathcal{J} . We will use the following fact.

Fact 4.12. *For every state \mathcal{J} and for every $q \leq p$, there exists $r \leq q$ such that $r \subseteq (A_{\mathcal{J}})_{\Delta_{|\mathcal{J}|}}$.*

Proof. Otherwise, by the pigeonhole principle, there would exist $r \leq q$ such that $r \subseteq (A_{\mathcal{J}})^c$. But then **I** could play r after the partial play \mathcal{J} , and **II** would answer, according to her strategy, by $x = \tau(\mathcal{J} \hat{\ } r)$ that should satisfy $x \triangleleft r$. Since $r \subseteq (A_{\mathcal{J}})^c$, this would imply that $x \in (A_{\mathcal{J}})^c$. But we also have, by the definition of $A_{\mathcal{J}}$, that $x \in A_{\mathcal{J}}$, a contradiction. \square

For two sequences $s, t \in X^{\leq \omega}$ of the same length, denote by $d(s, t) \leq \Delta$ the fact that for every $i < |s|$, we have $d(s_i, t_i) \leq \Delta_i$. Let $D \subseteq X$ be a countable dense set and let $(s_n)_{n \in \omega}$ be an enumeration of $D^{<\omega}$ such that if $s_m \subseteq s_n$, then $m \leq n$. We define, for some $n \in \omega$, a state \mathcal{J}_n realising a sequence t_n satisfying $d(s_n, t_n) \leq 2\Delta$, by induction in

the following way: $\mathcal{J}_0 = \emptyset$ and for $n \geq 1$, letting $s_n = s_m \hat{\ } y$ for some $m < n$ and some $y \in X$,

- if \mathcal{J}_m has been defined and if there exists $z \in X$ reachable from \mathcal{J}_m such that $d(y, z) \leq 2\Delta_{|\mathcal{J}_m|}$, then choose a $r \leq p$ such that $z = \tau(\mathcal{J}_m \hat{\ } r)$ and put $t_n = t_m \hat{\ } z$ and $\mathcal{J}_n = \mathcal{J}_m \hat{\ } (r, z)$,
- otherwise, \mathcal{J}_n is not defined.

Observe that if \mathcal{J}_n is defined and if $s_m \subseteq s_n$, then \mathcal{J}_m is defined, and we have $\mathcal{J}_m \subseteq \mathcal{J}_n$ and $t_m \subseteq t_n$.

We now define a \leq -decreasing sequence $(q_n)_{n \in \omega}$ of elements of P in the following way: $q_0 = p$ and

- if \mathcal{J}_n is defined, then q_{n+1} is the result of the application of Fact 4.12 to \mathcal{J}_n and q_n ;
- $q_{n+1} = q_n$ otherwise.

Finally, let $q \leq p$ be such that for every $n \in \omega$, $q \leq^* q_n$. We will show that **I** has a strategy in F_q to reach $(\mathcal{X})_{3\Delta}$. We describe this strategy on the following play of F_q :

$$\begin{array}{llll} \mathbf{I} & u_0 & u_1 & \dots \\ \mathbf{II} & x_0 & x_1 & \dots \end{array}$$

We moreover suppose that at the same time as this game is played, we build a sequence $(n_i)_{i \in \omega}$ of integers, with $n_0 = 0$ and n_i being defined during the i^{th} turn, such that $(s_{n_i})_{i \in \omega}$ is increasing and for every $i \in \omega$, $|s_{n_i}| = i$, \mathcal{J}_{n_i} is defined, and $d(s_{n_i}, (x_0, \dots, x_{i-1})) \leq \Delta$. This will be enough to conclude: indeed, $\bigcup_{i \in \omega} \mathcal{J}_{n_i}$ will be a total state realising the sequence $\bigcup_{i \in \omega} t_{n_i}$, showing that this sequence belongs to \mathcal{X} ; and since $d(\bigcup_{i \in \omega} t_{n_i}, (x_i)_{i \in \omega}) \leq d(\bigcup_{i \in \omega} t_{n_i}, \bigcup_{i \in \omega} s_{n_i}) + d(\bigcup_{i \in \omega} s_{n_i}, (x_i)_{i \in \omega}) \leq 3\Delta$, we will have that $(x_i)_{i \in \omega} \in (\mathcal{X})_{3\Delta}$.

Suppose that the i^{th} turn of the game has just been played, so the sequence (x_0, \dots, x_{i-1}) and the integers n_0, \dots, n_i has been defined. Then by construction of q_{n_i+1} , we have that $q_{n_i+1} \subseteq (A_{\mathcal{J}_{n_i}})_{\Delta_{|\mathcal{J}_{n_i}|}}$. We let **I** play some u_i such that $u_i \lesssim q$ and $u_i \leq q_{n_i+1}$. Then $u_i \subseteq (A_{\mathcal{J}_{n_i}})_{\Delta_{|\mathcal{J}_{n_i}|}}$. Now, suppose that **II** answers by x_i . Then we choose a $y_i \in D$ such that $d(x_i, y_i) \leq \Delta_i$ and we choose n_{i+1} in such a way that $s_{n_{i+1}} = s_{n_i} \hat{\ } y_i$. So we have that $y_i \in (A_{\mathcal{J}_{n_i}})_{2\Delta_{|\mathcal{J}_{n_i}|}}$; this shows that $\mathcal{J}_{n_{i+1}}$ has been defined. Moreover we have $d(s_{n_{i+1}}, (x_0, \dots, x_i)) \leq \Delta$ as wanted, what ends the proof. \square

Again, this proof can be done in $ZF + DC$, even if the space \mathcal{G} is not supposed effective.

5 Eliminating the asymptotic game

In this section, we provide a tool to deduce, from a statement of the form “player **I** has a strategy in F_p to reach \mathcal{X} ”, a conclusion of the form “in some subspace, every sequence satisfying some structural condition is in \mathcal{X} ”. This tool can be seen as a generalization of Lemma 3.7. It will allow us to get, from Ramsey results with game-theoretical conclusions, stronger results having the same form as Mathias–Silver’s theorem or Gowers’ theorem.

We will actually not add any structure on the set of points, but rather provide a tool enabling, in each concrete situation, to build this structure in the way we want. Our result could be stated in the setting of approximate Gowers spaces, but we prefer to state it in the more general setting of *approximate asymptotic spaces*, since it could be useful in itself in situations where we have no natural Gowers space structure.

Definition 5.1. An *approximate asymptotic space* is a quintuple $\mathcal{A} = \{P, X, d, \lesssim, \triangleleft\}$, where P is a nonempty set, (X, d) is a nonempty separable metric space, \lesssim is a quasiorder on P , and $\triangleleft \subseteq X \times P$ is a binary relation, satisfying the following properties:

1. for every $p, q, r \in P$, if $q \lesssim p$ and $r \lesssim p$, then there exists $u \in P$ such that $u \lesssim q$ and $u \lesssim r$;
2. for every $p \in P$, there exists $x \in X$ such that $x \triangleleft p$;
3. for every $x \in X$ and every $p, q \in P$, if $x \triangleleft p$ and $p \lesssim q$, then $x \triangleleft q$.

Every approximate Gowers space has a natural structure of approximate asymptotic space. In an approximate asymptotic space, define the notion of expansion, and the asymptotic game, in the same way as in an approximate Gowers space.

In the rest of this section, we fix $\mathcal{A} = \{P, X, d, \lesssim, \triangleleft\}$ an approximate asymptotic space. Recall that a subset of X is said to be *precompact* if its closure in X is compact. In what follows, for $K \subseteq X$ and $p \in P$, we abusively write $K \triangleleft p$ to say that the set $\{x \in K \mid x \triangleleft p\}$ is dense in K .

Definition 5.2. A *system of precompact sets* for \mathcal{A} is a set \mathcal{K} of precompact subsets of X , equipped with an associative binary operation \oplus , satisfying the following property: for every $p \in P$, and for every $K, L \in \mathcal{K}$, if $K \triangleleft p$ and $L \triangleleft p$, then $K \oplus L \triangleleft p$.

If (\mathcal{K}, \oplus) is a system of precompact sets for \mathcal{A} and if $(K_n)_{n \in \omega}$ is a sequence of elements of \mathcal{K} , then:

- for $A \subseteq \omega$ finite, denote by $\bigoplus_{n \in A} K_n$ the sum $K_{n_1} \oplus \dots \oplus K_{n_k}$, where n_1, \dots, n_k are the elements of A taken in increasing order;
- a *block sequence* of (K_n) is, by definition, a sequence $(x_i)_{i \in \omega} \in X^\omega$ for which there exists an increasing sequence of nonempty sets of integers $A_0 < A_1 < A_2 < \dots$ such that for every $i \in \omega$, we have $x_i \in \bigoplus_{n \in A_i} K_n$.

Denote by $\text{bs}((K_n)_{n \in \omega})$ the set of all block sequences of (K_n) .

We can already give some examples. For the Mathias–Silver space \mathcal{N} , let $\mathcal{K}_{\mathcal{N}}$ be the set of all singletons, and define the operation $\oplus_{\mathcal{N}}$ by $\{m\} \oplus_{\mathcal{N}} \{n\} = \{\max(m, n)\}$. Then $(\mathcal{K}_{\mathcal{N}}, \oplus_{\mathcal{N}})$ is a system of precompact sets. If $(m_i)_{i \in \omega}$ is an increasing sequence of integers, then the block sequences of $(\{m_i\})_{i \in \omega}$ are exactly the subsequences of (m_i) .

Now, for a Banach space E with a basis, consider the canonical approximate Gowers space \mathcal{G}_E . Let \mathcal{K}_E be the set of all unit spheres of finite-dimensional subspaces of E . Define the operation \oplus_E on \mathcal{K}_E by $S_F \oplus_E S_G = S_{F+G}$. Then $(\mathcal{K}_E, \oplus_E)$ is a system of precompact sets for \mathcal{G}_E . If $(x_n)_{n \in \omega}$ is a (normalized) block sequence of E , then for every n , $S_{\mathbb{R}x_n} = \{x_n, -x_n\}$ is in \mathcal{K}_E , and the block sequences of $(S_{\mathbb{R}x_n})_{n \in \omega}$ in the sense of \mathcal{K} are exactly the (normalized) block sequences of (x_n) in the Banach-theoretical sense. More generally, it is often useful to study the block sequences of sequences of the form $(S_{F_n})_{n \in \omega}$, where $(F_n)_{n \in \omega}$ is a FDD of a closed, infinite-dimensional subspace F of E (that is, a sequence such that every $x \in F$ can be written in a unique way as a sum $\sum_{n=0}^{\infty} x_n$, where for every n , $x_n \in F_n$).

In general, in an asymptotic space, a sequence $(K_n)_{n \in \omega}$ of elements of a system of precompact sets can be seen as another kind of subspace. Sometimes, some subspaces of the type $(K_n)_{n \in \omega}$ can be represented as elements of P ; that is, for example, the case in the Mathias–Silver space and in the canonical approximate Gowers space over a Banach space with a basis, as we just saw. We now introduce a theorem enabling us to build sequences $(K_n)_{n \in \omega}$ such that $\text{bs}((K_n)_{n \in \omega}) \subseteq \mathcal{X}$, knowing that player **I** has a strategy in an asymptotic game to reach \mathcal{X} . First, we have to define a new game.

Definition 5.3. Let (\mathcal{K}, \oplus) be a system of precompact sets for the space \mathcal{A} , and $p \in P$. The *strong asymptotic game below p* , denoted by SF_p , is defined as follows:

$$\begin{array}{llll} \text{I} & p_0 & p_1 & \dots \\ \text{II} & K_0 & K_1 & \dots \end{array}$$

where the K_n 's are elements of \mathcal{K} , and the p_n 's are elements of P . The rules are the following:

- for **I**: for all $n \in \omega$, $p_n \lesssim p$;
- for **II**: for all $n \in \omega$, $K_n \triangleleft p_n$.

The outcome of the game is the sequence $(K_n)_{n \in \omega} \in \mathcal{K}^{\omega}$.

Theorem 5.4. Let (\mathcal{K}, \oplus) be a system of precompact sets on the space \mathcal{A} , $p \in P$, $\mathcal{X} \subseteq X^{\omega}$, and Δ be a sequence of positive real numbers. Suppose that player **I** has a strategy in F_p to reach \mathcal{X} . Then he has a strategy in SF_p to build a sequence $(K_n)_{n \in \omega}$ such that $\text{bs}((K_n)_{n \in \omega}) \subseteq (\mathcal{X})_{\Delta}$.

Proof. For each $K \in \mathcal{K}$, each $q \in P$ such that $K \triangleleft q$, and each $i \in \omega$, let $\mathfrak{N}_{i,q}(K)$ be a Δ_i -net in K (that is, a finite subset of K such that $K \subseteq (\mathfrak{N}_{i,q}(K))_{\Delta_i}$), such that for every $x \in \mathfrak{N}_{i,q}(K)$, we have $x \triangleleft q$. We fix τ a strategy for **I** in F_p , enabling him to reach

\mathcal{X} . As in the proofs of Fact 3.4 and Lemma 3.7, we consider that in F_p , **II** is allowed to play against the rules, but that she immediately loses if she does; so we will view τ as a mapping from $X^{<\omega}$ to P , such that for every $s \in X^{<\omega}$, we have $\tau(s) \lesssim p$.

Let us describe a strategy for **I** in SF_p on a play $(p_0, K_0, p_1, K_1, \dots)$ of this game. Suppose that the first n turns have been played, so the p_j 's and the K_j 's, for $j < n$, are defined. Moreover suppose that the sequence $(p_j)_{j < n}$ is \lesssim -decreasing. Let $S_{(K_0, \dots, K_{n-1})} \subseteq X^{<\omega}$ be the set of all finite sequences (y_0, \dots, y_{k-1}) satisfying the following property: there exists an increasing sequence $A_0 < \dots < A_{k-1}$ of nonempty subsets of n such that for every $i < k$, we have $y_i \in \mathfrak{N}_{i, p_{\min(A_i)}}(\bigoplus_{j \in A_i} K_j)$. Then $S_{(K_0, \dots, K_{n-1})}$ is finite and for every $s \in S_{(K_0, \dots, K_{n-1})}$, we have $\tau(s) \lesssim p$, so by iterating the axiom 1. in the definition of an approximate asymptotic space, we can find $p_n \lesssim p$ such that for every $s \in S_{(K_0, \dots, K_{n-1})}$, we have $p_n \lesssim \tau(s)$. Moreover, if $n \geq 1$, we can choose p_n such that $p_n \lesssim p_{n-1}$. The strategy of **I** will consist in playing this p_n .

Now suppose that this play has been played completely; we show that $\text{bs}((K_n)_{n \in \omega}) \subseteq (\mathcal{X})_\Delta$. Let $(x_i)_{i \in \omega}$ be a block sequence of (K_n) and $A_0 < A_1 < \dots$ be a sequence of nonempty subsets of ω such that for every i , we have $x_i \in \bigoplus_{n \in A_i} K_n$. For every $i \in \omega$, we have $(\bigoplus_{n \in A_i} K_n) \triangleleft p_{\min(A_i)}$, so $\mathfrak{N}_{i, p_{\min(A_i)}}(\bigoplus_{n \in A_i} K_n)$ has been defined and we can choose a y_i in it such that $d(x_i, y_i) \leq \Delta_i$. We have to show that $(x_i)_{i \in \omega} \in (\mathcal{X})_\Delta$, so it is enough to show that $(y_i)_{i \in \omega} \in \mathcal{X}$. Knowing that τ is a strategy for **I** in F_p to reach \mathcal{X} , it is enough to show that, letting $q_i = \tau(y_0, \dots, y_{i-1})$ for all i , in the following play of F_p , **II** always respects the rules:

$$\begin{array}{cccc} \text{I} & q_0 & q_1 & \dots \\ \text{II} & y_0 & y_1 & \dots \end{array}$$

In other words, we have to show that for all $k \in \omega$, we have $y_k \triangleleft q_k$.

So let $k \in \omega$. We let $n_0 = \min A_k$. Since the sets A_0, \dots, A_{k-1} are subsets of n_0 , we have that $(y_0, \dots, y_{k-1}) \in S_{(K_0, \dots, K_{n_0-1})}$, and therefore $p_{n_0} \lesssim \tau(y_0, \dots, y_{k-1}) = q_k$. But $y_k \in \mathfrak{N}_{k, p_{n_0}}(\bigoplus_{n \in A_k} K_n)$, so $y_k \triangleleft p_{n_0}$, so $y_k \triangleleft q_k$, as wanted. \square

Again, under slight restrictions, we can prove Theorem 5.4 without using the full axiom of choice. Say that the approximate asymptotic space \mathcal{A} is *effective* if there exist a function $f : P^2 \rightarrow P$ such that for every $q, r \in P$, if there exist $p \in P$ such that $q \lesssim p$ and $r \lesssim p$, then we have $f(q, r) \lesssim q$ and $f(q, r) \lesssim r$. Effective approximate Gowers spaces, when seen as approximate asymptotic spaces, are effective. We will show that if \mathcal{A} is an effective approximate asymptotic space, if X is an analytic subset of a Polish space, if for every $p \in P$, the set $\{x \in X \mid x \triangleleft p\}$ is closed in X , and if every element of \mathcal{K} is compact, then Theorem 5.4 for \mathcal{A} and \mathcal{K} can be shown in $ZF + DC$. In the proof of Theorem 5.4, AC is only used:

- to choose p_n such that for every $s \in S_{(K_0, \dots, K_{n-1})}$, we have $p_n \lesssim \tau(s)$, and such that $p_n \lesssim p_{n-1}$ if $n \geq 1$;
- to choose the nets $\mathfrak{N}_{i, q}(K)$;

- and to choose $y_i \in \mathfrak{N}_{i, p_{\min(A_i)}} \left(\bigoplus_{n \in A_i} K_n \right)$ such that $d(x_i, y_i) \leq \Delta_i$.

The choice of the p_n 's can be done without AC as soon as the space \mathcal{A} is effective. For the choice of the nets and of the y_i 's, first, observe that, given $K \in \mathcal{K}$ and $q \in P$, since $\{x \in X \mid x \triangleleft q\}$ is closed in X , we have that $K \triangleleft q$ if and only if $K \subseteq \{x \in X \mid x \triangleleft q\}$; so $\mathfrak{N}_{i,q}(K)$ can actually be an arbitrary Δ_i -net in K , and does not need to depend on q . Thus, to be able to choose these nets and the y_i 's without AC , it is enough to show that we can choose, without AC , a Δ_i -net $\mathfrak{N}_i(K)$ in K and a wellordering $<_{i,K}$ on it, for every $K \in \mathcal{K}$ and every $i \in \omega$. This can be done in the following way. Let $\varphi : \omega^\omega \rightarrow X$ be a continuous surjection. If $K \in \mathcal{K}$, then $\varphi^{-1}(K)$ has the form $[T_K]$, where T_K is a pruned tree on ω . We can easily build, without choice, a countable dense subset of $[T_K]$, for example the set of all the u_s 's where for every $s \in T_K$, u_s is the leftmost branch of T_K satisfying $s \subseteq u_s$. Since T_K can naturally be wellordered, then this dense subset can also be wellordered. Pushing forward by φ , this enables us to get, for every $K \in \mathcal{K}$, a countable dense subset $D_K \subseteq K$ with a wellordering $<_K$. From this we can naturally wellorder the set of all finite subsets of D_K , take for $\mathfrak{N}_i(K)$ the least finite subset of D_K that is a Δ_i -net in K and take for $<_{i,K}$ the restriction of $<_K$ to $\mathfrak{N}_i(K)$.

Theorem 5.4, combined with the results of the last section and with the last remark, gives us the following corollary:

Corollary 5.5 (Abstract Gowers' theorem). *Let $\mathcal{G} = (P, X, d, \leq, \leq^*, \triangleleft)$ be an approximate Gowers space, equipped with a system of precompact sets (\mathcal{K}, \bigoplus) . Let $\mathcal{X} \subseteq X^\omega$, and suppose that we are in one of the following situations:*

- we work in ZFC , and \mathcal{X} is analytic;
- we work in ZFC , \mathcal{G} is analytic and \mathcal{X} is $\exists\Gamma$, for some suitable class Γ of subsets of Polish spaces such that every Γ -subset of \mathbb{R}^ω is determined;
- we work in $ZF + DC + AD_{\mathbb{R}}$, the space \mathcal{G} is effective, P is a subset of Polish space, for every $p \in P$, the set $\{x \in X \mid x \triangleleft p\}$ is closed in X , and every element of \mathcal{K} is compact.

Let $p \in P$ and Δ be a sequence of positive real numbers. Then there exists $q \leq p$ such that:

- either player **I** has a strategy in SF_q to build a sequence $(K_n)_{n \in \omega}$ such that $\text{bs}((K_n)_{n \in \omega}) \subseteq \mathcal{X}^c$;
- or player **II** has a strategy in G_q to reach $(\mathcal{X})_\Delta$.

Moreover, if \mathcal{G} satisfies the pigeonhole principle, then the second conclusion can be replaced with the following stronger one: player **I** has a strategy in SF_q to build a sequence $(K_n)_{n \in \omega}$ such that $\text{bs}((K_n)_{n \in \omega}) \subseteq (\mathcal{X})_\Delta$.

Now see how to deduce Mathias–Silver's theorem, Gowers' Theorem 1.2, and Gowers' theorem for c_0 (Theorem 1.5) from Corollary 5.5.

- For Mathias–Silver’s theorem, work in the Mathias–Silver space \mathcal{N} with the system $(\mathcal{K}_{\mathcal{N}}, \oplus_{\mathcal{N}})$ of precompact sets introduced before. Let M be an infinite set of integers, and $\mathcal{X} \subseteq [\omega]^\omega$ be analytic, that we will consider as a subset of ω^ω by identifying infinite subsets of ω with increasing sequences of integers. Applying Corollary 5.5 to \mathcal{X} , to M , and to the constant sequence equal to $\frac{1}{2}$, we get an infinite $N \subseteq M$ such that either **I** has a strategy in SF_N to build $(\{n_i\})_{i \in \omega}$ with $\text{bs}((\{n_i\})_{i \in \omega}) \subseteq \mathcal{X}$, or he has one to build $(\{n_i\})_{i \in \omega}$ with $\text{bs}((\{n_i\})_{i \in \omega}) \subseteq \mathcal{X}^c$. Observe that in SF_N , **II** can always play in such a way that the sequence $(n_i)_{i \in \omega}$ is increasing. So in the first case, we get an increasing sequence $(n_i)_{i \in \omega}$ of elements of N such that every block sequence of $(\{n_i\})_{i \in \omega}$ belongs to \mathcal{X} , or in other words, such that every infinite subset of $\{n_i \mid i \in \omega\}$ belongs to \mathcal{X} ; and in the second case, in the same way, we get an infinite subset of N every infinite subset of whose belongs to \mathcal{X}^c .
- For Gowers’ theorem, let E be a Banach space with a Schauder basis and work in the canonical approximate Gowers space \mathcal{G}_E with the system $(\mathcal{K}_E, \oplus_E)$ of precompact sets introduced before. Given $Y \in P$, in SF_Y , whatever **I** plays, **II** can always ensure that the outcome will have the form $(S_{\mathbb{R}y_n})_{n \in \omega}$, where $(y_n)_{n \in \omega}$ is a block sequence. So given $\mathcal{X} \subseteq [E]$ analytic, $X \subseteq E$ a block subspace, and Δ a sequence of positive real numbers, Corollary 5.5 gives us either a block sequence $(y_n)_{n \in \omega}$ in X such that $\text{bs}((S_{\mathbb{R}y_n})_{n \in \omega}) \subseteq \mathcal{X}^c$, or a block subspace $Y \subseteq X$ such that **II** has a strategy in G_Y to reach $(\mathcal{X})_{\frac{\Delta}{2}}$. In the first case, denoting by Y the block subspace generated by the sequence (y_n) , this precisely means that $[Y] \subseteq \mathcal{X}^c$. In the second case, we have to be careful because the Gowers’ game of the space \mathcal{G}_E is not exactly the same as this defined in the introduction: in the one of the introduction, player **II** is required to play vectors with finite support forming a block sequence, while in the one of \mathcal{G}_E , she can play any vector in the unit sphere of the subspace played by **I**. This is not a real problem as, by perturbing the vectors given by her strategy, player **II** can reach \mathcal{X}_Δ playing vectors with finite support; and without loss of generality, we can assume that the subspace Y_n played by **I** at the $(n+1)^{\text{th}}$ turn is chosen small enough to force **II** to play a y_n such that $\text{supp}(y_{n-1}) < \text{supp}(y_n)$.
- To deduce Gowers’ theorem for c_0 , the method is the same except that this time, \mathcal{G}_E satisfies the pigeonhole principle so Corollary 5.5 will give us a conclusion with a strong asymptotic game in both sides.

To finish this section, let us show on an example that the hypothesis “**I** has a strategy in F_p to reach \mathcal{X} ” does not always imply that for some subspace q , every sequence below q satisfying some natural structural condition (for instance, being block) is in \mathcal{X}_Δ . To see this, consider the Rosendal space $\mathcal{R}_K = (P, E \setminus \{0\}, \subseteq, \subseteq^*, \in)$ over a field K . We have the following fact:

Fact 5.6. *Suppose that K is a finite field. Let $\mathcal{X} \subseteq (E \setminus \{0\})^\omega$ and $X \in P$, and suppose that **I** has a strategy in F_X to reach \mathcal{X} . Then there exists a block subspace $Y \subseteq X$ such*

that every block sequence of Y is in \mathcal{X} .

Proof. Let \mathcal{K} be the set of all sets of the form $F \setminus \{0\}$, where F is a finite-dimensional subspace of E . Since the field K is finite, the elements of \mathcal{K} are finite too. For $F, G \subseteq E$ finite-dimensional, let $(F \setminus \{0\}) \oplus (G \setminus \{0\}) = (F + G) \setminus \{0\}$. Then (\mathcal{K}, \oplus) is a system of precompact sets. The conclusion follows from Theorem 5.4 applied to this system, using the same method as previously. \square

Observe that this proof does not work when K is infinite, and actually, this result is false. Let us give a counterexample. Let $(e_i)_{i < \omega}$ be the basis of E with respect to whose block subspaces are taken, and let $\varphi : K^* \rightarrow \omega$ be a bijection. For $x \in E \setminus \{0\}$, let $N(x)$ be the first nonzero coordinate of x . Let $\mathcal{Y} = \{(x, y) \in (E \setminus \{0\})^2 \mid \varphi(N(x)) < \min \text{supp}(y)\}$ and $\mathcal{X} = \{(x_n)_{n \in \omega} \in (E \setminus \{0\})^\omega \mid (x_0, x_1) \in \mathcal{Y}\}$. Then player **I** has a strategy in F_E to reach \mathcal{X} ; this strategy is illustrated on the following diagram:

$$\begin{array}{lll} \mathbf{I} & E & \text{span}(\{e_i \mid i > \varphi(N(x))\}) \\ \mathbf{II} & x & y \end{array}$$

But there is no block subspace Y of E such that every block sequence in Y belongs to \mathcal{X} . Indeed, given $Y \subseteq E$ a block subspace generated by a block sequence $(y_n)_{n \in \omega}$, we can take $\lambda \in K$ such that $\varphi(N(\lambda y_0)) = \min \text{supp}(y_1)$, and we have $(\lambda y_0, y_1, y_2, \dots) \notin \mathcal{X}$. Note that, just like the counterexample to the pigeonhole principle presented in section 3, this kind of counterexamples cannot be avoided by working in the projective Rosendal space: in this space, a similar construction works taking for $N(x)$ the quotient of the last nonzero coordinate of x by its first nonzero coordinate.

Therefore, in lots of cases, the “subspaces” of the form $(K_n)_{n \in \omega}$, where the K_n ’s are elements of a system of precompact sets, cannot always be identified with “genuine” subspaces (i.e. elements of P): we always need a form of compactness for that.

Acknowledgments. I would like to thank Stevo Todorčević who proposed me to work on the adversarial Ramsey principle and made me know Kastanas’ game, thereby giving me all the tools to prove it. I would also like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme “Mathematical, Foundational and Computational Aspects of the Higher Infinite” where work on this paper was undertaken. This work was supported by EPSRC grant numbers EP/K032208/1 and EP/R014604/1. In particular, I met there Jordi López-Abad who suggested me to give an approximate version of my result in a metric setting, what I do in section 4; I am grateful to him for that. I also thank Christian Rosendal who suggested me research directions in order to apply this result, and Valentin Ferenczi who gave me a very nice example of approximate Gowers space, that will be studied in a forthcoming paper with him and Wilson Cuellar Carrera. Finally, I would like to thank the anonymous referee for his useful comments that helped me much to improve the presentation of my results.

References

- [1] Argyros, S. A., Todorčević, S.: Ramsey methods in analysis. Adv. Courses in Math. CRM Barcelona, Birkhäuser Verlag, Basel (2005)

- [2] Bagaria, J., López-Abad, J.: Determinacy and weakly Ramsey sets in Banach spaces. *Trans. Amer. Math. Soc.* **354**, 1327–1349 (2002)
- [3] de Rancourt, N.: More on adversarially and strategically Ramsey sets. In preparation.
- [4] Ferenczi, V., Rosendal, C.: Banach spaces without minimal subspaces. *J. Funct. Anal.* **257**, 149–193 (2009)
- [5] Friedman, H. M.: Higher set theory and mathematical practice. *Ann. Math. Logic* **2**, 325–357 (1971)
- [6] Gowers, W. T.: Lipschitz functions on classical spaces. *Europ. J. Combin.* **13**, 141–151 (1992)
- [7] Gowers, W. T.: An infinite Ramsey theorem and some Banach-space dichotomies. *Ann. Math.* **156**, 797–833 (2002)
- [8] Harrington, L. A., Kechris, A. S.: On the determinacy of games on ordinals. *Ann. Math. Logic* **20**, 109–154 (1981)
- [9] Kastanas, I. G.: On the Ramsey property for sets of reals. *J. Symb. Logic* **48**, 1035–1045 (1983)
- [10] Kechris, A. S.: Classical descriptive set theory. Graduate Texts in Math. 156, Springer-Verlag, New York (1995)
- [11] Komorowski, R. A., Tomczak-Jaegermann, N.: Banach spaces without local unconditional structure. *Israel J. Math.* **89**, 205–226 (1995)
- [12] Mansfield, R.: A footnote to a theorem of Solovay on recursive encodability. In: *Logic Colloquium '77* (Macintyre, A., Pacholski, L., Paris, J., eds.), North-Holland, Amsterdam, 195–198 (1978)
- [13] Martin, D. A.: Measurable cardinals and analytic games. *Fund. Math.* **66**, 287–291 (1970)
- [14] Mathias, A. R. D.: On a generalization of Ramsey’s theorem. *Notices Amer. Math. Soc.* **15**, 931 (1968 (Abstract 68T-E19))
- [15] Milman, V. D.: Geometric theory of Banach spaces, part II: Geometry of the unit sphere. *Russian Math. Surveys* (6) **26**, 79–163 (1971)
- [16] Odell, E., Schlumprecht, T.: The distortion problem. *Acta Math.* **173**, 259–281 (1994)
- [17] Pelczar, A. M.: Subsymmetric sequences and minimal spaces. *Proc. Amer. Math. Soc.* **131**, 765–771 (2003)
- [18] Rosendal, C.: An exact Ramsey principle for block sequences. *Coll. Math.* **61**, 25–36 (2010)
- [19] Rosendal, C.: Determinacy of adversarial Gowers games. *Fund. Math.* **227**, 163–178 (2014)
- [20] Silver, J.: Every analytic set is Ramsey. *J. Symb. Logic* **35**, 60–64 (1970)

- [21] Todorčević, S.: Introduction to Ramsey spaces. Ann. Math. Studies 174, Princeton University Press, Princeton (2010)
- [22] Woodin, W. H.: On the consistency strength of projective uniformization. In: Proc. Herbrand Symposium. Logic Colloquium '81 (Stern, J., ed.), North-Holland, Amsterdam, 365–384 (1982)

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, UNIVERSITÉ PARIS DIDEROT,
BOÎTE COURRIER 7012, 75205 PARIS CEDEX 13, FRANCE

Current address: Universität Wien, Institut für Mathematik, Kurt Gödel Research Center, Augasse
2-6, UZA 1 – Building 2, 1090 Wien, AUSTRIA

Email address: `noe.de.rancourt@univie.ac.at`