



HAL
open science

Biomechanical evaluation of the ST-knot: a new suture for flexor tendon repair

Camille Brenac, Martine Pithioux, Sacha Tomczak, Marylene Lallemand,
Charlotte Jaloux, Jean Baptiste Bargemon

► **To cite this version:**

Camille Brenac, Martine Pithioux, Sacha Tomczak, Marylene Lallemand, Charlotte Jaloux, et al..
Biomechanical evaluation of the ST-knot: a new suture for flexor tendon repair. *Hand Surgery and
Rehabilitation*, In press. hal-04407010

HAL Id: hal-04407010

<https://hal.science/hal-04407010>

Submitted on 22 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hand Surgery and Rehabilitation

Biomechanical evaluation of the ST-knot: a new suture for flexor tendon repair

--Manuscript Draft--

Manuscript Number:	HANSUR-D-23-00396R2
Article Type:	Original Article
Keywords:	biomechanical; ex vivo; suture; tendon
Corresponding Author:	Camille Brenac Hospital Timone FRANCE
First Author:	Camille Brenac
Order of Authors:	Camille Brenac Martine Pithioux Sacha Tomczak Marylène Lallemand Charlotte Jaloux Jean Baptiste De Villeneuve Bargemon
Abstract:	<p>Purpose</p> <p>Although tendon lacerations are common injuries for the hand surgeon, there is currently no consensus regarding the choice of one suture over another. Easy and fast sutures that impart enough strength to allow mobilization are needed. This study compares the ex vivo biomechanical strength (force required to create a 2 mm tendon gap) of a new suture (ST-knot) with that of a conventional suture (double Kessler).</p> <p>Materials and methods</p> <p>Forty fresh deep flexor tendons from porcine forelimbs were used. Both repaired tendon ends were mounted on standard traction jaws of an axial traction machine at an initial distance of 40 mm for all tendons. A high-definition camera was used to determine the force forming a 2 mm gap. Ten tendons from group 1 (ST-knot) and ten tendons from group 2 (double Kessler) were prepared with PDS 4.0 (single thread for Kessler but double for ST-knot). Tendons in group 3 (ST-knot) and 4 (double Kessler) were repaired with PDS 1.0 using the same principle.</p> <p>Results</p> <p>There was no significant difference in the force required to form a 2 mm tendon gap between groups 1 and 2 and this trend was identical when using a stronger thread in groups 3 and 4. The maximum force before rupture, mode of repair failure, stress, and stiffness were also comparable, with no significant differences between groups 1 and 2, or between groups 3 and 4.</p> <p>Conclusions</p> <p>The ST-knot shows comparable results to the double-Kessler, regardless of the thread used. Because it involves fewer steps than conventional techniques, and it is easy to perform, the ST-knot may offer a therapeutic solution, particularly in complex traumas with multiple tendon injuries.</p>

Title: Biomechanical evaluation of the ST-knot: a new suture for flexor tendon repair

Authors : Camille Brenac, MD¹ ; Martine Pithioux, MD^{2,4,5}; Sacha Tomczak, MD⁶; Marylène Lallemand, MD^{3,4}; Charlotte Jaloux, MD, PhD⁷ ; Jean Baptiste de Villeneuve Bargemon, MD^{7,8}

Affiliations :

¹ *Hospices Civils de Lyon, Hôpital Croix Rousse, Service de Chirurgie Plastique, Esthétique et Réparatrice, France, Lyon F-69003*

² *Aix Marseille Univ, CNRS, ISM, 13009 Marseille, France*

³ *Ecole Centrale Marseille, 13013 Marseille, France*

⁴ *Aix Marseille Univ, APHM, CNRS, ISM, Mecabio Platform, Department of Orthopaedics and Traumatology, 13009 Marseille, France*

⁵ *Aix Marseille Univ, APHM, CNRS, ISM, Sainte-Marguerite Hospital, Institute for Locomotion, Department of Orthopaedics and Traumatology, 13009 Marseille, France*

⁶ *Plastic and Reconstructive Surgery Department, Hôpital Conception, AP-HM, Marseille, France*

⁷ *Hand surgery and limb reconstructive surgery, CHU de Timone, Aix-Marseille University, 264, rue Saint Pierre, 13005 Marseille*

⁸ *Hand, Wrist and Elbow Surgery, Saint Roch Private Hospital, Toulon, 99 avenue Saint Roch, 83100 Toulon, France*

Corresponding author:

Camille Brenac – Service de Chirurgie Plastique, Esthétique et Réparatrice

Hospices Civils Lyon, Hôpital Croix Rousse

103 Grande rue de la Croix Rousse, 69004 Croix Rousse, France

Tél +33 4 26 73 28 22

E-mail: camille.brenac@chu-lyon.fr

1 **Abstract**

2 **Purpose:** Although tendon lacerations are common injuries for the hand surgeon, there is currently no
3
4 3 consensus regarding the choice of one suture over another. Easy and fast sutures that impart enough
5
6 4 strength to allow mobilization are needed. This study compares the *ex vivo* biomechanical strength (force
7
8 5 required to create a 2 mm tendon gap) of a new suture (ST-knot) with that of a conventional suture
9
10 6 (double Kessler).

11 7 **Materials and methods:** Forty fresh deep flexor tendons from porcine forelimbs were used. Both
12
13 8 repaired tendon ends were mounted on standard traction jaws of an axial traction machine at an initial
14
15 9 distance of 40 mm for all tendons. A high-definition camera was used to determine the force forming a
16
17 10 2 mm gap. Ten tendons from group 1 (ST-knot) and ten tendons from group 2 (double Kessler) were
18
19 11 prepared with PDS 4.0 (single thread for Kessler but double for ST-knot). Tendons in group 3 (ST-knot)
20
21 12 and 4 (double Kessler) were repaired with PDS 1.0 using the same principle.

22 13 **Results:** There was no significant difference in the force required to form a 2 mm tendon gap between
23
24 14 groups 1 and 2 and this trend was identical when using a stronger thread in groups 3 and 4. The
25
26 15 maximum force before rupture, mode of repair failure, stress, and stiffness were also comparable, with
27
28 16 no significant differences between groups 1 and 2, or between groups 3 and 4.

29 17 **Conclusions:** The ST-knot shows comparable results to the double-Kessler, regardless of the thread
30
31 18 used. Because it involves fewer steps than conventional techniques, and it is easy to perform, the ST-
32
33 19 knot may offer a therapeutic solution, particularly in complex traumas with multiple tendon injuries.

34
35
36
37
38
39
40
41
42 20
43
44 21 **Keywords:** biomechanical study; *ex vivo*; suture; tendon
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

22 **Résumé**

23 **Objectif :** Bien que les sections tendineuses soient des blessures courantes pour le chirurgien de la
24 main, il n'existe actuellement aucun consensus concernant le choix d'une suture par rapport à une
25 autre. Des sutures simples et rapides, offrant une résistance suffisante pour permettre la mobilisation,
26 sont nécessaires. Cette étude compare la force biomécanique ex vivo (force nécessaire pour créer un
27 espace de 2 mm entre les deux extrémités suturées du tendon) d'une nouvelle suture (nœud ST) à celle
28 d'une suture conventionnelle (double Kessler).

29 **Matériels et méthodes :** Quarante tendons fléchisseurs profonds frais issus de membres antérieurs de
30 porcs ont été utilisés. Les deux extrémités du tendon réparé ont été fixées sur des mors de traction
31 standard d'une machine à traction axiale à une distance initiale de 40 mm pour tous les tendons. Une
32 caméra haute définition a été utilisée pour déterminer la force nécessaire à la formation d'un espace de
33 2 mm entre les deux extrémités suturées. Dix tendons du groupe 1 (nœud ST) et dix tendons du groupe
34 2 (double Kessler) ont été préparés avec du PDS 4.0 (simple pour le Kessler mais boucle pour le nœud
35 ST). Les tendons des groupes 3 (nœud ST) et 4 (double Kessler) ont été réparés avec du PDS 1.0 selon
36 le même principe.

37 **Résultats :** Aucune différence significative n'a été observée dans la force nécessaire pour former un
38 espace de 2 mm dans le tendon entre les groupes 1 et 2, et cette tendance était identique lors de
39 l'utilisation d'un fil plus résistant dans les groupes 3 et 4. La force maximale avant rupture, le mode de
40 rupture, la contrainte et la rigidité étaient également comparables, sans différence significative entre
41 les groupes 1 et 2, ni entre les groupes 3 et 4.

42 **Conclusion :** Le nœud ST montre des résultats biomécaniques comparables au double Kessler,
43 indépendamment du fil utilisé. Étant donné qu'il nécessite moins d'étapes que les techniques
44 conventionnelles et qu'il est facile à réaliser, le nœud ST pourrait offrir une solution thérapeutique,
45 notamment dans les traumatismes complexes avec de multiples lésions tendineuses.

46

47 **Mots clefs :** Étude biomécanique, ex-vivo, suture, tendon

48 **Introduction**

1
2 49 Tendon lacerations represent a common therapeutic challenge for hand surgeons [1]. Good outcomes
3
4 50 do not only depend on surgery, but also on early mobilization and physiotherapy [2–4]. To facilitate
5
6 51 early and optimal post-operative physiotherapy, the tendon suture must possess the capacity to endure
7
8 52 the stresses of mobilization. At present, a definitive consensus regarding the optimal selection of sutures
9
10 53 for employment in multi-strand repair techniques is lacking, and surgeon preferences frequently rely on
11
12 54 individual experience rather than robust scientific evidence [5]. It is currently understood that numerous
13
14 55 factors have a positive impact on repair strength. These include the quantity of strands traversing the
15
16 56 repair, slight crowding at the repair site, employment of a larger suture gauge, and configuring the
17
18 57 peripheral sutures appropriately [6,7]. In an *ex vivo* study, Savage et al. reported that six-strand repair
19
20 58 appears to be three times more potent than conventional two-strand repair (Kessler, Kleinert, Beckert,
21
22 59 Bunnel) [3]. Other more recent *ex vivo* studies however, seem to invalidate this theory [8]. Tang et al.
23
24 60 subsequently advocated for the adoption of loop sutures, which allows for the creation of two strands
25
26 61 per pass. This approach reduces tendon edge damage by minimizing the number of needle passes while
27
28 62 maintaining comparable strength to that of conventional sutures [7,9–11].
29
30

31
32
33 63 Nonetheless, no study has conclusively demonstrated the superiority of one 4-strand suture over
34
35 64 another. Some sutures present cumbersome configurations and involve multiple trans-tendinous
36
37 65 passages [12]. In order to aid the hand surgeon in the treatment of tendon injuries, it appears imperative
38
39 66 to employ a suture that is straightforward, swift, employs a single thread, and imparts sufficient strength
40
41 67 to facilitate mobilization. Moreover, in the various comparative studies of new sutures, the Kessler
42
43 68 suture seems to be the most widely used reference suture, with results equal to or better than those of
44
45 69 the new suture knots tested [3,12–17].
46
47

48
49 70 The aim of this study was to compare the mechanical properties of two suture techniques: the
50
51 71 four-strand double Kessler and the ST-Knot (which creates an S and a T pattern on the tendon upon
52
53 72 completion), a new suture technique. The primary objective was to evaluate the 2 mm gap-forming force
54
55 73 (N). The secondary objectives were to evaluate the maximum force before rupture (N_{max}), the mode of
56
57 74 repair failure (knot rupture, suture rupture, and suture pull-out), stress (ultimate tensile strength divided
58
59 75 by tendon cross-sectional area, in N/mm²), and stiffness (N/mm).
60
61
62
63
64
65

76 Materials and methods

77 *Surgical technique*

78 The ST-knot begins with the use of a 4/0 diameter looped suture. We prefer to use a polydioxanone
79 (PDS, Ethicon, Somerville, NJ), suture with a 19 mm needle. The first step consists in docking the
80 proximal tendon stump using Tsuge and Ikuta's stratagem [18] (**Figure 1A**). This allows the flexor
81 tendon, which tends to retract naturally, to be held in place, thus bypassing the use of a transtendinous
82 needle to reduce retraction forces. The suture continues by crossing intratendinously to reach the distal
83 tendon stump, thus initiating a transverse pass within the tendon (**Figure 1B and 1C**). The same
84 technique is used on the other side of the tendon. As the tendon is docked to the suture on the proximal
85 stump, this double anchoring in the distal stump brings the two tendon parts together, thereby reducing
86 the retraction forces on each tendon. As a result, the proximal stump is less likely to retract, and the two
87 tendon stumps are closer together (**Figure 1D**). When using an undocked suture, the suture slides
88 through the tendon, preventing the tendon from being approached. An intratendinous pass is then made
89 at the proximal stump on the contralateral side, to adjust the tendon suture axially and prevent
90 intratendinous torsion (**Figure 1E**). Note that this adjustment is simpler if the tendons are closer together
91 than if they are further apart. A final pass between the two passages secures the suture, forming a knot
92 without any retraction tension, thus eliminating the necessity for force to bring the two stumps into
93 proximity (**Figure 1F**). By applying an appropriate tension, this final pass enables the degree of
94 bunching up to be adjusted by 20 to 30%, as recommended by Tang JB et al. [7]. Following the method
95 outlined by Tsuge et al. a tension-free knot is made (**Figure 1G**)[18]. This produces a 6-strand
96 intratendinous suture resembles the M-Tang repair but employs a single thread, with an extra-tendinous
97 knot. The Kessler knot is executed using a PDS 4/0, a 26 mm needle, two strands, a single extratendinous
98 knot within the repair site, and four loops (**Figure 2**).

99

100 *Samples*

101 This study follows the concepts and sample sizes of other published biomechanical studies concerning
102 *ex vivo* flexor tendon repair techniques [5,19]. Forty fresh deep flexor tendons from porcine forelimbs
103 were used as experimental models. These tendons are similar in structure and size to the human flexor

104 digitorum profundus, and have been used in numerous studies [5,19–21]. The two deep flexor tendons
105 from each forelimb were harvested between pulleys A2 and A4. With an average cross-sectional area
106 of 21 mm² and a minimum length of 10 cm, these tendons were suitable for installation in the jaws of a
107 traction machine. All tendons were measured in width, thickness, and length using a calliper, and cross-
108 sectional area estimates were calculated using the following formula for the area of an ellipse [(small
109 diameter (mm)/2)*(large diameter (mm)/2)* π]. The tendons were maintained in a frozen state at -20°C
110 until 24 hours prior to testing. At that point, they were thawed and immersed in saline at -4°C.

112 *Repairs and equipment*

113 A 23 blade was used to make a clean cross-section through the middle of each tendon. The 40 tendons
114 were randomly divided into four groups of 10, repaired, and then tested in axial traction. All tendons
115 were repaired using PDS, which is a synthetic monofilament absorbable suture material commonly used
116 for tendon repair. The entry and exit points of the central sutures were located 7 mm from the cut end
117 of the tendon for all repairs with a 10% shortening to ensure the best tensile strength. This 7 mm distance
118 was systematically measured using the same metal ruler. This distance was chosen because it has been
119 shown that, although a distance of 4 mm or less results in significantly weaker repairs, increasing the
120 suture distance from 7 mm to 10 mm or 12 mm does not increase repair strength [21]. The reference
121 repair technique chosen for comparison in this study was the double Kessler technique or 4-strand locked
122 cross repair [22] (**Figure 2**). Groups 1 (ST1) and 3 (ST2) tendons were repaired with an ST-knot, while
123 groups 2 (K1) and 4 (K2) tendons were repaired using the double Kessler technique. For the central
124 sutures in groups 1 (ST1) and 2 (K1), a PDS 4.0 was used, including a loop PDS in group 1. Tendons
125 in groups 3 (ST2) and 4 (K2) were repaired with PDS 1.0. All tendon sutures were reinforced with a
126 peripheral PDS 5.0 peritendinous suture 2 mm from the transection site. The tendons were repaired by
127 two qualified surgeons.

129 *Biomechanical testing*

130 The tests were carried out using an Instron 5566A axial traction machine (Instron Corp; **Figure 3**). Both
131 mended tendon ends were affixed to standard traction mounts, with the initial distance uniformly set at

132 40 mm for all tendons. Subsequently, the repaired tendons underwent a tensile strength assessment.
1
2 133 After applying a preload of 1 newton (N), the tendons were drawn at a uniform rate of 10 mm/min. Data
3
4 134 points were captured at a frequency of 10 Hertz. The Bluehill Universal software (Instron Corp) was
5
6 135 employed to generate a force-versus-displacement curve, from which the ultimate tensile strength
7
8 136 (identified as the peak of the curve) was ascertained (**Figure 4**). Stiffness was calculated as the slope of
9
10 137 the initial linear section of the curve. A ruler was attached to the upper clamp next to the tendon, and
11
12 138 the test was monitored using a high-definition camera to determine the force forming a 2 mm gap
13
14 139 (**Figure 3**). The 2 mm benchmark was considered to have been reached as soon as a 2 mm gap had
15
16 140 formed over part of the repair site. The mode of failure was determined by visual examination of the
17
18 141 tendons by the surgeon, during and after the test.
19
20
21
22
23

24 143 *Statistical methods*

25
26 144 A Shapiro-Wilk test was used to ensure that tendon cross-sections were normally distributed. The non-
27
28 145 parametric Mann-Whitney test was used to compare numerical results (mechanical test results and cross-
29
30 146 sectional areas) between all groups.
31
32

33 147

35 148 **Results**

36
37 149 Regarding the primary endpoint, there was no significant difference in the occurrence of a tendon gap
38
39 150 at 2 mm between groups 1 and 2 (23.7N [17.5; 29.8] vs 24.2N [17.45; 30.9]; 95% CI) and this trend was
40
41 151 identical with the use of a stronger thread (groups 3 and 4) (55N [30.2; 79.8] vs 45.6N [30.8; 60.4]; 95%
42
43 152 CI). Other results (maximum force before rupture, mode of repair failure, stress and stiffness) were
44
45 153 comparable between groups 1 and 2, and between groups 3 and 4 (**Tables 1, 2, and 3**).
46
47
48

49 154 Regarding the mode of repair failure, 7 suture ruptures, 2 knot ruptures, and 1 suture pull-out
50
51 155 were observed in group 1 (ST1) while 7 suture ruptures and 3 suture pull-outs were observed in group
52
53 156 3 (ST2). The mode of repair failure in group 2 (K1) and 4 (K2) were almost exclusively due to suture
54
55 157 rupture, except for 1 suture pull-out which occurred in group 4 (**Table 1**).
56
57

58 158

60 159 **Discussion**

160 At present, although some recommendations were made over 30 years ago, there exists no unanimous
1
2 161 consensus regarding the treatment of flexor tendon injuries [23]. There appears to be a dispute
3
4 162 concerning the necessity of employing multiple intratendinous strands [3,8,10]. The present results
5
6 163 indicate that the number of passes does not appear to exert a notable influence on the maximum load or
7
8 164 the load required to induce a 2-mm gap. The findings herein are in line with previously published studies
9
10 demonstrating similar load levels [12,13,15,24,25]. Schädel-Höpfner et al. reported that the Kessler has
11 165
12 by far the best ultimate tensile strength when compared with 4 other sutures, likely explaining why this
13 166
14 knot is frequently found in comparative studies [13]. In 1999, McLarney demonstrated through a
15 167
16 biomechanical study that the cruciate four-strand flexor tendon repair exhibited a high failure load,
17 168
18 nearly double that of the Kessler knot [26]. However, no other biomechanical study of this knot without
19 169
20 a cross-lock was found in the literature. While forces up to 2.9N (2N [min; 0- max; 2.9]) might be
21 170
22 expected during unresisted active flexion of the wrist and up to 28.4N (18.6N [min; 1-max; 28.4]) during
23 171
24 unresisted active flexion of the long fingers, the ST-knot was consistent with these results showing a
25 172
26 mean resistance of 26N, a lower limit of 17.2N, and an upper limit of 34.9N when using PDS 4.0 [27].
27 173
28 The referenced study reported results in kilogram force (kgf), where 0.1 kgf corresponds to 1N. At
29 174
30 present, there is a wide disparity in the choice of suture between different centres and no single suture
31 175
32 appears to offer any real superiority [5,28,29]. In the management of flexor tendon injuries, it would
33 176
34 seem worthwhile to take a step back from the type of suture used and focus more on the practical and
35 177
36 technical use of these sutures. In the present study, the ST-knot yielded comparable results to the double-
37 178
38 Kessler technique, with no notable superiority observed for either method. The equivalence remains
39 179
40 unchanged with the use of a thicker suture.
41
42
43
44
45

46 181 According to various literature reviews and expert opinions, the key to successful tendon
47
48 suturing seems to be based not on the type of suture but on other multiple local factors such as pulley
49 182
50 preservation, adjusted tension between the two stumps, careful and limited soft tissue dissection
51 183
52 [6,28,29]. The ST-knot then takes its place as a practical addition to the suture. The tendon docking at
53 184
54 the very beginning of the suture enables the proximal stump to be mobilized solely by traction on the
55 185
56 suture thread as the M-Tang or Lim-Tsai suture when starting with the backwall of the running suture.
57 186
58 In addition, by bringing the two stumps together at an early stage and applying a slight tension at rest
59 187
60
61
62
63
64
65

188 between the two stumps, the use of tendon-damaging forceps is reduced, allowing intrinsic tendon
189 structures to be protected as much as possible. This is not possible when using the Kessler suture, as the
190 thread slips through the tendon if it is pulled along its axis. Moreover, the closing forces of the two
191 stumps are applied during the locking of the knot in a Kessler-type suture. In this situation, there is a
192 risk of suture rupture, due to the combination of the tension applied to lock the knot and the tension
193 needed to bring the two tendon stumps together. However, when the ST-Knot is performed, the two
194 tendon stumps are already brought together, reducing the shearing forces involved in locking the knot.
195 In addition, a double Kessler suture requires 12 different intratendinous passages, compared with 7 for
196 the ST-Knot. This is particularly important in the case of multiple tissue and tendon injuries, where
197 tendon repair time is paramount. Moreover, compared to other 6-strand sutures, the ST-knot enables a
198 6-strand suture to be produced using a single thread [11,30,31]. The strength of a suture must be
199 considered not only in terms of continuous axial traction, but also in terms of cyclic loading. Whether
200 the suture is 6- or 4-strand, previous studies showed that the transverse components of the suture are a
201 detrimental factor in the development of a tendon gap, however this aspect was not found in our study,
202 as the ST-knot has only one transverse component whereas the double Kessler has 4 transverse
203 components [32,33]. In our opinion, the ST-knot is not a stronger suture than the classic double Kessler,
204 but it does offer the surgeon greater comfort in suturing tendon injuries.

205 One criticism that can be made about the ST-knot is that the 6-strand tensions are neutralized
206 by a single knot. A recent study however, showed that the presence of 1 or 2 knots has no impact on the
207 functional outcome [34]. Furthermore, the presence of an extratendinous knot does not appear to have
208 a negative biomechanical impact compared with that of an intratendinous knot [11,35]. Indeed, Chang
209 et al. showed that the Lim/Tsai repair with extra-tendinous knot had a statistically significantly higher
210 ultimate strength and load to 2 mm gap force, as compared with the repair with intra-tendinous knot
211 [18,34,36]. However, suture knot location could affect glide especially in the digital canal. Noguchi et
212 al. showed in a *in vivo* study that the Tajima repair, whose knot is internal, exhibits significantly lower
213 gliding resistance than the modified Kessler which is identical except for an externally placed knot [37].
214 The contrary argument is that knot placement between the cuts could reduce the tendon end contact
215 surface area that is involved in healing [38].

216 Nonetheless, the present study is subject to several limitations, one of which is the exclusive
1 performance of in-axis traction. Novel modes of suture failure have indeed been shown when using
2 217
3 cyclic load tests [32,33]. As the ST-Knot has a transverse component, it would be interesting to explore
4 218
5 this weakness. In addition, although a total of 40 sutures were performed, each group contained only 10
6 219
7 sutures. Although this sample size is found in many comparative studies it may nevertheless represent
8 220
9 a potential limitation [13,25,32,33]. The failure loads observed in our study are relatively low for both
10 221
11 repairs (around 25 N) when using a 4.0 suture. The authors propose that a slightly lower strength is
12 222
13 observed, which may be attributed to the use of frozen tendons and the associated protocol,
14 223
15 distinguishing it from clinical findings. Furthermore, we claim that the ST-Knot appears to be easier
16 224
17 than a double Kessler suture, but unfortunately, we did not measure suture completion time, which could
18 225
19 be an interesting indicative marker to assert this ease. The completion time for an ST-knot should be
20 226
21 assessed on a cadaver finger, considering that one of the key features of this knot is the docking system.
22 227
23 In any case, suturing two tendon stumps ex-vivo does not appear to be representative of a real-life tendon
24 228
25 suturing clinical context. Moreover, other references than the Kessler knot could have been used in this
26 229
27 study. On the one hand, our choice to use the Kessler knot as a reference aligns with its widespread
28 230
29 utilization in recent biomechanical literature [14,19,24,39]. On the other hand, a comparative analysis
30 231
31 between the ST-knot and the cruciate four-strand flexor tendon repair would be of significant interest.
32 232
33
34
35
36
37
38
39

40 234 **Conclusion:**

41
42 235 The ST-knot seems to be a viable and secure suture, with biomechanical capacities that are comparable
43 236
44 to conventional sutures. Due to its fewer procedural steps and ease of execution compared to
45 237
46 conventional techniques, the ST-knot could provide a therapeutic option, especially in intricate traumas
47 238
48 involving multiple tendon injuries. To validate these hypotheses, *in vivo* studies are needed.
49 239
50

51 239 52 240 **Disclosure of interest**

53 241 The authors declare that they have no known competing financial or personal relationships that could
54 242
55 be viewed as influencing the work reported in this paper.
56 243
57
58
59
60
61
62
63
64
65

244 **Funding sources:**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

245 This research did not receive any specific grant from funding agencies in the public, commercial, or
246 not-for-profit sectors.

247
248 **Acknowledgements:** We thank Véréna Landel (Direction de la Recherche en Santé, Hospices
249 Civils de Lyon) for her help in manuscript preparation.

250 **References**

- 1
2 251 [1] Strickland JW. Development of flexor tendon surgery: twenty-five years of progress. *J Hand*
3 252 *Surg Am* 2000;25:214–35. <https://doi.org/10.1053/jhsu.2000.jhsu25a0214>.
- 4 253 [2] Sandow MJ, McMahon M. Active mobilisation following single cross grasp four-strand flexor
5 254 tenorrhaphy (Adelaide repair). *J Hand Surg Eur Vol* 2011;36:467–75.
6 255 <https://doi.org/10.1177/1753193411405937>.
- 7 256 [3] Savage R. In vitro studies of a new method of flexor tendon repair. *J Hand Surg Br*
8 257 1985;10:135–41. [https://doi.org/10.1016/0266-7681\(85\)90001-4](https://doi.org/10.1016/0266-7681(85)90001-4).
- 9 258 [4] Hung LK, Pang KW, Yeung PLC, Cheung L, Wong JMW, Chan P. Active mobilisation after
10 259 flexor tendon repair: comparison of results following injuries in zone 2 and other zones. *J Orthop Surg*
11 260 (Hong Kong) 2005;13:158–63. <https://doi.org/10.1177/230949900501300209>.
- 12 261 [5] Lawrence TM, Davis TRC. A biomechanical analysis of suture materials and their influence
13 262 on a four-strand flexor tendon repair. *J Hand Surg Am* 2005;30:836–41.
14 263 <https://doi.org/10.1016/j.jhsa.2005.03.011>.
- 15 264 [6] Ayhan E, Tuna Z, Oksuz C. Getting Better Results in Flexor Tendon Surgery and Therapy.
16 265 *Plast Reconstr Surg Glob Open* 2021;9:e3432. <https://doi.org/10.1097/GOX.0000000000003432>.
- 17 266 [7] Tang JB. Flexor Tendon Injuries. *Clinics in Plastic Surgery* 2019;46:295–306.
18 267 <https://doi.org/10.1016/j.cps.2019.02.003>.
- 19 268 [8] Uslu M, Isik C, Ozsahin M, Ozkan A, Yasar M, Orhan Z, et al. Flexor tendons repair: effect of
20 269 core sutures caliber with increased number of suture strands and peripheral sutures. A sheep model.
21 270 *Orthop Traumatol Surg Res* 2014;100:611–6. <https://doi.org/10.1016/j.otsr.2014.05.009>.
- 22 271 [9] Tang JB, Wang B, Chen F, Pan CZ, Xie RG. Biomechanical evaluation of flexor tendon repair
23 272 techniques. *Clin Orthop Relat Res* 2001:252–9. <https://doi.org/10.1097/00003086-200105000-00033>.
- 24 273 [10] Kang GH-Y, Wong Y-R, Lim RQ-R, Loke AM-K, Tay S-C. Cyclic Testing of the 6-Strand
25 274 Tang and Modified Lim-Tsai Flexor Tendon Repair Techniques. *J Hand Surg Am* 2018;43:285.e1-
26 275 285.e6. <https://doi.org/10.1016/j.jhsa.2017.08.014>.
- 27 276 [11] Wang B, Xie RG, Tang JB. Biomechanical analysis of a modification of Tang method of
28 277 tendon repair. *J Hand Surg Br* 2003;28:347–50. [https://doi.org/10.1016/s0266-7681\(03\)00019-6](https://doi.org/10.1016/s0266-7681(03)00019-6).
- 29 278 [12] Marrero-Amadeo IC, Chauhan A, Warden SJ, Merrell GA. Flexor tendon repair with a
30 279 knotless barbed suture: a comparative biomechanical study. *J Hand Surg Am* 2011;36:1204–8.
31 280 <https://doi.org/10.1016/j.jhsa.2011.04.003>.
- 32 281 [13] Schädel-Höpfner M, Windolf J, Lögters TT, Hakimi M, Celik I. Flexor tendon repair using a
33 282 new suture technique: a comparative in vitro biomechanical study. *Eur J Trauma Emerg Surg*
34 283 2011;37:79–84. <https://doi.org/10.1007/s00068-010-0019-8>.
- 35 284 [14] DüNDAR N, Güneri B, Uzel M, Doğaner A. Biomechanical comparison of Bunnell, modified
36 285 Kessler, and Tsuge tendon repair techniques using two suture types. *Acta Orthop Traumatol Turc*
37 286 2020;54:104–13. <https://doi.org/10.5152/j.aott.2020.01.411>.
- 38 287 [15] Yang W, Li J, Su Y, Liang W, Ren Y, Dong Y, et al. A Modified Flexor Tendon Suture
39 288 Technique Combining Kessler and Loop Lock Flexor Tendon Sutures. *Clinics (Sao Paulo)*
40 289 2021;76:e2358. <https://doi.org/10.6061/clinics/2021/e2358>.
- 41 290 [16] Zeplin PH, Zahn RK, Meffert RH, Schmidt K. Biomechanical evaluation of flexor tendon
42 291 repair using barbed suture material: a comparative ex vivo study. *J Hand Surg Am* 2011;36:446–9.
43 292 <https://doi.org/10.1016/j.jhsa.2010.11.031>.
- 44 293 [17] Low TH, Ahmad TS, Ng ES. Simplifying four-strand flexor tendon repair using double-
45 294 stranded suture: a comparative ex vivo study on tensile strength and bulking. *J Hand Surg Eur Vol*
46 295 2012;37:101–8. <https://doi.org/10.1177/1753193411409840>.
- 47 296 [18] Ikuta Y, Tsuge K. Postoperative results of looped nylon suture used in injuries of the digital
48 297 flexor tendons. *J Hand Surg Br* 1985;10:67–72. [https://doi.org/10.1016/s0266-7681\(85\)80021-8](https://doi.org/10.1016/s0266-7681(85)80021-8).
- 49 298 [19] Jordan MC, Schmitt V, Jansen H, Meffert RH, Hoelscher-Doht S. Biomechanical Analysis of
50 299 the Modified Kessler, Lahey, Adelaide, and Becker Sutures for Flexor Tendon Repair. *J Hand Surg*
51 300 *Am* 2015;40:1812–7. <https://doi.org/10.1016/j.jhsa.2015.05.032>.
- 52 301 [20] Croog A, Goldstein R, Nasser P, Lee SK. Comparative biomechanic performances of locked
53 302 cruciate four-strand flexor tendon repairs in an ex vivo porcine model. *J Hand Surg Am* 2007;32:225–
54 303 32. <https://doi.org/10.1016/j.jhsa.2006.11.009>.
- 55
56
57
58
59
60
61
62
63
64
65

- 304 [21] Tang JB, Zhang Y, Cao Y, Xie RG. Core suture purchase affects strength of tendon repairs. *J*
1 305 *Hand Surg Am* 2005;30:1262–6. <https://doi.org/10.1016/j.jhsa.2005.05.011>.
- 2 306 [22] Kessler I, Nissim F. Primary repair without immobilization of flexor tendon division within
3 307 the digital sheath. An experimental and clinical study. *Acta Orthop Scand* 1969;40:587–601.
4 308 <https://doi.org/10.3109/17453676908989524>.
- 5 309 [23] Strickland JW. Flexor tendon injuries. Part 2. Flexor tendon repair. *Orthop Rev* 1986;15:701–
6 310 21.
- 7 311 [24] Renner C, Corella F, Fischer N. Biomechanical evaluation of 4-strand flexor tendon repair
8 312 techniques, including a combined Kessler-Tsuge approach. *J Hand Surg Am* 2015;40:229–35.
9 313 <https://doi.org/10.1016/j.jhsa.2014.10.055>.
- 10 314 [25] Castoldi M, Solla F, Camuzard O, Pithioux M, Rampal V, Rosello O. A 3-Dimensional Suture
11 315 Technique for Flexor Tendon Repair: A Biomechanical Study. *J Hand Surg Am* 2023;48:194.e1-
12 316 194.e9. <https://doi.org/10.1016/j.jhsa.2021.09.030>.
- 13 317 [26] McLarney E, Hoffman H, Wolfe SW. Biomechanical analysis of the cruciate four-strand
14 318 flexor tendon repair. *J Hand Surg Am* 1999;24:295–301. <https://doi.org/10.1053/jhsu.1999.0295>.
- 15 319 [27] Schuind F, Garcia-Elias M, Cooney WP, An K-N. Flexor tendon forces: In vivo
16 320 measurements. *J Hand Surg Am* 1992;17:291–8. [https://doi.org/10.1016/0363-5023\(92\)90408-H](https://doi.org/10.1016/0363-5023(92)90408-H).
- 17 321 [28] Tang JB, Amadio PC, Boyer MI, Savage R, Zhao C, Sandow M, et al. Current practice of
18 322 primary flexor tendon repair: a global view. *Hand Clin* 2013;29:179–89.
19 323 <https://doi.org/10.1016/j.hcl.2013.02.003>.
- 20 324 [29] Klifto CS, Capo JT, Sapienza A, Yang SS, Paksima N. Flexor Tendon Injuries. *J Am Acad*
21 325 *Orthop Surg* 2018;26:e26–35. <https://doi.org/10.5435/JAAOS-D-16-00316>.
- 22 326 [30] Kozono N, Okada T, Takeuchi N, Shimoto T, Higaki H, Nakashima Y. A Biomechanical
23 327 Comparison Between Asymmetric Pennington Technique and Conventional Core Suture Techniques:
24 328 6-Strand Flexor Tendon Repair. *J Hand Surg Am* 2018;43:79.e1-79.e8.
25 329 <https://doi.org/10.1016/j.jhsa.2017.06.104>.
- 26 330 [31] Giesen T, Reissner L, Besmens I, Politikou O, Calcagni M. Flexor tendon repair in the hand
27 331 with the M-Tang technique (without peripheral sutures), pulley division, and early active motion. *J*
28 332 *Hand Surg Eur Vol* 2018;43:474–9. <https://doi.org/10.1177/1753193418758269>.
- 29 333 [32] Wu YF, Mao WF, Tang JB. The impact of transverse components on 4-strand tendon repairs.
30 334 *J Hand Surg Eur Vol* 2021;46:830–5. <https://doi.org/10.1177/17531934211018309>.
- 31 335 [33] Wu YF, Tang JB. The impact of transverse components on resistance and ultimate strength of
32 336 6-strand tendon repairs. *J Hand Surg Eur Vol* 2021;46:836–41.
33 337 <https://doi.org/10.1177/17531934211018327>.
- 34 338 [34] Gil JA, Skjong C, Katarincic JA, Got C. Flexor Tendon Repair With Looped Suture: 1 Versus
35 339 2 Knots. *J Hand Surg Am* 2016;41:422–6. <https://doi.org/10.1016/j.jhsa.2015.12.011>.
- 36 340 [35] Chen J, Fang Wu Y, Xing SG, Jun Pan Z. Suture knots between tendon stumps may not
37 341 benefit tendon repairs. *J Hand Surg Eur Vol* 2018;43:1005–6.
38 342 <https://doi.org/10.1177/1753193418788939>.
- 39 343 [36] Chang MK, Wong YR, Tay SC. Biomechanical comparison of modified Lim/Tsai tendon
40 344 repairs with intra- and extra-tendinous knots. *J Hand Surg Eur Vol* 2018;43:919–24.
41 345 <https://doi.org/10.1177/1753193418769804>.
- 42 346 [37] Noguchi M, Seiler JG, Gelberman RH, Sofranko RA, Woo SL. In vitro biomechanical
43 347 analysis of suture methods for flexor tendon repair. *J Orthop Res* 1993;11:603–11.
44 348 <https://doi.org/10.1002/jor.1100110415>.
- 45 349 [38] MASON ML, SHEARON CG. THE PROCESS OF TENDON REPAIR: AN
46 350 EXPERIMENTAL STUDY OF TENDON SUTURE AND TENDON GRAFT. *Archives of Surgery*
47 351 1932;25:615–92. <https://doi.org/10.1001/archsurg.1932.01160220003001>.
- 48 352 [39] Wongsiri S, Liawrungrueang W. Biomechanical study of a newly developed continuous
49 353 double knots technique compared with the 4-strand double-modified Kessler technique for flexor
50 354 tendon repair. *J Exp Orthop* 2021;8:79. <https://doi.org/10.1186/s40634-021-00404-4>.
- 51 355

356 **Figures legends**

1 357 **Figure 1: Illustration of the various stages of an ST-Knot.**

2
3 358 *A/ Docking of proximal tendon stump to PDS loop. B/ Intratendinous passage through proximal*
4
5 359 *stump, then lateral suture exit. C/ Transtendinous passage. D/ Intratendinous passage again. E/*
6 360 *Passage back into the proximal tendon stump. F/ Extratendinous passage in the distal stump,*
7
8 361 *returning to the proximal stump via an intratendinous passage. G/ Knot lock*

9
10 362
11 363 **Figure 2: Illustration of the 4-strand Kessler repair used in this study.**

12
13 364
14
15 365 **Figure 3: Installing the tendon in the Instron 5566A machine.**

16 366 *A/ Installation of tendon and camera*
17
18 367 *B/ Occurrence of peritendinous suture rupture after axial traction*

19
20 368
21 369 **Figure 4: Biomechanical test: displacement curve (mm) as a function of load (N).**

22
23 370 *ST-knot load-displacement graph with 10mm/min traction generated by Bluehill Universal software. A*
24
25 371 *deviation of 2mm generally occurred before rupture of the peritendinous suture.*

372 **Table 1. Biomechanical results according to the 4 different suture groups**

	ST1 (n=10)	K1 (n=10)	ST2 (n=10)	K2 (n=10)
Force (N)- 2mm gap, mean [95% CI]	23.7 [17.5 ; 29.8]	24.2 [17.45 ; 30.9]	55 [30.2 ; 79.8]	45.6 [30.8 ; 60.4]
Force (N) Max, mean [95% CI]	26 [17.2 ; 34.9]	29.8 [19.7 ; 39.9]	68 [35.6 ; 100.4]	60.7 [40.5 ; 80.9]
Stress (N/mm²), mean [95% CI]	1.3 [0.8 ; 1.8]	1.5 [0.9 ; 2]	2.9 [1.6 ; 4.2]	3 [1.7 ; 4.3]
Stiffness (N/mm), mean [95% CI]	4.2 [2.4 ; 6]	3.8 [2.5 ; 5.2]	4.8 [3.2 ; 6.5]	5.1 [3.2 ; 7]
Failure mode (n=40)	10	10	10	10
Suture rupture (n=33)	7	10	7	9
Knot rupture (n=2)	2	0	0	0
Suture pull-out (n=5)	1	0	3	1

374
 375 *Abbreviations: CI, Confidence interval; Max: maximum; N, Newton*

376 **Table 2. Statistical analysis for the comparison of biomechanical results from sutured tendons in**
 377 **groups 1 and 2.**

	ST1 (n=10) (Group 1)	K1 (n=10) (Group 2)	p-value
Force (N)- 2mm gap, mean [95% CI]	23.7 [17.5 ; 29.8]	24.2 [17.5 ; 30.9]	0.94
Force (N) Max, mean [95% CI]	26 [17.2 ; 34.9]	29.8 [19.7 ; 39.9]	0.57
Stress (N/mm²), mean [95% CI]	1.3 [0.8 ; 1.8]	1.5 [0.9 ; 2]	0.97
Stiffness (N/mm), mean [95% CI]	4.2 [2.4 ; 6]	3.8 [2.5 ; 5.2]	0.73

378

379 *Abbreviations: CI, Confidence interval; Max: maximum; N, Newton*

380 **Table 3. Statistical analysis for the comparison of biomechanical results from sutured tendons in**
 381 **groups 3 and 4.**

	ST2 (n=10) (Group 3)	K2 (n=10) (Group 4)	p-value
Force (N)- 2mm gap, mean [95% CI]	55 [30.2 ; 79.8]	45.6 [30.8 ; 60.4]	0.76
Force (N) Max, mean [95% CI]	68 [35.6 ; 100.4]	60.7 [40.5 ; 80.9]	0.97
Stress (N/mm²), mean [95% CI]	2.9 [1.6 ; 4.2]	3 [1.7 ; 4.3]	0.97
Stiffness (N/mm), mean [95% CI]	4.8 [3.2 ; 6.5]	5.1 [3.2 ; 7]	0.68

382
 383 *Abbreviations: CI, Confidence interval; Max, maximum; N, Newton*

POINT BY POINT RESPONSE TO REVIEWERS:

Dear Dr Dreant,

Thank you very much for sending us the reviewers' comments on the manuscript entitled "**Biomechanical evaluation of the ST-knot: a new suture for flexor tendon repair**". We would like to thank you for your interest in our work and for your contribution as experts in improving the quality of this submission.

We believe we have followed your advice and guidelines and addressed all comments the editor-in-chief had with our revised manuscript. The manuscript has been corrected in the text. We hope that you will judge the manuscript suitably revised for publication in *Hand Surgery and Rehabilitation* and remain at your disposal if necessary.

Submission:

Journal: *Hand Surgery and Rehabilitation*

Manuscript : HANSUR-D-23-00396

Type: Original article

Reviewer 1

1) Chers auteurs,

merci pour les corrections apportées à votre manuscrit.

Il reste encore quelques points avant de publier votre article.

Tout d'abord pourquoi ST knot? Le lecteur sera curieux de savoir à quoi correspondent ces deux lettres. Merci de le préciser dès la première apparition de ce terme dans votre manuscrit (pas dans le résumé ni dans le titre)

Auteurs : Le fil forme un S et un T sur le tendon lors de la réalisation du ST-knot, c'est pour cela que nous l'avons appelé de la sorte. Nous rajoutons la phrase suivante en accord avec votre commentaire page 3, ligne 71: (which creates an S and a T pattern on the tendon upon completion)

2) Merci d'écrire "et al." normalement, et non en italique.

Auteurs: La mise en forme de « et al. » a été corrigée dans l'ensemble du manuscrit.

3) La référence [1] fait allusion aux propriétés mécaniques du ligament scapho-lunaire. La référence [2] est par contre appropriée. Merci de supprimer la ref [1] qui vient mal à propos, surtout au tout début d'un article sur les tendons.

Auteurs : La référence [1] erronée a été supprimée.

4) Veuillez ré-écrire la bibliographie en faisant bien attention aux noms des revues.

Par exemple, dans la référence [2] le nom de la revue n'est pas "The journal of hand surgery" mais "J Hand Surg Am" comme il est écrit dans la base pubmed.

Auteurs : Nous avons vérifié et modifié l'ensemble des références erronées selon leur référencement pubmed.

5) Enfin, renvoyez seulement le manuscrit propre, sans corrections visibles.

Auteurs : Nous vous renvoyons uniquement le manuscrit corrigé.

Camille Brenac, MD

Service de Chirurgie Plastique, Esthétique et Réparatrice

Hospices Civils Lyon, Hôpital Croix Rousse

E-mail : camille.brenac@chu-lyon.fr

Dear Sir Dreant,

Editor-in-Chief

We would like to submit our manuscript entitled “Biomechanical evaluation of the ST-knot: a new suture for flexor tendon repair” for publication as an original article in *The Journal of Hand Surgery and Rehabilitation*


We believe that the teaching point of this article would be of great interest to the readers of *Journal of Hand Surgery and Rehabilitation*, as, in aiding the hand surgeon in the treatment of tendon lacerations, it appears imperative to employ a suture that is straightforward, swift, employs a single strand, and imparts sufficient strength to facilitate mobilization. In this biomechanical study, we demonstrate that ST-knots yield similar results in terms of resistance strength to double-Kessler sutures, irrespective of the thread size employed. Consequently, it emerges as a viable alternative for tendon repair.

All the authors have read the paper and agree with the content. No authors have conflicting interests. Neither the work nor any part of its essential substance, tables or figures have been, or will be, published or submitted to another scientific journal or are being considered for publication elsewhere. We are looking forward to reading your comments. Please feel free to contact me for any further questions.







Yours sincerely,

Camille Brenac on behalf of all authors

Figure 1 PDF



B [Click here to access/download;Figure;Figure 1 PDF.pdf](#)







Figure 3



A


[Click here to access/download;Figure;Figure 3 PDF pdf.pdf](#)



B


Figure 4

2 mm gap



Rupture of the peripheral suture

Suture breakage



Author responsibilities, integrity, ethics



This is an **editable** PDF form. It should be **saved to your computer, then completed** using Adobe reader or equivalent. Please **do NOT substitute** any other document (text file, scanned image, etc.).

Article title :

Human and animal rights

- The authors declare that the work described has been carried out in accordance with the [Declaration of Helsinki](#) of the World Medical Association revised in 2013 for experiments involving humans as well as in accordance with the EU Directive [2010/63/EU](#) for animal experiments.
- The authors declare that the work described has not involved experimentation on humans or animals.

Informed consent and patient details

- The authors declare that this report does not contain any [personal information](#) that could lead to the identification of the patient(s) and/or volunteers.
- The authors declare that they obtained a written [informed consent](#) from the patients and/or volunteers included in the article and that this report does not contain any [personal information](#) that could lead to their identification.
- The authors declare that the work described does not involve patients or volunteers.

Disclosure of interest

- The authors declare that they have no known [competing financial](#) or [personal relationships](#) that could be viewed as influencing the work reported in this paper.
- The authors declare the [following financial](#) or [personal relationships](#) that could be viewed as influencing the work reported in this paper:

Funding

- This work did not receive any [grant](#) from funding agencies in the public, commercial, or not-for-profit sectors.
- This work has been [supported](#) by:

Author contributions

- All authors attest that they meet the current International Committee of Medical Journal Editors ([ICMJE](#)) criteria for Authorship.
- All authors attest that they meet the current International Committee of Medical Journal Editors ([ICMJE](#)) criteria for Authorship. Individual author contributions are as follows:



Click here to access/download
e-component
Graphical Abstract.pdf

