



**HAL**  
open science

# Time-uniform convergence rates for diffusion Monte Carlo with applications to coupled harmonic oscillators

Michel Caffarel, Pierre Del Moral, Luc de Montella

► **To cite this version:**

Michel Caffarel, Pierre Del Moral, Luc de Montella. Time-uniform convergence rates for diffusion Monte Carlo with applications to coupled harmonic oscillators. 2024. hal-04409602v1

**HAL Id: hal-04409602**

**<https://hal.science/hal-04409602v1>**

Preprint submitted on 22 Jan 2024 (v1), last revised 15 Feb 2024 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Time-uniform convergence rates for diffusion Monte Carlo with applications to coupled harmonic oscillators

Michel Caffarel, Pierre Del Moral and Luc de Montella

January 22, 2024

## Abstract

The Diffusion quantum Monte Carlo method with constant number of walkers, also called Stochastic Reconfiguration as well as Sequential Monte Carlo, is a widely used genetic type Monte Carlo methodology for computing the ground-state energy and wave function of quantum systems. In this study, we present a general probabilistic framework with easily checked regularity conditions that ensure the uniform-in-time convergence of Diffusion Monte Carlo estimates towards the top of the spectrum of Schrödinger operators. We develop a novel powerful Lyapunov approach that applies to non necessarily compact state spaces, including linear diffusions evolving in quadratic absorbing potentials. We also illustrate the impact of these results in the context of coupled quantum harmonic oscillators, yielding what seems to be the first result of this type for this class of models.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Description of the models</b>	<b>3</b>
2.1	Free evolution semigroups . . . . .	3
2.2	Feynman-Kac semigroups . . . . .	4
2.3	Schrödinger semigroups . . . . .	6
2.4	Diffusion Quantum Monte Carlo . . . . .	8
<b>3</b>	<b>Statement of the main results</b>	<b>9</b>
3.1	Regularity conditions . . . . .	9
3.2	A time-uniform convergence Theorem . . . . .	10
3.3	Coupled harmonic oscillators . . . . .	11
<b>4</b>	<b>Stochastic interpolation</b>	<b>14</b>
4.1	Time varying semigroups . . . . .	14
4.2	Ground state estimates . . . . .	17

<b>5 Applications</b>	<b>17</b>
5.1 Coupled harmonic oscillators . . . . .	17
5.2 An Importance Sampling technique . . . . .	20
5.3 A divergence property . . . . .	22

## 1 Introduction

The many-body Schrödinger equation describes interacting quantum particles. In material sciences, these particles represent electrons and protons, as well as atoms in astrophysics. For realistic quantum systems, it is impossible to solve this equation analytically. The diffusion quantum Monte Carlo method (abbreviated DMC) provides a powerful stochastic approach to numerically approximate the ground state energy and wave function of Schrödinger operators.

The DMC methodology has a long and rich history, dating back to its first mention in 1949 by Ulam and Metropolis in [1]. The idea was first implemented by Donsker and Kac [2], and by Kalos [3] in the early 1960s. This methodology has been used to study fluxional molecules [4], to estimate the charge density of Lithium-ion battery [5], as well as to study the electronic structure of quantum dots [6]. For a more detailed discussion on the origins and the applications of these Monte Carlo techniques in physics we refer the reader to the more recent review article [7] as well as to [8, 9] and references therein.

Several variations of the method have been developed, see for example [10]. The version of interest here is the one with a fixed number of walkers, commonly known as *stochastic reconfiguration*; see the pioneering article by Hetherington [11], followed by Sorella and co-authors [12, 13] and by the first author and his co-workers in [14, 15, 16, 17]. Depending on the discipline, the methodology may be referred to by different names, such as genetic algorithm with selection and mutation, population Monte Carlo or sequential Monte Carlo [18, 19, 20, 21]. For a more thorough discussion on these application model areas we refer to the books [22, 23] and references therein.

These sequential Monte Carlo methods don't rely on biased variational techniques. They can be seen as a sophisticated genetic-type Monte Carlo methodology to simulate interacting quantum many-body systems. Various asymptotic results have been derived, including central limit theorems and large deviation principles, see for instance [24, 25] and [26, 27], as well as the books [28, 22, 23] for an overview.

Our work concerns less studied non-asymptotic and time-uniform problems. Recalling that the estimation of ground state energies relies on the limiting behavior of the walkers' evolution in the DMC method, it is therefore crucial to obtain uniform-in-time convergence estimates. Despite its importance, there is a notable gap in the literature and very few results have been proven in this respect. To the best of our knowledge, such uniform controls are mainly valid for compact state space models, see for instance [29, 22, 23] as well as [30]. Surprisingly, the theoretical efficiency of the method has never been verified even in basic linear-Gaussian scenarios such as the simple and well known harmonic oscillator.

In this paper, we address this gap by establishing the first uniform-in-time convergence estimates that apply to general state space models including the coupled harmonic oscillators presented in [31]. Our approach is partly based on recent developments on the stability of positive semigroups presented in [32, 33].

Our study leads us to conjecture that stable Markov transitions are necessary for the DMC method to be uniformly convergent. We thus prove this assumption in the simple linear Gaussian case. Additionally, we propose and, to some extent, establish the validity of an importance sampling transformation to overcome this difficulty. This type of technique, where a guiding wave function is used to direct the Monte Carlo moves, is commonly used to improve the efficiency of the DMC method [34, 35].

The paper is organised as follows: In Section 2, we provide a detailed description of the general framework in which our study is set, as well as the theoretical foundations on which our proof will be based.

Section 3 is devoted to the presentation of our main results. In section 4 we focus on proving convergence in the general framework.

Section 5 is devoted to the application of our convergence result to coupled harmonic oscillators [36, 37]. These models arise in various fields such as quantum optics [38], quantum cryptography [39] and photosynthesis [40]. In signal processing, the harmonic oscillator and the DMC methods coincides with the Kalman and the particle filter [41, 42].

## 2 Description of the models

### 2.1 Free evolution semigroups

Consider a Markov chain  $X_n$  indexed by  $n \in \mathbb{N}$  and taking values in a locally compact Polish space  $(E, \mathcal{E})$ , where  $\mathcal{E}$  is the Borel  $\sigma$ -field on  $E$ . Let  $\mathcal{C}(E)$  be the algebra of continuous measurable functions on  $E$ . We also define  $\mathcal{C}_b(E) \subset \mathcal{C}(E)$  as the sub-algebra of bounded measurable continuous functions endowed with the supremum norm  $\|\cdot\|$ . With a slight abuse of notation, we denote by 0 and 1 the null and unit scalars as well as the null and unit functions on  $E$  and we denote by  $I : x \in E \mapsto I(x) = x$  the identity function on  $E$ .

For  $n \in \mathbb{N}^*$ , we consider the Markov transitions  $P_n$  associated with  $X_n$ , and assume that they are Feller; in the sense that for any  $f \in \mathcal{C}_b(E)$  we have  $P_n(f) \in \mathcal{C}_b(E)$ , with the function  $P_n(f)$  defined for any  $x \in E$  by the integral operator

$$P_n(f)(x) := \int_E P_n(x, dy) f(y) = \mathbb{E}(f(X_n) \mid X_{n-1} = x)$$

Let  $\mathcal{C}_\infty(E) \subset \mathcal{C}(E)$  be the sub-algebra of uniformly positive continuous functions  $V$  that grow at infinity; that is, for any  $r \geq V_\star := \inf_E V > 0$ , the  $r$ -sub-level set  $\mathcal{V}(r) := \{V \leq r\} \subset E$  is a non-empty compact subset. We further assume that there exists a  $P$ -Lyapunov function  $V \in \mathcal{C}_\infty(E)$ ; in the sense there exists  $\epsilon \in [0, 1)$  and  $c \in \mathbb{R}$  such that for any  $n \in \mathbb{N}^*$  we have

$$P_n(V) \leq \epsilon V + c. \quad (1)$$

Let  $\mathcal{C}_V(E) \subset \mathcal{C}(E)$  be the sub-space of functions  $f \in \mathcal{C}(E)$  such that  $f/V$  is bounded, equipped with the norm  $\|f\|_V := \|f/V\|$ . The Markov semigroup associated with the Markov chain  $X_n$  is defined for any  $f \in \mathcal{C}_V(E)$  by

$$P_{k,n}(f)(x) := \mathbb{E}(f(X_n) \mid X_k = x).$$

Condition (1) ensures that  $P_{k,n}$  is  $V$ -Feller in the sense that for  $f \in \mathcal{C}_V(E)$  we have  $P_{k,n}(f) \in \mathcal{C}_V(E)$ . To ensure the semigroup  $P_{k,n}$  is exponentially stable [32], we assume the integral operator  $P_n(x, dy) = p_n(x, y)\nu(dy)$  has a density  $p_n$  w.r.t. some Radon measure  $\nu$  satisfying for some  $r_1 > 0$  and for any  $r \geq r_1$  the local minorization condition

$$0 < \inf_{n \in \mathbb{N}^*} \inf_{\mathcal{V}(r)^2} p_n \leq \sup_{n \in \mathbb{N}^*} \sup_{\mathcal{V}(r)^2} p_n < \infty \quad \text{and} \quad 0 < \nu(\mathcal{V}(r)) < \infty. \quad (2)$$

The  $V$ -norm semigroup contraction techniques developed in Section 8.2 in [43] (see also Lemma 2.3 in [33] and Theorem 2.2 in [32]), ensure that for any  $\mu \in \mathcal{P}_V(E)$ , there exists some parameters  $a, b > 0$  such that for any  $k \leq n$ , and any  $\mu_1, \mu_2 \in \mathcal{P}_V(E)$  we have

$$\|\mu_1 P_{k,n} - \mu_2 P_{k,n}\|_V \leq a e^{-b(n-k)} \|\mu_1 - \mu_2\|_V. \quad (3)$$

Note that the r.h.s condition in (2) is met as soon as  $V$  has compact sub-level sets with non empty interior and  $\nu$  is a Radon measure with full support; that is  $\nu$  is finite on compact sets and strictly positive on non-empty open sets. For time-homogeneous models, the l.h.s. minorization condition is satisfied as soon as  $(x, y) \in (E^\circ)^2 \mapsto p_n(x, y)$  is a continuous positive function on the interior  $E^\circ$  of the set  $E$ .

## 2.2 Feynman-Kac semigroups

We associate with a sequence of strictly positive functions  $(G_n)_{n \in \mathbb{N}} \in \mathcal{C}_V(E)^\mathbb{N}$  the discrete generation Feynman-Kac semigroups

$$Q_{k,n}(f)(x) = \mathbb{E} \left( f(X_n) \prod_{p=0}^{n-1} G_p(X_p) \mid X_k = x \right) \quad \text{and} \quad \widehat{Q}_{k,n}(f) := Q_{k,n}(G_n f).$$

To simplify notation, for  $k = (n - 1)$  sometimes we write  $Q_n$  instead of  $Q_{n-1,n}$ . We also use the convention  $Q_{n,n} = P_{n,n} = Id$ , the identity operator.

Let  $\mathcal{M}_b(E)$  be the set of bounded signed measures on  $E$ . Also, let  $\mathcal{P}(E) \subset \mathcal{M}_b(E)$  be the convex subset of probability measures on  $E$  and denote by  $\mathcal{P}_V(E)$  the convex set of probability measures  $\mu \in \mathcal{P}(E)$  such that  $\mu(V) < +\infty$ . The left action of  $Q_n$  on  $\mathcal{P}_V(E)$  is given for any  $(\eta, f) \in (\mathcal{P}_V(E), \mathcal{C}_V(E))$  by the formula

$$(\eta Q_n)(f) := \eta(Q_n(f)) = \int \eta(dx) Q_n(f)(x) = \int \eta(dx) Q_n(x, dy) f(y). \quad (4)$$

By Fubini's theorem, the integration order doesn't matter. Thus to simplify notation, we sometimes write  $\eta Q_n(f)$  instead of  $(\eta Q_n)(f)$  or  $\eta(Q_n(f))$ .

We denote by  $\mathcal{C}_0(E) := \{1/V : V \in \mathcal{C}_\infty(E)\} \subset \mathcal{C}_b(E)$  the sub-algebra of bounded continuous positive functions  $h$  that vanish at infinity; that is, for any  $0 < \epsilon \leq \|h\| < \infty$  the  $\epsilon$ -super-level set  $\{h \geq \epsilon\} \subset E$  is a non empty compact subset.

We further assume the Lyapunov function  $V$  introduced in (1) is a  $Q$ -Lyapunov function in the sense that (1) holds and there exists  $\Theta \in \mathcal{C}_0(E)$  and a compact subset  $K \subset E$  such that for any  $n \geq 1$  we have

$$Q_n(V)/V \leq \Theta \quad \text{and} \quad (G_{n-1}(x) - G_{n-1}(y))(\mathbb{1}_{E \setminus K}(x)V(x) - \mathbb{1}_{E \setminus K}(y)V(y)) \leq 0. \quad (5)$$

Note that the l.h.s. condition in (5) holds as soon as there exists  $G \in \mathcal{C}_0(E)$  such that for any  $G_n \leq G$ , for any  $n \geq 0$ . This condition ensures that for any positive function  $f \in \mathcal{C}_V(E)$  and  $n \geq 1$  we have

$$Q_n(f)/V \leq Q_n(V)/V \leq \Theta.$$

By (2) the integral operator  $Q_n(x, dy) = q_n(x, y) \nu_n(dy)$  also has a density given by  $q_n(x, y) = G_{n-1}(x)p_n(x, y)$  and for any  $r \geq r_1$  we have the local condition

$$0 < \inf_{\mathcal{V}(r)^2} q_n \leq \sup_{\mathcal{V}(r)^2} q_n < \infty.$$

Consider the normalized measure valued process  $\eta_n \in \mathcal{P}_V(E)$  starting at  $\eta_0 \in \mathcal{P}_V(E)$  defined for any  $n \geq 1$  by

$$\eta_{n+1} = \phi_{n+1}(\eta_n) := \psi_{G_n}(\eta_n)P_{n+1} \quad \text{and} \quad \hat{\eta}_n := \psi_{G_n}(\eta_n) \quad (6)$$

with the updated Boltzmann-Gibbs transformations  $\psi_{G_n}$  associated with the potential function  $G_n$  defined by

$$\psi_{G_n}(\eta_n)(dx) := \frac{1}{\eta(G_n)} G_n(x) \eta(dx).$$

We readily check that evolution semigroup  $\phi_{k,n} = \phi_{k+1,n} \circ \phi_k$  associated with the flow of measure  $\eta_n$  is given for any  $k \leq n$  by the formula

$$\phi_{k,n}(\eta_k) = \frac{\eta_k Q_{k,n}}{\eta_k Q_{k,n}(1)}$$

Note that for any  $\mu \in \mathcal{P}_V(E)$  we have the updating formula

$$\phi_{k,n}(\eta_k) = \psi_{H_{k,n}^\mu}(\eta_k) \bar{Q}_{k,n} \quad (7)$$

with the Markov operator

$$\bar{Q}_{k,n}(f) := Q_{k,n}(f)/Q_{k,n}(1) \quad \text{and} \quad H_{k,n}^\mu(x) := \frac{Q_{k,n}(1)(x)}{\phi_{0,k}(\mu)Q_{k,n}(1)}. \quad (8)$$

By Theorem 4.2 in [33], for any  $\mu \in \mathcal{P}_V(E)$ , there exists some parameters  $a, b > 0$  such that for any  $k \leq n$ , and any  $\mu_1, \mu_2 \in \mathcal{P}_V(E)$  we have

$$\|\mu_1 \bar{Q}_{k,n} - \mu_2 \bar{Q}_{k,n}\|_V \leq a e^{-b(n-k)} \|\mu_1 - \mu_2\|_{V/H_{k,n}^\mu}. \quad (9)$$

Note that for constant potential functions  $G_n(x) = G_n(y)$  we have  $\bar{Q}_{k,n} = P_{k,n}$  and the above contraction estimates resume to (3). For time homogeneous models  $Q_n = Q$ , Theorem 4.4 in [33] ensures the existence of a leading eigen triple  $(h, E_0, \eta_\infty) \in (C_V(E) \times \mathbb{R}_+^* \times \mathcal{P}_V(E))$ , such that

$$Q(h) = E_0 h \quad , \quad \eta_\infty Q = E_0 \eta_\infty \quad \text{and} \quad \eta_\infty(h) = 1.$$

### 2.3 Schrödinger semigroups

The objects defined in the previous subsection are core to a variety of physics problem. Indeed, consider an Hamiltonian differential operator  $\mathcal{H}$  given by the formula

$$\mathcal{H} := \mathcal{L} + V,$$

where  $V$  is a potential energy function from  $E$  to  $\mathbb{R}_+$ , and  $\mathcal{L}$  is a kinetic energy operator acting on a subset  $\mathcal{D}(\mathcal{L})$  of  $\mathcal{C}(E)$ . The time dependent Schrödinger equation and the imaginary time version associated with the hamiltonian  $\mathcal{H}$  are given, respectively, by the equations

$$i \partial_t \psi_t(x) = \mathcal{H}(\psi_t)(x) \quad \text{and} \quad -\partial_t \varphi_t(x) = \mathcal{H}(\varphi_t)(x),$$

with prescribed initial conditions  $(\psi_0, \varphi_0)$ . In the above display,  $i \in \mathbb{C}$  stands for the imaginary unit. The right-hand side equation is obtained via a formal time change by setting  $\varphi_t(x) = \psi_{-it}(x)$ , and can be equivalently written in the following form

$$\partial_t \varphi_t(x) = \mathcal{L}(\varphi_t)(x) - V(x) \varphi_t(x) \quad \text{with initial condition } \varphi_0. \quad (10)$$

For a twice differentiable function  $\varphi_0$ , the solution of (10) is given by the Feynman-Kac path integral formula

$$\begin{aligned} \varphi_t(x) &= Q_t(\varphi_0)(x) := \int Q_t(x, dy) \varphi_0(y) \\ &= \mathbb{E} \left( \varphi_0(X_t) \exp \left( - \int_0^t V(X_s) ds \right) \mid X_0 = x \right). \end{aligned} \quad (11)$$

In the above display,  $X_t$  stands for a time homogeneous stochastic process  $X_t$  on  $E$ , with generator  $\mathcal{L}$ . To facilitate the interpretation of the theoretical and numerical physics in the measure theoretical framework used in this article, we note that the Feynman-Kac propagator defined by the integral operator (11) is sometimes written in terms of the exponential of the Hamiltonian operator with the exponential-type symbol

$$Q_t := e^{-t\mathcal{H}} \quad \text{or in the bra-kets formalism} \quad Q_t(\varphi_0) = |e^{-t\mathcal{H}} \varphi_0\rangle.$$

The exponential notation is compatible with finite space models and the matrix notation of the continuous one-parameter semigroup for time homogenous models. The bra-ket (or Dirac) notation is also used to represent linear projection forms acting on Hilbert spaces associated with some reversible or some stationary measure, such as the Lebesgue measure for the harmonic oscillator.

The integral operator  $Q_t$  is sometimes called the Feynman-Kac propagator. For any  $s, t \geq 0$  the integral operators  $Q_t$  satisfy the semigroup property

$$Q_{s+t}(x, dz) = (Q_s Q_t)(x, dy) := \int Q_s(x, dz) Q_t(z, dy) \implies \varphi_{s+t} = Q_s(\varphi_t).$$

In terms of left action bra-kets, defining  $\mu_\varphi(dx) := \varphi(x)dx$ , Fubini's theorem yields

$$\begin{aligned} \langle \varphi | e^{-s\mathcal{H}} | \varphi_t \rangle &= \int dx \varphi(x) Q_s(x, dy) \varphi_t(dy) = (\mu_\varphi Q_s)(\psi_t) \\ &= \mu_\varphi((Q_s Q_t)(\varphi_0)) = \mu_\varphi Q_{s+t}(\varphi_0) = \langle \varphi | e^{-(s+t)\mathcal{H}} | \varphi_0 \rangle. \end{aligned}$$

The exponential notation is compatible with finite space models and the matrix notation of the continuous one-parameter semigroup for time homogeneous models. The bra-ket notation (a.k.a. Dirac notation) is also used to represents linear projection forms acting on Hilbert spaces associated with some reversible or some stationary measure, such as the Lebesgue measure for the harmonic oscillator.

The present article deals with different types of non necessarily stationary stochastic processes, including the free evolution process  $X_t$  discussed in (11). Apart in the reversible situation in which spectral theorems are stated on the Hilbert space associated with a reversible measure, the use of the exponential symbol or the use of the brackets formalism is clearly not adapted to represent different expectations with respect to different types of stochastic and non-necessarily reversible processes.

To analyze these general stochastic models, we have chosen to only use elementary and standard measure theory notation such as (4). The integral actions of a given integral operator  $Q_t$  on the right for functions and on the left for measures are clearly compatible with finite space models and matrix notation. The left action  $\mu \mapsto \mu Q_t$  maps measures into measures, while the right action  $f \mapsto Q_t(f)$  maps functions into functions.

The operator  $Q_t$  defined in (11) is sometimes called a Feynman-Kac propagator. However, despite its mathematical elegance, it can rarely be solved analytically.

The Feynman-Kac measure  $\eta_n$  introduced in (6) can be interpreted as the solution of a discrete-time approximation of the formula (11). Indeed, on a given time mesh  $t_n = n\lfloor t/n \rfloor$ , choosing  $G = e^{-V \Delta t}$  we have

$$\begin{aligned} &\mathbb{E} \left( f(X_{t_n}) \prod_{0 \leq k < n} G(X_{t_k}) \right) \\ &= \mathbb{E} \left( f(X_{t_n}) \exp \left\{ - \sum_{0 \leq t_k < t_n} V(X_{t_k})(t_{k+1} - t_k) \right\} \right) \simeq_{\Delta t \downarrow 0} \eta_0 Q_t(f). \end{aligned}$$



## 2.4 Diffusion Quantum Monte Carlo

The Diffusion Quantum Monte Carlo methodology relies on the fact that the flow of measures  $\eta_n$  introduced in (6) can be interpreted as the probability distributions  $\eta_n = \text{Law}(\bar{X}_n)$  of the random states  $\bar{X}_n$  of a nonlinear Markov chain  $\bar{X}_n$ . The choice of the Markov chain is far from unique. For instance, we have

$$\eta_{n+1} = \phi_{n+1}(\eta_n) = \eta_n K_{n+1, \eta_n}$$

with the local Markov transition

$$\begin{aligned} K_{n+1, \eta_n}(x, dz) &:= (S_{n, \eta_n} P_{n+1})(x, dz) \\ &:= \int S_{n, \eta_n}(x, dy) P_{n+1}(y, dz) = \mathbb{P}(\bar{X}_{n+1} \in dz \mid \bar{X}_n = x) \end{aligned}$$

In the above display,  $S_{n, \eta_n}$  stands for the Markov transition

$$S_{n, \eta_n}(x, dy) := \epsilon_n(\eta_n) G_n(x) \delta_x(dy) + (1 - \epsilon_n(\eta_n) G_n(x)) \psi_{G_n}(\eta_n)(dy).$$

for some tuning parameter  $\epsilon_n(\eta_n) \in [0, 1]$  chosen such that  $\epsilon_n(\eta_n) G_n(x) \in [0, 1]$ . For instance, for  $]0, 1[$ -valued potential functions, we can choose  $\epsilon_n(\eta_n) = 0$  as well as  $\epsilon_n(\eta_n) = 1$ . For more general models, we can also choose the inverse of the  $\eta_n$ -essential supremum of  $G_n$ .

Note that the transition  $\bar{X}_n \rightsquigarrow \bar{X}_{n+1}$  depends on the probability distributions  $\eta_n$  of the random states  $\bar{X}_n$ . In reference with similar nonlinear Markov chain models arising in fluid mechanics, the Markov chain  $\bar{X}_n$  is called a McKean interpretation of the flow of measures (6).

The mean field particle interpretation associated with a given McKean model is defined by a discrete-time system of  $N$  walkers  $\xi_n = (\xi_n^i)_{1 \leq i \leq N}$ . The system starts with  $N$  independent copies of a random variable  $\bar{X}_0 = X_0$  with distribution  $\eta_0$ . Given the system  $\xi_n$  at some time  $n \geq 0$ , we sample  $N$  conditionally independent walkers  $\xi_{n+1}^i$  with their respective distribution

$$K_{n+1, \eta_n^N}(\xi_n^i, dx) \quad \text{with} \quad \eta_n^N := \frac{1}{N} \sum_{1 \leq i \leq N} \delta_{\xi_n^i}$$

In other words, the DMC method consists of approximating the measure  $\eta_n$  by using the occupation measure  $\eta_n^N$  associated with a system of  $N$  walkers. The initial positions of the walkers are randomly chosen from the distribution  $\eta_0$ . The evolution of each walker follows then the following selection/mutation steps:

- **Selection:** We evaluate the current position  $\xi_n^i$  of a walker and its potential value  $G_n(\xi_n^i)$ . With probability  $(1 - \epsilon_n(\eta_n^N) G_n(\xi_n^i))$ ,  $\xi_n^i$  is killed and instantly replaced by another walker say  $\hat{\xi}_n^i = \xi_n^j$  with a probability proportional to  $G_n(\xi_n^j)$  and  $j \in \{1, \dots, N\}$ ; otherwise we keep it and set  $\hat{\xi}_n^i := \xi_n^i$ .

- Mutation: We move the selected walker  $\widehat{\xi}_n^i = x$  to a new location  $\xi_{n+1}^i = y$  using the transition kernel  $P_{n+1}(x, dy)$ .

The selection transition associated with the choice  $\epsilon_n(\eta_n^N) = 0$  coincides with the so-called proportional selection/reconfiguration. Note that the walker with the highest potential value is always selected when  $\epsilon_n(\eta_n^N)$  is the inverse of the  $\eta_n^N$ -essential supremum of  $G_n$ . For  $[0, 1]$ -valued potential functions  $G_n$  we can also choose  $\epsilon_n(\eta_n^N) = 1$ . In this situation, particles are killed at a geometric clock that depends on the potential function.

We expect that the occupation measures of the system approximate the solution of the measure-valued process (6); that is, in a sense to be given, for any time horizon  $n \geq 0$  we have

$$\eta_n^N = \frac{1}{N} \sum_{i=1}^N \delta_{\xi_n^i} \xrightarrow{N \rightarrow \infty} \eta_n$$

as well as

$$\psi_{G_n}(\eta_n^N) = \sum_{i=1}^N \frac{G_n(\xi_n^i)}{\sum_{1 \leq j \leq N} G_n(\xi_n^j)} \delta_{\xi_n^i} \xrightarrow{N \rightarrow \infty} \psi_{G_n}(\eta_n) = \widehat{\eta}_n.$$

### 3 Statement of the main results

#### 3.1 Regularity conditions

For  $f \in \mathcal{C}_b(E)$ , time-uniform  $L_p$ -convergence of the error made by the DMC method in estimating  $\eta_n(f)$  have been obtained (see for example [29]) under the assumption that there exists  $\epsilon_G \in \mathbb{R}_+^*$  such that

$$\forall (x, y, n) \in E^2 \times \mathbb{N}, \quad G_n(x) \geq \epsilon_G G_n(y)$$

A significant consequence of this assumption is a bound for any finite horizon  $n \in \mathbb{N}$  on the random potential function defined for some  $\gamma \in \mathcal{P}_V(E)$  and  $k \in \mathbb{N}$  by

$$G_{k,k+n}^\gamma : x \in E \mapsto \frac{1}{H_{k,k+n}^\gamma(x)}$$

In order to guarantee a time-uniform  $L_p$ -convergence, our framework requires not only the preservation of this bound, but also its uniformity in time. To do this, we first assume that there exists  $\gamma \in \mathcal{P}_V(E)$  such that

$$\sup_{n \in \mathbb{N}} \phi_{0,n}(\gamma)(G_n) < +\infty \tag{12}$$

For time-homogeneous models, without any further conditions on the potential, condition (12) is easily checked with  $\gamma = \eta_\infty$ . For this scenario, we will then consider in the rest that  $\gamma = \eta_\infty$ . Moreover, this hypothesis trivially holds if the functions  $G_n$  are bounded by some constant independent of  $n$ .

We assume that there exists  $W \in \mathcal{C}_V(E)$ ,  $\alpha \in ]0, 1]$  and a  $Q$ -Lyapunov function  $\bar{W} \in \mathcal{C}_V(E)$  such that

$$Q_n(W) \geq \bar{P} \times W \quad \text{and} \quad W^{-\alpha} \leq \bar{W}, \quad (13)$$

where

$$\bar{P} := \sup_{(n,x) \in \mathbb{N}^* \times E} P_n(G_n)(x).$$

For time homogeneous models, this condition can be relaxed into the following

$$Q(W) \geq \min \{ \bar{P}, E_0 \} \times W \quad \text{and} \quad W^{-\alpha} \leq \bar{W}. \quad (14)$$

Note that, for time-homogeneous models, the set of functions  $W \in \mathcal{P}_V(E)$  such that  $Q(W) \geq E_0 W$  is non-empty as it contains at least the ground state  $h$ . It is also worth noting that it is not necessary to know the exact value of  $\bar{P}$  nor the one of  $E_0$  in order to prove that (13) or (14) hold. Indeed, if one of these constants is less than some  $C \in \mathbb{R} \cup \{+\infty\}$ , then it is sufficient to prove that for any  $c < C$ , there exist  $W_c \in \mathcal{C}_V(E)$  and  $\alpha_c \in (0, 1]$  such that

$$Q(W_c) \geq c W_c \quad \text{and} \quad W_c^{-\alpha_c} \leq V.$$

Finally, we assume that there exists a  $Q$ -Lyapunov function  $\bar{V} \in \mathcal{C}(E)$  and  $\lambda \in 2\mathbb{N}$  such that

$$V^\lambda \leq \bar{V}. \quad (15)$$

Without further mention, we assume that  $V, \bar{V}$  and  $\bar{W}$  are integrable with respect to  $\eta_0$ , i.e.,  $\eta_0 \in \mathcal{P}_V(E) \cap \mathcal{P}_{\bar{W}}(E) \cap \mathcal{P}_{\bar{V}}(E)$ . Under conditions (12) and (13) or (14), it is then possible to obtain a time uniform bound on the random potential function  $G_{k,n}^\gamma$ .

**Lemma 1.** *Let  $\gamma$  and  $(\eta, \mu)$  be defined as in (12). There exists  $\bar{\beta} \in \mathbb{R}_+^*$  such that for any  $\beta \leq \bar{\beta}$  we have*

$$\sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[ \phi_q(\eta_{q-1}^N) (H_{q,n}^\gamma)^{-\beta} \right] < +\infty \quad \text{and} \quad \sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[ \eta_q^N (H_{q,n}^\gamma)^{-\beta} \right] < +\infty. \quad (16)$$

The proof of this pivotal Lemma is postponed to the appendix.

### 3.2 A time-uniform convergence Theorem

The main goal of this paper is to establish that in the context we described, the  $L_p$ -norm of the error made by the DMC method in approximating the Feynman-Kac measures  $\eta_m$  remains bounded in time and converges to zero as the number of particles increases. Our main result can be stated as follows

**Theorem 1.** For any  $p \in \mathbb{N}^*$ , there exists  $c \in \mathbb{R}$  and  $\beta \in (0, 1]$  such that for any  $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$  we have

$$\sup_{n \in \mathbb{N}} \mathbb{E} (|\eta_n(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}.$$

The proof of this theorem is provided in subsection 4.1. For time-homogeneous models, a direct consequence of Theorem 1 is a control over the estimation of the limit measure  $\eta_\infty$ .

**Corollary 1.** Let  $p \in \mathbb{N}^*$ , there exists  $(a, b, c) \in \mathbb{R}_+^{*3}$  and  $\beta \in (0, 1]$  such that for any  $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$  we have

$$\sup_{n \geq a+b \ln(N)} \mathbb{E} (|\eta_\infty(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}.$$

The proof the above Corollary is provided in subsection 4.2 .

Assuming that there exists a measure  $\mu \in \mathcal{P}(E)$  that is reversible for  $P$ , it becomes possible to obtain a re-normalized weak form of the ground state  $h$  and its associated eigenvalue from the limit measures  $\eta_\infty$  and  $\hat{\eta}_\infty$  of  $\eta_n$  and  $\hat{\eta}_n$ . Indeed, referring to Section 9.5.5 in [43] (see also [18] as well as Section 12.4 in [22]) :

$$\eta_\infty(G) = E_0 \quad , \quad \eta_\infty(f) = \mu(P(h)f)/\mu(P(h)) \quad \text{and} \quad \hat{\eta}_\infty(f) = \mu(hf)/\mu(h).$$

Those equalities, combined with the convergence stated in Corollary 1, guarantee the efficiency of the DMC method for approximating the ground-state energy and wave function of quantum systems.

**Corollary 2.** Let  $p \in \mathbb{N}^*$ , and assume that  $G \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ . There exists  $(a, b, c) \in \mathbb{R}_+^{*3}$  and  $\beta \in (0, 1]$  such that for any  $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$  and  $\hat{f} \in \mathcal{C}_{V^{\frac{\lambda}{4p}}/G}(E)$ , we have

$$\left\{ \begin{array}{l} \sup_{n \geq a+b \ln(N)} \mathbb{E} (|E_0 - \eta_n^N(G)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}} \\ \sup_{n \geq a+b \ln(N)} \mathbb{E} (|\eta_\infty(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}} \\ \sup_{n \geq a+b \ln(N)} \mathbb{E} (|E_0 \hat{\eta}_\infty(\hat{f}) - \eta_n^N(G\hat{f})|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}} \end{array} \right.$$

### 3.3 Coupled harmonic oscillators

To illustrate the practical applications of Theorem 1, we carry out an in-depth study of the coupled harmonic oscillator. First, we demonstrate its relevance by establishing easily verifiable sufficient conditions for time-uniform control of the DMC method in a framework that includes the harmonic oscillator.

We consider  $E = \mathbb{R}^d$  for some  $d \in \mathbb{N}^*$ . For any real definite positive  $d \times d$  matrices  $B$  and  $S$  and any a real  $d \times d$  matrix  $A$ , we denote by  $P_{A,B}$  and  $G_S$  the Markov kernel and the potential function defined as follows

$$P_{A,B}(x, dy) = \frac{1}{(2\pi)^{d/2} |B|^{1/2}} e^{-\frac{1}{2}(Ax-y)^\top B^{-1}(Ax-y)} \quad \text{and} \quad G_S(x) := \exp\left(-\frac{x^\top Sx}{2}\right).$$

Consider a family of Feller Markov transitions  $(P_n)_{n \in \mathbb{N}}$  with positive densities  $p_n$ , an initial distribution  $\eta_0 \in \mathcal{P}(E)$  and a family of positive functions  $(G_n)_{n \in \mathbb{N}} \in \mathcal{C}_0(\mathbb{R}^d)^{\mathbb{N}}$  that is uniformly bounded in time.

We assume the following, where  $(A, A')$  are real matrices,  $(B, B', S)$  real positive definite matrices,  $p_{A,B}$  the density of  $P_{A,B}$  and  $E_{A,B,S}$  the ground-state energy associated to the operators  $P_{A,B}$  and  $G_S$ .

- There exists  $c_1 \in \mathbb{R}$  such that for any  $(n, x, y) \in \mathbb{N} \times \mathbb{R}^{d^2}$  we have

$$p_n(x, y) \leq c_1 p_{A,B}(x, y). \quad (17)$$

- For any  $(n, x, y) \in \mathbb{N} \times \mathbb{R}^{d^2}$  we have

$$G_n(x) p_{n+1}(x, y) \geq \frac{\bar{P}}{E_{A',B',S}} \times G_S(x) p_{A',B'}(x, y) \quad (18)$$

- There exists a compact  $K' \subset \mathbb{R}^d$  such that for any  $(n, x, y) \in \mathbb{N} \times \mathbb{R}^{d^2}$  we have

$$[G_S(x) - G_S(y)][G_n^{-1}(x)\mathbb{1}_{\mathbb{R} \setminus K'}(x) - G_n^{-1}(y)\mathbb{1}_{\mathbb{R} \setminus K'}(y)] \geq 0. \quad (19)$$

- There exists  $c_3 \in \mathbb{R}_+^*$  such that  $G_S^{-c_3}$  is integrable with respect to  $\eta_0$ .

If  $P_n$  and  $G_n$  are time-independent, it is possible to replace  $\bar{P}$  in (18) by the ground state energy associated with  $P$  and  $G$ .

These conditions hold trivially for the coupled harmonic oscillator, i.e if  $P_n = P_{A,B}$  and  $G_n = G_S$ .

In this context, Theorem 1 leads to a simple sufficient matrix condition which guarantees the uniform convergence of the DMC method. The proof of the following corollary can be found in subsection 5.1.

**Corollary 3.** *Assume that  $A^\top S A < S$ . In this situation, for any  $p \in \mathbb{N}^*$ , there exist  $(\beta, \alpha, c) \in (0, 1] \times \mathbb{R}_+^{*2}$  such that for any function  $f \in \mathcal{C}_V(\mathbb{R}^d)$  we have*

$$\sup_{n \in \mathbb{N}} \mathbb{E} (|\eta_n(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}, \quad \text{with } V : x \in \mathbb{R}^d \mapsto \exp\left(\frac{\alpha}{2}x^\top Sx\right).$$

Shifting the focus to the approximation of the measures  $(\widehat{\eta}_n)_{n \in \mathbb{N}}$  within the coupled harmonic oscillator framework, the convergence condition given in the previous corollary can be overcome with a change of transition and selection in the DMC method. Considering  $P = P_{A,B}$ ,  $G = G_S$  as well as  $\eta_0 \sim \mathcal{N}(X_0, P_0)$  and assuming that  $A$ ,  $B$  and  $S$  can be diagonalized in the same basis, we recursively define the function  $\widehat{G}^{(k)} \in \mathcal{C}_V(E)$  and the Markov kernel  $P^{(k)}$  on  $E$  such that for all  $f \in \mathcal{C}_V(E)$  we have

$$\left\{ \begin{array}{l} \widehat{G}^{(1)} = P(G) \\ \widehat{P}^{(1)}(f) = \frac{P(fG)}{P(G)} \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} \widehat{G}^{(k)} = P(G \widehat{G}^{(k-1)}) \\ \widehat{P}^{(k)}(f) = \frac{\widehat{P}^{(k-1)}P(fG)}{\widehat{P}^{(k-1)}P(G)} \end{array} \right. .$$

We then have for  $f \in \mathcal{C}_V(E)$  the formula

$$\widehat{Q}_{0,k}(f) = \widehat{G}^{(k)} \widehat{P}^{(k)}(f). \quad (20)$$

For  $k \in \mathbb{N}^*$ , consider a system of walkers  $\widehat{\xi}_n^{(k)} = \left( \xi_n^{(k),i} \right)_{1 \leq i \leq N}$  associated to the DMC method with initial distribution  $\psi_G(\eta_0)$ , transitions  $\widehat{P}^{(k)}$  and selection function  $\widehat{G}^{(k)}$  as well as the empirical measures

$$\widehat{\eta}_n^{(k),N} := \frac{1}{N} \sum_{1 \leq i \leq N} \delta_{\widehat{\xi}_n^{(k),i}}. \quad (21)$$

This system of walkers offers an approximation of the measures  $\widehat{\eta}_n$  for any  $n \in k\mathbb{N}$ . This type of change in the approximation, based on an importance sampling transformation is analogous to using a guiding waves function to direct the Monte Carlo moves. Without any additional condition, Theorem 1 ensures the uniform convergence of the model. The details of the proof can be found in Subsection 5.1.

**Corollary 4.** *Let  $p \in \mathbb{N}^*$ , there exists  $\bar{k} \in \mathbb{N}$  such that for any  $k \geq \bar{k}$ , there exist  $(\beta, \alpha, c) \in (0, 1] \times \mathbb{R}_+^{*2}$  satisfying for any  $f \in \mathcal{C}_V(\mathbb{R})$*

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left( |\widehat{\eta}_{nk}(f) - \eta_n^{(k),N}(f)|^p \right)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}, \quad \text{with } V : x \in \mathbb{R}^d \mapsto \exp\left(\frac{\alpha}{2}x^T Sx\right).$$

Although the method does not provide an approximation for every time step, several strategies can be used to fill the gaps left by the approximation. A simple approach, though more computationally intensive, is to run independent systems of walkers for each time step in the interval  $\llbracket 0, k-1 \rrbracket$ . This method not only fills in the gaps, but also maintains the convergence property.

Our study concludes with a focus on the divergence of the DMC method when approximating the one-dimensional harmonic oscillator. This confirms that the stability condition stated in Corollary 3 is necessary and that, in some cases, the set of assumptions presented can closely approximate a sufficient and necessary condition. Additionally, it emphasizes the significance of the importance sampling method introduced in the

previous corollary. Specifically, in the one-dimensional context, the sufficient condition for uniform convergence of the DMC method is expressed by  $A^2 < 1$ . Proposition 1 establishes the divergence of the error made by the DMC method when  $A^2 > 1$ , leaving open only the case  $A = 1$ .

**Proposition 1.** *Assume that  $A^2 > 1$  and  $P_0 > 0$ . For any  $p \in \mathbb{N}^*$  we have*

$$\sup_{n \in \mathbb{N}} \mathbb{E} (|\eta_n(I) - \eta_n^N(I)|^p)^{\frac{1}{p}} = +\infty.$$

The proof of this proposition can be found in Subsection 5.3.

Note that all corollaries in this subsection can be extended to a control on the estimation of the limit measures, ground state, and eigenvalue using the same approach as presented in Corollary 2.

## 4 Stochastic interpolation

### 4.1 Time varying semigroups

In this subsection, we focus on proving Theorem 1. To take advantage of the conditional independence of the walkers, we structure our approach around the following decomposition of the difference between the Feynman-Kac measure and its empirical approximation, using the convention  $\eta_{-1}^N = \eta_0$ . Following [28, 29], we use the following stochastic interpolation formula

$$\eta_n^N - \eta_n = \sum_{q=0}^n [\phi_{q,n}(\eta_q^N) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))]. \quad (22)$$

Each term on the right-hand side represents the error that occurs when using the DMC approximation instead of the real propagator for a single extra time step. Combining the uniform bound given in Lemma 1 with the contraction property (9), the following Lemma establishes an exponentially decreasing control for these local errors.

**Lemma 2.** *For any  $p \in \mathbb{N}^*$ , there exists  $(c, \rho, \beta) \in \mathbb{R}_+^{*2} \times (0, 1]$  such that for any function  $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$  and any  $(N, q, n) \in \mathbb{N}^3$  with  $q \leq n$  we have*

$$\mathbb{E} \left[ |[\phi_{q,n}(\eta_q^N) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))](f)|^p \right]^{\frac{1}{p}} \leq ce^{-(n-q)\rho} N^{-\frac{\beta}{2}}. \quad (23)$$

**Proof:**

Let  $(\eta, \mu) \in \mathcal{P}(E)$  and let  $\gamma \in \mathcal{P}_V(E)$  be defined as in (12). Consider  $H_{q,n} := H_{q,n}^\gamma$  as defined in (8). Applying the updating formula (7), we obtain

$$\begin{aligned}
\phi_{q,n}(\eta)(f) - \phi_{q,n}(\mu)(f) &= (\psi_{H_{q,n}}(\eta)\bar{Q}_{q,n} - \psi_{H_{q,n}}(\mu)\bar{Q}_{q,n})(f) \\
&= \frac{1}{\eta(H_{q,n})}(\eta - \mu) (H_{q,n}\bar{Q}_{q,n}[f - \psi_{H_{q,n}}(\mu)\bar{Q}_{q,n}(f)]) \\
&= \frac{1}{\eta(H_{q,n})}(\eta - \mu)(F_{q,n}^\mu),
\end{aligned}$$

with

$$F_{q,n}^\mu(x) := H_{q,n}(x) \int_E \psi_{H_{q,n}}(\mu)(dy) [\bar{Q}_{q,n}(f)(x) - \bar{Q}_{q,n}(f)(y)].$$

Then, applying Hölder's inequality for any  $\beta \in [0, 1)$ , we obtain

$$\begin{aligned}
&\mathbb{E} (|\phi_{q,n}(\eta)(f) - \phi_{q,n}(\mu)(f)|^p)^{\frac{1}{p}} \\
&\leq \mathbb{E} \left( \eta(H_{q,n})^{-2p} |(\eta - \mu)(F_{q,n}^\mu)|^{2p(1-\beta)} \right)^{\frac{1}{2p}} \mathbb{E} \left( |(\eta - \mu)(F_{q,n}^\mu)|^{2\beta p} \right)^{\frac{1}{2p}} \\
&\leq \mathbb{E} \left( \eta(H_{q,n})^{-2\beta p} |\phi_{q,n}(\eta)(f) - \phi_{q,n}(\mu)(f)|^{2p(1-\beta)} \right)^{\frac{1}{2p}} \mathbb{E} \left( |(\eta - \mu)(F_{q,n}^\mu)|^{2\beta p} \right)^{\frac{1}{2p}} \\
&\leq \left[ \mathbb{E} \left( \phi_{q,n}(\eta)(V^\lambda) \right)^{\frac{1}{4p}} + \mathbb{E} \left( \phi_{q,n}(\mu)(V^\lambda) \right)^{\frac{1}{4p}} \right] \mathbb{E} \left( \eta(H_{q,n})^{-4\beta p} \right)^{\frac{1}{4p}} \mathbb{E} \left( |(\eta - \mu)(F_{q,n}^\mu)|^{2\beta p} \right)^{\frac{1}{2p}}.
\end{aligned}$$

From Lemmas 1 and 8, we deduce that, to conclude, it is enough to prove that, for some constant  $c \in \mathbb{R}_+^*$  independent of  $n, q$  and  $N$ , we have

$$\mathbb{E} \left[ |[\eta_q^N - \phi_q(\eta_{q-1}^N)](F_{q,n})|^{2\beta p} \right]^{\frac{1}{2p}} < ce^{-c_2(n-q)} N^{-\frac{\beta}{2}}, \quad (24)$$

where  $F_{q,n} := F_{q,n}^{\phi_q(\eta_{q-1}^N)} \in \mathcal{C}_{V^2}$ .

Let  $\beta' = \frac{2p\beta}{\lambda}$  and assume  $\beta$  small enough so that  $\beta' < 1/2$ .

For  $q > 0$ , the walkers  $(\xi_q^i)_{1 \leq i \leq N}$  are independent conditionally to  $\eta_{q-1}^N$ , and for any function  $f \in \mathcal{C}_{V^2}(E)$ , we have

$$[\eta_q^N - \phi_q(\eta_{q-1}^N)](f) = \eta_q^N \left( \frac{1}{N} \sum_{1 \leq i \leq N} f_i \right),$$

with  $f_i = f - S_{q-1, \eta_{q-1}^N} P_q(f)(\xi_{q-1}^i)$ .

For  $q = 0$ , the walkers are iid with common distribution  $\eta_0$ .

Using the convention  $E(X | \xi_{-1}) = \mathbb{E}(X)$ , we can thus apply Lemma 7.3.3 from [22], and deduce that there exists  $C \in \mathbb{R}$  such that



$$\begin{aligned}
\mathbb{E} \left[ \left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}) \right|^{2\beta p} \right]^{\frac{1}{2p}} &= \mathbb{E} \left[ \mathbb{E} \left( \left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}) \right|^{\lambda\beta'} \mid \xi_{q-1} \right) \right]^{\frac{1}{2p}} \\
&\leq \mathbb{E} \left[ \mathbb{E} \left( \left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}) \right|^\lambda \mid \xi_{q-1} \right)^{\beta'} \right]^{\frac{1}{2p}} \\
&\leq \frac{C}{N^{\beta/2}} \mathbb{E} \left[ \phi_q(\eta_{q-1}^N) \left( F_{q,n}^\lambda \right)^{\beta'} \right]^{1/2p}.
\end{aligned}$$

Applying contraction property property (9) with  $\mu = \delta_x$  and  $\eta = \delta_y$  we get the existence of  $(a, \rho) \in \mathbb{R}_+^2$  such that

$$|\bar{Q}_{q,n}(f)(x) - \bar{Q}_{q,n}(f)(y)| \leq ae^{-\rho(n-q)} \left( 1 + \frac{V(x)}{H_{q,n}(x)} \right) \left( 1 + \frac{V(y)}{H_{q,n}(y)} \right). \quad (25)$$

By substituting (25) into the definition of  $F_{q,n}$  and applying Hölder's inequality along with Jensen's inequality, we obtain, for some  $(a', \rho') \in \mathbb{R}_+^{*2}$

$$\begin{aligned}
&\mathbb{E} \left[ \phi_q(\eta_{q-1}^N) \left( F_{q,n}^\lambda \right)^{\beta'} \right]^{1/2p} \\
&\leq a' e^{-\rho'(n-q)} \mathbb{E} \left[ \phi_q(\eta_{q-1}^N) \{ (H_{q,n} + V)^\lambda \}^{\beta'} \phi_q(\eta_{q-1}^N) \{ (H_{q,n} + V) \}^{\lambda\beta'} \phi_q(\eta_{q-1}^N) (H_{q,n})^{-\lambda\beta'} \right]^{\frac{1}{2p}} \\
&\leq a' e^{-\rho'(n-q)} \mathbb{E} \left[ \phi_q(\eta_{q-1}^N) \{ (H_{q,n} + V)^\lambda \} \right]^{\frac{1}{4p}} \mathbb{E} \left[ \phi_q(\eta_{q-1}^N) (H_{q,n})^{-2\lambda\beta'} \right]^{\frac{1}{4p}}.
\end{aligned}$$

From our hypothesis on  $Q_n$ , we deduce from Lemma 3.2 in [33] that there exists a constant  $c$  such that for any  $(q', n') \in \mathbb{N}^2$ ,  $H_{q',n'} \leq cV$ . We can then conclude by choosing a small enough  $\beta'$  and using Lemmas 1 and 8.  $\square$

The proof of Theorem 1 is now relatively straightforward.

### Proof of Theorem 1:

Let  $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ . From the sub-additivity of the  $L_p$ -norm applied in (22), we have

$$E(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} \leq \sum_{q=0}^n \mathbb{E} \left( \left| \phi_{q,n}(\eta_q^N)(f) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))(f) \right|^p \right)^{\frac{1}{p}}.$$

By applying Lemma 2, we deduce that there exists  $(C, \rho, \beta) \in \mathbb{R} \times \mathbb{R}_+^* \times (0, 1]$  such that for any  $N \in \mathbb{N}^*$  we have

$$\mathbb{E}(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} \leq \frac{C}{N^{\frac{\beta}{2}}} \sum_{0 \leq l \leq n} e^{-(n-l)\rho} \leq \frac{C}{N^{\frac{\beta}{2}}(1 - e^{-\rho})}.$$

This ends the proof of the theorem.  $\square$

## 4.2 Ground state estimates

This subsection concentrates on proving Corollary 1. We consider thus the time-homogeneous model.

Let  $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ . Notice that we can decompose the error made by the DMC method in the following way:

$$\mathbb{E}(|\eta_n^N(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq \mathbb{E}(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} + \mathbb{E}(|\eta_n(f) - \eta_\infty(f)|^p)^{\frac{1}{p}}.$$

Theorem 1 implies that there exists  $(C_1, \beta) \in \mathbb{R}^* \times (0, 1]$  such that

$$\sup_{n \in \mathbb{N}} \mathbb{E}(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} \leq \frac{C_1}{N^{\beta/2}}$$

According to Theorem 4.3 in [33], there exists  $(C_2, \omega) \in \mathbb{R}_+^{*2}$  such that

$$\mathbb{E}(|\eta_n(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq C_2 e^{-\omega n}$$

Hence

$$\mathbb{E}(|\eta_n^N(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq \frac{C_1}{N^{\beta/2}} + C_2 e^{-\omega n}$$

Thus, letting  $a = \frac{1}{\omega} \ln(C_2/C_1)$  and  $b = \frac{\beta}{2\omega}$ , we deduce that there exists  $C \in \mathbb{R}$  such that

$$\sup_{n \geq a + b \ln(N)} \mathbb{E}(|\eta_n^N(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq \frac{C}{N^{\beta/2}}$$

This concludes the proof.  $\square$

## 5 Applications

### 5.1 Coupled harmonic oscillators

This subsection is dedicated to the proof of Corollary 3. Therefore we place ourselves within the framework associated with this corollary. We only consider the general case where  $P$  and  $G$  depend on a time parameter. If this is not the case, and  $\bar{P}$  is replaced by the ground state energy in (18), then the demonstration is completely analogous.

The Markov transition kernels  $P_n$  considered are Feller. Moreover, it is clear from Subsection 2.2 that proving the existence of a continuous  $P$ -Lyapunov function  $V \in \mathcal{C}_\infty(\mathbb{R}^d)$  makes (17) and (18) sufficient condition for (2) to hold. To guarantee the existence of an appropriate  $Q$ -Lyapunov function, we need a result obtained by Kato in [44]. We present it here using the formulation provided in [45].

**Lemma 3.** *Suppose that  $D \subset \mathbb{R}$  is an interval, and let  $A$  be a continuous function from  $D$  to the space of real  $d \times d$  matrices. In this case, there exist  $d$  eigenvalues (counted*

with algebraic multiplicities) of  $A(t)$  which can be parameterized as continuous functions  $\lambda_1(t), \dots, \lambda_d(t)$  from  $D$  to  $\mathbb{R}$ .

We can now ensure the existence of a  $Q$ -Lyapunov function under a simple matrix condition.

**Lemma 4.** *Assume that  $A^T S A < S$ . There exists  $\alpha \in \mathbb{R}_+^*$  such that the function*

$$V : x \in \mathbb{R}^d \mapsto \exp\left(\frac{\alpha}{2} x^T S x\right), \quad (26)$$

is integrable w.r.t  $\eta_0$  and  $Q$ -Lyapunov.

**Proof :**

From (19), we deduce that the r.h.s of (5) holds for  $V$  and  $G_n$ . Then, together with (17), we deduce that it is enough to prove that  $V$  is a Lyapunov function for  $P_{A,B}$  with  $\epsilon < 1/c$ . Let's then compute  $P_{A,B}(f)$  for any function of the form

$$f : x \in \mathbb{R}^d \mapsto \exp\left(\frac{1}{2} x^T F x\right),$$

where  $F$  is an invertible matrix such that  $B^{-1} - F$  is positive definite.

In this setting, the Woodbury matrix identity provides the following equality:

$$(B - F^{-1})^{-1} = B^{-1} - B^{-1} (B^{-1} - F)^{-1} B^{-1}.$$

We have then for any  $x \in \mathbb{R}^d$ , with  $\bar{B}_F := (B^{-1} - F)^{-1}$  :

$$\begin{aligned} P_{A,B}(f)(x) &= \frac{1}{(2\pi)^{d/2} \det(B)^{1/2}} \int_{\mathbb{R}^d} e^{-\frac{1}{2}[(Ax-y)^\top B^{-1}(Ax-y) - y^\top F y]} dy \\ &= f(x) \frac{\exp(\frac{1}{2} x^T (A^T B^{-1} \bar{B}_F B^{-1} A - A^T B^{-1} A - F)x)}{(2\pi)^{d/2} \det(B)^{1/2}} \\ &\quad \times \int_{\mathbb{R}^d} \exp\left(-\frac{1}{2} \left[ (\bar{B}_F B^{-1} A x - y)^\top \bar{B}_F^{-1} (\bar{B}_F B^{-1} A x - y) \right]\right) dy \\ &= \sqrt{\frac{\det(\bar{B}_F)}{\det(B)}} f(x) \exp\left(\frac{1}{2} x^T (A^T [B^{-1} (B^{-1} - F)^{-1} B^{-1} - B^{-1}] A - F)x\right) \\ &= \frac{1}{\sqrt{\det(I_d - BF)}} f(x) \exp\left(-\frac{1}{2} x^T (F + A^T (FB - I_d)^{-1} F A)x\right). \quad (27) \end{aligned}$$

From those calculations, we deduce that  $V$  is a Lyapunov function for  $P_{A,B}$  if the matrices  $B^{-1} - \alpha S$  and  $S - A^T (I_d - \alpha S B)^{-1} S A$  are positive definite.

Let  $\lambda_B$  be the greatest eigenvalue of  $B$  and  $\lambda_S$  be the greatest eigenvalue of  $S$ . It is clear that for  $\alpha \in (0, \frac{1}{\lambda_B \lambda_S})$ ,  $B^{-1} - \alpha S$  is positive definite.

Consider now the function

$$\psi : \alpha \in \left[0, \frac{1}{\lambda_B \lambda_S}\right) \mapsto sp(S - A^T(I_d - \alpha SB)^{-1}SA) \in \mathbb{R}^d.$$

Here,  $sp(M)$  represents the spectrum of a matrix  $M$  with multiplicity taken into account.

Given the hypotheses on  $A$  and  $S$ , we can conclude that  $\psi(0) \subset \mathbb{R}_+^{*d}$ . Furthermore, by Lemma 2, it is clear that  $\psi$  is a continuous function. Since  $\mathbb{R}_+^{*d}$  is an open set, there exists  $\bar{\alpha} \in \mathbb{R}$  such that for any  $\alpha \in (0, \bar{\alpha})$ ,  $\psi(\alpha) \subset \mathbb{R}_+^{*d}$ .

By choosing a sufficiently small value for  $\alpha$  to ensure that  $V$  is integrable w.r.t  $\eta_0$ , we can conclude.  $\square$

From this Lemma and the hypothesis on  $G_n$ , we deduce that the l.h.s of (5) holds as well. We can now focus on verifying that (14) holds by proving the following Lemma.

**Lemma 5.** *There exists a positive definite matrix  $H$  such that*

$$\forall n \in \mathbb{N}, Q_n(W) \geq \bar{P} \times W$$

with  $W(x) := \exp(-\frac{1}{2}x^T H x)$

**Proof :**

From (18), we have

$$Q_n(W) \geq \frac{\bar{P}}{E_{A',B',S}} \times G_S P_{A',B'}(W).$$

Using (27), we derive the following expression for  $G_S(x)P_{A',B'}(W)(x)$  with  $x \in \mathbb{R}^d$ :

$$\frac{1}{\sqrt{\det(I_d + B'H)}} W(x) \exp\left(\frac{1}{2}x^T (H - A'^T(HB' + I_d)^{-1}HA' - S)x\right).$$

Hence, choosing  $H$  as the solution to the Riccati equation

$$H - A'^T(HB' + I_d)^{-1}HA' - S = 0,$$

we deduce that

$$E_{A',B',S} = \frac{1}{\sqrt{\det(I_d + B'H)}}.$$

Thus:

$$Q_n(W) \geq \bar{P} \times W.$$

□

For  $\alpha$  sufficiently small,  $W^{-\alpha}$  is lower than  $V$ . The right-hand side of (5) is then also verified.

Under the condition  $A^T S A < S$ , we have confirmed that all the assumptions concerning  $P_n$ ,  $G_n$ , and  $\eta_0$  in Theorem 1 are satisfied. We can therefore apply it to conclude on the proof of Corollary 3.

## 5.2 An Importance Sampling technique

This subsection focuses on the study of the importance sampling described in Subsection 3.3 and on the proof of Corollary 4.

We consider the coupled harmonic oscillator, i.e, for some matrices  $A$ ,  $B$  and  $S$ , with  $B$  and  $S$  symmetric definite positive, we consider  $P = P_{A,B}$  and  $G = G_S$ . Up to this point, we have established that the  $L_p$ -norm of the error made by the DMC method is uniformly bounded in time, with a convergence rate of  $\frac{1}{N^{\beta/2}}$  for some  $\beta \in (0, 1]$  when  $A^T S A < A$ .

Our aim is now to use Theorem 1 to prove that the approximation of the measures  $\widehat{\eta}_n$  made by the DMC method - enhanced by the importance sampling scheme described in (21) - remains uniformly bounded in time, regardless of the value of  $A$  and  $S$ . However, there's a trade-off involved: we will only have access to the measures at specific times. Indeed, despite the converging property that we are about to prove, the sequence of empirical measures  $\widehat{\eta}_n^{(k)}$  only approximates the measures  $\widehat{\eta}_l$  for  $l \in k\mathbb{N}$ .

To proceed with our proof, we make the necessary assumption that the matrices  $A$ ,  $B$ , and  $S$  can all be diagonalized in the same basis.

Before presenting the central corollary of this subsection, we lay the foundation with a lemma that, using relation (20), proves that this scenario can be interpreted as another instance of the coupled harmonic oscillator approximated by the usual DMC method. We then proceed to compute the specific constants of this scenario. Consider the parameters

$$\lambda_1 := \frac{1}{\sqrt{\det(I + BS)}} \quad \& \quad S_1 := A^T(S^{-1} + B)^{-1}A$$

as well as

$$A_1 := (I + BS)^{-1}A \quad \& \quad B_1 := (B^{-1} + S)^{-1}$$

For any  $n \geq 0$  we also set

$$\lambda_{n+1} := \frac{\lambda_n}{\sqrt{\det(I + BS + BS_n)}} \quad \& \quad S_{n+1} := A^T(B + (S + S_n)^{-1})^{-1}A$$

as well as

$$\begin{aligned} A_{n+1} &:= (I + AB_n A^T S + BS)^{-1} A A_n \\ B_{n+1} &:= (I + AB_n A^T S + BS)^{-1} (AB_n A^T + B) \end{aligned}$$

**Lemma 6.** For any  $k \geq 0$  we have

$$\widehat{G}^{(k)}(x) = \lambda_k \exp\left(-\frac{1}{2}x^T S_k x\right) \quad \text{and} \quad \delta_x \widehat{P}^{(k)} \sim \mathcal{N}(A_k x, B_k)$$

The proof of this Lemma is relatively straightforward. However, it requires some technical calculations. It is therefore postponed to the Appendix.

We can now proceed to prove the central result of this subsection.

**Proof of Corollary 4 :**

Let  $k \in \mathbb{N}$ . From Corollary 3, to prove that the DMC method associated with  $\widehat{G}^{(k)}$  and  $\widehat{P}^{(k)}$  is uniformly converging toward the Feynman-Kac measures, it is enough to prove that

$$A_k^T S_k A_k < S_k.$$

Since all these matrices can be diagonalized in the same basis, proving that this criterion holds for the matrices  $A_k$  and  $S_k$  is equivalent to proving that all eigenvalues of  $A_k$  are in the interval  $(-1, 1)$ .

Let  $\mathcal{B}$  a basis in which the matrices  $A$ ,  $B$  and  $S$  are diagonal. We want to prove that for  $i \in \llbracket 1, d \rrbracket$ , the  $i$ -th eigenvalue of  $A$  is in the right interval. In the rest of the proof, we denote by  $M^{(i)}$  the  $i$ -th eigenvalue of a matrix  $M$  that can be diagonalized in  $\mathcal{B}$ .

Using the expression derived in Lemma 6, we obtain:

$$A_n^{(i)} = \prod_{1 \leq i \leq n} \frac{A^{(i)}}{1 + S^{(i)} t_n} \quad \text{with} \quad t_n := \begin{cases} A^{(i)^2} B_n^{(i)} + B^{(i)} & \text{if } n \geq 2 \\ B^{(i)} & \text{if } n = 1 \end{cases}.$$

We will establish that as  $n$  tends towards infinity, the limit of  $\left| \frac{A^{(i)}}{1 + S^{(i)} t_n} \right|$  is strictly less than 1. This result will consequently imply the convergence of the sequence  $A_n^{(i)}$  towards 0. For  $n \in \mathbb{N}^*$  :

$$t_{n+1} = A^{(i)^2} B_{n+1}^{(i)} + B^{(i)} = \frac{A^{(i)^2} t_n}{1 + S^{(i)} t_n} + B^{(i)} := \varphi(t_n). \quad (28)$$

Here,  $\varphi$  represents a Riccati operator defined as described in [46]. Using Equation (51) from the same article, we can derive that

$$\lim_{n \rightarrow +\infty} t_n > \frac{|A^{(i)}| - 1}{S^{(i)}}. \quad (29)$$

Hence  $\lim_{n \rightarrow +\infty} \left| \frac{A^{(i)}}{1 + S^{(i)} t_n} \right| < 1$ . For  $n$  large enough, the approximation made by the DMC method enhanced by importance sampling is then the same as the usual DMC approximation of an harmonic oscillator with a stable Markov transition. We can thus conclude using Theorem 1.  $\square$

### 5.3 A divergence property

In the previous subsections, we presented a simple sufficient condition for controlling the DMC method and introduced an importance sampling technique that satisfies this criterion. However, it is natural to question the robustness of this condition and whether it is necessary to use importance sampling. Specifically, for the uni-dimensional harmonic oscillator, the convergence condition reduced to  $A^2 < 1$ , and we will prove divergence of the DMC method when this stability condition is not met.

Within this framework, we can break down the evolution of the walkers into two distinct steps, a mutation transition and a selection transition

$$(\xi_n^i)_{i \in \llbracket 1, N \rrbracket} \in \mathbb{R}^N \xrightarrow{\text{selection}} (\widehat{\xi}_n^i)_{i \in \llbracket 1, N \rrbracket} \in \mathbb{R}^N \xrightarrow{\text{mutation}} (\xi_{n+1}^i)_{i \in \llbracket 1, N \rrbracket}.$$

The initial configuration  $(\xi_0^i)_{i \in \llbracket 1, N \rrbracket}$  is determined by sampling  $N$  independent random variables from the distribution  $\eta_0$ . The selection transition involves the sampling of  $N$  independent random variables  $(\widehat{\xi}_n^i)_{i \in \llbracket 1, N \rrbracket}$  using the weighted distributions

$$\epsilon_n (\eta_n^N) G_S(\xi_n^i) \delta_{\xi_n^i} + (1 - \epsilon_n (\eta_n^N) G_S(\xi_n^i)) \sum_{k \in \llbracket 1, N \rrbracket} \frac{e^{-\frac{S}{2} \xi_n^{k^2}}}{\sum_{j \in \llbracket 1, N \rrbracket} e^{-\frac{S}{2} \xi_n^{j^2}}} \delta_{\xi_n^k}.$$

The mutation transition is defined using a family of Gaussian random variables with zero-mean and unit variance  $(V_n^i)_{i \in \llbracket 1, N \rrbracket}$  such that

$$\xi_n^i = A \widehat{\xi}_{n-1}^i + \sqrt{B} V_n^i.$$

The measures  $(\eta_n)_{n \in \mathbb{N}}$  can be described exhaustively using the Kalman filter equations. It provides us with the mean and variances  $(X_n, P_n)$  of the Gaussian random variables  $\eta_n$  with the recurrent equations

$$\begin{cases} X_{n+1} = \frac{A}{1 + SP_n} X_n \\ P_{n+1} = \frac{A^2 P_n}{1 + SP_n} + B \end{cases}. \quad (30)$$

In this scenario, when the condition  $A^2 > 1$  is met, it is possible to prove that the DMC's error in approximating the Feynman-Kac measure does not admit a uniform-in-time bound. It is properly stated in Property 1, and we can now conduct its proof.

#### Proof of Proposition 1:

For any  $n \geq 2$ , we know from (30) that

$$\eta_n(I) := X_n = \frac{A^2}{(1 + SP_{n-1})(1 + SP_{n-2})} \leq A^2 X_{n-2}.$$

For any  $n \in \mathbb{N}^*$ , let  $\xi_n^* = \min_{i \in \llbracket 1, N \rrbracket} \xi_n^i$  and define the random variables  $V_n^*$  in the following way:

$$V_n^* = - \max_{i \in \llbracket 1, N \rrbracket} |V_n^i|.$$

By definition of the evolution of the walkers, there exists  $(i, j) \in \llbracket 1, N \rrbracket$  such that

$$\xi_{2n}^* = A^2 \xi_{2n-2}^j + \sqrt{B} V_{2n-1}^i + A \sqrt{B} V_{2n-2}^j \geq A^2 \xi_{2n-2}^* + \sqrt{B} V_{2n-1}^* + |A| \sqrt{B} V_{2n-2}^*.$$

Thus

$$\eta_{2n}^N(I) - \eta_{2n}(I) \geq A^2 (\xi_{2(n-1)}^* - X_{2(n-1)}) + \sqrt{B} V_{2n-1}^* + |A| \sqrt{B} V_{2n-2}^*.$$

Iterating the process, we obtain

$$\frac{\eta_{2n}^N(I) - \eta_{2n}(I)}{A^{2n}} \geq (\xi_0^* - X_0) + \sqrt{B} \sum_{1 \leq k \leq n} \frac{V_{2k-1}^*}{A^{2k}} + |A| \sqrt{B} \sum_{1 \leq k \leq n} \frac{V_{2(k-1)}^*}{A^{2k}}.$$

For any sequence of  $N$  independent centred Gaussian random variables  $U_i$  with unit variance, we have

$$\mathbb{E} \left[ \max_{1 \leq i \leq N} |U_i| \right] \leq \sqrt{2 \log(2N)}.$$

This inequality is obtained by using Jensen's inequality as follows, with  $t = \sqrt{2 \log(2N)}$

$$\exp \left[ t \mathbb{E} \left( \max_{1 \leq i \leq N} |U_i| \right) \right] \leq \mathbb{E} \left[ \exp \left( t \max_{1 \leq i \leq N} |U_i| \right) \right] \leq \sum_{i=1}^N \mathbb{E} [\exp(t|U_i|)],$$

and noticing that

$$\mathbb{E} [\exp(t|U_i|)] = 2 \int_0^{+\infty} \exp \left( -\frac{(x-t)^2 + t^2}{2} \right) dx \leq 2 \exp(t^2/2)$$

Then, on the event

$$\Omega_\epsilon := \left\{ \xi_0^* \geq \epsilon + X_0 + \frac{2\sqrt{B}(1+|A|)}{A^2-1} \sqrt{2 \log(2N)} \right\}, \quad (31)$$

with  $\epsilon \in \mathbb{R}_+$ , we have

$$\mathbb{E}[\eta_{2n}^N(I) - \eta_{2n}(I) | \xi_0^*] \geq \epsilon A^{2n} \xrightarrow{n \rightarrow +\infty} +\infty. \quad (32)$$

Integrating over  $\xi_0^*$  we deduce



$$\mathbb{E}[|\eta_{2n}^N(I) - \eta_{2n}(I)|] \geq \epsilon A^{2n} \mathbb{P}(\Omega_\epsilon). \quad (33)$$

We can then conclude by noticing

$$\mathbb{P}(\Omega_\epsilon) = \eta_0 \left\{ \left[ \epsilon + X_0 + \frac{2\sqrt{B}(1+|A|)}{A^2-1} \sqrt{2 \log(2N)}, +\infty \right) \right\}^N > 0. \quad (34)$$

□

For the case  $A = 1$ , we are not able to assert whether or not a uniform bound exists. To the best of our knowledge, the best result proved to date is a linear bound on the variance of the unnormalized measure when  $R = S = 1$  [47].

## Appendix

### Some technical Lemmas

**Lemma 7.** *Let  $(G, V) \in \mathcal{C}(E) \times \mathcal{C}_\infty(E)$  be positive functions and  $K \subset E$  be such that the right-hand side of (5) holds. There exists  $c \in \mathbb{R}_+$  such that for any probability measure  $\mu$  on  $E$  :*

$$\psi_G(\mu)(V) \leq \mu(V) + c$$

**Proof :** Let  $\bar{V} := \mathbb{1}_{\mathbb{R} \setminus K} V$ . For any  $(x, t) \in E^2$ , we have:

$$0 \geq (G(x) - G(y))(\bar{V}(x) - \bar{V}(y)) = G(x)\bar{V}(x) + G(y)\bar{V}(y) - G(y)\bar{V}(x) - G(x)\bar{V}(y).$$

Integrating with respect to the probability measure  $\mu$  over both  $x$  and  $y$ , we obtain:

$$\mu(G\bar{V}) - \mu(G)\mu(\bar{V}) \leq 0.$$

Hence

$$\psi_G(\mu)(\bar{V}) - \mu(\bar{V}) = \frac{\mu(G\bar{V}) - \mu(G)\mu(\bar{V})}{\mu(G)} \leq 0.$$

Thus

$$\psi_G(\mu)(V) \leq \psi_G(\mu)(\bar{V}) + \sup_K V \leq \mu(\bar{V}) + \sup_K V \leq \mu(V) + \sup_K V.$$

□

**Lemma 8.** *Let  $V \in \mathcal{C}_\infty(E)$  be a  $Q$ -Lyapunov function, then*

$$\sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} [\phi_{q,n}(\eta_q^N)(V)] < +\infty \quad \text{and} \quad \sup_{\substack{(q,n) \in \mathbb{N}^2 \\ q \leq n}} \mathbb{E} [\phi_{q,n}(\eta_q)(V)] < +\infty. \quad (35)$$

**Proof:** First, let's prove that for any  $\gamma \in \mathcal{P}_V(E)$ , we have, with  $(\epsilon, c) \in [0, 1) \times \mathbb{R}$  defined in (1),

$$\sup_{\substack{(q,n) \in \mathbb{N}^2 \\ q \leq n}} \phi_{q,n}(\gamma)(V) \leq \gamma(V) + \frac{c}{1 - \epsilon}. \quad (36)$$

To do so, we begin by using the  $Q$ -Lyapunov property of  $V$  as well as Lemma 7 for some  $l \in \mathbb{N}^*$ :

$$\phi_{q,n}(\gamma)(V) = \frac{\phi_{q,n-1}(\gamma)(G_n P_n(V))}{\phi_{q,n-1}(\gamma)(G_n)} \leq \epsilon \phi_{q,n-1}(\gamma)(V) + c'.$$

By iterating the process, we obtain (36). We now prove that

$$\sup_{n \in \mathbb{N}} \mathbb{E} [\eta_n^N(V)] \leq \eta_0(V) + \frac{c'}{1 - \epsilon}. \quad (37)$$

Notice first that

$$\mathbb{E} [\eta_n^N(V) \mid \eta_{n-1}^N] = \frac{1}{N} \sum_{i=1}^N \eta_{n-1}^N S_{n-1, \eta_{n-1}^N} P_n(V) = \psi_{G_{n-1}}(\eta_{n-1}^N)(P_n(V))$$

Then, using as previously the  $Q$ -Lyapunov property of  $V$  and Lemma 7 we obtain

$$\psi_{G_{n-1}}(\eta_{n-1}^N)(P_n(V)) \leq \epsilon \eta_{n-1}^N(V) + c'.$$

By iterating the process, we obtain

$$\mathbb{E} [\eta_n^N(V)] \leq \epsilon^n \eta_0(V) + c' \sum_{i=0}^{n-1} \epsilon^i \leq \eta_0(V) + \frac{c'}{1 - \epsilon}.$$

Thus, by combining (36) and (37), we can conclude regarding the first part of (35). The second part is obtained by proceeding in a strictly analogous way.  $\square$

## Proof of Lemma 1

We first consider the case where only (13) holds and prove that

$$\phi_{0,q}(\gamma)Q_{q,n}(1) \leq C\bar{P}^{n-q-1}. \quad (38)$$

For  $q \leq n + 2$ , we have

$$\begin{aligned} Q_{q,n}(1)(x) &= Q_{q,n-2}(G_{n-1}P_n(G_n))(x) \\ &\leq \bar{P}Q_{q,n-2}(G_{n-1})(x) \\ &\leq \bar{P}Q_{q,n-1}(1)(x). \end{aligned}$$

Iterating the process, we deduce

$$Q_{q,n}(1) \leq \bar{P}^{n-q-1}G_{q-1}.$$

Using (12) we deduce (38).

For  $q \geq 1$  and  $\mu \in \{\eta_q^N, \phi_q(\eta_{q-1}^N)\}$ , we have then

$$\frac{1}{\mu(H_{q,n}^\gamma)} = \phi_{q,n}(\mu)(W) \times \frac{\phi_{0,q}(\gamma)Q_{q,n}(1)}{\mu(Q_{q,n}(W))} \leq C \frac{\bar{P}^{n-q-1}\phi_{q,n}(\mu)(W)}{\mu(Q_{q,n}(W))}.$$

From Holder's inequality, Jensen's inequality and the hypothesis on  $W$ , we get :

$$\begin{aligned} \mathbb{E} \left[ \mu(H_{q,n}^\gamma)^{-\beta} \right] &\leq C^\beta \mathbb{E} \left[ \phi_{q,n}(\mu)(W)^{2\beta} \right]^{\frac{1}{2}} \mathbb{E} \left[ \bar{P}^{2\beta(n-q-1)} \mu(Q_{q,n}(W))^{-2\beta} \right]^{\frac{1}{2}} \\ &\leq \frac{C^\beta}{\bar{P}^\beta} \mathbb{E} \left[ \phi_{q,n}(\mu)(V)^{2\beta} \right]^{\frac{1}{2}} \mathbb{E} \left[ \mu(W^{-2\beta}) \right]^{\frac{1}{2}}. \end{aligned}$$

Then, by choosing  $\beta$  small enough such that  $W^{-2\beta} \leq \bar{W}$  and  $V^{2\beta} \leq \bar{V}$ , where  $\bar{W}$  and  $\bar{V}$  are  $Q$ -Lyapunov functions, we obtain

$$\sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[ \mu(H_{q,n}^\gamma)^{-\beta} \right] \leq \frac{C^\beta}{\bar{E}_0^\beta} \sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[ \mu(\bar{V}) \right]^{\frac{1}{2}} \times \sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[ \mu(\bar{W}) \right]^{\frac{1}{2}}$$

We can conclude using Lemma 8.

The demonstration for time-homogeneous models with  $E_0 < \bar{P}$  is analogous. Indeed the equivalent of (38) is obtained by noticing that

$$\phi_{0,q}(\eta_\infty)Q_{q,n}(1) = \eta_\infty Q_{q,n-1}(G) = \eta_\infty Q_{q,n-1}(1)\eta_\infty(G) = E_0 \times \eta_\infty Q_{q,n-1}(1) = E_0^{n-q}$$

The rest of the proof follows the same arguments, thus it is skipped. This ends the proof of the lemma.  $\square$

## Proof of Lemma 6

Let us begin by proving the result for  $k = 1$ . Take  $x \in \mathbb{R}^d$ :

$$\widehat{G}^{(1)}(x) = \frac{1}{\sqrt{\det(I + BS)}} \exp\left(-\frac{1}{2}x^T A^T (S^{-1} + B)^{-1} Ax\right),$$

Applying the Woodbury matrix identity we get:

$$\begin{aligned} & \sqrt{(2\pi)^d \det([B^{-1} + S]^{-1})} \times \widehat{P}^{(1)}(x, dy) \\ &= \exp\left(-\frac{1}{2}[y^T Sy + (y - Ax)^T B^{-1}(y - Ax) - x^T A^T (S^{-1} + B)^{-1} Ax]\right) dy \\ &= \exp\left(-\frac{1}{2}(y - (I + BS)^{-1} Ax)^T (B^{-1} + S)(y - (I + BS)^{-1} Ax)\right) dy. \end{aligned}$$

Assume the property true for  $k \in \mathbb{N}^*$ . We let  $\bar{S}_k := S_k + S$ . In this notation, we have

$$\widehat{G}^{(k+1)}(x) = P(G\widehat{G}^{(k)})(x) = \lambda_k P\left(z \mapsto \exp\left(-\frac{1}{2}z^T \bar{S}_k z\right)\right)(x)$$

This implies that

$$\begin{aligned} & \widehat{G}^{(k+1)}(x) \\ &= \frac{\lambda_k}{\sqrt{(2\pi)^d \det(B)}} \int_{\mathbb{R}^d} \exp\left(-\frac{1}{2}[z \bar{S}_k z^T + (z - Ax)^T B^{-1}(z - Ax)]\right) dz \\ &= \frac{\lambda_k}{\sqrt{(2\pi)^d \det(B)}} \exp\left(-\frac{1}{2}[x^T A^T B^{-1} Ax - x^T A^T B^{-1}(\bar{S}_k + B^{-1})^{-1} B^{-1} Ax]\right) \\ & \times \int_{\mathbb{R}^d} \exp\left(-\frac{1}{2}(z - (\bar{S}_k + B^{-1})^{-1} B^{-1} Ax)^T (\bar{S}_k + B^{-1})(z - (\bar{S}_k + B^{-1})^{-1} B^{-1} Ax)\right) dz \end{aligned}$$

from which we check that

$$\widehat{G}^{(k+1)}(x) = \frac{\lambda_k}{\sqrt{\det(I + BS + BS_k)}} \exp\left(-\frac{1}{2}x^T A^T (B + (S + S_k)^{-1})^{-1} Ax\right).$$

Moreover, since  $\delta_x \widehat{P}^{(k)} \sim \mathcal{N}(A_k x, B_k^2)$  we also have

$$\delta_x \widehat{P}^{(k)} P \sim \mathcal{N}(AA_k x, AB_k A^T + B).$$

According the Gaussian update formula (see Proposition 4.5.2 in [43] for example), if  $\eta = \mathcal{N}(m, \Sigma)$ , then  $\psi_{G_S}(\eta) = \mathcal{N}[(I + \Sigma S)^{-1} m, (I + \Sigma S)^{-1} \Sigma]$ . Then we have

$$\psi_G(\delta_x \widehat{P}^{(k+1)}) \sim \mathcal{N}((I + AB_k A^T S + BS)^{-1} AA_k x, (I + AB_k A^T S + BS)^{-1} (AB_k A^T + B)).$$

This concludes the proof.  $\square$

## References

- [1] Nicholas Metropolis and Stanislaw Ulam. The monte carlo method. *Journal of the American statistical association*, 44(247):335–341, 1949.
- [2] MD Donsker and M Kac. A sampling method for determining the lowest eigenvalue and the principal eigenfunction of schrödinger’s equation. *Journal of Research of the National Bureau of Standards*, 44(50):551–557, 1950.
- [3] Malvin H Kalos. Monte carlo calculations of the ground state of three-and four-body nuclei. *Physical Review*, 128(4):1791, 1962.
- [4] Ryan J DiRisio, Jacob M Finney, and Anne B McCoy. Diffusion monte carlo approaches for studying nuclear quantum effects in fluxional molecules. *Wiley Interdisciplinary Reviews: Computational Molecular Science*, 12(6):e1615, 2022.
- [5] Kayahan Saritas, Eric R Fadel, Boris Kozinsky, and Jeffrey C Grossman. Charge density and redox potential of linio2 using ab initio diffusion quantum monte carlo. *The Journal of Physical Chemistry C*, 124(11):5893–5901, 2020.
- [6] J Shumway, LRC Fonseca, J-P Leburton, Richard M Martin, and DM Ceperley. Electronic structure of self-assembled quantum dots: comparison between density functional theory and diffusion quantum monte carlo. In *Physical Models for Quantum Dots*, pages 633–649. Jenny Stanford Publishing, 2021.
- [7] Michel Mareschal. The early years of quantum monte carlo (1): the ground state. *The European Physical Journal H*, 46(1):11, 2021.
- [8] Yao Ma, Lu Meng, Yan-Ke Chen, and Shi-Lin Zhu. Ground state baryons in the flux-tube three-body confinement model using diffusion monte carlo. *Physical Review D*, 107(5):054035, 2023.
- [9] WMC Foulkes, Lubos Mitas, RJ Needs, and Guna Rajagopal. Quantum monte carlo simulations of solids. *Reviews of Modern Physics*, 73(1):33, 2001.
- [10] Jindřich Kolorenč and Lubos Mitas. Applications of quantum monte carlo methods in condensed systems. *Reports on Progress in Physics*, 74(2):026502, 2011.
- [11] JH Hetherington. Observations on the statistical iteration of matrices. *Physical Review A*, 30(5):2713, 1984.
- [12] Sandro Sorella. Green function monte carlo with stochastic reconfiguration. *Physical review letters*, 80(20):4558, 1998.
- [13] Sandro Sorella and Luca Capriotti. Green function monte carlo with stochastic reconfiguration: An effective remedy for the sign problem. *Physical Review B*, 61(4):2599, 2000.

- [14] Michel Caffarel. Stochastic methods in quantum mechanics. In *Numerical Determination of the Electronic Structure of Atoms, Diatomic and Polyatomic Molecules*, pages 85–105. Springer, 1989.
- [15] Michel Caffarel and Pierre Claverie. Development of a pure diffusion quantum monte carlo method using a full generalized feynman–kac formula. i. formalism. *The Journal of chemical physics*, 88(2):1088–1099, 1988.
- [16] Michel Caffarel and Pierre Claverie. Treatment of the schrödinger equation through a monte carlo method based upon the generalized feynman–kac formula. *Journal of Statistical Physics*, 43:797–801, 1986.
- [17] Roland Assaraf, Michel Caffarel, and Anatole Khelif. Diffusion monte carlo methods with a fixed number of walkers. *Physical Review E*, 61(4):4566, 2000.
- [18] Pierre Del Moral and Laurent Miclo. Particle approximations of lyapunov exponents connected to schrödinger operators and feynman–kac semigroups. *ESAIM: Probability and Statistics*, 7:171–208, 2003.
- [19] Pierre Del Moral and Arnaud Doucet. Particle motions in absorbing medium with hard and soft obstacles. *Stochastic Analysis and Applications*, 22(5):1175–1207, 2004.
- [20] Mathias Rousset. On the control of an interacting particle estimation of schrödinger ground states. *SIAM J. Math. Anal.*, 38:824–844, 2006.
- [21] Eric Cancés, Benjamin Jourdain, and Tony Lelièvre. Quantum monte carlo simulations of fermions. a mathematical analysis of the xed-no de approximation. *Mathematical Models and Methods in Applied Sciences*, 16, 11 2011.
- [22] Pierre Del Moral. *Feynman-Kac Formulae: Genealogical and Interacting Particle Systems With Applications*, volume 100. 05 2004.
- [23] Pierre Del Moral. *Mean field simulation for Monte Carlo integration*. CRC press, 2013.
- [24] Pierre Del Moral and Alice Guionnet. Large deviations for interacting particle systems: applications to non-linear filtering. *Stochastic processes and their applications*, 78(1):69–95, 1998.
- [25] Donald A Dawson and Pierre Del Moral. Large deviations for interacting processes in the strong topology. In *Statistical modeling and analysis for complex data problems*, pages 179–208. Springer, 2005.
- [26] Pierre Del Moral and Alice Guionnet. Central limit theorem for nonlinear filtering and interacting particle systems. *Annals of Applied Probability*, pages 275–297, 1999.

- [27] Nicolas Chopin. Central limit theorem for sequential monte carlo methods and its application to bayesian inference. 2004.
- [28] P. Del Moral and L. Miclo. *Branching and interacting particle systems approximations of feynman-kac formulae with applications to non-linear filtering*, pages 1–145. Springer Berlin Heidelberg, Berlin, Heidelberg, 2000.
- [29] Pierre Del Moral and Alice Guionnet. On the stability of interacting processes with applications to filtering and genetic algorithms. In *Annales de l’Institut Henri Poincaré (B) Probability and Statistics*, volume 37, pages 155–194. Elsevier, 2001.
- [30] Nick Whiteley. Sequential monte carlo samplers: error bounds and insensitivity to initial conditions. *Stochastic Analysis and Applications*, 30(5):774–798, 2012.
- [31] P Del Moral and E Horton. Coupled quantum harmonic oscillators and feynman–kac path integrals for linear diffusive particles. *Communications in Mathematical Physics*, 402(2):2079–2127, 2023.
- [32] Marc Arnaudon, Pierre Del Moral, and El Maati Ouhabaz. A lyapunov approach to stability of positive semigroups: An overview with illustrations. *Stochastic Analysis and Applications*, pages 1–80, 2023.
- [33] Pierre Del Moral, Emma Horton, and Ajay Jasra. On the stability of positive semigroups. *arXiv preprint arXiv:2112.03751*, 2021.
- [34] M. H. Kalos. Monte carlo calculations of the ground state of three- and four-body nuclei. *Phys. Rev.*, 128:1791–1795, Nov 1962.
- [35] M. H. Kalos, D. Levesque, and L. Verlet. Helium at zero temperature with hard-sphere and other forces. *Phys. Rev. A*, 9:2178–2195, May 1974.
- [36] A. Messiah. *Quantum Mechanics*. Number vol. 2 in Quantum Mechanics. Elsevier Science, 1961.
- [37] David J. Griffiths and Darrell F. Schroeter. *Introduction to quantum mechanics*. Cambridge University Press, Cambridge ; New York, NY, third edition edition, 2018.
- [38] Dmitry N. Makarov. Quantum entanglement and reflection coefficient for coupled harmonic oscillators. *Phys. Rev. E*, 102:052213, Nov 2020.
- [39] Artur K. Ekert. Quantum cryptography based on bell’s theorem. *Phys. Rev. Lett.*, 67:661–663, Aug 1991.
- [40] Elisabet Romero, Ramunas Augulis, Vladimir Novoderezhkin, Marco Ferretti, Jos Thieme, Donatas Zigmantas, and Rienk van Grondelle. Quantum coherence in photosynthesis for efficient solar energy conversion. *Nature Physics*, 10, 07 2014.

- [41] Pierre Del Moral. Non linear filtering: Interacting particle solution. *Markov Processes and Related Fields*, 2:555–580, 03 1996.
- [42] Rudolf E. Kálmán and Richard S. Bucy. New results in linear filtering and prediction theory. *Journal of Basic Engineering*, 83:95–108, 1961.
- [43] Pierre del Moral and Spiridon Penev. *Stochastic Processes: From Applications to Theory* . December 2016.
- [44] Tosio Kato. *Perturbation Theory for Linear Operators*. 1966.
- [45] Chi-Kwong Li and Fuzhen Zhang. Eigenvalue continuity and gersgorin’s theorem. *Electronic Journal of Linear Algebra*, 35:619–625, 12 2019.
- [46] Pierre Del Moral and Emma Horton. A theoretical analysis of one-dimensional discrete generation ensemble kalman particle filters, 07 2021.
- [47] Nick Whiteley, Nikolas Kantas, and Ajay Jasra. Linear variance bounds for particle approximations of time-homogeneous feynman–kac formulae. *Stochastic Processes and their Applications*, 122(4):1840–1865, 2012.