



HAL
open science

On the Mathematical foundations of Diffusion Monte Carlo

Michel Caffarel, Pierre Del Moral, Luc de Montella

► **To cite this version:**

Michel Caffarel, Pierre Del Moral, Luc de Montella. On the Mathematical foundations of Diffusion Monte Carlo. 2024. hal-04409602v3

HAL Id: hal-04409602

<https://hal.science/hal-04409602v3>

Preprint submitted on 5 Feb 2024 (v3), last revised 15 Feb 2024 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

On the Mathematical foundations of Diffusion Monte Carlo

Michel Caffarel¹, Pierre Del Moral² and Luc de Montella²

¹University of Toulouse, Lab. de Chimie et Physique Quantiques, Toulouse , 31062, FR. E-Mail: `caffarel@irsamc.ups-tlse.fr`

²Centre de Recherche Inria Bordeaux Sud-Ouest, Talence, 33405, FR. E-Mail: `pierre.del-moral@inria.fr,luc.de-montella@inria.fr`

February 5, 2024

Abstract

The Diffusion Monte Carlo method with constant number of walkers, also called Stochastic Reconfiguration as well as Sequential Monte Carlo, is a widely used Monte Carlo methodology for computing the ground-state energy and wave function of quantum systems. In this study, we present the first mathematically rigorous analysis of this class of stochastic methods on non necessarily compact state spaces, including linear diffusions evolving in quadratic absorbing potentials, yielding what seems to be the first result of this type for this class of models. We present a novel and general mathematical framework with easily checked Lyapunov stability conditions that ensure the uniform-in-time convergence of Diffusion Monte Carlo estimates towards the top of the spectrum of Schrödinger operators. For transient free evolutions, we also present a divergence blow up of the estimates w.r.t. the time horizon even when the asymptotic fluctuation variances are uniformly bounded. We also illustrate the impact of these results in the context of generalized coupled quantum harmonic oscillators with non necessarily reversible nor stable diffusive particle and a quadratic energy absorbing well associated with a semi-definite positive matrix force.

1 Introduction

The many-body Schrödinger equation describes interacting quantum particles. Depending on the domain of application, these particles may represent electrons in solid-state physics or quantum chemistry, nucleons in nuclear physics, atoms in quantum liquid physics, or coupled modes of oscillators in molecular spectroscopy, among the main applications. Except for trivial quantum systems, it is impossible to solve this equation analytically. The diffusion Monte Carlo method (abbreviated DMC) provides a powerful stochastic approach to numerically approximate the ground state energy and wave function of Schrödinger operators.

The DMC methodology has a long and rich history, dating back to its first mention in 1949 by Ulam and Metropolis in [1]. The idea was first implemented by Donsker and Kac [2], and by Kalos [3] in the early 1960s. Over the years, the physics community has proposed numerous variants of Diffusion Monte Carlo, known by various names such as Green’s function Monte Carlo,[3, 4] Fixed-Node Diffusion Monte Carlo,[5], Pure Diffusion Monte Carlo, [6, 7] Stochastic Reconfiguration Monte Carlo,[8, 9, 10, 11] and Reptation Monte Carlo,[12] to cite the main ones. Despite their apparent diversity, all these approaches are fundamentally based, in one way or another, on the stochastic simulation of a specific implementation of the Feynman-Kac formula with importance sampling.

For a more detailed discussion on the origins and the applications of these Monte Carlo techniques in physics we refer the reader to the recent review article [13] as well as to [14, 15] and references therein.

The version of interest employed here is the DMC method with a fixed number of walkers, commonly known in physics as *Stochastic Reconfiguration Monte Carlo*; see the pioneering article by Hetherington [8], followed by Sorella and co-authors [9, 10] and by the first author and his co-workers in [11].

In mathematics, the methodology may also be referred to by different names, such as genetic algorithm with selection and mutation, population Monte Carlo or sequential Monte Carlo [16, 17, 18, 19, 20]. For a more thorough discussion on these application model areas we refer to the books [21, 22] and references therein.

These sequential Monte Carlo methods do not rely on biased variational techniques. They can be seen as a sophisticated genetic-type Monte Carlo methodology to simulate interacting quantum many-body systems. Various asymptotic results have been derived, including central limit theorems and large deviation principles, see for instance [23, 24] and [25, 26], as well as the books [27, 21, 22] for an overview.

Our work concerns less studied non-asymptotic and time-uniform problems. Recalling that the estimation of ground state energies relies on the limiting behavior of the walkers’ evolution in the DMC method, it is therefore crucial to obtain uniform-in-time convergence estimates. Despite its importance, there is a notable gap in the literature and very few results have been proven in this respect. To the best of our knowledge, such uniform controls are mainly valid for compact state space models, see for instance [28, 21, 22] as well as [29]. Surprisingly, the theoretical efficiency of the DMC method has never been verified rigorously even in basic linear-Gaussian scenarios such as the simple and well known harmonic oscillator. In this paper, we address this gap by establishing the first uniform-in-time convergence estimates that apply to general state space models including the coupled harmonic oscillators presented in [30].

Our approach is partly based on recent developments on the stability of positive semigroups presented in [31, 32], see also the analysis of generalized coupled harmonic oscillators presented in [30]. In the present article, we provide a natural Lyapunov condition that ensures the exponential stability of possibly time varying positive semigroups on non necessarily compact state spaces (cf. (5) and the local conditions (6)). In the context of time homogeneous positive semigroups, these conditions ensure the existence

of an unique leading eigen-triple (see for instance (11)). We underline that these results do not rely on any reversibility-type condition, nor on some spectral theorem. They can be seen as an extended version of Perron-Frobenius and Krein-Rutman theorems for possibly time varying positive operators.

We present a nonlinear Markov chain interpretation of the DMC methodologies. In this interpretation, the genetic type evolution of the walkers can be seen as a mean field particle simulation of a nonlinear Markov chain (see Section 2.4). In this context, we present an auxiliary Lyapunov condition that depends on the potential function and the free evolution of the walkers that ensures the time uniform performance of the DMC methodology (cf. condition (5), as well as Theorem 1 and Corollary 2). We illustrate this condition in the context of generalized coupled harmonic oscillators for a linear diffusive-type particle and a quadratic energy absorbing well associated with a semi-definite positive matrix force. In this context, we also show that the DMC methodology may diverge when the free evolution of the walkers is unstable *for any fixed number of walkers, even if the asymptotic variance of the Central Limit Theorem is uniformly bounded* with respect to the time parameter (see Proposition 1, as well as Section 5.3 and Proposition 2).

In the context of absorbing wells centered at the origin, this study leads us to conjecture that stable free evolution transitions is a necessary and sufficient condition for the DMC method to be uniformly convergent w.r.t. the time horizon.

Additionally, we propose and, to some extent, establish the validity of an importance sampling transformation to overcome this difficulty. This type of technique is related but not identical to the use of guiding wave functions in physics to direct the Monte Carlo moves to improve the efficiency of the DMC method [33, 34]. In contrast with conventional guiding waves techniques our approach is based on conditional free evolutions transitions and survival weight potential functions (cf. Section 2.4 and Section 5.2).

The rest of the article is organized as follows: In Section 2, we provide a detailed description of the general framework in which our study is set, as well as the theoretical foundations on which our proof will be based.

Section 3 is devoted to the presentation of our main results. Section 4 is mainly concerned the detailed proofs of time-uniform estimates.

Section 5 is devoted to the application of our convergence result to generalized coupled harmonic oscillators [30, 35, 36]. These models arise in various fields such as molecular spectroscopy [37], quantum optics [38], quantum cryptography [39] and photosynthesis [40]. In signal processing, the harmonic oscillator and the DMC methods coincides with the Kalman and the particle filter [41, 42].

2 Description of the models

2.1 Free evolution semigroups

Consider a Markov chain X_n indexed by $n \in \mathbb{N}$ and taking values in a locally compact Polish space (E, \mathcal{E}) , where \mathcal{E} is the Borel σ -field on E . Let $\mathcal{C}(E)$ be the algebra of

continuous measurable functions on E . We also define $\mathcal{C}_b(E) \subset \mathcal{C}(E)$ as the sub-algebra of bounded measurable continuous functions endowed with the supremum norm $\|\cdot\|$. With a slight abuse of notation, we denote by 0 and 1 the null and unit scalars as well as the null and unit functions on E and we denote by $I : x \in E \mapsto I(x) = x$ the identity function on E .

For $n \in \mathbb{N}^*$, we consider the Markov transitions P_n associated with X_n , and assume that they are Feller; in the sense that for any $f \in \mathcal{C}_b(E)$ we have $P_n(f) \in \mathcal{C}_b(E)$, with the function $P_n(f)$ defined for any $x \in E$ by the integral operator

$$P_n(f)(x) := \int_E P_n(x, dy) f(y) = \mathbb{E}(f(X_n) \mid X_{n-1} = x).$$

Let $\mathcal{C}_\infty(E) \subset \mathcal{C}(E)$ be the sub-algebra of uniformly positive continuous functions V that grow at infinity; that is, for any $r \geq V_\star := \inf_E V > 0$, the r -sub-level set $\mathcal{V}(r) := \{V \leq r\} \subset E$ is a non-empty compact subset. We further assume that there exists a P -Lyapunov function $V \in \mathcal{C}_\infty(E)$; in the sense that $V(E) \subset [1, \infty)$ and there exists $\epsilon \in [0, 1)$ and $c \in \mathbb{R}$ such that for any $n \in \mathbb{N}^*$ we have

$$P_n(V) \leq \epsilon V + c. \quad (1)$$

Let $\mathcal{C}_V(E) \subset \mathcal{C}(E)$ be the sub-space of functions $f \in \mathcal{C}(E)$ such that f/V is bounded, equipped with the norm $\|f\|_V := \|f/V\|$. The Markov semigroup associated with the Markov chain X_n is defined for any $f \in \mathcal{C}_V(E)$ by

$$P_{k,n}(f)(x) := \mathbb{E}(f(X_n) \mid X_k = x).$$

Condition (1) ensures that $P_{k,n}$ is V -Feller in the sense that for $f \in \mathcal{C}_V(E)$ we have $P_{k,n}(f) \in \mathcal{C}_V(E)$. To ensure the semigroup $P_{k,n}$ is exponentially stable [31], we assume the integral operator

$$P_n(x, dy) = p_n(x, y) \nu(dy)$$

has a density p_n w.r.t. some Radon measure ν satisfying for some $r_1 > 0$ and for any $r \geq r_1$ the local minorization condition

$$0 < \inf_{n \in \mathbb{N}^*} \inf_{\mathcal{V}(r)^2} p_n \leq \sup_{n \in \mathbb{N}^*} \sup_{\mathcal{V}(r)^2} p_n < \infty \quad \text{and} \quad 0 < \nu(\mathcal{V}(r)) < \infty. \quad (2)$$

The V -norm semigroup contraction techniques developed in Section 8.2 in [43] (see also Lemma 2.3 in [32] and Theorem 2.2 in [31]), ensure that for any $\mu \in \mathcal{P}_V(E)$, there exists some parameters $a, b > 0$ such that for any $k \leq n$, and any $\mu_1, \mu_2 \in \mathcal{P}_V(E)$ we have

$$\|\mu_1 P_{k,n} - \mu_2 P_{k,n}\|_V \leq a e^{-b(n-k)} \|\mu_1 - \mu_2\|_V. \quad (3)$$

Note that the r.h.s condition in (2) is met as soon as V has compact sub-level sets with non empty interior and ν is a Radon measure with full support; that is ν is finite on compact sets and strictly positive on non-empty open sets. For time-homogeneous models, the l.h.s. minorization condition is satisfied as soon as $(x, y) \in (E^\circ)^2 \mapsto p_n(x, y)$ is a continuous positive function on the interior E° of the set E .

2.2 Feynman-Kac semigroups

We associate with a sequence of strictly positive functions $(G_n)_{n \in \mathbb{N}} \in \mathcal{C}_V(E)^\mathbb{N}$ the discrete generation Feynman-Kac semigroups

$$Q_{k,n}(f)(x) = \mathbb{E} \left(f(X_n) \prod_{p=k}^{n-1} G_p(X_p) \mid X_k = x \right) \quad \text{and} \quad \widehat{Q}_{k,n}(f) := G_k^{-1} Q_{k,n}(G_n f),$$

with $G_k^{-1} := 1/G_k$. To simplify notation, for $k = (n-1)$ sometimes we write Q_n and \widehat{Q}_n instead of $Q_{n-1,n}$ and $\widehat{Q}_{n-1,n}$. In this notation, we have

$$Q_n(f)(x) = G_{n-1}(x) P_n(f)(x) \quad \text{and} \quad \widehat{Q}_n(f)(x) = P_n(G_n f)(x).$$

We also use the convention $Q_{n,n} = P_{n,n} = Id$, the identity operator.

Let $\mathcal{M}_b(E)$ be the set of bounded signed measures on E . Also, let $\mathcal{P}(E) \subset \mathcal{M}_b(E)$ be the convex subset of probability measures on E and denote by $\mathcal{P}_V(E)$ the convex set of probability measures $\mu \in \mathcal{P}(E)$ such that $\mu(V) < +\infty$. The left action of Q_n on $\mathcal{P}_V(E)$ is given for any $(\eta, f) \in (\mathcal{P}_V(E), \mathcal{C}_V(E))$ by the formula

$$(\eta Q_n)(f) := \eta(Q_n(f)) = \int \eta(dx) Q_n(f)(x) = \int \eta(dx) Q_n(x, dy) f(y). \quad (4)$$

By Fubini's theorem, the integration order doesn't matter. Thus to simplify notation, we sometimes write $\eta Q_n(f)$ instead of $(\eta Q_n)(f)$ or $\eta(Q_n(f))$.

We denote by $\mathcal{C}_0(E) := \{1/V : V \in \mathcal{C}_\infty(E)\} \subset \mathcal{C}_b(E)$ the sub-algebra of bounded continuous positive functions h that vanish at infinity; that is, for any $0 < \epsilon \leq \|h\| < \infty$ the ϵ -super-level set $\{h \geq \epsilon\} \subset E$ is a non empty compact subset.

We further assume the Lyapunov function V introduced in (1) is a Q -Lyapunov function in the sense that (1) holds and there exists $\Theta \in \mathcal{C}_0(E)$ and a compact subset $K \subset E$ such that for any $n \geq 1$ we have

$$Q_n(V)/V \leq \Theta \quad \text{and} \quad (G_{n-1}(x) - G_{n-1}(y))(\mathbb{1}_{E \setminus K}(x)V(x) - \mathbb{1}_{E \setminus K}(y)V(y)) \leq 0. \quad (5)$$

Note that the l.h.s. condition in (5) holds as soon as there exists $G \in \mathcal{C}_0(E)$ such that for any $G_n \leq G$, for any $n \geq 0$. This condition ensures that for any positive function $f \in \mathcal{C}_V(E)$ and $n \geq 1$ we have

$$Q_n(f)/V \leq Q_n(V)/V \leq \Theta.$$

By (2) the integral operator $Q_n(x, dy) = q_n(x, y) \nu(dy)$ also has a density given by $q_n(x, y) = G_{n-1}(x) p_n(x, y)$ and for any $r \geq r_1$ we have the local condition

$$0 < \inf_{\nu(r)^2} q_n \leq \sup_{\nu(r)^2} q_n < \infty. \quad (6)$$

Consider the normalized measure valued process $\eta_n \in \mathcal{P}_V(E)$ starting at $\eta_0 \in \mathcal{P}_V(E)$ defined for any $n \geq 1$ by

$$\eta_{n+1} = \phi_{n+1}(\eta_n) := \psi_{G_n}(\eta_n)P_{n+1} \quad \text{and} \quad \widehat{\eta}_n := \psi_{G_n}(\eta_n), \quad (7)$$

with the updated Boltzmann-Gibbs transformations ψ_{G_n} associated with the potential function G_n defined by

$$\psi_{G_n}(\eta_n)(dx) := \frac{1}{\eta(G_n)} G_n(x) \eta(dx).$$

We readily check that the evolution semigroup $\phi_{k,n} = \phi_{k+1,n} \circ \phi_k$ associated with the flow of measure η_n is given for any $k \leq n$ by the formula

$$\phi_{k,n}(\eta_k) = \frac{\eta_k Q_{k,n}}{\eta_k Q_{k,n}(1)}.$$

Note that for any $\mu \in \mathcal{P}_V(E)$ we have the updating formula

$$\phi_{k,n}(\eta_k) = \psi_{H_{k,n}^\mu}(\eta_k) \bar{Q}_{k,n}, \quad (8)$$

with the Markov operator

$$\bar{Q}_{k,n}(f) := Q_{k,n}(f)/Q_{k,n}(1) \quad \text{and} \quad H_{k,n}^\mu(x) := \frac{Q_{k,n}(1)(x)}{\phi_{0,k}(\mu)Q_{k,n}(1)}. \quad (9)$$

By Theorem 4.2 in [32] (see also Theorem 1 in [44] in the context of nonlinear filtering), for any $\mu \in \mathcal{P}_V(E)$, there exists some parameters $a, b > 0$ such that for any $k \leq n$, and any $\mu_1, \mu_2 \in \mathcal{P}_V(E)$ we have

$$\|\mu_1 \bar{Q}_{k,n} - \mu_2 \bar{Q}_{k,n}\|_V \leq a e^{-b(n-k)} \|\mu_1 - \mu_2\|_{V/H_{k,n}^\mu}. \quad (10)$$

Note that for constant potential functions $G_n(x) = G_n(y)$ we have $\bar{Q}_{k,n} = P_{k,n}$ and the above contraction estimates resume to (3).

For time homogeneous models $Q_n = Q$, Theorem 4.4 in [32] ensures the existence of a leading eigen triple $(h, E_0, \eta_\infty) \in (C_V(E) \times \mathbb{R}_+^* \times \mathcal{P}_V(E))$, such that

$$Q(h) = E_0 h \quad , \quad \eta_\infty Q = E_0 \eta_\infty \quad \text{and} \quad \eta_\infty(h) = 1. \quad (11)$$

2.3 Schrödinger semigroups

The objects defined in the previous subsection are core to a variety of physics problem. Indeed, consider an Hamiltonian differential operator \mathcal{H} given by the formula

$$\mathcal{H} := -\mathcal{L} + U,$$

where U is a potential energy function from E to \mathbb{R}_+ , and \mathcal{L} is a kinetic energy operator acting on a subset $\mathcal{D}(\mathcal{L})$ of $\mathcal{C}(E)$. The time dependent Schrödinger equation and the

imaginary time version associated with the hamiltonian \mathcal{H} are given, respectively, by the equations

$$i \partial_t \psi_t(x) = \mathcal{H}(\psi_t)(x) \quad \text{and} \quad -\partial_t \varphi_t(x) = \mathcal{H}(\varphi_t)(x),$$

with prescribed initial conditions (ψ_0, φ_0) . In the above display, $i \in \mathbb{C}$ stands for the imaginary unit. The right-hand side equation is obtained via a formal time change by setting $\varphi_t(x) = \psi_{-it}(x)$, and can be equivalently written in the following form

$$\partial_t \varphi_t(x) = \mathcal{L}(\varphi_t)(x) - U(x)\varphi_t(x) \quad \text{with initial condition } \varphi_0. \quad (12)$$

For a twice differentiable function φ_0 , the solution of (12) is given by the Feynman-Kac path integral formula

$$\begin{aligned} \varphi_t(x) &= \mathcal{Q}_t(\varphi_0)(x) := \int \mathcal{Q}_t(x, dy) \varphi_0(y) \\ &= \mathbb{E} \left(\varphi_0(\mathcal{X}_t) \exp \left(- \int_0^t U(\mathcal{X}_s) ds \right) \mid \mathcal{X}_0 = x \right). \end{aligned} \quad (13)$$

In the above display, \mathcal{X}_t stands for a time homogeneous stochastic process \mathcal{X}_t on E , with generator \mathcal{L} . To facilitate the interpretation of the theoretical and numerical physics in the measure theoretical framework used in this article, we note that the Feynman-Kac propagator defined by the integral operator (13) is sometimes written in terms of the exponential of the Hamiltonian operator with the exponential-type symbol

$$\mathcal{Q}_t := e^{-t\mathcal{H}} \quad \text{or in the bra-kets formalism} \quad \mathcal{Q}_t(\varphi_0) = |e^{-t\mathcal{H}}|\varphi_0\rangle.$$

The integral operator \mathcal{Q}_t is sometimes called the Feynman-Kac propagator. For any $s, t \geq 0$ the integral operators \mathcal{Q}_t satisfy the semigroup property

$$\mathcal{Q}_{s+t}(x, dz) = (\mathcal{Q}_s \mathcal{Q}_t)(x, dy) := \int \mathcal{Q}_s(x, dz) \mathcal{Q}_t(z, dy) \implies \varphi_{s+t} = \mathcal{Q}_s(\varphi_t).$$

In terms of left action bra-kets, defining $\mu_\varphi(dx) := \varphi(x)dx$, Fubini's theorem yields

$$\begin{aligned} \langle \varphi | e^{-s\mathcal{H}} | \varphi_t \rangle &= \int dx \varphi(x) \mathcal{Q}_s(x, dy) \varphi_t(dy) = (\mu_\varphi \mathcal{Q}_s)(\psi_t) \\ &= \mu_\varphi((\mathcal{Q}_s \mathcal{Q}_t)(\varphi_0)) = \mu_\varphi \mathcal{Q}_{s+t}(\varphi_0) = \langle \varphi | e^{-(s+t)\mathcal{H}} | \varphi_0 \rangle. \end{aligned}$$

The exponential notation is compatible with finite space models and the matrix notation of the continuous one-parameter semigroup for time homogeneous models. The bra-ket notation (a.k.a. Dirac notation) is also used to represent linear projection forms acting on Hilbert spaces associated with some reversible or some stationary measure, such as the Lebesgue measure for the harmonic oscillator.

The present article deals with different types of non necessarily stationary stochastic processes, including the free evolution process X_t discussed in (13). Apart in the reversible situation in which spectral theorems are stated on the Hilbert space associated

with a reversible measure, the use of the exponential symbol or the use of the brackets formalism is clearly not adapted to represent different expectations with respect to different types of stochastic and non-necessarily reversible processes.

To analyze these general stochastic models, we have chosen to only use elementary and standard measure theory notation such as (4). The integral actions of a given integral operator Q_t on the right for functions and on the left for measures are clearly compatible with finite space models and matrix notation. The left action $\mu \mapsto \mu Q_t$ maps measures into measures, while the right action $f \mapsto Q_t(f)$ maps functions into functions.

$$(\mu Q_t)(dy) := \int_E \mu(dx) Q_t(x, dy) \quad \text{and} \quad Q_t(f)(x) := \int_E Q_t(x, dy) f(y).$$

In this context the normalized measures are defined for any $s \leq t$ by the flow of measures

$$\mu_t(f) := \Phi_{s,t}(\mu_s)(f) := \mu_s Q_{t-s}(f) / \mu_s Q_{t-s}(1), \quad (14)$$

where μ_0 stands for the distribution of the initial random state \mathcal{X}_0 . Note that

$$\begin{aligned} \partial_t \mu_0 Q_t(1) &= -\mathbb{E} \left(U(\mathcal{X}_t) \exp \left(- \int_0^t U(\mathcal{X}_s) ds \right) \right) = -\mu_0 Q_t(U) \\ \implies \partial_t \log \mu_0 Q_t(1) &= -\mu_t(U) \implies \mu_0 Q_t(1) = \exp \left(- \int_0^t \mu_s(U) ds \right). \end{aligned} \quad (15)$$

The operator Q_t defined in (13) is sometimes called a Feynman-Kac propagator. However, despite its mathematical elegance, it can rarely be solved analytically. Under some regularity conditions (cf. for instance [32]) the flow of measures μ_t converge as $t \rightarrow \infty$ to some limiting fixed point measure $\mu_\infty = \Phi_{s,t}(\mu_\infty)$ (a.k.a. quasi-invariant measure). In this case, choosing $\mu_0 = \mu_\infty$ in (15) we have

$$\mu_\infty Q_t(1) = e^{-\lambda_0 t} \quad \text{with} \quad \lambda_0 := \mu_\infty(U).$$

Whenever it exists, the ground state h_0 is the leading eigen-function associated with λ_0 ; that is, we have

$$Q_t(h_0) = e^{-\lambda_0 t} h_0.$$

On a given time mesh $t_n = n\delta$, with time step $t_n - t_{n-1} = \delta > 0$ we clearly have

$$\mu_{t_n}(f) = \frac{\mu_{t_{n-1}} Q_\delta(f)}{\mu_{t_{n-1}} Q_\delta(1)} \quad \text{and} \quad Q_\delta(x, dy) = G_\delta(x) P_\delta(x, dy),$$

with the function

$$G_\delta(x) := Q_\delta(1)(x) \quad \text{and the Markov transition} \quad P_\delta(x, dy) := \frac{Q_\delta(x, dy)}{Q_\delta(1)(x)}.$$

Note that

$$\mu_\infty(G_\delta) = \mu_\infty Q_\delta(1) = e^{-\lambda_0 \delta} \quad \text{as well as} \quad Q_\delta(h_0) = e^{-\lambda_0 \delta} h_0 \quad \text{and} \quad \mu_\infty Q_\delta = e^{-\lambda_0 \delta} \mu_\infty.$$

In other words, choosing $(G, P) = (\mathcal{G}_\delta, \mathcal{P}_\delta)$ in the time-homogeneous Feynman-Kac model discussed in (7) and (11), we obtain the leading triple

$$(\eta_\infty, E_0, h) = (\mu_\infty, e^{-\lambda_0 \delta}, h_0).$$

This yields the formula

$$-\frac{1}{\delta} \log \mu_\infty(\mathcal{G}_\delta) = \lambda_0 = \mu_\infty(U).$$

Unfortunately, with the notable exception of coupled harmonic models (cf. [30] as well as Proposition 7.1 in [31]), the potential function \mathcal{G}_δ can rarely be evaluated and the Markov transition \mathcal{P}_δ cannot be sampled. The Feynman-Kac measure η_n introduced in (7) can also be interpreted as the solution of a discrete-time approximation of the formula (13). Indeed, consider a discrete time approximation X_{t_n} of the process \mathcal{X}_{t_n} and let

$$Q(x, dy) = G(x)P(x, dy),$$

with

$$G(x) = \exp(-U(x)\delta) \quad \text{and} \quad P(x, dy) = \mathbb{P}(X_{t_n}^\delta \in dy \mid X_{t_{n-1}}^\delta = x). \quad (16)$$

In this situation, choosing $n = \lfloor t/\delta \rfloor$ we have

$$\begin{aligned} \eta_0 Q_{0,n}(f) &= \mathbb{E} \left(f(X_{t_n}^\delta) \prod_{0 \leq k < n} G(X_{t_k}^\delta) \right) \\ &= \mathbb{E} \left(f(X_{t_n}^\delta) \exp \left\{ - \sum_{0 \leq k < n} U(X_{t_k}^\delta)(t_{k+1} - t_k) \right\} \right) \simeq_{\delta \downarrow 0} \eta_0 \mathcal{Q}_t(f). \end{aligned}$$

The leading triple (η_∞, E_0, h) associated with the discrete time Feynman-Kac approximation model now depends on the time step δ and we have

$$(\eta_\infty, h) \simeq_{\delta \downarrow 0} (\mu_\infty, h_0) \quad \text{and} \quad -\frac{1}{\delta} \log \eta_\infty(G) = -\frac{1}{\delta} \log \eta_\infty(e^{-U\delta}) \simeq_{\delta \downarrow 0} \lambda_0 = \mu_\infty(U).$$

It is clearly out of the scope of this article to analyze the bias introduced by the discrete time approximation discussed above.

2.4 Diffusion Quantum Monte Carlo

The Diffusion Quantum Monte Carlo methodology relies on the fact that the flow of measures η_n introduced in (7) can be interpreted as the probability distributions $\eta_n = \text{Law}(\bar{X}_n)$ of the random states \bar{X}_n of a nonlinear Markov chain \bar{X}_n . The choice of the Markov chain is far from unique. For instance, we have

$$\eta_{n+1} = \phi_{n+1}(\eta_n) = \eta_n K_{n+1, \eta_n},$$

with the local Markov transition

$$\begin{aligned} K_{n+1,\eta_n}(x, dz) &:= (S_{n,\eta_n}P_{n+1})(x, dz) \\ &:= \int S_{n,\eta_n}(x, dy) P_{n+1}(y, dz) = \mathbb{P}(\bar{X}_{n+1} \in dz \mid \bar{X}_n = x). \end{aligned}$$

In the above display, S_{n,η_n} stands for the Markov transition

$$S_{n,\eta_n}(x, dy) := \epsilon_n(\eta_n)G_n(x)\delta_x(dy) + (1 - \epsilon_n(\eta_n)G_n(x)) \psi_{G_n}(\eta_n)(dy).$$

for some tuning parameter $\epsilon_n(\eta_n) \in [0, 1]$ chosen such that $\epsilon_n(\eta_n)G_n(x) \in [0, 1]$. For instance, for $]0, 1]$ -valued potential functions, we can choose $\epsilon_n(\eta_n) = 0$ as well as $\epsilon_n(\eta_n) = 1$. For more general models, we can also choose the inverse of the η_n -essential supremum of G_n .

Note that the transition $\bar{X}_n \rightsquigarrow \bar{X}_{n+1}$ depends on the probability distributions η_n of the random states \bar{X}_n . In reference with similar nonlinear Markov chain models arising in fluid mechanics, the Markov chain \bar{X}_n is called a McKean interpretation of the flow of measures (7).

The mean field particle interpretation associated with a given McKean model is defined by a discrete-time system of N walkers $\xi_n = (\xi_n^i)_{1 \leq i \leq N}$. The system starts with N independent copies of a random variable $\bar{X}_0 = X_0$ with distribution η_0 . Given the system ξ_n at some time $n \geq 0$, we sample N conditionally independent walkers ξ_{n+1}^i with their respective distribution

$$K_{n+1,\eta_n^N}(\xi_n^i, dx) \quad \text{with} \quad \eta_n^N := \frac{1}{N} \sum_{1 \leq i \leq N} \delta_{\xi_n^i}.$$

In other words, the DMC method consists of approximating the measure η_n by using the occupation measure η_n^N associated with a system of N walkers. The initial positions of the walkers are randomly chosen from the distribution η_0 . The evolution of each walker follows then the following selection/mutation steps:

- Selection: We evaluate the current position ξ_n^i of a walker and its potential value $G_n(\xi_n^i)$. With probability $(1 - \epsilon_n(\eta_n^N)G_n(\xi_n^i))$, ξ_n^i is killed and instantly replaced by another walker say $\hat{\xi}_n^i = \xi_n^j$ with a probability proportional to $G_n(\xi_n^j)$ and $j \in \{1, \dots, N\}$; otherwise we keep it and set $\hat{\xi}_n^i := \xi_n^i$.
- Mutation: We move the selected walker $\hat{\xi}_n^i = x$ to a new location $\xi_{n+1}^i = y$ using the transition kernel $P_{n+1}(x, dy)$.

The selection transition associated with the choice $\epsilon_n(\eta_n^N) = 0$ coincides with the so-called proportional selection/reconfiguration. Note that the walker with the highest potential value is always selected when $\epsilon_n(\eta_n^N)$ is the inverse of the η_n^N -essential supremum of G_n .

For $[0, 1]$ -valued potential functions G_n we can also choose $\epsilon_n(\eta_n^N) = 1$. In this situation, particle are killed at a geometric clock that depends on the potential function.

In the context of the discrete time approximating Feynman-Kac models discussed in (16), when the time step δ tends to 0, these geometric killing-rates converges to an exponential killing rate. The limiting DMC scheme in continuous time consists of a system of N walkers. Between killing times, walkers explore the space with a free evolution with generator \mathcal{L} ; at rate U the walkers are killed and instantly another walker in the pool duplicates (see for instance [16, 27, 45, 46, 47]). *We underline that without the geometric killing rates, the variance of the proportional reconfiguration associated with the choice $\epsilon_n(\eta_n^N) = 0$ blows up when the time step tends to 0.*

We expect that the occupation measures of the system approximate the solution of the measure-valued process (7); that is, in a sense to be given, for any time horizon $n \geq 0$ we have

$$\eta_n^N = \frac{1}{N} \sum_{i=1}^N \delta_{\xi_n^i} \xrightarrow{N \rightarrow \infty} \eta_n,$$

as well as

$$\psi_{G_n}(\eta_n^N) = \sum_{i=1}^N \frac{G_n(\xi_n^i)}{\sum_{1 \leq j \leq N} G_n(\xi_n^j)} \delta_{\xi_n^i} \xrightarrow{N \rightarrow \infty} \psi_{G_n}(\eta_n) = \hat{\eta}_n.$$

The Lyapunov drift condition (1) combined with the local minorization condition (2) ensures that the free evolution of the walkers is stable, in the sense that it forgets exponentially fast its initial condition (see (3)). This condition is not satisfied for linear gaussian processes with an unstable drift matrix. In this context, as shown in Proposition 1 and Proposition 2 (see also Section 5.3) *the DMC method diverges for any fixed number of walkers, even if the asymptotic variance of the Central Limit Theorem stated in Lemma 7 is uniformly bounded w.r.t. the time horizon.*

Note that the stability regions and the Lyapunov functions are connected to the potential function by (5). Importance sampling techniques and twisted guiding waves can be used to design stable-like free evolutions. For instance, using the decomposition

$$\hat{Q}_{n+1}(f)(x) = \hat{G}_n \hat{P}_{n+1}(f),$$

with the potential function

$$\hat{G}_n := P_{n+1}(G_{n+1}) \quad \text{and} \quad \hat{P}_{n+1}(f) := \frac{P_{n+1}(G_{n+1}f)}{P_{n+1}(G_{n+1})},$$

we readily check the evolution equation

$$\hat{\eta}_{n+1} = \hat{\phi}_{n+1}(\hat{\eta}_n) := \psi_{\hat{G}_n}(\hat{\eta}_n) \hat{P}_{n+1}.$$

This shows that the updated measures evolve as in (7) by replacing (G_n, P_n) by (\hat{G}_n, \hat{P}_n) . The DMC associated with these objects is a genetic-type Monte Carlo sampler with selection fitness functions \hat{G}_n and mutation transitions \hat{P}_n . From the mathematical viewpoint, this model coincides with the one discussed in (7). Nevertheless, the free evolution of the walkers associated with (\hat{G}_n, \hat{P}_n) is now driven by the potential function.

This local conditioning importance sampling strategy is often used to turn infinite energy absorbing wells (a.k.a. hard obstacles) into soft ones [17, 48].

More generally, for any given time mesh $k_n \leq k_{n+1}$ we have

$$\widehat{Q}_{k_n, k_{n+1}}(f) = \widehat{G}_{k_n} \widehat{P}_{k_n, k_{n+1}}(f),$$

with the potential function

$$\widehat{G}_{k_n} := \widehat{Q}_{k_n, k_{n+1}}(1) \quad \text{and} \quad \widehat{P}_{k_n, k_{n+1}}(f) := \frac{\widehat{Q}_{k_n, k_{n+1}}(f)}{\widehat{Q}_{k_n, k_{n+1}}(1)}.$$

This yields the formula

$$\widehat{\eta}_{k_{n+1}}(f) = \widehat{\phi}_{k_n, k_{n+1}}(\widehat{\eta}_{k_n}) = \psi_{\widehat{G}_{k_n}}(\widehat{\eta}_{k_n}) \widehat{P}_{k_n, k_{n+1}}.$$

This shows that the updated measures $\widehat{\eta}_{k_n}$ evolve as in (7) by replacing (G_n, P_n) by $(\widehat{G}_{k_n}, \widehat{P}_{k_n, k_{n+1}})$. The DMC associated with these objects is a genetic-type Monte Carlo sampler with selection fitness functions \widehat{G}_{k_n} and mutation transitions $\widehat{P}_{k_n, k_{n+1}}$.

As shown in Section 5.2 (see also Corollary 4) in the context of coupled harmonic oscillators there exists a time mesh for which the mutation transitions $\widehat{P}_{k_n, k_{n+1}}$ and the potential functions \widehat{G}_{k_n} satisfy the required stability properties. For more general models, these objects do not have an analytic form. In this context, we can use the unbiased Monte Carlo methodologies discussed in [49], see also Section 2.3.2 in [27], and Section 11.5 in [21].

3 Statement of the main results

3.1 Some regularity conditions

For $f \in \mathcal{C}_b(E)$, time-uniform L_p -convergence of the error made by the DMC method in estimating $\eta_n(f)$ have been obtained (see for example [27, 28], as well as Chapter 4 in [21], Chapter 12 in [22] and the more recent article [32]) under the strong mixing assumption that there exist $\epsilon_P \in \mathbb{R}_+$ and $\epsilon_G \in \mathbb{R}_+$ such that

$$\forall (x_1, x_2, n) \in E^2 \times \mathbb{N}, \quad G_n(x_1) \geq \epsilon_G G_n(x_2) \quad \text{and} \quad P_n(x_1, dy) \geq \epsilon_P P_n(x_2, dy).$$

A significant consequence of this assumption is a time uniform bound on the potential function defined for some $\gamma \in \mathcal{P}_V(E)$ and $k \in \mathbb{N}$ by

$$G_{k, k+n}^\gamma : x \in \mathbb{E} \mapsto G_{k, k+n}^\gamma(x) := 1/H_{k, k+n}^\gamma(x).$$

Unfortunately, these uniform minorization and majorization conditions are rarely satisfied when E is non-compact.

In order to guarantee a time-uniform L_p -convergence for more general models including coupled harmonic oscillators, our framework requires to estimate uniformly in

time the inverse moments of $\eta_q^N(H_{q,n}^\gamma)$. To do this, we first assume that there exists $\gamma \in \mathcal{P}_V(E)$ such that

$$\sup_{n \in \mathbb{N}} \phi_{0,n}(\gamma)(G_n) < +\infty. \quad (17)$$

For time-homogeneous models, without any further conditions on the potential, condition (17) is easily checked with $\gamma = \eta_\infty$. For this scenario, we will then consider in the rest that $\gamma = \eta_\infty$. Moreover, this hypothesis trivially holds if the functions G_n are bounded by some constant independent of n .

We assume that there exists $W \in \mathcal{C}_V(E)$, $\alpha \in]0, 1]$ and a Q -Lyapunov function $\bar{W} \in \mathcal{C}_V(E)$ such that

$$Q_n(W) \geq \chi \times W \quad \text{and} \quad W^{-\alpha} \leq \bar{W}, \quad (18)$$

where

$$\chi := \sup_{(n,x) \in (\mathbb{N}^* \times E)} P_n(G_n)(x).$$

For time homogeneous models, this condition can be relaxed into the following

$$Q(W) \geq \min\{\chi, E_0\} \times W \quad \text{and} \quad W^{-\alpha} \leq \bar{W}. \quad (19)$$

Note that, for time-homogeneous models, the set of functions $W \in \mathcal{P}_V(E)$ such that $Q(W) \geq E_0 W$ is non-empty as it contains at least the ground state h . It is also worth noting that it is not necessary to know the exact value of χ nor the one of E_0 in order to prove that (18) or (19) hold. Indeed, if one of these constants is less than some $C \in \mathbb{R} \cup \{+\infty\}$, then it is sufficient to prove that for any $c < C$, there exist $W_c \in \mathcal{C}_V(E)$ and $\alpha_c \in (0, 1]$ such that

$$Q(W_c) \geq c W_c \quad \text{and} \quad W_c^{-\alpha_c} \leq V.$$

Finally, we assume that there exists a Q -Lyapunov function $\bar{V} \in \mathcal{C}(E)$ and $\lambda \in 2\mathbb{N}$ such that

$$V^\lambda \leq \bar{V}. \quad (20)$$

Without further mention, we assume that V, \bar{V} and \bar{W} are integrable with respect to η_0 , i.e., $\eta_0 \in \mathcal{P}_V(E) \cap \mathcal{P}_{\bar{W}}(E) \cap \mathcal{P}_{\bar{V}}(E)$. Under conditions (17) and (18) or (19), it is then possible to obtain a time uniform bound on the random potential function $G_{k,n}^\gamma$.

Lemma 1. *Let γ and (η, μ) be defined as in (17). There exists $\bar{\beta} \in \mathbb{R}_+^*$ such that for any $\beta \leq \bar{\beta}$ we have*

$$\sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[\phi_q(\eta_{q-1}^N)(H_{q,n}^\gamma)^{-\beta} \right] < +\infty \quad \text{and} \quad \sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[\eta_q^N(H_{q,n}^\gamma)^{-\beta} \right] < +\infty. \quad (21)$$

The proof of this pivotal Lemma is postponed to the appendix.

3.2 A time-uniform convergence Theorem

The main goal of this paper is to establish that in the context we described, the L_p -norm of the error made by the DMC method in approximating the Feynman-Kac measures η_n remains bounded in time and converges to zero as the number of particles increases. Our main result can be stated as follows

Theorem 1. *For any $p \in \mathbb{N}^*$, there exists $c \in \mathbb{R}$ and $\beta \in (0, 1]$ such that for any $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ we have*

$$\sup_{n \in \mathbb{N}} \mathbb{E} (|\eta_n(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}.$$

The proof of this theorem is provided in subsection 4.1. For time-homogeneous models, a direct consequence of Theorem 1 is a control over the estimation of the limiting quasi-invariant measure η_∞ .

Corollary 1. *For any $p \in \mathbb{N}^*$, there exists $(a, b, c) \in \mathbb{R}_+^{*3}$ and $\beta \in (0, 1]$ such that for any $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ we have*

$$\sup_{n \geq a + b \ln(N)} \mathbb{E} (|\eta_\infty(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}.$$

The proof the above Corollary is provided in subsection 4.2 .

Assuming that there exists a measure $\mu \in \mathcal{P}(E)$ that is reversible for P , it becomes possible to obtain a re-normalized weak form of the ground state h and its associated eigenvalue from the limit measures η_∞ and $\widehat{\eta}_\infty$ of η_n and $\widehat{\eta}_n$.

Indeed, referring to Section 9.5.5 in [43] (see also [16] as well as Section 12.4 in [21]), we have

$$\eta_\infty(G) = E_0 \quad \text{and} \quad \eta_\infty(f) = \frac{\mu(P(h)f)}{\mu(P(h))} = \frac{\mu(hP(f))}{\mu(h)} = \psi_h(\mu)P(f),$$

Note that

$$\eta_\infty(f) = \frac{\mu(Q(h)f/G)}{\mu(h)} = E_0 \psi_h(\mu)(f/G).$$

In the reverse angle, we have the updated limiting measures

$$\widehat{\eta}_\infty(f) = \psi_G(\eta_\infty)(f) = \psi_h(\mu)(f) = \frac{\mu(hf)}{\mu(h)} \implies \eta_\infty = \widehat{\eta}_\infty P$$

The existence of a reversible measure is actually not required to express the ground-state energy using the limit measure; indeed, we always have

$$\eta_\infty(h) = \phi(\eta_\infty)(h) = \frac{\eta_\infty(GP(h))}{\eta_\infty(G)} = E_0 \frac{\eta_\infty(h)}{\eta_\infty(G)} \implies \eta_\infty(G) = E_0.$$

Those equalities, combined with the convergence stated in Corollary 1, guarantee the efficiency of the DMC method for approximating the ground-state energy and wave function of quantum systems.

Corollary 2. *Let $p \in \mathbb{N}^*$, and assume that $G \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$. There exists $(a, b, c) \in \mathbb{R}_+^{*3}$ and $\beta \in (0, 1]$ such that for any $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ we have*

$$\left\{ \begin{array}{l} \sup_{n \geq a+b \ln(N)} \mathbb{E} (|E_0 - \eta_n^N(G)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}} \\ \sup_{n \geq a+b \ln(N)} \mathbb{E} (|\psi_h(\mu)(P(f)) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}} \\ \sup_{n \geq a+b \ln(N)} \mathbb{E} (|E_0 \psi_h(\mu)(f/G) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}} \end{array} \right. .$$

3.3 Coupled harmonic oscillators

To illustrate the practical applications of Theorem 1, we carry out an in-depth study of the generalized coupled harmonic oscillator [30]. First, we demonstrate its relevance by establishing easily verifiable sufficient conditions for time-uniform control of the DMC method in a framework that includes the harmonic oscillator.

We consider $E = \mathbb{R}^d$ for some $d \in \mathbb{N}^*$. For any real definite positive $d \times d$ matrices B and S and any a real $d \times d$ matrix A , we denote by $P_{A,B}$ and G_S the Markov kernel and the potential function defined as follows

$$P_{A,B}(x, dy) = \frac{1}{(2\pi)^{k/2} |B|^{1/2}} e^{-\frac{1}{2}(Ax-y)^\top B^{-1}(Ax-y)} \quad \text{and} \quad G_S(x) := \exp\left(-\frac{x^\top Sx}{2}\right).$$

Note that the matrix A may not be symmetric nor a stable (also called Hurwitz) matrix. In addition, the transition $P_{A,B}$ is not necessarily reversible, unless $AB = BA^T$, where A^T stands for the transposition of the matrix A .

Consider a family of Feller Markov transitions $(P_n)_{n \in \mathbb{N}}$ with positive densities p_n , an initial distribution $\eta_0 \in \mathcal{P}(E)$ and a family of positive functions $(G_n)_{n \in \mathbb{N}} \in \mathcal{C}_0(\mathbb{R}^d)^{\mathbb{N}}$ that is uniformly bounded in time.

We assume the following, where (A, A') are real matrices, (B, B', S) real positive definite matrices, $p_{A,B}$ the density of $P_{A,B}$ and $E_{A,B,S}$ the ground-state energy associated to the operators $P_{A,B}$ and G_S .

- There exists $c_1 \in \mathbb{R}$ such that for any $(n, x, y) \in \mathbb{N} \times \mathbb{R}^{d^2}$ we have

$$p_n(x, y) \leq c_1 p_{A,B}(x, y). \quad (22)$$

- For any $(n, x, y) \in \mathbb{N} \times \mathbb{R}^{d^2}$ we have

$$G_n(x) p_{n+1}(x, y) \geq \frac{\chi}{E_{A',B',S}} \times G_S(x) p_{A',B'}(x, y). \quad (23)$$

- There exists a compact $K \subset \mathbb{R}^d$ such that for any $(n, x, y) \in \mathbb{N} \times \mathbb{R}^{d^2}$ we have

$$[G_S(x) - G_S(y)][G_n^{-1}(x)\mathbb{1}_{\mathbb{R}^d \setminus K}(x) - G_n^{-1}(y)\mathbb{1}_{\mathbb{R}^d \setminus K}(y)] \geq 0. \quad (24)$$

- There exists $c_3 \in \mathbb{R}_+^*$ such that $G_S^{-c_3}$ is integrable with respect to η_0 .

If P_n and G_n are time-independent, it is possible to replace χ in (23) by the ground state energy associated with P and G .

These conditions hold trivially for the coupled harmonic oscillator, i.e if $P_n = P_{A,B}$, $G_n = G_S$ and η_0 is a normal distribution.

In this context, Theorem 1 leads to a simple sufficient matrix condition which guarantees the uniform convergence of the DMC method. The proof of the following corollary can be found in subsection 5.1.

Corollary 3. *Assume that $A^T S A < S$. In this situation, for any $p \in \mathbb{N}^*$, there exist $(\beta, \alpha, c) \in (0, 1] \times \mathbb{R}_+^{*2}$ such that for any function $f \in \mathcal{C}_V(\mathbb{R}^d)$ we have*

$$\sup_{n \in \mathbb{N}} \mathbb{E} (|\eta_n(f) - \eta_n^N(f)|^p)^{\frac{1}{p}} \leq c N^{-\frac{\beta}{2}}, \quad \text{with } V : x \in \mathbb{R}^d \mapsto \exp\left(\frac{\alpha}{2} x^T S x\right).$$

Shifting the focus to the approximation of the measures $(\hat{\eta}_n)_{n \in \mathbb{N}}$ within the coupled harmonic oscillator framework, the convergence condition given in the previous corollary can be overcome with a change of transition and selection in the DMC method. Considering $P = P_{A,B}$, $G = G_S$ as well as $\eta_0 \sim \mathcal{N}(X_0, P_0)$ and assuming that A , B and S can be diagonalized in the same basis, we recursively define for any $k \geq 1$ the function $\hat{G}^{(k)} \in \mathcal{C}_V(E)$ and the Markov kernel $P^{(k)}$ on E such that for all $f \in \mathcal{C}_V(E)$ we have

$$\hat{G}^{(k)} = P(G \hat{G}^{(k-1)}) \quad \text{and} \quad \hat{P}^{(k)}(f) = \frac{\hat{P}^{(k-1)}(P(fG))}{\hat{P}^{(k-1)}(P(G))},$$

with the convention $\hat{G}^{(0)} = 1$ and $\hat{P}^{(0)} = Id$ for $k = 0$. Equivalently, for any $f \in \mathcal{C}_V(E)$ and $k \geq 1$ we have the formula

$$\hat{Q}_{0,k}(f) = \hat{G}^{(k)} \hat{P}^{(k)}(f) \quad \text{with} \quad \hat{G}^{(k)} := \hat{Q}_{0,k}(1) \quad \text{and} \quad \hat{P}^{(k)}(f) := \hat{Q}_{0,k}(f) / \hat{Q}_{0,k}(1). \quad (25)$$

For $k \in \mathbb{N}^*$, consider a system of walkers $\hat{\xi}_n^{(k)} = \left(\xi_n^{(k),i} \right)_{1 \leq i \leq N}$ associated to the DMC method with initial distribution $\psi_G(\eta_0)$, transitions $\hat{P}^{(k)}$ and selection function $\hat{G}^{(k)}$ as well as the empirical measures

$$\hat{\eta}_n^{(k),N} := \frac{1}{N} \sum_{1 \leq i \leq N} \delta_{\hat{\xi}_n^{(k),i}}. \quad (26)$$

This system of walkers offers an approximation of the measures $\hat{\eta}_n$ for any $n \in k\mathbb{N}$. This type of change in the approximation, based on an importance sampling transformation is analogous to using a guiding waves function to direct the Monte Carlo moves. Without any additional condition, Theorem 1 ensures the uniform convergence of the model. The details of the proof can be found in Subsection 5.1.

Corollary 4. *Let $p \in \mathbb{N}^*$, there exists $\bar{k} \in \mathbb{N}$ such that for any $k \geq \bar{k}$, there exist $(\beta, \alpha, c) \in (0, 1] \times \mathbb{R}_+^{*2}$ satisfying for any $f \in \mathcal{C}_V(\mathbb{R})$*

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left(|\widehat{\eta}_{nk}(f) - \eta_n^{(k),N}(f)|^p \right)^{\frac{1}{p}} \leq cN^{-\frac{\beta}{2}}, \quad \text{with } V : x \in \mathbb{R}^d \mapsto \exp\left(\frac{\alpha}{2}x^T Sx\right).$$

Although the method does not provide an approximation for every time step, several strategies can be used to fill the gaps left by the approximation. A simple approach, though more computationally intensive, is to run independent systems of walkers for each time step in the interval $\llbracket 0, k-1 \rrbracket$. This method not only fills in the gaps, but also maintains the convergence property.

Our study concludes with a focus on the divergence of the DMC method when approximating the one-dimensional harmonic oscillator. This confirms that the stability condition stated in Corollary 3 is necessary and that, in some cases, the set of assumptions presented can closely approximate a sufficient and necessary condition. Additionally, it emphasizes the significance of the importance sampling method introduced in the previous corollary. Specifically, in the one-dimensional context, the sufficient condition for uniform convergence of the DMC method is expressed by $A^2 < 1$. Proposition 1 establishes the divergence of the error made by the DMC method when $A^2 > 1$, leaving open only the case $A = 1$.

Proposition 1. *Assume that $A^2 > 1$ and $P_0 > 0$. For any $p \in \mathbb{N}^*$ we have*

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left(|\eta_n(I) - \eta_n^N(I)|^p \right)^{\frac{1}{p}} = +\infty.$$

The proof of this proposition can be found in Subsection 5.3.

Note that all corollaries in this subsection can be extended to a control on the estimation of the limit measures, ground state, and eigenvalue using the same approach as presented in Corollary 2.

4 Stochastic interpolation

4.1 Time varying semigroups

In this subsection, we focus on proving Theorem 1. To take advantage of the conditional independence of the walkers, we structure our approach around the following decomposition of the difference between the Feynman-Kac measure and its empirical approximation, using the convention $\eta_{-1}^N = \eta_0$. Following [27, 28], we use the following stochastic interpolation formula

$$\eta_n^N - \eta_n = \sum_{q=0}^n [\phi_{q,n}(\eta_q^N) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))]. \quad (27)$$

Each term on the right-hand side represents the error that occurs when using the DMC approximation instead of the real propagator for a single extra time step. Combining the uniform bound given in Lemma 1 with the contraction property (10), the following Lemma establishes an exponentially decreasing control for these local errors.

Lemma 2. *For any $p \in \mathbb{N}^*$, there exists $(c, \rho, \beta) \in \mathbb{R}_+^{*2} \times (0, 1]$ such that for any function $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$ and any $(N, q, n) \in \mathbb{N}^3$ with $q \leq n$ we have*

$$\mathbb{E} \left[\left| [\phi_{q,n}(\eta_q^N) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))](f) \right|^p \right]^{\frac{1}{p}} \leq ce^{-(n-q)\rho} N^{-\frac{\beta}{2}}. \quad (28)$$

Proof:

Let $(\eta, \mu) \in \mathcal{P}(E)$ and let $\gamma \in \mathcal{P}_V(E)$ be defined as in (17). Consider $H_{q,n} := H_{q,n}^\gamma$ as defined in (9). Applying the updating formula (8), we obtain

$$\begin{aligned} \phi_{q,n}(\eta)(f) - \phi_{q,n}(\mu)(f) &= (\psi_{H_{q,n}}(\eta)\bar{Q}_{q,n} - \psi_{H_{q,n}}(\mu)\bar{Q}_{q,n})(f) \\ &= \frac{1}{\eta(H_{q,n})}(\eta - \mu)(H_{q,n}\bar{Q}_{q,n}[f - \psi_{H_{q,n}}(\mu)\bar{Q}_{q,n}(f)]). \end{aligned}$$

This yields the formula

$$\phi_{q,n}(\eta)(f) - \phi_{q,n}(\mu)(f) = \frac{1}{\eta(H_{q,n})}(\eta - \mu)(F_{q,n}^\mu), \quad (29)$$

with the function

$$F_{q,n}^\mu(x) := H_{q,n}(x) \int_E \psi_{H_{q,n}}(\mu)(dy) [\bar{Q}_{q,n}(f)(x) - \bar{Q}_{q,n}(f)(y)].$$

Then, applying Hölder's inequality for any $\beta \in [0, 1)$, we obtain the estimate

$$\begin{aligned} &\mathbb{E} \left(\left| \phi_{q,n}(\eta_q^N)(f) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))(f) \right|^p \right)^{\frac{1}{p}} \\ &\leq \mathbb{E} \left(\eta_q^N(H_{q,n})^{-2p} \left| (\eta_q^N - \phi_q(\eta_{q-1}^N))(F_{q,n}^N) \right|^{2p(1-\beta)} \right)^{\frac{1}{2p}} \mathbb{E} \left(\left| (\eta_q^N - \phi_q(\eta_{q-1}^N))(F_{q,n}^N) \right|^{2\beta p} \right)^{\frac{1}{2p}}, \end{aligned}$$

with $F_{q,n}^N := F_{q,n}^{\phi_q(\eta_{q-1}^N)}$. Using (29) this yields the estimate

$$\begin{aligned} &\mathbb{E} \left(\left| \phi_{q,n}(\eta_q^N)(f) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))(f) \right|^p \right)^{\frac{1}{p}} \\ &\leq \mathbb{E} \left(\eta_q^N(H_{q,n})^{-2\beta p} \left| \phi_{q,n}(\eta_q^N)(f) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))(f) \right|^{2p(1-\beta)} \right)^{\frac{1}{2p}} \\ &\quad \times \mathbb{E} \left(\left| (\eta_q^N - \phi_q(\eta_{q-1}^N))(F_{q,n}^N) \right|^{2\beta p} \right)^{\frac{1}{2p}} \end{aligned}$$

Recalling that $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$, this implies that

$$\begin{aligned} & \mathbb{E} \left(\left| \phi_{q,n}(\eta_q^N)(f) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))(f) \right|^p \right)^{\frac{1}{p}} \\ & \leq \left[\mathbb{E} \left(\phi_{q,n}(\eta_q^N)(V^\lambda) \right)^{\frac{1}{4p}} + \mathbb{E} \left(\phi_{q,n}(\phi_q(\eta_{q-1}^N))(V^\lambda) \right)^{\frac{1}{4p}} \right] \\ & \quad \times \mathbb{E} \left(\eta_q^N (H_{q,n})^{-4\beta p} \right)^{\frac{1}{4p}} \mathbb{E} \left(\left| (\eta_q^N - \phi_q(\eta_{q-1}^N))(F_{q,n}^N) \right|^{2\beta p} \right)^{\frac{1}{2p}}. \end{aligned}$$

From Lemmas 1 and 9, we deduce that, to conclude, it is enough to prove that, for some constant $c \in \mathbb{R}_+^*$ independent of n, q and N , we have

$$\mathbb{E} \left[\left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}^N) \right|^{2\beta p} \right]^{\frac{1}{2p}} < c e^{-c_2(n-q)} N^{-\frac{\beta}{2}}. \quad (30)$$

Let $\beta' = \frac{2p\beta}{\lambda}$ and assume β small enough so that $\beta' < 1/4$.

For $q > 0$, the walkers $(\xi_q^i)_{1 \leq i \leq N}$ are independent conditionally to η_{q-1}^N we have

$$[\eta_q^N - \phi_q(\eta_{q-1}^N)](f) = \eta_q^N \left(\frac{1}{N} \sum_{1 \leq i \leq N} h_i \right) \quad \text{with } h_i := f - S_{q-1, \eta_{q-1}^N} P_q(f)(\xi_{q-1}^i).$$

Moreover, we have

$$\phi_q(\eta_{q-1}^N) = \frac{1}{N} \sum_{1 \leq i \leq n} \mu_i \quad \text{with } \mu_i = \delta_{\xi_{q-1}^i} S_{q-1, \eta_{q-1}^N} P_q.$$

Since for any $i \in \llbracket 1, N \rrbracket$, we have $\mu_i(h_i) = 0$, we can apply Lemma 7.3.3 from [21], and deduce that there exists $C \in \mathbb{R}$ such that

$$\begin{aligned} \mathbb{E} \left[\left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}^N) \right|^{2\beta p} \right]^{\frac{1}{2p}} &= \mathbb{E} \left[\mathbb{E} \left(\left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}^N) \right|^{\lambda\beta'} \mid \xi_{q-1} \right) \right]^{\frac{1}{2p}} \\ &\leq \mathbb{E} \left[\mathbb{E} \left(\left| [\eta_q^N - \phi_q(\eta_{q-1}^N)] (F_{q,n}^N) \right|^\lambda \mid \xi_{q-1} \right)^{\beta'} \right]^{\frac{1}{2p}} \\ &\leq \frac{C}{N^{\beta/2}} \mathbb{E} \left[\phi_q(\eta_{q-1}^N) \left(|F_{q,n}^N|^\lambda \right)^{\beta'} \right]^{1/2p}. \end{aligned}$$

For $q = 0$, the walkers are iid with common distribution η_0 . The previous reasoning therefore holds using the convention $E(X \mid \xi_{-1}) = E(X)$.

Applying the contraction property (10) with $\mu = \delta_x$ and $\eta = \delta_y$ we get the existence of $(a, \rho) \in \mathbb{R}_+^2$ such that

$$|\bar{Q}_{q,n}(f)(x) - \bar{Q}_{q,n}(f)(y)| \leq a e^{-\rho(n-q)} \left(1 + \frac{V(x)}{H_{q,n}(x)} \right) \left(1 + \frac{V(y)}{H_{q,n}(y)} \right). \quad (31)$$

By substituting (31) into the definition of $F_{q,n}$ and applying Hölder's inequality along with Jensen's inequality, we obtain, for some $(a', \rho') \in \mathbb{R}_+^{*2}$

$$\begin{aligned} & \mathbb{E} \left[\phi_q(\eta_{q-1}^N) \left(|F_{q,n}^N|^\lambda \right)^{\beta'} \right]^{1/2p} \\ & \leq a' e^{-\rho'(n-q)} \mathbb{E} \left[\phi_q(\eta_{q-1}^N) \{ (H_{q,n} + V)^\lambda \}^{\beta'} \phi_q(\eta_{q-1}^N) \{ (H_{q,n} + V) \}^{\lambda\beta'} \phi_q(\eta_{q-1}^N) (H_{q,n})^{-\lambda\beta'} \right]^{\frac{1}{2p}} \\ & \leq a' e^{-\rho'(n-q)} \mathbb{E} \left[\phi_q(\eta_{q-1}^N) \{ (H_{q,n} + V)^\lambda \} \right]^{\frac{1}{4p}} \mathbb{E} \left[\phi_q(\eta_{q-1}^N) (H_{q,n})^{-2\lambda\beta'} \right]^{\frac{1}{4p}}. \end{aligned}$$

From our hypothesis on Q_n , we deduce from Lemma 3.2 in [32] that there exists a constant c such that for any $(q', n') \in \mathbb{N}^2$, $H_{q',n'} \leq cV$. We can then conclude by choosing a small enough β' and using Lemmas 1 and 9. \square

The proof of Theorem 1 is now relatively straightforward.

Proof of Theorem 1:

Let $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$. From the sub-additivity of the L_p -norm applied in (27), we have

$$E(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} \leq \sum_{q=0}^n \mathbb{E} \left(|\phi_{q,n}(\eta_q^N)(f) - \phi_{q,n}(\phi_q(\eta_{q-1}^N))(f)|^p \right)^{\frac{1}{p}}.$$

By applying Lemma 2, we deduce that there exists $(C, \rho, \beta) \in \mathbb{R} \times \mathbb{R}_+^* \times (0, 1]$ such that for any $N \in \mathbb{N}^*$ we have

$$E(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} \leq \frac{C}{N^{\frac{\beta}{2}}} \sum_{0 \leq l \leq n} e^{-(n-l)\rho} \leq \frac{C}{N^{\frac{\beta}{2}}(1 - e^{-\rho})}.$$

This ends the proof of the theorem. \square

4.2 Ground state estimates

This subsection concentrates on proving Corollary 1. We consider thus the time-homogeneous model.

Let $f \in \mathcal{C}_{V^{\frac{\lambda}{4p}}}(E)$. Notice that we can decompose the error made by the DMC method in the following way:

$$\mathbb{E}(|\eta_n^N(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq \mathbb{E}(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} + |\eta_n(f) - \eta_\infty(f)|.$$

Theorem 1 implies that there exists $(C_1, \beta) \in \mathbb{R}^* \times (0, 1]$ such that

$$\sup_{n \in \mathbb{N}} \mathbb{E}(|\eta_n^N(f) - \eta_n(f)|^p)^{\frac{1}{p}} \leq \frac{C_1}{N^{\beta/2}}.$$

According to Theorem 4.3 in [32], there exists $(C_2, \omega) \in \mathbb{R}_+^{*2}$ such that

$$|\eta_n(f) - \eta_\infty(f)| \leq C_2 e^{-\omega n}.$$

Hence

$$\mathbb{E}(|\eta_n^N(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq \frac{C_1}{N^{\beta/2}} + C_2 e^{-\omega n}.$$

Thus, letting $a = \frac{1}{\omega} \ln(C_2/C_1)$ and $b = \frac{\beta}{2\omega}$, we deduce that there exists $C \in \mathbb{R}$ such that

$$\sup_{n \geq a + b \ln(N)} \mathbb{E}(|\eta_n^N(f) - \eta_\infty(f)|^p)^{\frac{1}{p}} \leq \frac{C}{N^{\beta/2}}.$$

This concludes the proof. \square

5 Coupled harmonic oscillators

5.1 Lyapunov functions

This subsection is dedicated to the proof of Corollary 3. Therefore we place ourselves within the framework associated with this corollary. We only consider the general case where P and G depend on a time parameter. If this is not the case, and χ is replaced by the ground state energy in (23), then the demonstration is completely analogous.

The Markov transition kernels P_n considered are Feller. Moreover, it is clear from Subsection 2.2 that proving the existence of a continuous P -Lyapunov function $V \in \mathcal{C}_\infty(\mathbb{R}^d)$ makes (22) and (23) sufficient condition for (2) to hold. To guarantee the existence of an appropriate Q -Lyapunov function, we need a result obtained by Kato in [50]. We present it here using the formulation provided in [51].

Lemma 3. *Suppose that $D \subset \mathbb{R}$ is an interval, and let A be a continuous function from D to the space of real $d \times d$ matrices. In this case, there exist d eigenvalues (counted with algebraic multiplicities) of $A(t)$ which can be parameterized as continuous functions $\lambda_1(t), \dots, \lambda_d(t)$ from D to \mathbb{R} .*

We can now ensure the existence of a Q -Lyapunov function under a simple matrix condition.

Lemma 4. *Assume that $A^T S A < S$. There exists $\alpha \in \mathbb{R}_+^*$ such that the function*

$$V : x \in \mathbb{R}^d \mapsto \exp\left(\frac{\alpha}{2} x^T S x\right), \quad (32)$$

is a Q -Lyapunov function and it is integrable w.r.t η_0 .

Proof :

From (24), we deduce that the r.h.s of (5) holds for V and G_n . Then, together with (22), we deduce that it is enough to prove that V is a Lyapunov function for $P_{A,B}$ with $\epsilon < 1/c$. Let's then compute $P_{A,B}(f)$ for any function of the form

$$f : x \in \mathbb{R}^d \mapsto \exp\left(\frac{1}{2}x^T F x\right),$$

where F is an invertible matrix such that $B^{-1} - F$ is positive definite.

In this setting, the Woodbury matrix identity provides the following equality:

$$(B - F^{-1})^{-1} = B^{-1} - B^{-1}(B^{-1} - F)^{-1}B^{-1}.$$

We have then for any $x \in \mathbb{R}^d$, with $\bar{B}_F := (B^{-1} - F)^{-1}$:

$$\begin{aligned} P_{A,B}(f)(x) &= \frac{1}{(2\pi)^{d/2} \det(B)^{1/2}} \int_{\mathbb{R}^d} e^{-\frac{1}{2}[(Ax-y)^\top B^{-1}(Ax-y) - y^T F y]} dy \\ &= f(x) \frac{\exp(\frac{1}{2}x^T (A^T B^{-1} \bar{B}_F B^{-1} A - A^T B^{-1} A - F)x)}{(2\pi)^{d/2} \det(B)^{1/2}} \\ &\quad \times \int_{\mathbb{R}^d} \exp\left(-\frac{1}{2}\left[(\bar{B}_F B^{-1} Ax - y)^\top \bar{B}_F^{-1} (\bar{B}_F B^{-1} Ax - y)\right]\right) dy. \end{aligned}$$

This yields the formulae

$$\begin{aligned} &P_{A,B}(f)(x) \\ &= \sqrt{\frac{\det(\bar{B}_F)}{\det(B)}} f(x) \exp\left(\frac{1}{2}x^T (A^T [B^{-1}(B^{-1} - F)^{-1}B^{-1} - B^{-1}]A - F)x\right) \quad (33) \\ &= \frac{1}{\sqrt{\det(I_d - BF)}} f(x) \exp\left(-\frac{1}{2}x^T (F + A^T (FB - I_d)^{-1}FA)x\right). \end{aligned}$$

From those calculations, we deduce that V is a Lyapunov function for $P_{A,B}$ if the matrices $B^{-1} - \alpha S$ and $S - A^T (I_d - \alpha SB)^{-1} SA$ are positive definite.

Let λ_B be the greatest eigenvalue of B and λ_S be the greatest eigenvalue of S . It is clear that for $\alpha \in (0, \frac{1}{\lambda_B \lambda_S})$, $B^{-1} - \alpha S$ is positive definite.

Consider now the function

$$\psi : \alpha \in \left[0, \frac{1}{\lambda_B \lambda_S}\right) \mapsto sp(S - A^T (I_d - \alpha SB)^{-1} SA) \in \mathbb{R}^d.$$

Here, $sp(M)$ represents the spectrum of a matrix M with multiplicity taken into account.

Given the hypotheses on A and S , we can conclude that $\psi(0) \subset \mathbb{R}_+^{*d}$. Furthermore, by Lemma 2, it is clear that ψ is a continuous function. Since \mathbb{R}_+^{*d} is an open set, there exists $\bar{\alpha} \in \mathbb{R}$ such that for any $\alpha \in (0, \bar{\alpha})$, $\psi(\alpha) \subset \mathbb{R}_+^{*d}$.

By choosing a sufficiently small value for α to ensure that V is integrable w.r.t η_0 , we can conclude. \square

From this Lemma and the hypothesis on G_n , we deduce that the l.h.s of (5) holds as well. We can now focus on verifying that (19) holds by proving the following Lemma.

Lemma 5. *There exists a positive definite matrix H such that*

$$\forall n \in \mathbb{N}, Q_n(W) \geq \chi \times W \quad \text{with} \quad W(x) := \exp\left(-\frac{1}{2}x^T H x\right).$$

Proof :

From (23), we have

$$Q_n(W) \geq \frac{\chi}{E_{A',B',S}} \times G_S P_{A',B'}(W).$$

Using (33), we derive the following expression for $G_S(x)P_{A',B'}(W)(x)$ with $x \in \mathbb{R}^d$:

$$\frac{1}{\sqrt{\det(I_d + B'H)}} W(x) \exp\left(\frac{1}{2}x^T (H - A'^T (HB' + I_d)^{-1} HA' - S)x\right).$$

Hence, choosing H as the solution to the Riccati equation

$$H - A'^T (HB' + I_d)^{-1} HA' - S = 0,$$

we deduce that

$$E_{A',B',S} = \frac{1}{\sqrt{\det(I_d + B'H)}}.$$

Thus:

$$Q_n(W) \geq \chi \times W.$$

\square

For α sufficiently small, $W^{-\alpha}$ is lower than V . The right-hand side of (19) is then also verified.

Under the condition $A^T S A < S$, we have confirmed that all the assumptions concerning P_n , G_n , and η_0 in Theorem 1 are satisfied. We can therefore apply it to conclude on the proof of Corollary 3.

5.2 Conditional free evolutions

This subsection focuses on the study of the importance sampling described in Section 2.4, Section 3.3 and on the proof of Corollary 4.

We consider the coupled harmonic oscillator, i.e, for some matrices A , B and S , with B and S symmetric definite positive, we consider $P = P_{A,B}$ and $G = G_S$. Up to this point, we have established that the L_p -norm of the error made by the DMC method is uniformly bounded in time, with a convergence rate of $\frac{1}{N^{\beta/2}}$ for some $\beta \in (0, 1]$ when $A^T S A < A$.

Our aim is now to use Theorem 1 to prove that the approximation of the measures $\hat{\eta}_n$ made by the DMC method - enhanced by the importance sampling scheme described in (26) - remains uniformly bounded in time, regardless of the value of A and S . However, there's a trade-off involved: we will only have access to the measures at specific times. Indeed, despite the converging property that we are about to prove, the sequence of empirical measures $\hat{\eta}_n^{(k)}$ only approximates the measures $\hat{\eta}_l$ for $l \in k\mathbb{N}$.

To proceed with our proof, we make the necessary assumption that the matrices A , B , and S can all be diagonalized in the same basis.

Before presenting the central corollary of this subsection, we lay the foundation with a lemma that, using relation (25), proves that this scenario can be interpreted as another instance of the coupled harmonic oscillator approximated by the usual DMC method. We then proceed to compute the specific constants of this scenario. Consider the parameters

$$\lambda_1 := \frac{1}{\sqrt{\det(I + BS)}} \quad \& \quad S_1 := A^T(S^{-1} + B)^{-1}A,$$

as well as

$$A_1 := (I + BS)^{-1}A \quad \& \quad B_1 := (B^{-1} + S)^{-1}.$$

For any $n \geq 0$ we also set

$$\lambda_{n+1} := \frac{\lambda_n}{\sqrt{\det(I + BS + BS_n)}} \quad \& \quad S_{n+1} := A^T(B + (S + S_n)^{-1})^{-1}A,$$

as well as

$$\begin{aligned} A_{n+1} &:= (I + AB_k A^T S + BS)^{-1} A A_n \\ B_{n+1} &:= (I + AB_n A^T S + BS)^{-1} (AB_n A^T + B). \end{aligned}$$

Lemma 6. *For any $k \geq 0$ we have*

$$\hat{G}^{(k)}(x) = \lambda_k \exp\left(-\frac{1}{2}x^T S_k x\right) \quad \text{and} \quad \delta_x \hat{P}^{(k)} \sim \mathcal{N}(A_k x, B_k).$$

The proof of this Lemma is relatively straightforward. However, it requires some technical calculations. It is therefore postponed to the Appendix.

We can now proceed to prove the central result of this subsection.

Proof of Corollary 4: Let $k \in \mathbb{N}$. From Corollary 3, to prove that the DMC method associated with $\widehat{G}^{(k)}$ and $\widehat{P}^{(k)}$ is uniformly converging toward the Feynman-Kac measures, it is enough to prove that

$$A_k^T S_k A_k < S_k.$$

Since all these matrices can be diagonalized in the same basis, proving that this criterion holds for the matrices A_k and S_k is equivalent to proving that all eigenvalues of A_k are in the interval $(-1, 1)$.

Let \mathcal{B} a basis in which the matrices A , B and S are diagonal. We want to prove that for $i \in \llbracket 1, d \rrbracket$, the i -th eigenvalue of A is in the right interval. In the rest of the proof, we denote by $M^{(i)}$ the i -th eigenvalue of a matrix M that can be diagonalized in \mathcal{B} .

Using the expression derived in Lemma 6, we obtain:

$$A_n^{(i)} = \prod_{1 \leq i \leq n} \frac{A^{(i)}}{1 + S^{(i)} t_n} \quad \text{with} \quad t_n := \begin{cases} A^{(i)^2} B_n^{(i)} + B^{(i)} & \text{if } n \geq 2 \\ B^{(i)} & \text{if } n = 1 \end{cases}.$$

We will establish that as n tends towards infinity, the limit of $\left| \frac{A^{(i)}}{1 + S^{(i)} t_n} \right|$ is strictly less than 1. This result will consequently imply the convergence of the sequence $A_n^{(i)}$ towards 0. For $n \in \mathbb{N}^*$:

$$t_{n+1} = A^{(i)^2} B_{n+1}^{(i)} + B^{(i)} = \frac{A^{(i)^2} t_n}{1 + S^{(i)} t_n} + B^{(i)} := \varphi(t_n). \quad (34)$$

Here, φ represents a Riccati operator defined as described in [52]. Using Equation (51) from the same article, we can derive that

$$\lim_{n \rightarrow +\infty} t_n > \frac{|A^{(i)}| - 1}{S^{(i)}}. \quad (35)$$

Hence $\lim_{n \rightarrow +\infty} \left| \frac{A^{(i)}}{1 + S^{(i)} t_n} \right| < 1$. For n large enough, the approximation made by the DMC method enhanced by importance sampling is then the same as the usual DMC approximation of an harmonic oscillator with a stable Markov transition. We can thus conclude using Theorem 1. \square

5.3 Divergence and fluctuation estimates

In the previous subsections, we presented a simple sufficient condition for controlling the DMC method and introduced an importance sampling technique that satisfies this criterion. However, it is natural to question the robustness of this condition and whether it is necessary to use importance sampling. Specifically, for the uni-dimensional harmonic oscillator, the convergence condition reduced to $A^2 < 1$, and we will prove divergence of the DMC method when this stability condition is not met.

Within this framework, we can break down the evolution of the walkers into two distinct steps, a mutation transition and a selection transition

$$\left(\xi_n^i\right)_{i \in \llbracket 1, N \rrbracket} \in \mathbb{R}^N \xrightarrow{\text{selection}} \left(\widehat{\xi}_n^i\right)_{i \in \llbracket 1, N \rrbracket} \in \mathbb{R}^N \xrightarrow{\text{mutation}} \left(\xi_{n+1}^i\right)_{i \in \llbracket 1, N \rrbracket}.$$

The initial configuration $\left(\xi_0^i\right)_{i \in \llbracket 1, N \rrbracket}$ is determined by sampling N independent random variables from the distribution η_0 . The selection transition involves the sampling of N independent random variables $\left(\widehat{\xi}_n^i\right)_{i \in \llbracket 1, N \rrbracket}$ using the weighted distributions

$$\epsilon_n(\eta_n^N)G_S(\xi_n^i)\delta_{\xi_n^i} + (1 - \epsilon_n(\eta_n^N)G_S(\xi_n^i)) \sum_{k \in \llbracket 1, N \rrbracket} \frac{e^{-\frac{S}{2}\xi_n^{k^2}}}{\sum_{j \in \llbracket 1, N \rrbracket} e^{-\frac{S}{2}\xi_n^{j^2}}} \delta_{\xi_n^k}.$$

The mutation transition is defined using a family of Gaussian random variables with zero-mean and unit variance $(V_n^i)_{i \in \llbracket 1, N \rrbracket}$ such that

$$\xi_n^i = A\widehat{\xi}_{n-1}^i + \sqrt{B}V_n^i.$$

The measures $(\eta_n)_{n \in \mathbb{N}}$ can be described exhaustively using the Kalman filter equations. It provides us with the mean and variances (m_n, σ_n^2) of the Gaussian random variables η_n with the recurrent equations

$$\begin{cases} m_{n+1} = \frac{A}{1 + S\sigma_n^2} m_n \\ \sigma_{n+1}^2 = \frac{A^2\sigma_n^2}{1 + S\sigma_n^2} + B \end{cases}. \quad (36)$$

In this scenario, when the condition $A^2 > 1$ is met, it is possible to prove that the DMC's error in approximating the Feynman-Kac measure does not admit a uniform-in-time bound. It is properly stated in Property 1, and we can now conduct its proof.

Proof of Proposition 1:

For any $n \geq 2$, we know from (36) that

$$\eta_n(I) := m_n = \frac{A^2}{(1 + S\sigma_{n-1}^2)(1 + S\sigma_{n-2}^2)} m_{n-2} \leq A^2 m_{n-2}.$$

For any $n \in \mathbb{N}^*$, let $\xi_n^* = \min_{i \in \llbracket 1, N \rrbracket} \xi_n^i$ and define the random variables V_n^* in the following way:

$$V_n^* = - \max_{i \in \llbracket 1, N \rrbracket} |V_n^i|.$$

By definition of the evolution of the walkers, there exists $(i, j) \in \llbracket 1, N \rrbracket$ such that

$$\xi_{2n}^* = A^2 \xi_{2n-2}^j + \sqrt{B} V_{2n-1}^i + A \sqrt{B} V_{2n-2}^j \geq A^2 \xi_{2n-2}^* + \sqrt{B} V_{2n-1}^* + |A| \sqrt{B} V_{2n-2}^*.$$

Thus

$$\eta_{2n}^N(I) - \eta_{2n}(I) \geq A^2 (\xi_{2(n-1)}^* - m_{2(n-1)}) + \sqrt{B} V_{2n-1}^* + |A| \sqrt{B} V_{2n-2}^*.$$

Iterating the process, we obtain

$$\frac{\eta_{2n}^N(I) - \eta_{2n}(I)}{A^{2n}} \geq (\xi_0^* - m_0) + \sqrt{B} \sum_{1 \leq k \leq n} \frac{V_{2k-1}^*}{A^{2k}} + |A| \sqrt{B} \sum_{1 \leq k \leq n} \frac{V_{2(k-1)}^*}{A^{2k}}.$$

For any sequence of N independent centred Gaussian random variables U_i with unit variance, we have

$$\mathbb{E} \left[\max_{1 \leq i \leq N} |U_i| \right] \leq \sqrt{2 \log(2N)}.$$

This inequality is obtained by using Jensen's inequality as follows, with $t = \sqrt{2 \log(2N)}$

$$\exp \left[t \mathbb{E} \left(\max_{1 \leq i \leq N} |U_i| \right) \right] \leq \mathbb{E} \left[\exp \left(t \max_{1 \leq i \leq N} |U_i| \right) \right] \leq \sum_{i=1}^N \mathbb{E} [\exp(t|U_i|)],$$

and noticing that

$$\mathbb{E} [\exp(t|U_i|)] = 2 \int_0^{+\infty} \exp \left(-\frac{(x-t)^2 + t^2}{2} \right) dx \leq 2 \exp(t^2/2).$$

Then, on the event

$$\Omega_\epsilon := \left\{ \xi_0^* \geq \epsilon + m_0 + \frac{2\sqrt{B}(1+|A|)}{A^2-1} \sqrt{2 \log(2N)} \right\}, \quad (37)$$

with $\epsilon \in \mathbb{R}_+^*$, we have

$$\mathbb{E}[\eta_{2n}^N(I) - \eta_{2n}(I) | \xi_0^*] \geq \epsilon A^{2n} \xrightarrow{n \rightarrow +\infty} +\infty. \quad (38)$$

Integrating over ξ_0^* we deduce

$$\mathbb{E}[|\eta_{2n}^N(I) - \eta_{2n}(I)|] \geq \epsilon A^{2n} \mathbb{P}(\Omega_\epsilon). \quad (39)$$

We can then conclude by noticing

$$\mathbb{P}(\Omega_\epsilon) = \eta_0 \left\{ \left[\epsilon + X_0 + \frac{2\sqrt{B}(1+|A|)}{A^2-1} \sqrt{2 \log(2N)}, +\infty \right) \right\}^N > 0. \quad (40)$$

□

For the case $A = 1$, we are not able to assert whether or not a uniform bound exists. To the best of our knowledge, the best divergence-type result proved to date is a linear bound on the variance of the unnormalized measure when $R = S = 1$ [53].

The divergence result in Proposition 1 highlights the importance of studying non-asymptotic uniform convergence results rather than relying solely on central limit theorems (CLTs). Extensive research has been devoted to CLTs and, under appropriate assumptions. We quote the first studies in this field [25, 27] mainly based on uniformly bounded potential and test functions.

More general fluctuation theorems that apply to more general models including diffusion-type processes with Lipschitz drift and diffusion functions as well as test functions with at most polynomial growth are discussed in [54, 55]. We can formulate the following result :

Lemma 7. *In the context of proportional selection/reconfiguration, for any $n \geq 1$, we have the following convergence in law as N tends toward $+\infty$*

$$\sqrt{N}[\eta_n(I) - \eta_n^N(I)] \xrightarrow[N \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, \sigma_n^2),$$

with the asymptotic variance

$$\sigma_n^2 := \sum_{p=0}^n \eta_p \left[\left(\frac{Q_{p,n}(1)}{\eta_p Q_{p,n}(1)} \bar{Q}_{p,n}(I - \eta_n(I)) \right)^2 \right].$$

Related asymptotic variance formulae and comparisons are discussed in [48] in the context of random walks with absorbing barriers, including geometric killing rates and local reflection moves.

In [56], Nick Withley also obtains a uniform time bound on the asymptotic variance. However, our next proposition shows that, for a given system, the DMC method can have a uniform time bound on its asymptotic variance, despite the fact that its non-asymptotic variance is unbounded.

Proposition 2. *Let $G : x \in \mathbb{R} \mapsto e^{-\frac{x^2}{2}S}$, and P such that $\delta_x P \sim \mathcal{N}(Ax, B)$ for some $(A, B, S) \in (1, +\infty) \times \mathbb{R}_+^{*2}$. There exists an initial distribution η_0 such that*

$$\sup_{n \in \mathbb{N}} \sigma_n^2 < \infty \quad \text{and} \quad \sup_{n \in \mathbb{N}} \mathbb{E} \left(|\eta_n(I) - \eta_n^N(I)|^p \right)^{\frac{1}{p}} = +\infty, \quad (41)$$

where σ_n^2 is defined as in Lemma 7.

Proof :

Considering $\eta_0 \sim \mathcal{N}(m_0, \sigma_0^2)$ and using the Kalman filter's equations, we are able to fully describe the measures η_n . For $n \in \mathbb{N}^*$ we have

$$\eta_n \sim \mathcal{N}(m_n, \sigma_n^2) \quad \text{with} \quad \begin{cases} m_{n+1} = \frac{A}{1+S\sigma_n^2} m_n \\ \sigma_{n+1}^2 = \frac{A^2}{1+S\sigma_n^2} + B \end{cases}.$$

The limit measure is given by $\eta_\infty = \mathcal{N}(0, \sigma_\infty^2)$ where σ_∞^2 is the fixed point of the function $x \in \mathbb{R}_+ \mapsto \frac{A^2}{1+Sx} + B$. From now on, we assume that $\eta_0 = \eta_\infty$. We have then

$$\sigma_n^2 = \sum_{p=0}^n \frac{\eta_\infty(Q_{p,n}(I)^2)}{(\eta_\infty Q_{p,n}(1))^2}.$$

Using the calculations and notations from Lemma 10, we have then

$$\begin{cases} \eta_\infty(Q_{p,n}(1)) &= \frac{\lambda_{n-p}}{\sqrt{2\pi\sigma_\infty^2}} \int_{\mathbb{R}} e^{-\frac{y^2}{2\sigma_\infty^2}} e^{-\frac{y^2}{2} q_{n-p}} dy = \frac{\lambda_{n-p}}{\sqrt{q_{n-p}\sigma_\infty^2 + 1}} \\ \eta_\infty(Q_{p,n}(I)^2) &= \frac{\mu_{n-p}^2}{\sqrt{2\pi\sigma_\infty^2}} \int_{\mathbb{R}} y^2 e^{-\frac{y^2}{2\sigma_\infty^2}} e^{-y^2 q_{n-p}} dy = \frac{\mu_{n-p}^2 \sigma_\infty^2}{(2q_{n-p}\sigma_\infty^2 + 1)^{3/2}} \end{cases},$$

and

$$\sigma_n^2 = \sigma_\infty^2 \sum_{p=0}^n \frac{\mu_{n-p}^2 (q_{n-p}\sigma_\infty^2 + 1)^{3/2}}{\lambda_{n-p}^2 (2q_{n-p}\sigma_\infty^2 + 1)^{3/2}}.$$

As a solution to a Riccati equation, q_n converges towards some $q_\infty \in \mathbb{R}_+^*$ such that

$$q_\infty = \frac{A^2 q_\infty}{1 + q_\infty B} + S.$$

Since $A > 1$, we obtain that $1 + q_\infty B = A^2 + S/q_\infty > A$. Thus, there exists $C \in (0, 1)$ such that, for n large enough, we have

$$\mu_{n+1} \leq C \frac{\mu_n}{\sqrt{1 + q_n B}}$$

Thus, comparing the definition of λ_n and the previous bound, we deduce that there exists $\alpha \in \mathbb{R}$ such that we have

$$\frac{\mu_{n-p}^2 (q_{n-p}\sigma_\infty^2 + 1)^{3/2}}{\lambda_{n-p}^2 (2q_{n-p}\sigma_\infty^2 + 1)^{3/2}} \leq \alpha C^{2(n-p)}.$$

We can thus conclude on the right-hand side. of (41). The left-hand side is a direct application of Proposition 1. \square

Acknowledgments

MC thanks the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 863481) for financial support.

Appendix

Some technical Lemmas

Lemma 8. *Let $(G, V) \in \mathcal{C}(E) \times \mathcal{C}_\infty(E)$ be positive functions and $K \subset E$ be such that the right-hand side of (5) holds. There exists $c \in \mathbb{R}_+$ such that for any probability measure μ on E :*

$$\psi_G(\mu)(V) \leq \mu(V) + c.$$

Proof : Let $\bar{V} := \mathbb{1}_{\mathbb{R} \setminus K} V$. For any $(x, t) \in E^2$, we have:

$$0 \geq (G(x) - G(y))(\bar{V}(x) - \bar{V}(y)) = G(x)\bar{V}(x) + G(y)\bar{V}(y) - G(y)\bar{V}(x) - G(x)\bar{V}(y).$$

Integrating with respect to the probability measure μ over both x and y , we obtain:

$$\mu(G\bar{V}) - \mu(G)\mu(\bar{V}) \leq 0.$$

Hence

$$\psi_G(\mu)(\bar{V}) - \mu(\bar{V}) = \frac{\mu(G\bar{V}) - \mu(G)\mu(\bar{V})}{\mu(G)} \leq 0.$$

Thus

$$\psi_G(\mu)(V) \leq \psi_G(\mu)(\bar{V}) + \sup_K V \leq \mu(\bar{V}) + \sup_K V \leq \mu(V) + \sup_K V.$$

□

Lemma 9. *Let $V \in \mathcal{C}_\infty(E)$ be a Q -Lyapunov function, then*

$$\sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} [\phi_{q,n}(\eta_q^N)(V)] < +\infty \quad \text{and} \quad \sup_{\substack{(q,n) \in \mathbb{N}^2 \\ q \leq n}} \phi_{q,n}(\eta_q)(V) < +\infty. \quad (42)$$

Proof: First, let's prove that for any $\gamma \in \mathcal{P}_V(E)$, we have, with $(\epsilon, c) \in [0, 1) \times \mathbb{R}$ defined in (1),

$$\sup_{\substack{(q,n) \in \mathbb{N}^2 \\ q \leq n}} \phi_{q,n}(\gamma)(V) \leq \gamma(V) + \frac{c}{1 - \epsilon}. \quad (43)$$

To do so, we begin by using the Q -Lyapunov property of V as well as Lemma 8 for some $l \in \mathbb{N}^*$ to deduce that

$$\phi_{q,n}(\gamma)(V) = \frac{\phi_{q,n-1}(\gamma)(G_n P_n(V))}{\phi_{q,n-1}(\gamma)(G_n)} \leq \epsilon \phi_{q,n-1}(\gamma)(V) + c'.$$

By iterating the process, we obtain (43). We now prove that

$$\sup_{n \in \mathbb{N}} \mathbb{E} [\eta_n^N(V)] \leq \eta_0(V) + \frac{c'}{1 - \epsilon}. \quad (44)$$

Notice first that

$$\mathbb{E} [\eta_n^N(V) \mid \eta_{n-1}^N] = \frac{1}{N} \sum_{i=1}^N \eta_{n-1}^N S_{n-1, \eta_{n-1}^N} P_n(V) = \psi_{G_{n-1}}(\eta_{n-1}^N)(P_n(V)).$$

Then, using as previously the Q -Lyapunov property of V and Lemma 8 we obtain

$$\psi_{G_{n-1}}(\eta_{n-1}^N)(P_n(V)) \leq \epsilon \eta_{n-1}^N(V) + c'.$$

By iterating the process, we obtain

$$\mathbb{E} [\eta_n^N(V)] \leq \epsilon^n \eta_0(V) + c' \sum_{i=0}^{n-1} \epsilon^i \leq \eta_0(V) + \frac{c'}{1 - \epsilon}.$$

Thus, by combining (43) and (44), we can conclude regarding the first part of (42). The second part is obtained by proceeding in a strictly analogous way. \square

Proof of Lemma 1

We first consider the case where only (18) holds and prove that

$$\phi_{0,q}(\gamma)(Q_{q,n}(1)) \leq C \chi^{n-q-1}. \quad (45)$$

For $q \leq n + 2$, we have

$$\begin{aligned} Q_{q,n}(1)(x) &= Q_{q,n-2}(G_{n-1} P_n(G_n))(x) \\ &\leq \chi Q_{q,n-2}(G_{n-1})(x) \leq \chi Q_{q,n-1}(1)(x). \end{aligned}$$

Iterating the process, we deduce

$$Q_{q,n}(1) \leq \chi^{n-q-1} G_{q-1}.$$

Using (17) we deduce (45).

For $q \geq 1$ and $\mu \in \{\eta_q^N, \phi_q(\eta_{q-1}^N)\}$, we have then

$$\frac{1}{\mu(H_{q,n}^\gamma)} = \phi_{q,n}(\mu)(W) \times \frac{\phi_{0,q}(\gamma)Q_{q,n}(1)}{\mu(Q_{q,n}(W))} \leq C \frac{\chi^{n-q-1}\phi_{q,n}(\mu)(W)}{\mu(Q_{q,n}(W))}.$$

From Holder's inequality, Jensen's inequality and the hypothesis on W , we get :

$$\begin{aligned} \mathbb{E} \left[\mu(H_{q,n})^{-\beta} \right] &\leq C^\beta \mathbb{E} \left[\phi_{q,n}(\mu)(W)^{2\beta} \right]^{\frac{1}{2}} \mathbb{E} \left[\chi^{2\beta(n-q-1)} \mu(Q_{q,n}(W))^{-2\beta} \right]^{\frac{1}{2}} \\ &\leq \frac{C^\beta}{\chi^\beta} \mathbb{E} \left[\phi_{q,n}(\mu)(V)^{2\beta} \right]^{\frac{1}{2}} \mathbb{E} \left[\mu(W^{-2\beta}) \right]^{\frac{1}{2}}. \end{aligned}$$

Then, by choosing β small enough such that $W^{-2\beta} \leq \bar{W}$ and $V^{2\beta} \leq \bar{V}$, where \bar{W} and \bar{V} are Q -Lyapunov functions, we obtain

$$\sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[\mu(H_{q,n}^\gamma)^{-\beta} \right] \leq \frac{C^\beta}{\chi^\beta} \sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[\mu(\bar{V}) \right]^{\frac{1}{2}} \times \sup_{\substack{(q,n,N) \in \mathbb{N}^3 \\ q \leq n}} \mathbb{E} \left[\mu(\bar{W}) \right]^{\frac{1}{2}}.$$

We can conclude using Lemma 9.

The demonstration for time-homogeneous models with $E_0 < \chi$ is analogous. Indeed the equivalent of (45) is obtained by noticing that

$$\phi_{0,q}(\eta_\infty)Q_{q,n}(1) = \eta_\infty Q_{q,n-1}(G) = \eta_\infty Q_{q,n-1}(1)\eta_\infty(G) = E_0 \times \eta_\infty Q_{q,n-1}(1) = E_0^{n-q}.$$

The rest of the proof follows the same arguments, thus it is skipped. This ends the proof of the lemma. \square

Proof of Lemma 6

Let us begin by proving the result for $k = 1$. Take $x \in \mathbb{R}^d$:

$$\widehat{G}^{(1)}(x) = \frac{1}{\sqrt{\det(I + BS)}} \exp \left(-\frac{1}{2} x^T A^T (S^{-1} + B)^{-1} Ax \right),$$

Applying the Woodbury matrix identity we get:

$$\begin{aligned} &\sqrt{(2\pi)^d \det([B^{-1} + S]^{-1})} \times \widehat{P}^{(1)}(x, dy) \\ &= \exp \left(-\frac{1}{2} [y^T S y + (y - Ax)^T B^{-1} (y - Ax) - x^T A^T (S^{-1} + B)^{-1} Ax] \right) dy \\ &= \exp \left(-\frac{1}{2} (y - (I + BS)^{-1} Ax)^T (B^{-1} + S) (y - (I + BS)^{-1} Ax) \right) dy. \end{aligned}$$

Assume the property true for $k \in \mathbb{N}^*$. We let $\bar{S}_k := S_k + S$. In this notation, we have

$$\widehat{G}^{(k+1)}(x) = P(G\widehat{G}^{(k)})(x) = \lambda_k P \left(z \mapsto \exp\left(-\frac{1}{2}z^T \bar{S}_k z\right) \right) (x).$$

This implies that

$$\begin{aligned} & \widehat{G}^{(k+1)}(x) \\ &= \frac{\lambda_k}{\sqrt{(2\pi)^d \det(B)}} \int_{\mathbb{R}^d} \exp \left(-\frac{1}{2} [z \bar{S}_k z^T + (z - Ax)^T B^{-1} (z - Ax)] \right) dz \\ &= \frac{\lambda_k}{\sqrt{(2\pi)^d \det(B)}} \exp \left(-\frac{1}{2} [x^T A^T B^{-1} Ax - x^T A^T B^{-1} (\bar{S}_k + B^{-1})^{-1} B^{-1} Ax] \right) \\ & \times \int_{\mathbb{R}^d} \exp \left(-\frac{1}{2} (z - (\bar{S}_k + B^{-1})^{-1} B^{-1} Ax)^T (\bar{S}_k + B^{-1}) (z - (\bar{S}_k + B^{-1})^{-1} B^{-1} Ax) \right) dz \end{aligned}$$

from which we check that

$$\widehat{G}^{(k+1)}(x) = \frac{\lambda_k}{\sqrt{\det(I + BS + BS_k)}} \exp \left(-\frac{1}{2} x^T A^T (B + (S + S_k)^{-1})^{-1} Ax \right).$$

Moreover, since $\delta_x \widehat{P}^{(k)} \sim \mathcal{N}(A_k x, B_k^2)$ we also have

$$\delta_x \widehat{P}^{(k)} P \sim \mathcal{N}(AA_k x, AB_k A^T + B).$$

According the Gaussian update formula (see Proposition 4.5.2 in [43] for example), if $\eta = \mathcal{N}(m, \Sigma)$, then $\psi_{G_S}(\eta) = \mathcal{N}[(I + \Sigma S)^{-1} m, (I + \Sigma S)^{-1} \Sigma]$. Then we have

$$\psi_G(\delta_x \widehat{P}^{(k+1)}) \sim \mathcal{N}((I + AB_k A^T S + BS)^{-1} AA_k x, (I + AB_k A^T S + BS)^{-1} (AB_k A^T + B)).$$

This concludes the proof. \square

Lemma 10. Let $G : x \in \mathbb{R} \mapsto e^{-\frac{x^2}{2}S}$, and P such that $\delta_x P \sim \mathcal{N}(Ax, B)$ for some $(A, B, S) \in \mathbb{R} \times \mathbb{R}_+^{*2}$. Denoting by Q^n the n -times composition of the operator Q , we have

$$Q^n(1)(x) = \lambda_n e^{-\frac{x^2}{2}q_n} \quad \text{and} \quad Q^n(I)(x) = \mu_n x e^{-\frac{x^2}{2}q_n}, \quad (46)$$

with the parameters $(q_0, \lambda_0, \mu_0) = (S, 1, A)$ and

$$\begin{cases} q_{n+1} = \frac{A^2 q_n}{1 + q_n B} + S \\ \lambda_{n+1} = \frac{\lambda_n}{\sqrt{1 + q_n B}} \quad \text{and} \quad \mu_{n+1} = \frac{A \mu_n}{(1 + q_n B)^{3/2}} \end{cases}.$$

Proof :

For $n = 0$ the result is immediate. Assume that it holds for some $n \in \mathbb{N}$. In this situation, we have

$$\begin{aligned}
Q^{n+1}(1)(x) &= Q(Q^n(1))(x) = \lambda_n Q \left[y \mapsto e^{-\frac{y^2}{2} q_n} \right] (x) \\
&= \frac{\lambda_n}{\sqrt{2\pi B}} e^{-\frac{x^2}{2} S} \int_{\mathbb{R}} e^{-\frac{y^2}{2} q_n - \frac{(Ax-y)^2}{2B}} dy \\
&= \frac{\lambda_n}{\sqrt{2\pi B}} e^{-\frac{x^2}{2} \left(\frac{A^2}{B} - \frac{A^2}{B(1+q_n B)} + S \right)} \int_{\mathbb{R}} e^{-\frac{(1+q_n B) \left(\frac{A}{1+q_n B} x - y \right)^2}{2B}} dy \\
&= \frac{\lambda_n}{\sqrt{1+q_n B}} e^{-\frac{x^2}{2} \left(\frac{A^2 q_n}{1+q_n B} + S \right)}.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
Q^{n+1}(I)(x) &= Q(Q^n(I))(x) = \mu_n Q \left[y \mapsto ye^{-\frac{y^2}{2} q_n} \right] (x) \\
&= \frac{\mu_n}{\sqrt{2\pi B}} e^{-\frac{x^2}{2} \left(\frac{A^2 q_n}{1+q_n B} + S \right)} \int_{\mathbb{R}} ye^{-\frac{(1+q_n B) \left(\frac{A}{1+q_n B} x - y \right)^2}{2B}} dy \\
&= \frac{A\mu_n}{(1+q_n B)^{3/2}} xe^{-\frac{x^2}{2} \left(\frac{A^2 q_n}{1+q_n B} + S \right)}.
\end{aligned}$$

This ends the proof. □

References

- [1] Nicholas Metropolis and Stanislaw Ulam. The Monte Carlo method. *Journal of the American statistical association*, 44(247):335–341, 1949.
- [2] Monroe D. Donsker and Mark Kac. A sampling method for determining the lowest eigenvalue and the principal eigenfunction of schrödinger’s equation. *Journal of Research of the National Bureau of Standards*, 44(50):551–557, 1950.
- [3] Malvin H Kalos. Monte Carlo calculations of the ground state of three-and four-body nuclei. *Physical Review*, 128(4):1791, 1962.
- [4] Malvin H. Kalos D. M. Ceperley. *Monte Carlo Methods in Statistical Physics*. Springer-Verlag Berlin, 1986.
- [5] Peter J. Reynolds, David M. Ceperley, Berni J. Alder, and Jr. Lester, William A. Fixed-node quantum Monte Carlo for molecules. *The Journal of Chemical Physics*, 77(11):5593–5603, 12 1982.

- [6] Michel Caffarel and Pierre Claverie. Treatment of the Schrödinger equation through a Monte Carlo method based upon the generalized Feynman-Kac formula. *Journal of Statistical Physics*, 43:797–801, 1986.
- [7] Michel Caffarel and Pierre Claverie. Development of a pure diffusion quantum Monte Carlo method using a full generalized Feynman-Kac formula. i. formalism. *The Journal of chemical physics*, 88(2):1088–1099, 1988.
- [8] Jack H. Hetherington. Observations on the statistical iteration of matrices. *Physical Review A*, 30(5):2713, 1984.
- [9] Sandro Sorella. Green function Monte Carlo with stochastic reconfiguration. *Physical review letters*, 80(20):4558, 1998.
- [10] Sandro Sorella and Luca Capriotti. Green function Monte Carlo with stochastic reconfiguration: An effective remedy for the sign problem. *Physical Review B*, 61(4):2599, 2000.
- [11] Roland Assaraf, Michel Caffarel, and Anatole Khelif. Diffusion Monte Carlo methods with a fixed number of walkers. *Physical Review E*, 61(4):4566, 2000.
- [12] Baroni Stefano and Moroni Saverio. Reptation Quantum Monte Carlo: A method for unbiased ground-state averages and imaginary-time correlations. *Phys. Rev. Lett.*, 82:4745, 1999.
- [13] Michel Mareschal. The early years of quantum Monte Carlo (1): the ground state. *The European Physical Journal H*, 46(1):11, 2021.
- [14] Yao Ma, Lu Meng, Yan-Ke Chen, and Shi-Lin Zhu. Ground state baryons in the flux-tube three-body confinement model using diffusion Monte Carlo. *Physical Review D*, 107(5):054035, 2023.
- [15] Matthew WMC Foulkes, Lubos Mitas, RJ Needs, and Guna Rajagopal. Quantum Monte carlo simulations of solids. *Reviews of Modern Physics*, 73(1):33, 2001.
- [16] Pierre Del Moral and Laurent Miclo. Particle approximations of Lyapunov exponents connected to Schrödinger operators and Feynman-Kac semigroups. *ESAIM: Probability and Statistics*, 7:171–208, 2003.
- [17] Pierre Del Moral and Arnaud Doucet. Particle motions in absorbing medium with hard and soft obstacles. *Stochastic Analysis and Applications*, 22(5):1175–1207, 2004.
- [18] Mathias Rousset. On the control of an interacting particle estimation of schrödinger ground states. *SIAM J. Math. Anal.*, 38:824–844, 2006.
- [19] Eric Cancés, Benjamin Jourdain, and Tony Lelièvre. Quantum Monte Carlo simulations of fermions. A mathematical analysis of the fixed-node approximation. *Mathematical Models and Methods in Applied Sciences*, 16, 11 2011.

- [20] Mohamed El Makrini, Benjamin Jourdain, and Tony Lelièvre. Diffusion monte carlo method: Numerical analysis in a simple case. *ESAIM: Mathematical Modelling and Numerical Analysis*, 41(2):189–213, 2007.
- [21] Pierre Del Moral. *Feynman-Kac Formulae: Genealogical and Interacting Particle Systems With Applications*, volume 100. 05 2004.
- [22] Pierre Del Moral. *Mean field simulation for Monte Carlo integration*. CRC press, 2013.
- [23] Pierre Del Moral and Alice Guionnet. Large deviations for interacting particle systems: applications to non-linear filtering. *Stochastic processes and their applications*, 78(1):69–95, 1998.
- [24] Donald A Dawson and Pierre Del Moral. Large deviations for interacting processes in the strong topology. In *Statistical modeling and analysis for complex data problems*, pages 179–208. Springer, 2005.
- [25] Pierre Del Moral and Alice Guionnet. Central limit theorem for nonlinear filtering and interacting particle systems. *Annals of Applied Probability*, pages 275–297, 1999.
- [26] Nicolas Chopin. Central limit theorem for sequential Monte Carlo methods and its application to bayesian inference. 2004.
- [27] P. Del Moral and L. Miclo. *Branching and interacting particle systems approximations of Feynman-Kac formulae with applications to non-linear filtering*, pages 1–145. Springer Berlin Heidelberg, Berlin, Heidelberg, 2000.
- [28] Pierre Del Moral and Alice Guionnet. On the stability of interacting processes with applications to filtering and genetic algorithms. In *Annales de l’Institut Henri Poincaré (B) Probability and Statistics*, volume 37, pages 155–194. Elsevier, 2001.
- [29] Nick Whiteley. Sequential Monte Carlo samplers: error bounds and insensitivity to initial conditions. *Stochastic Analysis and Applications*, 30(5):774–798, 2012.
- [30] P Del Moral and E Horton. Coupled quantum harmonic oscillators and Feynman-Kac path integrals for linear diffusive particles. *Communications in Mathematical Physics*, 402(2):2079–2127, 2023.
- [31] Marc Arnaudon, Pierre Del Moral, and El Maati Ouhabaz. A lyapunov approach to stability of positive semigroups: An overview with illustrations. *Stochastic Analysis and Applications*, pages 1–80, 2023.
- [32] Pierre Del Moral, Emma Horton, and Ajay Jasra. On the stability of positive semigroups. *The Annals of Applied Probability*, 33(6A):4424–4490, 2023.
- [33] Malvin H. Kalos. Monte Carlo calculations of the ground state of three- and four-body nuclei. *Phys. Rev.*, 128:1791–1795, Nov 1962.

- [34] Malvin H. Kalos, Dominique Levesque, and Loup Verlet. Helium at zero temperature with hard-sphere and other forces. *Phys. Rev. A*, 9:2178–2195, May 1974.
- [35] Albert Messiah. *Quantum Mechanics*. Number vol. 2 in Quantum Mechanics. Elsevier Science, 1961.
- [36] David J. Griffiths and Darrell F. Schroeter. *Introduction to quantum mechanics*. Cambridge University Press, Cambridge ; New York, NY, third edition edition, 2018.
- [37] Gerhard Herzberg. *Molecular Spectra and Molecular Structure: Infrared and Raman of Polyatomic Molecules*. D. Van Nostrand Company, Inc., 1945.
- [38] Dmitry N. Makarov. Quantum entanglement and reflection coefficient for coupled harmonic oscillators. *Phys. Rev. E*, 102:052213, Nov 2020.
- [39] Artur K. Ekert. Quantum cryptography based on bell’s theorem. *Phys. Rev. Lett.*, 67:661–663, Aug 1991.
- [40] Elisabet Romero, Ramunas Augulis, Vladimir Novoderezhkin, Marco Ferretti, Jos Thieme, Donatas Zigmantas, and Rienk van Grondelle. Quantum coherence in photosynthesis for efficient solar energy conversion. *Nature Physics*, 10, 07 2014.
- [41] Pierre Del Moral. Non linear filtering: Interacting particle solution. *Markov Processes and Related Fields*, 2:555–580, 03 1996.
- [42] Rudolf E. Kálmán and Richard S. Bucy. New results in linear filtering and prediction theory. *Journal of Basic Engineering*, 83:95–108, 1961.
- [43] Pierre Del Moral and Spiridon Penev. *Stochastic Processes: From Applications to Theory*. December 2016.
- [44] Nick Whiteley. Stability properties of some particle filters. *The Annals of Applied Probability*, 23(6):2500 – 2537, 2013.
- [45] P.Del Moral and L. Miclo. A Moran particle system approximation of Feynman-Kac formulae. *Stochastic Processes and their Applications*, 86(2):193–216, 2000.
- [46] Marc Arnaudon and Pierre Del Moral. A duality formula and a particle Gibbs sampler for continuous time Feynman-Kac measures on path spaces. *Electronic Journal of Probability*, 25(none):1 – 54, 2020.
- [47] Mathias Rousset. On the control of an interacting particle estimation of Schrödinger ground states. *SIAM Journal on Mathematical Analysis*, 38(3):824–844, 2006.
- [48] Pierre Del Moral and Ajay Jasra. A note on random walks with absorbing barriers and sequential monte carlo methods. *Stochastic Analysis and Applications*, 36(3):413–442, 2018.

- [49] Pierre Del Moral. Measure-valued processes and interacting particle systems. application to nonlinear filtering problems. *The Annals of Applied Probability*, 8(2):438–495, 1998.
- [50] Tosio Kato. *Perturbation Theory for Linear Operators*. 1966.
- [51] Chi-Kwong Li and Fuzhen Zhang. Eigenvalue continuity and gersgorin’s theorem. *Electronic Journal of Linear Algebra*, 35:619–625, 12 2019.
- [52] Pierre Del Moral and Emma Horton. A theoretical analysis of one-dimensional discrete generation Ensemble Kalman particle filters, 07 2021.
- [53] Nick Whiteley, Nikolas Kantas, and Ajay Jasra. Linear variance bounds for particle approximations of time-homogeneous Feynman-Kac formulae. *Stochastic Processes and their Applications*, 122(4):1840–1865, 2012.
- [54] Pierre Del Moral and Jean Jacod. The Monte-Carlo method for filtering with discrete time observations. central limit theorems. In *Proceedings Workshop on Numerical Methods and Stochastics. The Fields Institute 1999*, editors T.J. Lyons and T.S Salisbury. American Mathematical Society, 2002.
- [55] Pierre Del Moral and J.Jacod. Interacting particle filtering with discrete-time observations: asymptotic behaviour in the gaussian case. In Kunita H. Rajput B. Watanabe S. Hida T., Karandikar R. and Xiong J., editors, *Stochastics in finite/infinite dimensions*, pages 101–123. Trends in Mathematics, Birkhauser, 2001.
- [56] Nick Whiteley. Stability properties of some particle filters. *The Annals of Applied Probability*, 23(6), dec 2013.