



Evaluation of the biodegradability of chlordecone during mesophilic and thermophilic organic waste treatment via anaerobic digestion

Perla Alnajjar, Déborah Martin, Claire Soligot-Hognon, Delphine Muselet, Hussein Kanso, Stéphane Pacaud, Cyril Feidt, Pierre-Loïc Saaidi, Yves Le Roux

► To cite this version:

Perla Alnajjar, Déborah Martin, Claire Soligot-Hognon, Delphine Muselet, Hussein Kanso, et al.. Evaluation of the biodegradability of chlordecone during mesophilic and thermophilic organic waste treatment via anaerobic digestion. 7 ISEBE - Seventh International Symposium on Environmental Biotechnology and Engineering, May 2023, Marseille, France. hal-04410185

HAL Id: hal-04410185

<https://hal.science/hal-04410185>

Submitted on 22 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evaluation of the biodegradability of chlordecone during mesophilic and thermophilic organic waste treatment via anaerobic digestion

Perla Alnajjar^{1,2}, Déborah Martin³, Claire Soligot-Hognon¹, Delphine Muselet³, Hussein Kanso³, Stéphane Pacaud⁴, Cyril Feidt¹, Pierre-Loïc Saaidi³, Yves Le Roux^{1,4}

- 1 URAFPA, Université de Lorraine, 2 Avenue de la Forêt de Haye, 54505 Vandœuvre-Lès-Nancy, France
- 2 Ecole doctorale en Sciences et Technologie, Université Libanaise, Tripoli, Liban
- 3 UMR 8030 Génomique Métabolique, CEA, Institut de Biologie François Jacob, Genoscope, Université d'Evry Val d'Essonne, Université Paris-Saclay, Evry, France
- 4 Agrométhéa, Chaire industrielle-ENSAIA, 54280 Champenoux, France

Anaerobic digestion (AD) has long been studied as an effective treatment strategy for organic wastes contaminated with recalcitrant pollutants. The bioremediation potential of AD was tested on digestates supplemented with chlordecone (CLD, $C_{10}Cl_{10}H_2O_2$), a persistent organic pollutant that has been applied in the French West Indies (1972-1993) against the banana black weevil *Cosmopolites sordidus*. Its application has resulted in an extensive long-term environmental contamination; concentrations between 0.03-24.2 mg.kg⁻¹ of soils are still recorded¹. CLD has been shown to accumulate in the food chain² triggering harmful acute and chronic toxicities in humans³.

Anaerobic digestates from animal and plant origins were spiked with 20 mg.L⁻¹ of CLD and incubated, in batch mode, under mesophilic (37.5°C) and thermophilic (55°C) conditions, respectively. This concentration was chosen to facilitate the detection and/or quantification of transformation products (TPs) at early stages as well as the quantification of CLD after 40 incubation days. A sacrificial approach was followed in which the bottles were stopped at different incubation days (0, 5, 20, and 40), extracted with a developed QuEChERS⁴ protocol, and analyzed with a combined GC-MS and LC-HRMS approach.

The Biochemical Methane Potential (BMP), followed throughout the entire experiment, showed that a concentration of CLD as high as 20 mg.L⁻¹ did not alter the production of biogas, as evidenced by the absence of a statistically significant difference between the bottles filled with CLD-treated digestates and the control bottles. Moreover, a redox potential lower than -350mV was maintained, providing thus sufficient reducing conditions reported to be needed for efficient CLD degradation⁵. Indeed, a significant reduction of around 87 +/- 4% of CLD under

thermophilic conditions, after 40 days, was observed. Under mesophilic conditions, the reduction of CLD was rather moderate with a decrease in CLD concentration of 32 +/- 7%. Various TPs, with a similar profile to that obtained with previous microbiological degradation studies^{6,7}, appeared in a concomitant manner with CLD's removal. Hydrochlordecones ($C_{10}Cl_{10-n}O_2H_{2+n}$, $n=1,2$), arising from reductive dechlorination, as well as polychloroindenes ($C_9Cl_{6-n}H_{2+n}$, $n=1,2$), formed after ring-opening, were quantified. Polychloroindenecarboxylic acids ($C_{10}Cl_{5-n}H_{3+n}O_2$, $n=1,2$) were also detected.

The anaerobic digestion, in our experimental conditions, has therefore provided the adequate conditions needed not only to dechlorinate CLD but also to open its ring structure which is partially responsible for its persistence in the environment. AD could thus possibly be considered as an effective treatment technique for CLD-contaminated organic wastes. The fate of the obtained TPs as well as their possible toxicity will be explored in subsequent works.

References:

- 1: Crabit, A., Cattani, P., Colin, F., & Voltz, M. (2016). Soil and river contamination patterns of chlordecone in a tropical volcanic catchment in the French West Indies (Guadeloupe). *Environmental Pollution*, 212, 615–626. <https://doi.org/10.1016/j.envpol.2016.02.055>
- 2: Dubuisson, C., Héraud, F., Leblanc, J.-C., Gallotti, S., Flamand, C., Blateau, A., Quenel, P., & Volatier, J.-L. (2007). Impact of subsistence production on the management options to reduce the food exposure of the Martinican population to Chlordecone. *Regulatory Toxicology and Pharmacology: RTP*, 49(1), 5–16. <https://doi.org/10.1016/j.yrtph.2007.04.008>
- 3: Maudouit, M., & Rochoy, M. (2019). [Systematic review of the impact of chlordecone on human health in the French West Indies]. *Thérapie*, 74(6), 611–625. <https://doi.org/10.1016/j.therap.2019.01.010>
- 4: Déborah Martin, Delphine Muselet, Perla Alnajjar, Stéphane Pacaud, Yves Le Roux, Sébastien Bristeau, Jennifer Hellal, Cyril Feidt et Pierre-Loïc Saaidi. Protocole d'extraction QuEChERS pour l'analyse de la chlordécone et ses produits de transformation dans des procédés de (bio)remédiation et des échantillons de sol. Poster présenté dans le cadre du « GFP 2022 », 18-20 Mai 2022, Namur, Belgique.
- 5: Dictor M.C., Mercier A., Lereau L., Amalric L., Bristeau S., Mouvet C., avec la collaboration de Auger P., Béchu E., Breeze D., Touzelet S., Tris H. et Henry B. (2011). Décontamination de sols pollués par la chlordécone. Validation de procédés de dépollution physico-chimique et biologique, étude des produits de dégradation et amélioration de la sensibilité analytique pour la chlordécone dans les sols. Rapport final. BRGM/RP-59481-FR, 201p., 70 fig., 42 tabl., 6 ann.
- 6: Chevallier, M. L., Della-Negra, O., Chaussonnerie, S., Barbance, A., Muselet, D., Lagarde, F., Darii, E., Ugarte, E., Lescop, E., Fonknechten, N., Weissenbach, J., Woignier, T., Gallard, J.-F., Vuilleumier, S., Imfeld, G., Le Paslier, D., & Saaidi, P.-L. (2019). Natural Chlordecone Degradation Revealed by Numerous Transformation Products Characterized in Key French West Indies Environmental Compartments. *Environmental Science & Technology*, 53(11), 6133–6143. <https://doi.org/10.1021/acs.est.8b06305>
- 7: Lomheim, L., Laquitaine, L., Rambinaising, S., Flick, R., Starostine, A., Jean-Marius, C., Edwards, E. A., & Gaspard, S. (2020). Evidence for extensive anaerobic dechlorination and transformation of the pesticide chlordecone ($C_{10}Cl_{10}O$) by indigenous microbes in microcosms from Guadeloupe soil. *PloS One*, 15(4), e0231219.