



**HAL**  
open science

# Rotating polygons : Numerical and experimental study of the base flow

Wen Yang, Ivan Delbende, Yann Fraigneau, Laurent Martin Witkowski

► **To cite this version:**

Wen Yang, Ivan Delbende, Yann Fraigneau, Laurent Martin Witkowski. Rotating polygons : Numerical and experimental study of the base flow. 12th European Fluid Mechanics Conference, Sep 2018, Vienna, Austria. hal-04410239

**HAL Id: hal-04410239**

**<https://hal.science/hal-04410239v1>**

Submitted on 22 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Rotating polygons : Numerical and experimental study of the base flow

W. Yang<sup>\*†</sup>, I. Delbende<sup>‡§</sup>, Y. Fraigneau<sup>§</sup> and L. Martin Witkowski<sup>‡§</sup>

Rotating polygons, as shown in Fig. 1 (right), have challenged the curiosity of various research groups<sup>12</sup>. The flow is induced by a disk rotating at the bottom of a cylindrical tank partially filled with a working fluid. The polygons appear above a threshold that breaks the symmetry of the base flow initially axisymmetric. Instability mechanisms have been proposed based on simplified base flow model<sup>3</sup>. The aim of our study is to characterize in details the base flow using numerical techniques – computation of steady solutions or mean unsteady two-dimensional and three-dimensional direct simulations – as well as Laser-Doppler Velocimetry measurements.

Numerical results averaged in time and space reveal the structure of the toroidal flow located at the periphery of the central solid body rotation region. When viewed in a meridional plane, this flow cell is found bordered by four layers, in particular by a layer at the free surface leading to a better understanding of the complete flow structure. The Froude number associated to surface deformation and the Reynolds number if sufficiently large are shown to have little effect on the flow structure. Experimental measurements corroborate the azimuthal velocity distribution used in the simplified models. But experimental and numerical results also reveal a large robust modal structure as seen in Fig. 1 (left) with the same azimuthal wave number as the corners' number of the rotating polygon. This structure is observed while the free surface remains almost axisymmetric and thus before the rotating polygon settles in.

---

\*Sorbonne Université, Collège Doctoral, F-75005 Paris

†FAST UMR 7608, Parc-Club Orsay Université, F-91405 Orsay, France

‡Sorbonne Université, Faculté des Sciences et Ingénierie, UFR d'Ingénierie, F-75005 Paris

§LIMSI-CNRS, Bât 508, rue John von Neumann, F-91405 Orsay Cedex, France

<sup>1</sup>Vatistas, *J. Fluid Mech.* **217**, 241 (1990).

<sup>2</sup>Jansson et al., *Phys. Rev. Lett.* **96**, 174502 (2006).

<sup>3</sup>Mougel et al., *J. Fluid Mech.* **824**, 230 (2017)




Figure 1: (Left) Visualization with kalliroscope of a mode 3 below threshold for the rotating polygon, (Right) Rotating polygon once it is settled.