



HAL
open science

Mechanism involved of post-exercise cold water immersion: Blood redistribution and increase in energy expenditure during rewarming

Dorian Giraud, Laura Pomportes, Caroline Nicol, Denis Bertin, Jean-Laurent Gardarein, Arnaud Hays

► To cite this version:

Dorian Giraud, Laura Pomportes, Caroline Nicol, Denis Bertin, Jean-Laurent Gardarein, et al.. Mechanism involved of post-exercise cold water immersion: Blood redistribution and increase in energy expenditure during rewarming. *Temperature*, In press, 10 (1), pp.3-8. 10.1080/23328940.2024.2303332 . hal-04411334v1

HAL Id: hal-04411334

<https://hal.science/hal-04411334v1>

Submitted on 23 Jan 2024 (v1), last revised 8 Feb 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Mechanism involved of post-exercise cold water immersion: Blood redistribution and increase in energy expenditure during rewarming

Dorian Giraud, Laura Pomportes, Caroline Nicol, Denis Bertin, Jean-Laurent Gardarein & Arnaud Hays

To cite this article: Dorian Giraud, Laura Pomportes, Caroline Nicol, Denis Bertin, Jean-Laurent Gardarein & Arnaud Hays (15 Jan 2024): Mechanism involved of post-exercise cold water immersion: Blood redistribution and increase in energy expenditure during rewarming, Temperature, DOI: [10.1080/23328940.2024.2303332](https://doi.org/10.1080/23328940.2024.2303332)

To link to this article: <https://doi.org/10.1080/23328940.2024.2303332>



© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



View supplementary material [↗](#)



Accepted author version posted online: 15 Jan 2024.



Submit your article to this journal [↗](#)



Article views: 82



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis & Informa UK Limited, trading as Taylor & Francis Group

Journal: Temperature

DOI: 10.1080/23328940.2023.2179820

Mechanism involved of post-exercise cold water immersion: Blood redistribution and increase in energy expenditure during rewarming

Dorian Giraud^{1,2}, Laura Pomportes³, Caroline Nicol³, Denis Bertin^{1,3}, Jean-Laurent Gardarein², Arnaud Hays¹

1 – Aix-Marseille University, HIPE Human Lab, Marseille, France

2 - Aix-Marseille University, CNRS, IUSTI, Marseille, France

3 - Aix-Marseille University, CNRS, ISM, Marseille, France

Corresponding author : dorian.giraud@univ-amu.fr

Abstract

Thermogenesis is well understood, but the relationships between cold water immersion (CWI), the post-CWI rewarming and the associated physiological changes are not. This study investigated muscle and systemic oxygenation, cardiorespiratory and haemodynamic responses, and gastrointestinal temperature during and after CWI. 21 healthy men completed randomly 2 protocols. Both protocols consisted of a 48 minutes heating cycling exercise followed by 3 recovery periods (R1-R3), but they differed in R2. R1 lasted 20 minutes in a passive semi-seated position on a physiotherapy table at ambient room temperature. Depending on the protocol, R2 lasted 15 minutes at either ambient condition (R2_AMB) or in a CWI condition at 10 °C up to the iliac crest (R2_CWI). R3 lasted 40 minutes at AMB while favouring rewarming after R2_CWI. This was followed by 10 minutes of cycling. Compared to R2_AMB, R2_CWI ended at higher $\dot{V}O_2$ in the non-immersed body part due to thermogenesis (7.16(2.15) vs. 4.83(1.62) ml.min⁻¹.kg⁻¹) and lower femoral artery blood flow (475(165) vs. 704(257) ml.min⁻¹) (p<0.001). Only after CWI, R3 showed a progressive decrease in vastus and gastrocnemius medialis O₂ saturation, significant after 34 minutes (p<0.001). As blood flow did not differ from the AMB protocol, this indicated local thermogenesis in the immersed part of the body. After CWI, a lower gastrointestinal temperature on resumption of cycling compared to AMB (36.31(0.45) vs. 37.30(0.49) °C, p<0.001) indicated incomplete muscle thermogenesis. In conclusion, the rewarming period after CWI was non-linear and metabolically costly. Immersion and rewarming should be considered as a continuum rather than separate events.

Keywords:

- Thermoregulation
- Cold-water immersion
- Muscle oxygenation
- Oxygen uptake
- Core temperature
- Cardiocirculatory system
- Metabolic kinetics

Abbreviation

AMB = Ambient temperature (°C)

ATP = Adenosine triphosphate

BAT = Brown adipose tissue

CWI = Cold-water immersion

FAD = Femoral Artery Diameter (mm)

FABF = Femoral Artery Blood Flow ($\text{ml}\cdot\text{min}^{-1}$)

FABV = Femoral Artery Blood Velocity ($\text{cm}\cdot\text{s}^{-1}$)

T_{GI} = Gastrointestinal temperature (°C)

NST = Non-shivering thermogenesis

\dot{Q}_c = Cardiac output ($\text{L}\cdot\text{min}^{-1}$)

RER = Respiratory exchange ratio

MSO_2GM = Muscle oxygen saturation of the Gastrocnemius Medialis muscle (%)

MSO_2VM = Muscle oxygen saturation of the Vastus Medialis muscle (%)

ST = shivering thermogenesis

UCP1 = uncoupling protein 1

\dot{V}_E = Ventilation ($\text{L}\cdot\text{min}^{-1}$)

$\dot{V}\text{CO}_2$ = Relative carbon dioxide output ($\text{ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$)

$\dot{V}\text{O}_2$ = Relative oxygen uptake ($\text{ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$)

VT1 = First ventilatory threshold

VT2 = Second ventilatory threshold

Introduction

Cold-water Immersion (CWI) is a common recovery method used in various sports, such as contact sports [1], sports with high mechanical stress [2] and endurance sports [3]. Recent research [4] found that 86 % of a panel of 111 athletes, coaches and support practitioners used CWI, and 78 % found it effective for recovery. Several mechanisms are proposed to explain the beneficial effects of CWI on recovery including hydrostatic pressure and cold-induced analgesic effect and vasoconstriction [5]. However, controversial results remain in the literature, largely due to the variability in CWI protocols between studies [4]. Environmental conditions (e.g., room temperature, humidity) are another source of discrepancies between studies. The recommendation for or against the use of CWI is influenced by a number of factors, including the duration and temperature of CWI, the degree of hyperthermia prior to CWI, the duration of the post-CWI event, the control protocol used, the potential benefit on inflammation, and the type of exercise [6]. The depth of immersion also plays an important role in the CWI protocol [7]. On the other hand, the energy expenditure due to thermoregulation accounts for a significant proportion of our total energy expenditure under normal conditions [8]. As the cooling/rewarming couple appears to be inseparable, a number of studies have focused on temperature changes during different rewarming protocols. It has been shown that body temperature continues to fall after leaving a cold bath, known as the 'afterdrop'. This has been observed at three measurement points with similar kinetics but different timing in rectal, oesophageal and ear canal temperatures [9,10]. It was initially proposed that afterdrop was due to increased perfusion of the extremities as a result of post-immersion vasodilation, increasing the amount of cooled blood returning to cool the core, but this hypothesis was refuted by a study focusing on blood flow variation during rewarming [10]. Instead, this study suggested that this phenomenon was related to heat conduction along a thermal gradient towards a cold inner shell (including convective heat exchange between tissues of different temperatures) and a reduction in heat production due to shivering inhibition. The conduction and convection phenomena were validated by Romet [9]. Giesbrecht et al. [11] reported a much lower energy expenditure during shivering suppression, starting 5 min after injection of the suppressant (mepederine) and continuing between 30 and 35 min after immersion. Another study investigated different modalities of rewarming [12] and found approximately the same kinetics of metabolic rate (calculated from oxygen consumption, $\dot{V}O_2$) as Giesbrecht et al. [11]. This demonstrates the significant metabolic impact of the rewarming period. It is therefore interesting to quantify the energetic cost of a cold bath and its subsequent rewarming. Indeed, as part of a recovery strategy, it may be important to compensate for the energy deficit caused by CWI with an adequate energy intake.

Few studies have focused on the energy costs of CWI. Four studies have found an increase in $\dot{V}O_2$ during CWI [11,13–15], reflecting an increase in body heat production. Three thermoregulatory mechanisms can explain this heat production. The first two mechanisms refer to shivering thermogenesis (ST) and non-shivering thermogenesis (NST). ST can be defined as involuntary rhythmic muscle contractions controlled by the sympathetic nervous system to produce heat [16]. The ATP production required to sustain shivering comes from glucose, lipids and proteins, depending on the nutritional status of the participant prior to cold exposure, the shivering intensity and variation in muscle fibre recruitment [17,18]. Giesbrecht et al. [11] highlighted the energy cost of ST, during CWI and during the rewarming period. NST is divided into two parts: non-shivering muscle thermogenesis (NST-M) and non-shivering thermogenesis due to the use of brown (and beige) adipose tissue (NST-B). NST-M is based on the activity of the SERCA protein in the sarcoplasmic reticulum of muscle fibres, which allows the active transport of Ca^{2+} from the cytosol to the lumen of the sarcoplasmic reticulum [19]. In the case of NST-M, sarcolipin induces a structural change in the Ca^{2+} transport site that prevents Ca^{2+} from being transported during each catalytic cycle and pushes it back into the cytosol. ATP is hydrolysed without Ca^{2+} transport, thereby producing only heat [20]. Another poorly understood mechanism is mitochondrial proton leak, which could be mediated by the UCP3 protein or by changes in mitochondrial morphology [21]. Concerning NST-B, brown adipose tissue (BAT) contains mitochondria and uncoupling protein 1 (UCP1) [22]. When this protein is activated, the mitochondrial proton gradient converts the energy of fat oxidation into heat rather than creating ATP [23,24]. However, $\dot{V}O_2$ changes due to cold water immersion must be interpreted with caution, as cold shock responses occur within the first 30 sec and stabilise after 3 min in most cases [25]. These responses include an inspiratory gasp of 2 and 3 litres, depending on water temperature [26]. Increased ventilation due to cold shock causes hypocapnia [27], which is partially offset by an increase in metabolic rate, leading to an increase in CO_2 release [26]. This increase in metabolic rate logically leads to an increase in $\dot{V}O_2$. At the same time, there is an increase in cardiac output [28] due to tachycardia and the increase of cutaneous vasoconstriction [26]. All these mechanisms are due to the cold shock response and are not thermoregulatory mechanisms.

Vasoconstriction is another important thermoregulatory mechanism to conserve the heat, as blood plays an essential role in transporting heat to tissues, both to regulate deep body temperature and to rewarm tissues. Vasoconstriction leads to a reduction in the transport of heat by the blood, resulting in a drop in muscle and skin temperature. This event has 2 effects: First, less cold blood flows back to the central core, helping to protect deep body temperature. Second, it reduces the heat flow between the body and the water, thereby reducing the heat loss from the body by reducing the temperature difference between the muscles and the skin with water. Superficial and

deep body tissues continue to cool by conduction, but the body reduces heat delivery to the submerged limb, preserving heat for the central core.

These thermoregulatory mechanisms are primarily activated by the transient receptor potential cation channel subfamily M (melastatin) member 8 (TRMP8), which transmits information to the preoptic area. TRMP8 is a cation channel whose conductance is activated by mild cooling (<28 °C) [29]. A feedback loop related to the deep body temperature involves detection of brain, spinal cord and visceral temperature [30]. During acute cold stress, these different sensors elicit different responses according to environmental and deep body temperature. The periods of exposure to cold and heat are likely to result in different energy expenditure responses. Therefore, to understand the impact of CWI, it is necessary to understand the time course of thermoregulatory mechanisms during the exposure and rewarming periods.

The present study was designed to investigate the specific cooling-rewarming pair after a heating exercise, which reflects a common type of recovery in high performance sports. In this context, we investigated the time course of central or local heat production and vasoconstriction during the subsequent CWI and rewarming period after a heating cycling exercise. We hypothesised that the different thermoregulatory mechanisms that occur during rewarming compared to CWI may be an additional cause of the different effects reported in the literature on increased energy expenditure during and after CWI. To test this hypothesis, several physiological parameters were measured pre-, per- and post CWI.

Methods

Twenty-one male volunteers (mean(SD); age: 24(5) years, body mass: 68(8) kg; height: 177(6) cm; $\dot{V}O_{2max}$: 55(6) ml.kg⁻¹.min⁻¹, maximal aerobic power: 300(50) W, body fat percentage: 14.7(3) %) who regularly participated in endurance sports such as cycling or trail running, were recruited for this study. All participants were fully informed of the potential risks and requirements of the study and provided written informed consent. The study was approved by the national ethical committee for STAPS (Science and Technique of Physical activity and Sport) Research (number IRB00012476-2022-21-04-176). To be eligible, participants were required to have completed an incremental cycling exercise test providing data on their power at the first and second ventilatory thresholds (P_{VT1} and P_{VT2} , respectively), maximal aerobic power and $\dot{V}O_{2max}$. Any participant who failed to complete the test protocol or sustained an injury during the process was excluded from the study.

Incremental exercise test

The incremental protocol consisted of a 2 min warm-up at 100 W, followed by increments of 30 W every 2 min. The first ventilatory threshold (VT1) was defined as the time at which an initial deviation

from expiratory volume ($\dot{V}E$) linearity was observed, and when a systematic increase in the ventilatory equivalent for O_2 ($\dot{V}E/\dot{V}O_2$) and the fraction of expired O_2 first appeared. The second ventilatory threshold (VT2) was defined based on a secondary increase in $\dot{V}E$ and $\dot{V}E/\dot{V}O_2$, and a marked increase in the ventilatory equivalent for CO_2 ($\dot{V}E/\dot{V}CO_2$) combined with a decrease in the fraction of expired CO_2 [31]. Participants were required to abstain from alcohol (48 h), caffeine, tea or guarana shot (24 h) and strenuous exercise (24 h) before and during the experiment. Additionally, participants were requested to arrive well-hydrated.

Experimental design

To ensure familiarity with the protocol for measuring cardiac output ($\dot{Q}c$), all participants underwent a familiarisation session. This was followed by two experimental protocols, randomised 48 hours apart.

As shown in Figure 1, both protocols session started with a 5 min warm-up cycle exercise (WU), followed by a 48 min heating cycling exercise (H), 3 recovery periods (R1-R3), and a final cycling exercise (FI). The protocols only differed in R2. More precisely, WU started at 80 % of the power at VT1, followed by 4x30 s at the power at VT1 plus 90 % of the difference between the power at VT1 and the power at VT2, interspaced by 1 min passive recovery [32]. It ended with 2 all-out isokinetic sprints at a mechanical cadence of 120 rpm [33]. The 48 min H exercise consisted of 8 sets of 5 min at a fixed cycling workload (90 % of the power reach at VT1) with 1 min recovery at 75 W. Cognitive tasks were performed during sets 1, 2, 4, 5, 7 and 8. These data were used in another study investigating cognitive decision making before, during and after exercise [34]. As the data for this period are not covered in a way that is consistent with the topic of this article, you will find the data for the heating period in the supplementary material. All recovery periods (R1-R3) were carried out in a standardised semi-seated position at 20.8(1.2) °C room temperature and 40.7(8.9) % humidity. In R1, participants rested passively on a physiotherapy table for 25 min while some measurements were taken and some equipment was placed on the participant (as described in the Measurement section), as in the Peiffer et al. [35] study. During R1, the recorded physiological variables returned to relatively stable values, while rectal temperature continued to fall thereafter [36]. Depending on the protocol, R2 lasted 15 min in either ambient condition (R2_AMB) or cold water immersion (R2_CWI) conditions, which determined the name of the protocol (either CWI or AMB) (Figure 1). The order between these 2 protocols was randomised and well-balanced between participants. More precisely, in the R2_CWI condition, the participant sat down in a bathtub filled-up with water at 10 °C for 15 min [37]. The instruction was to sit down as soon as possible on a stool adjusted so that water would rise up to the iliac crest. The bathtub was located 1 metre from the physiotherapy table where the passive R1 recovery was completed. In the R2_AMB condition, participants mimicked getting in and

out of the bathtub to ensure that the observed effects were related to cold and not to movement. For both protocols, in R3 the participant sat on the physiotherapy table at ambient room temperature for a further 40 min [37]. In the protocol including the R2_CWI condition, the participant dried off with their towel, kept their bathing suit and put on a dry T-shirt. For them only, R3 corresponded to a rewarming period. Both protocols ended with a FI similar to the exercise used in WU to examine the changes in deep body temperature during a cycling activity following CWI.




Figure 1: Test protocols and measurement schedule. CWI, cold-water immersion at 10(0.5) °C; AMB, ambient room condition at 20.8(1.2) °C.

Measurements

1h15 before the start of the CWI, participants were asked to swallow a pill (e-Celsius Performance, BodyCap, France) which was connected to their monitor via Bluetooth to record their gastrointestinal temperature (T_{GI}) every 30 s throughout the experiment. For ease of reading, we have set $T=0$ as the start of the R2(CWI and AMB) period (Figure 1). Cardiorespiratory variables including $\dot{V}O_2$, $\dot{V}CO_2$, $\dot{V}E$ and respiratory exchange ratio (RER) were recorded with breath-by-breath acquisition and cardiac output (\dot{Q}_c) by rebreathing [38] from -7 to 55 min (end of R3) using the Innocor system (Cosmed, Italy). To reflect the local energy expenditure of the targeted muscles, muscle oxygen saturation of the vastus medialis (MSO_2VM) and gastrocnemius medialis (MSO_2GM) of the left lower limb were recorded with a NIRS system (MOXY, Moxymonitor, United States) at 2 Hz as previously described [39,40]. To monitor the vasoconstriction mechanisms, the changes in superficial femoral artery blood flow (FABF) complemented the \dot{Q}_c recordings. FABF was measured using a Doppler ultrasound device (Echowave, Teled, Italy) at each of the measurement times described in Figure 1. The probe was 6 cm wide and the sampling frequency set at 9 MHz (HL9.0/60/128Z-2, Teled, Italy). FABF Data were

collected at an insonation angle of 60 ° [41]. The room humidity and temperature were measured with one thermo-hygro-barometer (RS-1161, RS Pro, United Kingdom). The water temperature was continuously monitored and kept at 10(0.5) °C using a water temperature controller (CryopackPerf, Cryo control, Castanet Tolosan, France). The water was not stirred manually, but the cooling system necessarily involved suction and discharge of water into the basin, resulting in a movement of water in the bath.

Data Management

FABF was determined from superficial femoral artery diameter (FAD) and blood velocity (FABV) data using an Echowave (Echo wave II, version 4.1.0, Telemed, Italy). FAD was defined as the average of measurements in 4 directions and FABV was the average peak blood velocity during 5 heartbeats before the FAD image was taken. FABF was calculated as follows:

$$FABF = \frac{\pi \times \frac{FAD^2}{2}}{1000} \times \frac{FABV}{100} \times 10^5$$

where FAD is measured in mm, FABV in $\text{cm}\cdot\text{s}^{-1}$ and FABF in $\text{ml}\cdot\text{min}^{-1}$.

The recorded T_{GI} , MSO_2 and cardiorespiratory data were imported into Matlab (R2020a, MathWorks, United states). Gas exchange data were smoothed with 11-point rolling average (between $n-5$ and $n+5$), and then averaged over the 30-s period prior to each \dot{Q}_c (the 9 recording times are shown in Figure 1). T_{GI} was measured at 5 min intervals from the end of the last heating exercise intensity until the first \dot{Q}_c measurement, then at each \dot{Q}_c measurement. Finally, data were collected after 5 min of WU after R3, and at the end of WU (12 min cycling).

Statistics

A linear mixed model was performed using the lme4 [42] and lmerTest [43] packages in R. To assess the protocol effect as a function of time, protocol and time were used as fixed effects and the random effects included test day and subject. Time corresponds to each measurement instant as described earlier in the method. A pairwise comparison with Holm's adjustment was then used for the interaction. The following data were included in the analysis: $\dot{V}O_2$, $\dot{V}CO_2$, $\dot{V}E$, RER, T_{GI} , \dot{Q}_c and FABF. The statistical method was similar to that used in the study by Macchi et al. [44]. The effect size (ES) of mean differences was determined using Cohen's d coefficient, with effect sizes equal to or greater than 0.2, 0.6, and 1.2 interpreted as small, moderate, and large effects, respectively [45]. Partial eta squared (η_p^2) was calculated for each interaction. For MSO_2VM and MSO_2GM , a one-dimension statistical parametric mapping (SPM) [46] was used to examine the influence of the protocol. This method applied the test directly to the original (recorded) curves and was analysed using the open-source spm1d code (version 0.4.8, www.spm1d.org) in Matlab (R2020a, The Mathworks Inc, Natick, USA). Normality was checked using an SPM Shapiro Wilk test before analysing

the NIRS data. The statistical calculations are described in Pataky [47]. A 5% risk was used for all statistical tests.

Results

As all the parameters studied, except muscle O_2 saturation, showed a significant interaction between the effects of the protocol (CWI vs. AMB) and the time of measurement, only the interactions are detailed in their case.

Cardio-Respiratory parameters

$\dot{V}O_2$, $\dot{V}E$, $\dot{V}CO_2$ and RER showed a significant time x protocol (CWI vs. AMB) interaction ($p < 0.001$ for all variables; $\dot{V}O_2 \eta_p^2 = 0.09$; $\dot{V}CO_2 \eta_p^2 = 0.18$; $\dot{V}E \eta_p^2 = 0.30$; RER $\eta_p^2 = 0.30$). The group mean values (\pm SD) are shown for each protocol in figure 2 together with the statistics related to the protocol effect. The time effect statistics are detailed (p -values and effect sizes) in Tables 1 and 2.

Time effect in each protocol

Only the CWI protocol showed a significant time effect, due to an overall increase in cardiorespiratory parameters during the R2_CWI condition compared to the R1 and R3 recovery periods.

During R2_CWI, $\dot{V}O_2$ at 30 sec was significantly higher than in R1 and R3 periods. $\dot{V}O_2$ then decreased up to 9 min before increasing again at 14 min. At 4 min $\dot{V}O_2$ was still higher than in R3 (up to 45 min), at 9 min not different from R1 and R3, and at 14 min significantly higher than in R3.

$\dot{V}CO_2$ clearly peaked at 30 sec, being significantly higher than at any other time point in R1, R2 and R3. Due to its subsequent decrease, at 4 min $\dot{V}CO_2$ was only higher than in R3 (up to 45 min). A slight increase in both $\dot{V}O_2$ and $\dot{V}CO_2$ was observed at 55 min in R3.

$\dot{V}E$ also peaked at 30 sec before decreasing progressively. At both 30 sec and 4 min $\dot{V}E$ was significantly higher than at any other time point in R1, R2 and R3. At 9 min, $\dot{V}E$ was not different from R1 and R3. At 14 min, due to a slight increase, $\dot{V}E$ was again higher than in R3.

RER also peaked at 30 sec before decreasing. At 30 sec RER was higher than at any other time point in R1, R2 and R3. At 4 min, RER was still higher than at 9 min and 14 min in R2, and at 20 min in R3. Compared to R1, RER was also higher at 9 min and 14 min in R2, and at 55 min in R3.




Figure 2: Influence of the protocol (CWI vs. AMB) on the time course of the relative oxygen uptake ($\dot{V}O_2$), relative carbon dioxide output ($\dot{V}CO_2$), ventilation ($\dot{V}E$) and respiratory exchange ratio (RER) during the 3 recovery periods (R1-R3) following a heating cycling exercise. Time was set at 0 at the beginning of R2. The black and grey lines correspond to the protocol with the R2 recovery in either cold water immersion (CWI) or at ambient room temperature (AMB). The statistical values of the time effect in each protocol are detailed in Tables 1 and 2. * $p < 0.05$, ** $p < 0.01$ between protocols, *** $p < 0.001$ between protocols.

Table 2: P-values (p) and effect size (Cohen's d value) for the significant time effect on $\dot{V}E$ and RER.

			R1	R2				R3			
			-7 min	30 sec	4 min	9min	14 min	20 min	30 min	45 min	55 min
$\dot{V}E$	R1	p		<0.001	<0.001						
		d		2.18	1.10						
	R2	30 sec	p		<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
			d		0.91	1.81	1.74	2.76	2.69	2.55	2.37
		4 min	p			<0.001	0.009	<0.001	0.015	<0.001	<0.001
			d			0,88	0.73	1,64	1.62	1.51	1.41
	14 min	p					0.011	0.004	0.027	0.011	
		d					1.03	1.03	0.91	0.82	
	R3	20 min	p								
			d								
		30 min	p								
			d								
45 min		p									
		d									
55 min	p										
d											
RER	R1	p		<0.001		0.001	<0.001	0.004			0.036
		d		1.84		1.07	1.29	0.93			1.14
	R2	30 sec	p		<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
			d		1.73	2.59	2.77	2.58	2.66	2.28	2.64
		4 min	p			0.006	0.004	0.021			
			d			0.86	1.03	0.86			
	9 min	p									
		d									
	14 min	p									
		d									
	R3	20 min	p								
			d								
30 min		p									
		d									
45 min		p									
		d									
55 min	p										
d											

The table only shows the statistically significant differences in ventilation ($\dot{V}E$) and respiratory exchange ratio (RER) between different measurement times in each protocol. The upper right (smoothed) corresponds to the CWI protocol, and the lower left (hatched) corresponds to the AMB protocol. R1, R2 and R3 correspond to the first, second and third recovery periods after a heating cycling exercise. The two protocols only differed in R2, which was performed in either cold water immersion (CWI) or at ambient room temperature (AMB). From lightest to darkest coloring: from no difference or small effect size, to moderate effect size, and large effect size for the darkest.

Hemodynamic parameters

A significant interaction time x protocol (CWI vs. AMB) was found for FABF and $\dot{Q}c$ ($p < 0.001$; FABF $\eta_p^2 = 0.06$; $\dot{Q}c$ $\eta_p^2 = 0.09$). The group mean values (\pm SD) are shown in figure 3 together with the statistics related to the protocol effect. The time effect statistics are detailed in Table 3.




Figure 3: Influence of the protocol (CWI vs. AMB) on the time course of the femoral artery blood flow and cardiac output during the 3 recovery periods (R1-R3) following a heating cycling exercise. Time was set at 0 at the beginning of R2. The black and grey lines correspond to the protocol with the R2 recovery in either cold water immersion (CWI) or at ambient room temperature (AMB). * $p < 0.05$, *** $p < 0.001$ between protocols.

Time effect in each protocol

Only the CWI protocol showed a significant time effect on FABF and on \dot{Q}_c :

During R2-CWI, FABF decreased sharply and remained significantly lower than in R1 throughout. During the subsequent rewarming period R3, FABF was not significantly different from R1 or R2, except at 30 min, which was significantly higher than at 14 min in R2.

\dot{Q}_c peaked at 30 sec, being significantly higher than at any other time point in R1, R2 and R3. It then decreased progressively until the beginning of R3 at 20 min, being below R1 and R2 up to 4 min. A slight decrease occurred during R3 at 40 min resulted in a difference with R1 and R2 at 30 sec.

Protocol Effect

As shown in Figure 3, the R2_CWI condition was characterised by lower FABF values than the R2_AMB condition (at 30 sec: $p < 0.001$, $d = 0.80$; 4 min: $p < 0.001$, $d = 0.95$; 9 min: $p < 0.001$, $d = 0.97$; 14 min: $p < 0.001$, $d = 1.06$) while \dot{Q}_c was only initially higher (at 30 sec: $p < 0.001$, $d = 0.84$ and 4 min: $p = 0.048$, $d = 0.55$). During the subsequent rewarming period R3, FABF was again significantly lower than in the AMB protocol at 45 min ($p = 0.049$, $d = 0.52$) and 55 min ($p = 0.017$, $d = 0.57$).

Table 3: P-values (p) and effect size (Cohen's d value) for the significant time effect on femoral artery blood flow and cardiac output.

			R1	R2				R3			
			-7 min	30 sec	4 min	9min	14 min	20 min	30 min	45 min	55 min
Femoral artery blood flow	R1	-7 min	p	<0.001	<0.001	<0.001	<0.001				
		d	0.91	0.99	0.92	1.15					
	R2	30 sec	p								
		d									
		4 min	p								
		d									
	R3	9 min	p								
		d									
		14 min	p						0.043		
		d							0.89		
R3	20 min	p									
	d										
	30 min	p									
	d										
R3	45 min	p									
	d										
	55 min	p									
	d										
CWI Protocol											
Cardiac Output	R1	-7 min	p	0.002				0.005		0.042	
		d	0.71				0.91		0.72		
	R2	30 sec	p		<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
		d		0.78	1.03	1.26	1.48	1.29	1.31	1.15	
		4 min	p					0.024			
		d					0.72				
	R3	9 min	p								
		d									
		14 min	p								
		d									
R3	20 min	p									
	d										
	30 min	p									
	d										
R3	45 min	p									
	d										
	55 min	p									
	d										
AMB Protocol											

The table only shows the statistically significant differences in the femoral artery blood flow and cardiac output between different measurement times in each protocol. The upper right (smoothed) corresponds to the CWI protocol, and the lower left (hatched) corresponds to the AMB protocol. R1, R2 and R3 correspond to the first, second and third recovery periods after a heating cycling exercise. The two protocols only differed in R2, which was performed in either cold water immersion (CWI) or at ambient room temperature (AMB). From lightest to darkest coloring: from no difference or small effect size, to moderate effect size, and large effect size for the darkest.

Gastrointestinal temperature

A significant interaction time x protocol (CWI vs. AMB) was found for T_{GI} ($p < 0.001$; $\eta_p^2 = 0.25$). The group mean values (\pm SD) are shown in figure 4 together with the statistics related to the protocol effect. The time effect statistics are detailed in Table 4.




Figure 4: Influence of the protocol (CWI vs. AMB) on the time course of stomach temperature during the 3 recovery periods (R1-R3) following a heating cycling exercise. Time was set at 0 at the beginning of R2. The black and grey lines correspond to the protocol with the R2 recovery in either cold water immersion (CWI) or at ambient room temperature (AMB).

*** $p < 0.001$ between protocols.

Time effect in each protocol

During R1, in both protocols, T_{GI} dropped during the second half of R1 (being lower at -10 and -7 min and thus 15 to 18 min after the end of the heating exercise). During the subsequent CWI and R3 periods, T_{GI} remained rather stable, decreasing only in the CWI protocol at 55 min in R3. During FI, T_{GI} changed in an opposite way in the two protocols: In the CWI protocol, it decreased, being significantly lower than at any other time point in R1, R2 and R3. In the AMB protocol, it was lower compared to the 10 first min of R1, but its value after 12 min of cycling was significantly higher than at 45 min and 55 min in R3.

Protocol effect

As shown in figure 4, T_{GI} varied similarly in both recovery protocols throughout the recovery period. However, the final cycling exercise was associated with significantly different T_{GI} changes: in the CWI protocol, T_{GI} decreased so that it was significantly lower than in the AMB protocol after 5 min ($p < 0.001$, $d = 2.05$) and 12 min ($p < 0.001$, $d = 2.73$) of cycling.

Table 4: P-value (p) and effect size (Cohen's d value) of the time effect on gastrointestinal temperature.

		R1					R2				R3				FI		
		-25 min	-20 min	-15 min	-10 min	-7 min	30 sec	4 min	9 min	14 min	20 min	30 min	45 min	55 min	5 min cycling	12 min cycling	
Gastrointestinal temperature	R1	-25 min p					<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	
		-25 min d					1.74	1.62	1.58	1.41	1.65	1.77	1.93	2.39	2.73	3.82	3.48
		-20 min p			0.044	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
		-20 min d			0.71	1.77	1.68	1.63	1.47	1.69	1.81	1.94	2.37	2.68	3.74	3.44	
		-15 min p			<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
	-15 min d			0.97	2.23	2.06	2.03	1.87	2.13	2.27	2.42	2.90	3.23	4.28	3.91		
	-10 min p	0.047		0.017			0.013				0.041	0.007	<0.001	<0.001	<0.001	<0.001	
	-10 min d	0.69		0.88			0.80				0.81	0.93	1.32	1.63	2.75	2.53	
	-7 min p	<0.001	<0.001	<0.001	<0.001									<0.001	0.021	<0.001	
	-7 min d	1.61	1.47	2.21	1.27									1.30	2.82	2.47	
	30 sec p	<0.001	<0.001	<0.001	0.002										<0.001	<0.001	
	30 sec d	1.58	1.44	2.17	1.21										2.34	2.09	
	4 min p	<0.001	<0.001	<0.001	0.016									0.013	<0.001	<0.001	
	4 min d	1.45	1.31	1.95	1.02									1.15	2.58	2.30	
	9 min p	<0.001	<0.001	<0.001										<0.001	<0.001	<0.001	
	9 min d	1.38	1.24	1.86										1.52	2.95	2.62	
14 min p	<0.001	<0.001	<0.001	0.011									0.007	<0.001	<0.001		
14 min d	1.52	1.38	2.10	1.12									1.34	2.84	2.51		
20 min p	<0.001	<0.001	<0.001	0.004									0.025	<0.001	<0.001		
20 min d	1.60	1.45	2.24	1.23									1.26	2.81	2.47		
30 min p	<0.001	<0.001	<0.001	<0.001										<0.001	<0.001		
30 min d	1.75	1.60	2.45	1.46										2.70	2.37		
45 min p	<0.001	<0.001	<0.001	<0.001										<0.001	<0.001		
45 min d	1.96	1.81	2.80	1.79										2.23	1.94		
55 min p	<0.001	<0.001	<0.001	<0.001										<0.001	<0.001		
55 min d	1.95	1.81	2.75	1.78										1.76	1.52		
5 min cycling p	<0.001	<0.001	<0.001	<0.001													
5 min cycling d	1.62	1.49	2.24	1.32													
12 min cycling p	<0.001	<0.001	<0.001									0.018	0.007				
12 min cycling d	0.91	0.79	1.17									1.90	1.85				

CWI protocol

AMB Protocol

The table only shows the statistically significant differences in the gastrointestinal temperature between different measurement times in each protocol. The upper right (smoothed) corresponds to the CWI protocol, and the lower left (hatched) corresponds to the AMB protocol. R1, R2 and R3 correspond to the first, second and third recovery periods after a heating cycling exercise. The two protocols only differed in R2, which was performed in either cold water immersion (CWI) or at ambient room temperature (AMB). From lightest to darkest colouring: from no difference or small effect size, to moderate effect size, and large effect size for the darkest.

Muscle O₂ saturation

As the statistical test is different, the results are presented in a different way. We will only discuss the difference between CWI and AMB measurements over time.

During R2, MSO₂VM et MSO₂GM did not differ significantly between protocols. During R3 in the CWI protocol, MSO₂VM and MSO₂GM decreased progressively after 22 min (7 min after the cold bath) (Figure 4A, 4B). Compared to the AMB protocol (Figure 4C), in the CWI protocol MSO₂VM was significantly lower from 31 min (16 min after the cold bath) to the end of R3, and MSO₂GM from 38 min (23 min after the cold bath) to the end of R3.




Figure 5: Muscle saturation in O₂ of vastus medialis (A) and gastrocnemius medialis (B) and t-test results including the raw SPM{t} with the random field theory computed critical t threshold (C). Muscle O₂ saturation in the CWI protocol is significantly lower than in the AMB protocol when its SPM{t} is higher than the significance threshold (horizontal cut line). CWI: Cold Water Immersion protocol; AMB: Ambient protocol.

Discussion

The aim of the study was to define the time course of central or local heat production and vasoconstriction during the subsequent CWI and rewarming period after a heating cycling exercise. The observation of an increase in $\dot{V}O_2$ with associated peripheral vasoconstriction during CWI, followed by a decrease in MSO₂ (indicating increased O₂ consumption in lower limb muscles) during

rewarming, supports our hypothesis that different thermoregulatory mechanisms may contribute to the increase in energy expenditure during and after CWI. Specifically, we distinguished non-thermoregulatory responses during early CWI from different thermoregulatory responses during and after CWI. The physiological thermoregulatory changes over time demonstrated that the rewarming period as an integral part of the recovery protocol using CWI could potentially explain some of the contradictions found in the literature. As most studies in the literature have examined each of these variables separately, we will discuss them in turn before discussing their potential relationships.

With regard to the changes in body temperature during CWI, most studies have recorded rectal temperature [35–37,41] and, more recently, gastrointestinal temperature using a pill [48,49]. Among them, Crampton et al. [48] showed a decrease in intestinal temperature from the beginning to the end of a 30 min CWI in 15 °C water. However, the intestinal temperature of their participants had already decreased during the first 5 min of the CWI, which differs from the current results where the gastrointestinal temperature remained stable. However, their participants' intestinal temperature at the beginning of the CWI was around 38 °C (above the ideal temperature of the human body), whereas ours was 37.21 °C. Another explanatory factor may be the different location of the temperature measurement: In Crampton et al. [48] study, the pill was swallowed 3 hours before immersion, whereas our participants took it 1 hour before. As a result, we measured temperature changes higher up in the body than they did. A third and most likely reason is that their participants were immersed in the bath 5 min after the end of the heating exercise. In the current study, immersion did not take place until 25 min after the heating exercise. During this intermediate (R1) recovery period at ambient room temperature, a decrease in gastrointestinal temperature began between 10 to 15 min later. Under ambient conditions, Peiffer et al. [36] also reported a decrease in rectal temperature 5 to 10 min later. Therefore, it is possible that the decrease in intestinal temperature reported by Crampton et al. [48] during immersion was due to the effect of CWI combined with natural recovery post-exercise. Similarly, these authors also reported a decrease in intestinal temperature during exercise after leaving the cold bath. In a more recent study [49], intestinal temperature was measured during a 10 min immersion protocol at 8, 15 and 20 °C without prior exercise. Consistent with our results, they found no significant difference in intestinal temperature between the beginning and end of the immersion, regardless of water temperature. A maintained gastrointestinal temperature during CWI can be attributed to two phenomena: A vasoconstrictive effect that reduces convective heat loss from the immersed limb by lowering their muscle and skin temperature [48–50] and/or a production of heat to counteract cooling.

In our study, we observed a decrease in FABF during CWI, which is supported by the findings of Vaile et al. [37] at the end of CWI. However, our FABF values were higher than theirs, possibly due to the use of different measurement techniques (venous occlusion plethysmography vs. doppler

ultrasound device). During whole-body cold exposure, a significant decrease in arterial blood velocity has been reported due to blood redistribution to the deep circulation [5,37]. This may be due to an increase in capillary vasoconstriction and hydrostatic pressure due to CWI. Their combination with the increase in heart rate is expected to explain the increase in \dot{Q}_c at 30 sec and 4 min in the CWI protocol [25,26]. These results suggest that during the CWI the muscles cool down, reducing the exchange with the blood and maintaining a stable deep body temperature. This is supported by the large decrease in gastrointestinal temperature when cycling resumed. This is due to the cold being transferred to the deep body by increased blood flow from the cold muscles.

Several studies have reported an increase in $\dot{V}O_2$ during CWI. This response is initially neural in origin and is referred to as the cold shock response. In this line, we observed a 35 % increase in $\dot{V}O_2$ at 30 sec during the R2_CWI period. However, it contrasted with the 400 % increase reported during the first minute of immersion by Mekjavic & Bligh [14]. One reason for this could be that their study involved immersion up to the neck, whereas in our study it was only involved up to the iliac crest. Indeed, Ran et al. [51] have shown an increase in the neuronal response in the spinal cord as the number of cold receptors stimulated increased with decreasing temperature. We can therefore hypothesise that an increase in the number of activated receptors due to an increase in the area of immersion will also lead to an increase in neuronal activity, and therefore the intensity of the cold shock response. In addition, our participants first placed their calves in the water before sitting on the stool, which may have attenuated the reflex increase of $\dot{V}O_2$ [25]. Staged immersion has been shown to reduce the cold shock response as evidenced by a smaller increase in $\dot{V}E$ compared to non-staged immersion [52]. As observed in our study, cold shock responses are reported to reach a peak within 30 sec of immersion and adapt over the first 3 min of immersion in most individuals [25]. Similarly, the increase in RER observed 30 sec after entering the cold bath was attributed to the reflex increase in $\dot{V}E$, since the induced increase in pulmonary air exchange should lead to an increase in expired $\dot{V}CO_2$ associated with a decrease of alveolar CO_2 [27]. This agrees with the hypocapnia reported when entering a cold bath [25]. Therefore, it is suggested that the current increase in RER was due to non-metabolic phenomena.

In a second phase, to maintain its deep body temperature in CWI and after it, the body produces heat through two primary mechanisms: Non-shivering thermogenesis (NST) and shivering thermogenesis (ST), both of which consume O_2 [21,24]. In our study we found a 25 % increase in $\dot{V}O_2$ after 14 min of immersion at 10 °C vs. 130 % reported after 15 min at 14 °C by Šrámek [15]. In the latter study, immersion was again up to the neck. It appears that $\dot{V}O_2$ increases as the surface area of the body exposed to cold increases, probably due to the increased energy flow exchanged when more of the body is in contact with the water in addition to the neural response. This requires the body to produce more heat, resulting in increased O_2 consumption. In addition, Eyolfson et al. [13]

observed a gradual increase in $\dot{V}O_2$ during immersion in colder water (from 20 °C to 8 °C) over 15 min. The plateau in $\dot{V}O_2$ observed in our study during the entire CWI is probably due to ST. Indeed, it can increase $\dot{V}O_2$ up to 40 % of $\dot{V}O_{2max}$ [17]. Nevertheless, as indicated by the stability of MSO_2GM and MSO_2VM , this metabolic phenomenon is unlikely to be due to ST or NST in the immersed lower limbs. Moreover, it has been shown that ST suppression during a cold bath results in a decrease in $\dot{V}O_2$ compared to a protocol without suppression [11]. Since brown fat is mainly found in different parts of the trunk [53] and inhibition of shivering would only be effective at a core temperature below 31 °C [54], ST and NST could have occurred in the trunk and upper limbs, but unfortunately we did not measure them. At this stage, it is concluded that the increase in $\dot{V}O_2$ during CWI, first by neural phenomena and then by metabolic phenomena, with a return to the same value as in the AMB protocol during R3, suggests that different mechanisms were involved during CWI, but also after during the rewarming period.

Regarding muscle O_2 consumption, only two studies have examined MSO_2 in the immersed muscles during the CWI period and one during the rewarming period. Therefore, it is difficult to draw conclusions based on the comparison with their results. However, our results during the CWI period are consistent with some findings in the literature (e.g. Roberts et al [55]), which reported no change in MSO_2 in the vastus lateralis during CWI up to the umbilical level at 10 °C for 10 min. In contrast, an increase in MSO_2 in the vastus lateralis has been reported during leg immersion at 10 °C for 15 min (Ihsan et al. [56]). This discrepancy is attributed to differences in the test protocol. In the latter study, only one leg was immersed up to the level of the gluteal fold, covered with a polyethylene bag to avoid contact between the NIRS and the water, and the participants were instructed to distribute their weight evenly while standing. Given all these differences in protocols, it seems difficult to compare our results with theirs. If the result of Roberts et al. [55] are closer to ours during CWI, they reported no change in MSO_2 in the vastus lateralis during the rewarming period, unlike our. However, their participants walked to another experimentation room to perform an isometric maximum voluntary contraction. This may explain why Roberts et al. [55] did not observe a decrease in MSO_2 during the rewarming period, due to the increase in muscle temperature caused by the muscle activity through increased heat generation and blood flow. However, it is important to note that one of the limitations of this measurement is the change in muscle temperature during and after immersion. Indeed, it has been shown that the NIRS signal can be used to measure the in vivo temperature of water-containing tissue [57]. Nevertheless, several studies have investigated the evolution of muscle temperature at depths of 1,2,3 cm [41,58]. In 10 min their immersion protocols at 8 °C, muscle temperature at 2 cm has already decreased by 3 to 5 °C 1 min after exiting the cold bath. We can therefore hypothesise that during 15 min immersion at 10 °C, muscle temperature at 2 cm depth starts to decrease during immersion. The NIRS signal is measured at a muscle depth

corresponding to 50-60 % of transmitter -receiver distance [59]. The farthest receiver on our NIRS was 25 mm from the transmitter, so we measured oxygenation at a depth of about 12 mm. As the current NIRS signal remained extremely stable throughout the immersion and decreased regularly 7 min after exiting, we believe that the observed variation can't be explained by the muscle temperature alone. Nevertheless, this remains a limitation of the current paper because this part of the discussion is partially based on the hypothesis that muscle temperature changes during CWI. The current decrease of MSO_2VM and MSO_2GM during the passive rewarming period could be rather explained by ST or the NST of the muscle (NST-M) since our participants did not perform any muscular activity after CWI but oxygen seems to be consumed. Unfortunately, neither EMG electrodes nor accelerometer were placed on the vastus medialis and gastrocnemius medialis muscles, which would have allowed us to differentiate between ST and NST-M [25]. At the present stage it can only be said that NST-M and/or ST of the immersed muscles may have occurred more than 10 min after leaving the cold bath.

Combining all the results after the heating exercise, i.e. the increase in $\dot{V}O_2$ during the cold water immersion with peripheral vasoconstriction, the decrease in MSO_2 during the subsequent rewarming during R3, and the decrease in gastrointestinal temperature during the final cycling period, reveals a continuum of different responses during CWI and rewarming. The first phase after the onset of immersion, the cold shock response, is thought to be neural rather than thermoregulatory in origin. This phase, which lasts approximately 3 min, was observed in our study with reflex vasoconstriction and increased $\dot{V}O_2$, $\dot{V}CO_2$, RER and \dot{Q}_c in agreement with the literature. The first thermoregulatory responses then occur, with $\dot{V}O_2$ stabilising at a higher value than in the AMB condition and RER decreasing. These phenomena are well described in the literature. The higher $\dot{V}O_2$ indicates an increase in energy expenditure of the non-submerged part of the body, probably related to ST or NST. Indeed, this does not seem to be due to an increase in metabolic activity of the lower limb muscles as shown by the stagnation of MSO_2VM and MSO_2GM , but rather of the trunk and/or upper limb muscles. There seems to be a transitional period when the participant comes out of the cold bath. Indeed, $\dot{V}O_2$ has already returned to the same level as in AMB 5 min after the end of immersion, while peripheral vasodilation also seems to relax, as shown by the return of FABF to a lower but not significantly different level than in AMB. However, during this period of stagnation, MSO_2GM and MSO_2VM start to decrease 7 min after the end of the cold bath, but do not differ significantly from the AMB protocol until 31 min, i.e. 16 min after the end of immersion. At this point, a new phase in the fight against cold seems to emerge. Indeed, the gastrointestinal temperature drops after 30 min. This well-known mechanism is called afterdrop. However, it is interesting to note the stability of gas exchange values, but a slight relapse in FABF may indicate renewed peripheral vasoconstriction. The decrease in MSO_2 tends to be explained by an increase in thermogenesis at the

muscle level, although the effect of temperature on the NIRS signal may be a limitation. These different phases are schematised in Figure 6, in relation to the evolution of the different measured variables useful for our discussion. It is also important to note that thermogenesis in the lower limb muscles did not completely warm them. Indeed, the decrease in gastrointestinal temperature on resumption of cycling could be due to an increase in convection link to the increase in blood flow to the legs, which cools on contact with the cold muscles and thus cool the deep body. The main limitation of this work is the lack of measurements to differentiate between ST, NST-B and NST-M. This would have allowed us to deepen our discussion, particularly by expressing the parts of the body where they occur during and after immersion. The second factor to be investigated in future research is the evolution of muscle temperature during the rewarming period.




Figure 6: schematic evolution of oxygen consumption ($\dot{V}O_2$), gastrointestinal temperature (T_{GI}), muscle O_2 saturation (MSO_2) and arterial blood flow (FABF) during the CWI and the rewarming.

Conclusion and perspective

This study represents one of the first attempts to combine measurements of stomach temperature, cardiac and arterial output with systemic and local oxygen consumption in immersed muscles during the immersion and rewarming periods, compared to passive recovery in ambient conditions, post cycling exercise. It's important to note that the participants were immersed up to the waist, and the responses might have been different if they hadn't exercised beforehand or if they had been immersed up to the neck. Our results demonstrated the existence of different responses during and after cold immersion, which can be distinguished by gastric temperature kinetics, haemodynamic and metabolic variables. Our main result was that thermogenesis during CWI seemed to take place

initially in the non-immersed zones, before shifting to the immersed limbs during the rewarming period. Nevertheless, at 55 min post-CWI, thermogenesis in the lower limbs was still insufficient, given the drop in deep body temperature on resumption of physical activity. This underlines the importance of recording core, muscle and skin temperature changes in parallel, considering immersion and rewarming as a continuum rather than as separate events. Our results also highlight the importance of studying the rewarming state after cold immersion in order to understand the whole effects of this last.

Additional information

DATA accessibility

All the data are available to the following link:

<https://dataverse.harvard.edu/privateurl.xhtml?token=d5a43e33-8082-48e5-a3cc-cfa68acd7696>

If the article is accepted, a permanent link will be created.

Acknowledgement

First of all, we would like to thank the company Cryocontrol for the loan of the equipment and the technical assistance during the experiment. We would also like to thank Robin Macchi for his invaluable help with the statistical processing and Pr Jerome Laurin for checking the manuscript.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

Funding

Experimental material was bought by Aix-Marseille University. Mr Dorian Giraud was supported by a PhD grant. Other authors were paid by the University of Aix-Marseille.

References

- [1] Pointon M, Duffield R. Cold Water Immersion Recovery after Simulated Collision Sport Exercise. *Med Sci Sports Exerc.* 2012;44:206–216. doi: 10.1249/MSS.0b013e31822b0977.
- [2] Vieira A, Siqueira A, Ferreira-Junior J, et al. The Effect of Water Temperature during Cold-Water Immersion on Recovery from Exercise-Induced Muscle Damage. *Int J Sports Med.* 2016;37:937–943. doi: 10.1055/s-0042-111438.
- [3] De Paula F, Escobar K, Ottone V, et al. Post-exercise cold-water immersion improves the performance in a subsequent 5-km running trial. *Temperature.* 2018;5:359–370. doi: 10.1080/23328940.2018.1495023.
- [4] Allan R, Akin B, Sinclair J, et al. Athlete, coach and practitioner knowledge and perceptions of post-exercise cold-water immersion for recovery: a qualitative and quantitative exploration. *Sport Sci Health.* 2022;18:699–713. doi: 10.1007/s11332-021-00839-3.


- [5] Ihsan M, Watson G, Abbiss CR. What are the Physiological Mechanisms for Post-Exercise Cold Water Immersion in the Recovery from Prolonged Endurance and Intermittent Exercise? *Sports Med.* 2016;46:1095–1109. doi: 10.1007/s40279-016-0483-3.
- [6] Tipton MJ, Collier N, Massey H, et al. Cold water immersion: kill or cure?: Cold water immersion: kill or cure? *Exp Physiol.* 2017;102:1335–1355. doi: 10.1113/EP086283.
- [7] Lee DT, Toner MM, McArdle WD, et al. Thermal and metabolic responses to cold-water immersion at knee, hip, and shoulder levels. *J Appl Physiol.* 1997;82:1523–1530. doi: 10.1152/jappl.1997.82.5.1523.
- [8] Landsberg L. Core temperature: a forgotten variable in energy expenditure and obesity? *Obes Rev.* 2012;13:97–104. doi: 10.1111/j.1467-789X.2012.01040.x.
- [9] Romet TT. Mechanism of afterdrop after cold water immersion. *J Appl Physiol.* 1988;65:1535–1538. doi: 10.1152/jappl.1988.65.4.1535.
- [10] Savard GK, Cooper KE, Veale WL, et al. Peripheral blood flow during rewarming from mild hypothermia in humans. *J Appl Physiol.* 1985;58:4–13. doi: 10.1152/jappl.1985.58.1.4.
- [11] Giesbrecht GG, Goheen MSL, Johnston CE, et al. Inhibition of shivering increases core temperature afterdrop and attenuates rewarming in hypothermic humans. *J Appl Physiol.* 1997;83:1630–1634. doi: 10.1152/jappl.1997.83.5.1630.
- [12] Williams AB. Rewarming of healthy volunteers after induced mild hypothermia: a healthy volunteer study. *Emerg Med J.* 2005;22:182–184. doi: 10.1136/emj.2003.007963.
- [13] Eyolfson DA, Tikuisis P, Xu X, et al. Measurement and prediction of peak shivering intensity in humans. *Eur J Appl Physiol.* 2001;84:100–106. doi: 10.1007/s004210000329.
- [14] Mekjavic IB, Bligh J. The increased oxygen uptake upon immersion: The raised external pressure could be a causative factor. *Eur J Appl Physiol.* 1989;58:556–562. doi: 10.1007/BF02330712.
- [15] Šrámek P, Šimečková M, Janský L, et al. Human physiological responses to immersion into water of different temperatures. *Eur J Appl Physiol.* 2000;81:436–442.
- [16] Hemingway A. Shivering. *Physiol Rev.* 1963;43:397–422. doi: 10.1152/physrev.1963.43.3.397.
- [17] Haman F. Metabolic requirements of shivering humans. *Front Biosci.* 2010;S2:1155–1168. doi: 10.2741/s124.
- [18] Haman F, Blondin DP. Shivering thermogenesis in humans: Origin, contribution and metabolic requirement. *Temperature.* 2017;4:217–226. doi: 10.1080/23328940.2017.1328999.
- [19] Blondin DP, Haman F. Shivering and nonshivering thermogenesis in skeletal muscles. *Handb Clin Neurol* [Internet]. Elsevier; 2018 [cited 2022 Oct 10]. p. 153–173. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780444639127000102> doi: 10.1016/B978-0-444-63912-7.00010-2.
- [20] Autry JM, Thomas DD, Espinoza-Fonseca LM. Sarcolipin Promotes Uncoupling of the SERCA Ca²⁺ Pump by Inducing a Structural Rearrangement in the Energy-Transduction Domain. *Biochemistry.* 2016;55:6083–6086. doi: 10.1021/acs.biochem.6b00728.

- [21] Townsend LK, Wang D, Wright DC, et al. Skeletal muscle, not adipose tissue, mediates cold-induced metabolic benefits. *Nat Metab* [Internet]. 2023 [cited 2023 Jul 11]; Available from: <https://www.nature.com/articles/s42255-023-00837-4> doi: 10.1038/s42255-023-00837-4.
- [22] Lagarde D. Rôle des flux de lactate dans le métabolisme des tissus adipeux beiges et bruns [PhD Thesis]. Toulouse 3; 2020.
- [23] Fedorenko A, Lishko PV, Kirichok Y. Mechanism of Fatty-Acid-Dependent UCP1 Uncoupling in Brown Fat Mitochondria. *Cell*. 2012;151:400–413. doi: 10.1016/j.cell.2012.09.010.
- [24] Holets HM, Kerna NA, Chawla S, et al. Brown Adipose Tissue: A Comprehensive Review. 2022;26–40.
- [25] Datta A, Tipton M. Respiratory responses to cold water immersion: neural pathways, interactions, and clinical consequences awake and asleep. *J Appl Physiol*. 2006;100:2057–2064. doi: 10.1152/jappphysiol.01201.2005.
- [26] Tipton MJ. The Initial Responses to Cold-Water Immersion in Man. *Clin Sci*. 1989;77:581–588. doi: 10.1042/cs0770581.
- [27] Keatinge WR, Evans M. The respiratory and cardiovascular response to immersion in cold and warm water. *Q J Exp Physiol Cogn Med Sci*. 1961;46:83–94. doi: 10.1113/expphysiol.1961.sp001519.
- [28] Keatinge WR, McIlroy MB, Goldfien A. Cardiovascular responses to ice-cold showers. *J Appl Physiol*. 1964;19:1145–1150. doi: 10.1152/jappl.1964.19.6.1145.
- [29] Peier AM, Moqrich A, Hergarden AC, et al. A TRP Channel that Senses Cold Stimuli and Menthol. *Cell*. 2002;108:705–715. doi: 10.1016/S0092-8674(02)00652-9.
- [30] Morrison SF, Nakamura K. Central Mechanisms for Thermoregulation. *Annu Rev Physiol*. 2019;81:285–308. doi: 10.1146/annurev-physiol-020518-114546.
- [31] McLellan T. Ventilatory and Plasma Lactate Response with Different Exercise Protocols: A Comparison of Methods. *Int J Sports Med*. 1985;06:30–35. doi: 10.1055/s-2008-1025809.
- [32] Hays A, Nicol C, Bertin D, et al. Physiological and Mechanical Indices Serving the New Cross-Country Olympic Mountain Bike Performance. *Int J Sports Physiol Perform*. 2021;16:1008–1013. doi: 10.1123/ijsp.2020-0319.
- [33] Brughelli M, Van Leemputte M. Reliability of Power Output During Eccentric Sprint Cycling. *J Strength Cond Res*. 2013;27:76–82. doi: 10.1519/JSC.0b013e31824f2055.
- [34] Davranche K, Giraud D, Hays A, et al. High-intensity physical activity enhances cognitive decision processes [Internet]. *Neuroscience*; 2023 [cited 2023 Oct 16]. Available from: <http://biorxiv.org/lookup/doi/10.1101/2023.02.14.528466> doi: 10.1101/2023.02.14.528466.
- [35] Peiffer JJ, Abbiss CR, Nosaka K, et al. Effect of cold water immersion after exercise in the heat on muscle function, body temperatures, and vessel diameter. *J Sci Med Sport*. 2009;12:91–96. doi: 10.1016/j.jsams.2007.10.011.


- [36] Peiffer JJ, Abbiss CR, Watson G, et al. Effect of cold-water immersion duration on body temperature and muscle function. *J Sports Sci.* 2009;27:987–993. doi: 10.1080/02640410903207424.
- [37] Vaile J, O’Hagan C, Stefanovic B, et al. Effect of cold water immersion on repeated cycling performance and limb blood flow. *Br J Sports Med.* 2011;45:825–829. doi: 10.1136/bjsm.2009.067272.
- [38] Dong L, Wang J, Jiang C. Validation of the use of foreign gas rebreathing method for non-invasive determination of cardiac output in heart disease patients. *J Zhejiang Univ Sci B.* 2005;6:1157–1162. doi: 10.1631/jzus.2005.B1157.
- [39] Jones B, Hesford CM, Cooper CE. The Use of Portable NIRS to Measure Muscle Oxygenation and Haemodynamics During a Repeated Sprint Running Test. In: Van Huffel S, Naulaers G, Caicedo A, et al., editors. *Oxyg Transp Tissue XXXV* [Internet]. New York, NY: Springer New York; 2013 [cited 2022 Jul 8]. p. 185–191. Available from: http://link.springer.com/10.1007/978-1-4614-7411-1_26 doi: 10.1007/978-1-4614-7411-1_26.
- [40] Wang B, Xu G, Tian Q, et al. Differences between the vastus lateralis and gastrocnemius lateralis in the assessment ability of breakpoints of muscle oxygenation for aerobic capacity indices during an incremental cycling exercise. 2012;8.
- [41] Mawhinney C, Jones H, Joo CH, et al. Influence of cold-water immersion on limb and cutaneous blood flow after exercise. *Med Sci Sports Exerc.* 2013;45:2277–2285. doi: 10.1249/MSS.0b013e31829d8e2e.
- [42] Bates D, Maechler M, Bolker B, et al. lme4: Linear mixed-effects models using “Eigen” and S4 classes. *R Package Version.* 2020;1:15.
- [43] Kuznetsova A, Brockhoff PB, Christensen RH. lmerTest package: tests in linear mixed effects models. *J Stat Softw.* 2017;82:1–26.
- [44] Macchi R, Santuz A, Hays A, et al. Sex influence on muscle synergies in a ballistic force-velocity test during the delayed recovery phase after a graded endurance run. *Heliyon.* 2022;8:e09573. doi: 10.1016/j.heliyon.2022.e09573.
- [45] Hopkins WG, Marshall SW, Batterham AM, et al. Progressive Statistics for Studies in Sports Medicine and Exercise Science. *Med Sci Sports Exerc.* 2009;41:3–12. doi: 10.1249/MSS.0b013e31818cb278.
- [46] Friston K, Ashburner J, Kiebel S, et al., editors. *Statistical Parametric Mapping* [Internet]. London: Academic Press; 2007. Available from: <https://www.sciencedirect.com/science/article/pii/B9780123725608500000> doi: <https://doi.org/10.1016/B978-012372560-8/50000-0>.
- [47] Pataky TC. One-dimensional statistical parametric mapping in Python. *Comput Methods Biomech Biomed Engin.* 2012;15:295–301. doi: 10.1080/10255842.2010.527837.
- [48] Crampton D, Egaña M, Donne B, et al. Including arm exercise during a cold water immersion recovery better assists restoration of sprint cycling performance: Active immersion recovery for sprint cycling. *Scand J Med Sci Sports.* 2014;24:e290–e298. doi: 10.1111/sms.12169.

- [49] Mawhinney C, Heinonen I, Low DA, et al. Changes in quadriceps femoris muscle perfusion following different degrees of cold-water immersion. *J Appl Physiol*. 2020;128:1392–1401. doi: 10.1152/jappphysiol.00833.2019.
- [50] Otte JW, Merrick MA, Ingersoll CD, et al. Subcutaneous adipose tissue thickness alters cooling time during cryotherapy. *Arch Phys Med Rehabil*. 2002;83:1501–1505. doi: 10.1053/apmr.2002.34833.
- [51] Ran C, Hoon MA, Chen X. The coding of cutaneous temperature in the spinal cord. *Nat Neurosci*. 2016;19:1201–1209. doi: 10.1038/nn.4350.
- [52] Hayward JS, French CD. Hyperventilation response to cold water immersion: reduction by staged entry. *Aviat Space Environ Med*. 1989;60:1163–1165.
- [53] Sacks H, Symonds ME. Anatomical Locations of Human Brown Adipose Tissue. *Diabetes*. 2013;62:1783–1790. doi: 10.2337/db12-1430.
- [54] Parsons KC. Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort, and performance. 2nd ed. London: Taylor & Francis; 2006.
- [55] Roberts LA, Muthalib M, Stanley J, et al. Effects of cold water immersion and active recovery on hemodynamics and recovery of muscle strength following resistance exercise. *Am J Physiol-Regul Integr Comp Physiol*. 2015;309:R389–R398. doi: 10.1152/ajpregu.00151.2015.
- [56] Ihsan M, Watson G, Lipski M, et al. Influence of Postexercise Cooling on Muscle Oxygenation and Blood Volume Changes. *Med Sci Sports Exerc*. 2013;45:876–882. doi: 10.1249/MSS.0b013e31827e13a2.
- [57] Raiko J, Koskensalo K, Sainio T. Imaging-based internal body temperature measurements: The journal *Temperature* toolbox. *Temperature*. 2020;7:363–388. doi: 10.1080/23328940.2020.1769006.
- [58] Mawhinney C, Jones H, Low DA, et al. Influence of cold-water immersion on limb blood flow after resistance exercise. *Eur J Sport Sci*. 2017;17:519–529. doi: 10.1080/17461391.2017.1279222.
- [59] Bhambhani YN. Muscle Oxygenation Trends During Dynamic Exercise Measured by Near Infrared Spectroscopy. *Can J Appl Physiol*. 2004;29:504–523. doi: 10.1139/h04-033.


[[Figure 1]]




[[Figure 2]]




[[Figure 3]]




[[Figure 4]]



[[Figure 5]]



[[Figure 6]]



Methods

Twenty-one male volunteers (mean(SD); age: 24(5) years, body mass: 68(8) kg; height: 177 (6) cm; $\dot{V}O_{2max}$: 55(6) ml.kg⁻¹.min⁻¹, maximal aerobic power: 300(50) W, body fat percentage: 14.7(3)%) who regularly participated in endurance sports such as cycling or trail running, were recruited for this study. All participants were fully informed of the potential risks and requirements of the study and provided written informed consent. The study was approved by the national ethical committee for STAPS (Science and Technique of Physical activity and Sport) Research (number IRB00012476-2022-21-04-176). To be eligible, participants were required to have completed an incremental cycling exercise test providing data on their power at the first and second ventilatory thresholds (P_{VT1} and P_{VT2} , respectively), maximal aerobic power and $\dot{V}O_{2max}$. Any participant who failed to complete the test protocol or sustained an injury during the process was excluded from the study.

Incremental exercise test

The incremental protocol consisted of a 2 min warm-up at 100 W, followed by increments of 30 W every 2 min. The first ventilatory threshold (VT1) was defined as the time at which an initial deviation from expiratory volume ($\dot{V}E$) linearity was observed, and when a systematic increase in the ventilatory equivalent for O₂ ($\dot{V}E/\dot{V}O_2$) and the fraction of expired O₂ first appeared. The second ventilatory threshold (VT2) was defined based on a secondary increase in $\dot{V}E$ and $\dot{V}E/\dot{V}O_2$, and a marked increase in the ventilatory equivalent for CO₂ ($\dot{V}E/\dot{V}CO_2$) combined with a decrease in the fraction of expired CO₂ (McLellan, 1985). Participants were required to abstain from alcohol (48 h), caffeine, tea or guarana shot (24 h) and strenuous exercise (24 h) before and during the experiment. Additionally, participants were requested to arrive well-hydrated.

Experimental design

As shown in Figure 1, both protocols session started with a 5 min warm-up cycle exercise (WU), followed by a 48 min heating cycling exercise (H), 3 recovery periods (R1-R3), and a final cycling exercise (FI). The protocols only differed in R2. More precisely, WU started at 80 % of the power at VT1, followed by 4x30 s at the power at VT1 plus 90 % of the difference between the power at VT1 and the power at VT2, interspaced by 1 min passive recovery. It ended with 2 all-out isokinetic sprints at a mechanical cadence of 120 rpm. The 48 min H exercise consisted of 8 sets of 5 min at a fixed cycling workload (90 % of the power reach at VT1) with 1 min recovery at 75 W. Cognitive tasks were performed during sets 1, 2, 4, 5, 7 and 8. These data were used in another study investigating cognitive decision making before, during and after exercise (Davranche et al., 2023).

Measurements

Upon arrival, participants were asked to swallow a pill (e-Celsius Performance, BodyCap, France) which was connected to their monitor via Bluetooth to record their stomach temperature (T_s) every 30 s throughout the experiment. For ease of reading, we have set $T=0$ as the start of the R2(CWI and AMB) period. Gas exchange measurements were obtained using breath-by-breath acquisition for the last 10 seconds of each intensity and the subsequent 20 seconds during recovery. Collection of data began from the end of the first sprint to the end of the last sprint while in the heating period and 7 minutes before cold water immersion. We collected temperature data from the start of the warm-up and continued throughout the heating period until the end of the experimental session.

Data Management

Stomach temperature data were averaged during the minute surrounding the end of each sprint (from $n-30$ to $n+30$). Additionally, stomach temperature was averaged one minute before the commencement of the warm-up. After the heating phase, T_s data were averaged every five minutes for a period of one minute until "- 7 min" data were obtained, which correspond with the same data presented in the main article. For $\dot{V}O_2$ data, the average of the three highest data points (excluding the maximum) was calculated at the end of each sprint. Subsequently, the "post-heating" data from the main article were retrieved.

Result

$\dot{V}O_2$

The $\dot{V}O_2$ data are shown in Figure 1.




Figure 7 : Time evolution of relative oxygen uptake ($\dot{V}O_2$) during the heating phase and the recovery phase (R1). The dark curve includes the R2_CWI condition and the clear curve the R2_AMB condition. The dotted lines represent the evolution of each individual in each condition.

$\dot{V}CO_2$

The $\dot{V}CO_2$ data are shown in Figure 2.




Figure 8: Time evolution of relative oxygen uptake ($\dot{V}CO_2$) during the heating phase and the recovery phase (R1). The dark curve includes the R2_CWI condition and the clear curve the R2_AMB condition. The dotted lines represent the evolution of each individual in each condition.

Temp

The stomach temperature data are shown in Figure 3. Large standard deviations are noticeable for this variable due to big temperature drops, as can be seen from the individual results. The permissible drinking of participants during the experiment caused these changes. The measurements were taken at stomach level, and any instance of drinking caused a temporary temperature decrease that later rose again to its original level. However, it is observed that in R1, as soon as the participants were forbidden to drink, the temperature stabilised with smaller standard deviations

and gradually decreased until the "- 7 min" measurement.




Figure 9: Time evolution of stomach temperature during the heating phase and the recovery phase (R1). The dark curve includes the R2_CWI condition and the clear curve the R2 AMB condition. The dotted lines represent the evolution of each individual in each condition.