



HAL
open science

Stability of the $P_{nc^1 - (P^0 + P^1)}$ element

Erell Jamelot, Patrick Ciarlet, Stefan Sauter

► **To cite this version:**

Erell Jamelot, Patrick Ciarlet, Stefan Sauter. Stability of the $P_{nc^1 - (P^0 + P^1)}$ element. ENUMATH 2023 - The European Conference on Numerical Mathematics and Advanced Applications, Sep 2023, Lisbonne, Portugal. hal-04414894v2

HAL Id: hal-04414894

<https://hal.science/hal-04414894v2>

Submitted on 12 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Stability Of The $\mathbf{P}_{nc}^1 - (P^0 + P^1)$ Element

Erell Jamelot^[0000-0002-9919-0793] and
Patrick Ciarlet, Jr.^[0000-0003-1198-5063] and
Stefan Sauter^[0000-0002-1403-1874]

Abstract We solve the Stokes problem numerically. We analyse the $\mathbf{P}_{nc}^1 - (P^0 + P^1)$ mixed finite element method, which exhibits interesting numerical features. However, only a partial proof of the stability is available. Our goal here is to gather evidence regarding the inf-sup condition.

1 Stokes Problem

A popular low-order method for solving the Stokes problem is the nonconforming Crouzeix-Raviart mixed finite element method [1, Example 4] (cf. [2, 3] for higher orders). The discrete velocity is piecewise affine, continuous in the barycentre of the interfaces between elements, but not globally continuous, \mathbf{P}_{nc}^1 ; and the discrete pressure is piecewise constant, P^0 . This method introduces some consistency error on the discrete velocity, which may lead to an unphysical solution when solving the Navier-Stokes equations, especially when the source field is a strong gradient [4, 5]. To recover consistency, the discrete pressure space can be generated by P^1 -continuous plus P^0 basis functions, cf. the TrioCFD code [6, 7]. This leads to the $\mathbf{P}_{nc}^1 - (P^0 + P^1)$ mixed finite element method, which exhibits interesting numerical features. However, only a partial proof of the stability is available in [8, 9]. Below we gather evidence regarding the inf-sup condition. Consider Stokes problem set in

Erell Jamelot
Université Paris-Saclay, CEA, Service de Thermo-hydraulique et de Mécanique des Fluides, 91191
Gif-sur-Yvette cedex, France, e-mail: erell.jamelot@cea.fr

Patrick Ciarlet, Jr
POEMS, CNRS, INRIA, ENSTA Paris, Institut Polytechnique de Paris, 91120 Palaiseau, France
e-mail: patrick.ciarlet@ensta-paris.fr

Stefan Sauter
Institut für Mathematik, Universität Zürich, Winterthurerstr 190, CH-8057 Zürich, Switzerland
e-mail: stas@math.uzh.ch

an open, bounded, connected subset Ω of \mathbb{R}^d , $d = 2, 3$, with a Lipschitz boundary:

$$\text{Find } (\mathbf{u}, p) \text{ such that } -\nu \Delta \mathbf{u} + \nabla p = \mathbf{f} \text{ and } \operatorname{div} \mathbf{u} = 0 \text{ in } \Omega, \quad (1)$$

with boundary condition $\mathbf{u} = 0$ on $\partial\Omega$ and a normalisation condition $\int_{\Omega} p = 0$. Given D an open subset of \mathbb{R}^d , let us set $\mathbf{L}^2(D) = (L^2(D))^d$, and $L^2_{zmv}(D) = \{q \in L^2(D) \mid \int_D q = 0\}$. We denote by $(\cdot, \cdot)_D$ the $L^2(D)$ or $\mathbf{L}^2(D)$ scalar product, and $\|\cdot\|_D$ the associated norm. Let $\mathbf{H}_0^1(D) = (H_0^1(D))^d$, and $\mathbf{H}^{-1}(D) = (H^{-1}(D))^d$ its dual space. Due Poincaré-Steklov inequality [10, Lemma 3.24], the semi-norm in $H_0^1(\Omega)$ is equivalent to the natural norm, so that $(v, w)_{H_0^1(\Omega)} = (\nabla v, \nabla w)_{\Omega}$ and $\|v\|_{H_0^1(\Omega)} = \|\nabla v\|_{\Omega}$. Let $\mathbf{v} \in \mathbf{H}_0^1(\Omega)$. We denote by $(v_i)_{i=1}^d$ the components of \mathbf{v} , and we set $\underline{\nabla} \mathbf{v} = (\partial_j v_i)_{i,j=1}^d \in [L^2(\Omega)]^{d \times d}$. Let us set $\mathbf{V} = \{\mathbf{v} \in \mathbf{H}_0^1(\Omega) \mid \operatorname{div} \mathbf{v} = 0\}$ which is a closed subset of $\mathbf{H}_0^1(\Omega)$. We denote by \mathbf{V}^{\perp} the orthogonal of \mathbf{V} in $\mathbf{H}_0^1(\Omega)$.

Proposition 1 *The operator $\operatorname{div} : \mathbf{H}_0^1(\Omega) \rightarrow L^2(\Omega)$ is an isomorphism of \mathbf{V}^{\perp} onto $L^2_{zmv}(\Omega)$. We call $C_{\operatorname{div}} > 1$ the constant such that: $\forall p \in L^2_{zmv}(\Omega)$, $\exists! \mathbf{v}_p \in \mathbf{V}^{\perp} \mid \operatorname{div} \mathbf{v}_p = p$ and $\|\mathbf{v}_p\|_{\mathbf{H}_0^1(\Omega)} \leq C_{\operatorname{div}} \|p\|_{\Omega}$ [11, Theorem 3.1 (e)].*

Let $\mathbf{f} \in \mathbf{H}^{-1}(\Omega)$. The variational formulation of Problem (1) reads:
Find $(\mathbf{u}, p) \in \mathbf{H}_0^1(\Omega) \times L^2_{zmv}(\Omega)$ such that $\forall (\mathbf{v}, q) \in \mathbf{H}_0^1(\Omega) \times L^2_{zmv}(\Omega)$:

$$\nu(\mathbf{u}, \mathbf{v})_{\mathbf{H}_0^1(\Omega)} - (p, \operatorname{div} \mathbf{v})_{\Omega} - (q, \operatorname{div} \mathbf{u})_{\Omega} = \langle \mathbf{f}, \mathbf{v} \rangle_{\mathbf{H}_0^1(\Omega)}. \quad (2)$$

One can prove that Problem (2) is well-posed with the T-coercivity theory [12, 13, 14], using Poincaré-Steklov inequality [10, Lemma 3.24] and Proposition 1.

2 Discretization

Consider $(\mathcal{T}_h)_h$ a family of simplicial triangulations of Ω . For a given $\mathcal{T}_h := \{K\}$, we call $\mathcal{F}_h := \{F\}$ its set of $(d-1)$ -simplices, called facets (faces if $d = 3$, edges if $d = 2$), and $\mathcal{V}_h := \{S\}$ its set of vertices. In the bounds, we will use \lesssim and \gtrsim for (hidden) constants independent of \mathcal{T}_h .

- For all $K \in \mathcal{T}_h$, we call h_K and ρ_K the diameters of K and its inscribed sphere respectively, and we let: $\sigma_K = \frac{h_K}{\rho_K}$. When the $(\mathcal{T}_h)_h$ is a shape-regular triangulation family (see e.g. [10, def. 11.2]), there exists a constant $\sigma > 1$, called the shape regularity parameter, such that for all h , for all $K \in \mathcal{T}_h$, $\sigma_K \leq \sigma$.
- Let $\mathcal{F}_h = \mathcal{F}_h^i \cup \mathcal{F}_h^b$, where $\mathcal{F}_h^b := \{F \in \mathcal{F}_h \mid F \subset \partial\Omega\}$, $\mathcal{F}_h^i = \mathcal{F}_h \setminus \mathcal{F}_h^b$. For all $K \in \mathcal{T}_h$, we set $\mathcal{F}_K^i = \{F \in \mathcal{F}_h^i \mid F \subset \partial K\}$.

For all $F \in \mathcal{F}_h$, M_F denotes the barycentre of F , and \mathbf{n}_F its unit normal (outward oriented if $F \subset \partial\Omega$). For all $K \in \mathcal{T}_h$ and $F \subset \partial K$, we denote by $\mathbf{n}_{F,K}$ the normal vector of F outgoing from K . Let \hat{K} be the reference simplex. For all $K \in \mathcal{T}_h$, we denote by $T_K : \hat{K} \rightarrow K$ the geometric mapping such that $\forall \hat{\mathbf{x}} \in \hat{K}$, $\mathbf{x}_{|K} = T_K(\hat{\mathbf{x}}) =$

$\mathbb{B}_K \hat{\mathbf{x}} + \mathbf{b}_K$, where $\mathbb{B}_K \in \mathbb{R}^{d \times d}$ and $\mathbf{b}_K \in \mathbb{R}^d$. Let us set $J_K = \det(\mathbb{B}_K)$. Recall that [10, Lemma 11.1]: $|J_K| = |K|/|\hat{K}|$, $\|\mathbb{B}_K\| \leq h_K/\rho_{\hat{K}}$, $\|\mathbb{B}_K^{-1}\| \leq h_{\hat{K}}/\rho_K$. For all $v \in L^2(\Omega)$, we set: $v_K = v|_K$, $\underline{v}_K = \int_K v/|K|$, $\tilde{v}_K = v_K - \underline{v}_K$, $\hat{v}_K = v \circ T_K$. By changing the variable, we get: $\|v\|_K^2 = |J_K| \|\hat{v}_K\|_{\hat{K}}^2$. Let $v \in H^1(K)$. By changing the variable, we get: $\nabla v|_K = (\mathbb{B}_K^{-1})^T \nabla_{\hat{\mathbf{x}}} \hat{v}$, and $\|\nabla v\|_K^2 \leq \|\mathbb{B}_K^{-1}\|^2 |J_K| \|\nabla_{\hat{\mathbf{x}}} \hat{v}\|_{\hat{K}}^2$, $\|\nabla_{\hat{\mathbf{x}}} \hat{v}\|_{\hat{K}}^2 \leq \|\mathbb{B}_K\|^2 |J_K|^{-1} \|\nabla v\|_K^2$.

We set $\mathcal{P}_h H^1 = \{v \in L^2(\Omega); \forall K \in \mathcal{T}_h, v|_K \in H^1(K)\}$, endowed with: $(v, w)_h := \sum_{K \in \mathcal{T}_h} (\nabla v, \nabla w)_K$, $\|v\|_h^2 = \sum_{K \in \mathcal{T}_h} \|\nabla v\|_K^2$. We define the operator ∇_h such that: $\forall q \in \mathcal{P}_h H^1$, $\nabla_h q \in \mathbf{L}^2(\Omega)$ is such that $\forall \mathbf{v} \in \mathbf{L}^2(\Omega)$, $(\nabla_h q, \mathbf{v})_\Omega = \sum_{K \in \mathcal{T}_h} (\nabla q, \mathbf{v})_K$. Let $K \in \mathcal{T}_h$ and $F \subset \partial K$, we denote by $T_{F,K}$ the geometric mapping such that $\forall \mathbf{x} \in F$, $T_{F,K}(T_K^{-1}(\mathbf{x})) = \mathbf{x}$. We let $\hat{F}_{F,K} = T_K^{-1}(F)$. For all $v \in \mathcal{P}_h H^1$, for all $K \in \mathcal{T}_h$ and for all $F \subset \partial K$, we set: $v_{F,K} = (v|_K)|_F$, $\underline{v}_{F,K} = \int_F v|_K/|F|$, $\tilde{v}_{F,K} = v|_K - \underline{v}_{F,K}$, $\hat{v}_{F,K} = v \circ T_{F,K}$. By changing the variable, we get $\|v_{F,K}\|_F^2 = (|F|/|\hat{F}_{F,K}|) \|\hat{v}_{F,K}\|_{\hat{F}_{F,K}}^2$. According to Equation [1, (3.17)], it holds:

$$\forall K \in \mathcal{T}_h, \forall F \subset \partial K, \quad |F| |K|^{-1} \lesssim \rho_K^{-1}. \quad (3)$$

We set $\mathcal{P}_h \mathbf{H}^1 = [\mathcal{P}_h H^1]^d$ and for all $\mathbf{v} \in \mathcal{P}_h \mathbf{H}^1$: $\underline{\nabla}_h \mathbf{v} = ((\nabla_h v_i)_{i=1}^d) \in [L^2(\Omega)]^{d \times d}$.

Let $F \in \mathcal{F}_h^i$ such that $F = \partial K_L \cap \partial K_R$ and $\mathbf{n}_F = \mathbf{n}_{F,K_L}$. The jump of $v \in \mathcal{P}_h H^1$ across F is defined by: $[v]_F := v|_{K_L} - v|_{K_R}$. For $F \in \mathcal{F}_h^b$, we set: $[v]_F := v|_F$.

For a subset D of \mathbb{R}^d , for $k \in \mathbb{N}$, we call $P^k(D)$ the set of degree k polynomials on D , $\mathbf{P}^k(D) = (P^k(D))^d$, and $\mathcal{P}_{disc}^k(\mathcal{T}_h) = \{q \in L^2(\Omega); \forall K \in \mathcal{T}_h, q|_K \in P^k(K)\}$. For $k > 0$ we define $\mathcal{P}^k(\mathcal{T}_h) = C^0(\bar{\Omega}) \cap \mathcal{P}_{disc}^k(\mathcal{T}_h)$. We call $\pi_0 : L^2(\Omega) \mapsto \mathcal{P}_{disc}^0(\mathcal{T}_h)$ the L^2 -orthogonal projection onto $\mathcal{P}_{disc}^0(\mathcal{T}_h)$, such that $\forall q \in L^2(\Omega)$, $\pi_0(q) = \sum_{K \in \mathcal{T}_h} \pi_0(q)(K) \mathbb{1}_K$ where $\forall K \in \mathcal{T}_h$, $\pi_0(q)(K) = \int_K q/|K|$.

For $K \in \mathcal{T}_h$ and a vertex S of K , $\lambda_{S,K}$ denotes the barycentric coordinate related to S . We can endow $\mathcal{P}^1(\mathcal{T}_h)$ with the basis $(\phi_S)_{S \in \mathcal{V}_h}$ such that: $\forall K \in \mathcal{T}_h$, $\phi_S|_K = \lambda_{S,K}$ if $S \in K$, zero otherwise, so that $\mathcal{P}^1(\mathcal{T}_h) = \text{vect}((\phi_S)_{S \in \mathcal{V}_h})$.

Let $k \in \mathbb{N}^*$, $v \in H^1(K) \cap P^k(K)$. Using the discrete trace inequality [15, Thm. 5] and Poincaré-Steklov inequality [10, Lemma 12.11], we have:

$$\|(\widehat{\tilde{v}_{F,K}})_{F,K}\|_{\hat{F}_{F,K}}^2 \lesssim \|(\widehat{\tilde{v}_{F,K}})_K\|_{\hat{K}}^2 \lesssim \|\nabla_{\hat{\mathbf{x}}} \hat{v}_K\|_{\hat{K}}^2. \quad (4)$$

Proposition 2 For all $K \in \mathcal{T}_h$, $\forall v \in H^1(K) \cap P^1(K)$, $\|\nabla v\|_K \lesssim (\rho_K)^{-1} \|\tilde{v}\|_K$.

Proof. We follow the proof of [16, Lemma 2]: changing the variable, we have $\|\nabla v\|_{L^2(K)} \lesssim \|\mathbb{B}_K^{-1}\| |K|^{1/2} |\int_{\hat{K}} \nabla_{\hat{\mathbf{x}}} \hat{v}_K|$. Integrating by parts in the reference element, it holds: $|\int_{\hat{K}} \nabla_{\hat{\mathbf{x}}} \hat{v}_K| = |\int_{\partial \hat{K}} \hat{v}_K \hat{\mathbf{n}}|_{\partial \hat{K}}|$. We get the result using Cauchy-Schwarz, [15, Thm. 5] and changing the variable again. \square

We set $\mathcal{P}_h \mathbf{H}(\text{div}) = \{\mathbf{v} \in \mathbf{L}^2(\Omega); \forall K \in \mathcal{T}_h, \mathbf{v}|_K \in \mathbf{H}(\text{div}; K)\}$.

We define the operator div_h such that: $\forall \mathbf{v} \in \mathcal{P}_h \mathbf{H}(\text{div})$, $\text{div}_h \mathbf{v} \in L^2(\Omega)$ is such that $\forall q \in L^2(\Omega)$, $(\text{div}_h \mathbf{v}, q)_\Omega = \sum_{K \in \mathcal{T}_h} (\text{div } \mathbf{v}, q)_K$.

Let $\mathbf{X}_{0,h}$ be the space of the discrete velocity of the nonconforming Crouzeix-Raviart mixed finite element method, defined by:

$$\begin{aligned} \mathbf{X}_h &= (X_h)^d, \quad X_h = \{v_h \in \mathcal{P}_{disc}^1(\mathcal{T}_h); \quad \forall F \in \mathcal{F}_h^i, \int_F [v_h] = 0\}; \\ \mathbf{X}_{0,h} &= (X_{0,h})^d, \quad X_{0,h} = \{v_h \in X_h; \quad \forall F \in \mathcal{F}_h^b, \int_F v_h = 0\}. \end{aligned} \quad (5)$$

The condition on the jumps of v_h on the inner facets is often called the patch-test condition. Due to this condition, one can prove that [1, Lemma 2]:

Proposition 3 *The broken norm $v_h \rightarrow \|v_h\|_h$ is a norm over $X_{0,h}$.*

For all $K \in \mathcal{T}_h$, for all $F \subset \partial K$, $\lambda_{F,K}$ denotes the barycentric coordinate related to the vertex $S_{F,K}$ opposite to facet F in K . We can endow X_h with the basis $(\psi_F)_{F \in \mathcal{F}_h}$ such that for all $K \in \mathcal{T}_h$, $\psi_F|_K = 1 - d\lambda_{F,K}$ if $F \subset \partial K$ and zero otherwise. We then have $\psi_F|_F = 1$, so that $[\psi_F]_F = 0$ if $F \in \mathcal{F}_h^i$, and for all $F' \neq F$, $\int_{F'} \psi_F = 0$. We have: $X_h = \text{vect}((\psi_F)_{F \in \mathcal{F}_h})$ and $X_{0,h} = \text{vect}((\psi_F)_{F \in \mathcal{F}_h^i})$.

Let $\Pi_h : \mathbf{H}^1(\Omega) \mapsto \mathbf{X}_h$ be the Crouzeix-Raviart interpolation operator, such that: $\forall \mathbf{v} \in \mathbf{H}^1(\Omega)$, $\Pi_h(\mathbf{v}) = \sum_{F \in \mathcal{F}_h} \Pi_F \mathbf{v} \psi_F$, $\Pi_F \mathbf{v} = \underline{\mathbf{v}}_F = \int_F \mathbf{v} / |F|$.

Proposition 4 *The interpolation operator Π_h is such that [16, Lemma 2]: $\forall \mathbf{v}_h \in \mathbf{H}_0^1(\Omega)$, $\|\Pi_h(\mathbf{v}_h)\|_h \leq \|\mathbf{v}_h\|_{\mathbf{H}_0^1(\Omega)}$.*

However, due to the consistency error on the discrete velocity, it is well-known [17, 18] that choosing the space of the discrete pressures equal to $\mathcal{P}_{disc}^0(\mathcal{T}_h) \cap L_{zmv}^2(\Omega)$ yields inaccurate results when solving Navier-Stokes equations. Starting from [19], a new element is proposed in [8]. The space of the discrete pressures is then: $Q_h = Q_{0,h} + Q_{1,h}$, where $Q_{0,h} := \mathcal{P}_{disc}^0(\mathcal{T}_h) \cap L_{zmv}^2$ and for $k > 0$, $Q_{k,h} := \mathcal{P}^k(\mathcal{T}_h) \cap L_{zmv}^2(\Omega)$. One can build an L^2 -orthogonal sum letting $\tilde{Q}_{1,h} = \{q_h \in Q_h \mid \forall K \in \mathcal{T}_h, \int_K q_h = 0\}$.

Proposition 5 *We have: $Q_h = Q_{0,h} \oplus \tilde{Q}_{1,h}$. Let $q_h \in Q_h$ and $\tilde{q}_{1,h} := q_h - \pi_0(q_h)$, it holds: $\|q_h\|_{\Omega}^2 = \|\pi_0(q_h)\|_{\Omega}^2 + \|\tilde{q}_{1,h}\|_{\Omega}^2$.*

To prove the discrete inf-sup condition of $\mathbf{X}_{0,h} \times Q_h$, we need some Lemmas.

Lemma 1 *Let $k \in \mathbb{N}^*$ and $(\mathbf{v}_h, q_{k,h}) \in \mathbf{X}_{0,h} \times Q_{k,h}$. We then have:*

$$|\sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\mathbf{v}_h \cdot \mathbf{n}_{F,K}, q_{k,h})_F| \lesssim \sigma \|\mathbf{v}_h\|_h \|\tilde{q}_{k,h}\|_{\Omega}. \quad (6)$$

Proof. We start with the proof of [19, Lemma 3.1]. Using the patch-test, one obtains:

$$\sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\mathbf{v}_h \cdot \mathbf{n}_{F,K}, q_{k,h}|_K)_F = \sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\tilde{\mathbf{v}}_{F,K} \cdot \mathbf{n}_{F,K}, \tilde{q}_{k,h}|_K)_F.$$

We can compute the following estimates, using (4) twice and (3):

$$|(\tilde{\mathbf{v}}_{F,K} \cdot \mathbf{n}_{F,K}, \tilde{q}_{k,h}|_K)_F| \lesssim |F| \|\widehat{\tilde{\mathbf{v}}}_{F,K}\|_{\hat{F}_{F,K}} \|\widehat{(\tilde{q}_{k,h})}_K\|_{\hat{F}_{F,K}} \lesssim \sigma_K \|\underline{\mathbf{v}}_h\|_K \|\tilde{q}_{k,h}\|_K.$$

Summing and using the discrete Cauchy-Schwarz inequality, we obtain (6). \square

Let us set $\mathbf{X}_{N,h} := \{\mathbf{v}_h \in \mathbf{X}_{0,h} \mid \forall F \in \mathcal{F}_h^i, \mathbf{v}_h \times \mathbf{n}_F = 0\}$. We can build an explicit right inverse of div_h from Q_h to $\mathbf{X}_{N,h}$.

Lemma 2 *For all $q_h \in Q_h$, there exists $\mathbf{v}_{N,h} \in \mathbf{X}_{N,h}$ such that:*

$$-(\text{div}_h \mathbf{v}_{N,h}, \pi_0(q_h))_\Omega = \nu^{-1} \|\pi_0(q_h)\|_\Omega^2, \quad (7)$$

$$\|\mathbf{v}_{N,h}\|_h \leq C_{\text{div}} \nu^{-1} \|\pi_0(q_h)\|_\Omega. \quad (8)$$

Proof. Let $q_h \in Q_h$. From Proposition 1, $\exists \mathbf{v}_N \in \mathbf{H}_0^1(\Omega) \mid \text{div } \mathbf{v}_N = -\nu^{-1} \pi_0(q_h)$ and $\|\mathbf{v}_N\|_{\mathbf{H}_0^1(\Omega)} \leq C_{\text{div}} \nu^{-1} \|\pi_0(q_h)\|_\Omega$. Let $\mathbf{v}_{N,h} \in \mathbf{X}_{N,h} \mid \forall F \in \mathcal{F}_h^i, \mathbf{v}_{N,h}(M_F) = (\Pi_h(\mathbf{v}_N)(M_F) \cdot \mathbf{n}_F) \mathbf{n}_F$. Integrating by part twice, we get: $\forall K \in \mathcal{T}_h, \int_K \text{div } \mathbf{v}_{N,h} = \int_K \text{div } \mathbf{v}_N$, hence (7). Using Proposition 4, we get (8). \square

We can also build an explicit right inverse of the discrete divergence operator from $Q_{1,h}$ to $\mathbf{X}_{T,h} := \{\mathbf{v}_h \in \mathbf{X}_{0,h} \mid \forall F \in \mathcal{F}_h^i, \mathbf{v}_h \cdot \mathbf{n}_F = 0\}$, if [19, Hyp. 4.1] holds:

Hypothesis 1. *We suppose that \mathcal{T}_h is such that $\partial\Omega$ contains at most one edge for $d = 2$ and at most two facets for $d = 3$, of the same element $K \in \mathcal{T}_h$.*

Lemma 3 *Assuming Hypothesis 1, $\forall q_{1,h} \in Q_{1,h}, \exists \mathbf{v}_{T,h} \in \mathbf{X}_{T,h}$ such that:*

$$(\nabla q_{1,h}, \mathbf{v}_{T,h})_\Omega \gtrsim \nu^{-1} \sigma^{-2(d-2)} \|\tilde{q}_{1,h}\|_\Omega^2, \quad (9)$$

$$(\text{div } \mathbf{v}_{T,h}, q_{0,h})_K = 0 \quad \forall q_{0,h} \in Q_{0,h}, \quad (10)$$

$$\|\mathbf{v}_{T,h}\|_h \lesssim \nu^{-1} \sigma^2 \|\tilde{q}_{1,h}\|_\Omega. \quad (11)$$

Proof. Let us consider $q_{1,h} \in Q_{1,h}$. For all $K \in \mathcal{T}_h$, we let $q_K := q_{1,h}|_K$ and $\hat{q}_K = q_K \circ T_K$. Since $q_{1,h} \in H^1(\Omega)$ we have (see e.g. [20, Prop. 2.2.10]): $\forall F \in \mathcal{F}_h^i \mid F = K \cap K', \nabla q_K \times \mathbf{n}_F = \nabla q_{K'} \times \mathbf{n}_F$. We construct $\mathbf{v}_{T,h} \in \mathbf{X}_{T,h} \mid \forall F \in \mathcal{F}_h^i, \mathbf{v}_{T,h}(M_F) = \nu^{-1} h_F^2 \mathbf{n}_F \times (\nabla q_{1,h} \times \mathbf{n}_F)$, where h_F is the diameter of F . We have:

$$\forall K \in \mathcal{T}_h \quad (\nabla q_{1,h}, \mathbf{v}_{T,h})_K = \nu^{-1} (d+1)^{-1} |K| \sum_{F \in \mathcal{F}_K^i} h_F^2 |\nabla q_K \times \mathbf{n}_F|^2. \quad (12)$$

Let $\mathbf{n}_{\hat{F}_{F,K}}$ be the unit normal vector of the facet $\hat{F}_{F,K}$ outgoing from \hat{K} . We have:

$$\mathbf{n}_{F,K} = |K| (\mathbb{B}_K^{-1})^T |\hat{F}_{F,K}| |F|^{-1} \mathbf{n}_{\hat{F}_{F,K}}.$$

Consider $d = 2$. Changing the variable, assuming Hypothesis 1, we get:

$$\sum_{F \in \mathcal{F}_K^i} h_F^2 |\nabla q_K \times \mathbf{n}_F|^2 = \sum_{F \in \mathcal{F}_K^i} |\hat{F}_{F,K}|^2 |\nabla_{\hat{\mathbf{x}}} \hat{q}_K \times \mathbf{n}_{\hat{F}_{F,K}}|^2 \gtrsim |\nabla_{\hat{\mathbf{x}}} \hat{q}_K|^2. \quad (13)$$

Consider $d = 3$. It holds that $|\nabla q_K \times \mathbf{n}_F| = |\hat{F}_{F,K}| |F|^{-1} |\mathbb{B}_K \nabla_{\hat{\mathbf{x}}} \hat{q}_K \times \mathbf{n}_{\hat{F}_{F,K}}|$ changing the variable. Hence: $|\nabla q_K \times \mathbf{n}_F| \gtrsim |F|^{-1} \|\mathbb{B}_K^{-1}\|^{-1} |\nabla_{\hat{\mathbf{x}}} \hat{q}_K \times \mathbf{n}_{\hat{F}_{F,K}}|$. Notice that $h_F |F|^{-1} \|\mathbb{B}_K^{-1}\|^{-1} \gtrsim \sigma_K^{-1}$. Assuming Hypothesis 1, we then have:

$$\sum_{F \in \mathcal{F}_K^i} h_F^2 |\nabla q_K \times \mathbf{n}_F|^2 \gtrsim \sigma_K^{-2} |\nabla_{\hat{\mathbf{x}}} \hat{q}_K|^2. \quad (14)$$

Using Poincaré-Steklov inequality [10, Lemma 12.11] in \hat{K} , we have: $|\nabla_{\hat{\mathbf{x}}} \hat{q}_K|^2 \gtrsim \|\hat{q}_K - \underline{\hat{q}}_K\|_{\hat{K}}^2$. Using (13) and (14) in (12), it holds:

$$(\nabla q_{1,h}, \mathbf{v}_{T,h})_K \gtrsim \nu^{-1} \sigma_K^{-2(d-2)} |K| |\nabla_{\hat{\mathbf{x}}} \hat{q}_K|^2 \gtrsim \nu^{-1} \sigma_K^{-2(d-2)} \|\tilde{q}_K\|_K^2. \quad (15)$$

We obtain (9) by summation. Remark that $\nabla \psi_{F|K} = |K|^{-1} |F| \mathbf{n}_{F,K}$. Let us prove (10). Since $\nabla \psi_{F|K} \perp \mathbf{v}_{T,h}(M_F)$, $(\operatorname{div} \mathbf{v}_{T,h}, q_{0,h})_K = |K| q_{0,h}(K) \sum_{F \in \mathcal{F}_K^i} \nabla \psi_{F|K} \cdot \mathbf{v}_{T,h}(M_F) = 0$. Let us prove (11). For all $K \in \mathcal{T}_h$, $\underline{\nabla} \mathbf{v}_{T,h|K} = \sum_{F \in \mathcal{F}_K^i} \mathbf{v}_{T,h}(M_F) \otimes \nabla \psi_{F|K}$. Hence: $\|\underline{\nabla} \mathbf{v}_{T,h}\|_K^2 \lesssim \nu^{-2} \sum_{F \in \mathcal{F}_K^i} |F|^2 |K|^{-1} h_F^4 |\nabla q_K \times \mathbf{n}_F|^2$. Using (3), and assuming Hypothesis 1, we get: $\|\underline{\nabla} \mathbf{v}_{T,h}\|_K^2 \lesssim \nu^{-2} \sigma_K^2 h_K^2 |K| \sum_{F \in \mathcal{F}_K^i} |\nabla q_K \times \mathbf{n}_F|^2 \lesssim \nu^{-2} \sigma_K^2 h_K^2 \|\nabla q_K\|_K^2$. From Proposition 2: $\|\underline{\nabla} \mathbf{v}_{T,h}\|_K \lesssim \nu^{-1} \sigma_K^2 \|\tilde{q}_K\|_K$. \square

Let $b_h : \mathbf{X}_h \times \mathcal{Q}_h \mapsto \mathbb{R} \mid \forall (\mathbf{v}_h, q_h) \in \mathbf{X}_h \times \mathcal{Q}_h, b_h(\mathbf{v}_h, q_h) = -(\operatorname{div}_h \mathbf{v}_h, q_{0,h})_\Omega + (\mathbf{v}_h, \nabla q_{1,h})_\Omega$ with $q_h = q_{0,h} + q_{1,h}$ such that $(q_{0,h}, q_{1,h}) \in \mathcal{Q}_{0,h} \times \mathcal{Q}_{1,h}$.

Theorem 1 *The following continuity property holds:*

$$\forall (\mathbf{v}_h, q_h) \in \mathbf{X}_{0,h} \times \mathcal{Q}_h, \quad |b_h(\mathbf{v}_h, q_h)| \lesssim \sigma \|\mathbf{v}_h\|_h \|q_h\|_\Omega. \quad (16)$$

Assuming Hypothesis 1, the following inf-sup condition holds:

$$\forall q_h \in \mathcal{Q}_h, \exists \mathbf{v}_h \in \mathbf{X}_{0,h}, \quad b_h(\mathbf{v}_h, q_h) \gtrsim \frac{1}{\sqrt{2}} C_{\operatorname{div}}^{-2} \sigma^{-2d} \|\mathbf{v}_h\|_h \|q_h\|_\Omega. \quad (17)$$

Proof. Let us prove (16). Let $(\mathbf{v}_h, q_h) \in \mathbf{X}_{0,h} \times \mathcal{Q}_h$. Integrating by parts, we have:

$$b_h(\mathbf{v}_h, q_h) = -(\operatorname{div}_h \mathbf{v}_h, \pi_0(q_h))_\Omega + \sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\mathbf{v}_h \cdot \mathbf{n}_{F,K}, q_{1,h})_F. \quad (18)$$

Using Cauchy-Schwarz, we have: $|(\operatorname{div}_h \mathbf{v}_h, \pi_0(q_h))_\Omega| \leq \sqrt{d} \|\mathbf{v}_h\|_h \|\pi_0(q_h)\|_\Omega$.

Using (6) in (18), we obtain (16) from Proposition 5. Let us prove (17), starting from the proof of [19, Lemma 4.2]. Let $q_h \in \mathcal{Q}_h \setminus \{0\}$, where $q_h = q_{0,h} + q_{1,h}$ is such that $(q_{0,h}, q_{1,h}) \in \mathcal{Q}_{0,h} \times \mathcal{Q}_{1,h}$. Let $\mathbf{v}_{N,h} := \mathbf{v}_{N,h}(q_h)$ be like in Lemma 2. Using (18) with $\mathbf{v}_h = \mathbf{v}_{N,h}$, (6) and (7), we have:

$$b_h(\mathbf{v}_{N,h}, q_h) \gtrsim \nu^{-1} \|\pi_0(q_h)\|_\Omega^2 - \sigma \|\mathbf{v}_{N,h}\|_h \|\tilde{q}_{1,h}\|_\Omega. \quad (19)$$

Using (8) and Young inequality, we have for all $\varepsilon > 0$:

$$-\|\mathbf{v}_{N,h}\|_h \|\tilde{q}_{1,h}\|_\Omega \geq -\frac{1}{2} C_{\operatorname{div}} \nu^{-1} (\varepsilon \|\pi_0(q_h)\|_\Omega^2 + \varepsilon^{-1} \|\tilde{q}_{1,h}\|_\Omega^2). \quad (20)$$

We now insert the bound (20) in (19) to get:

$$b_h(\mathbf{v}_{N,h}, q_h) \gtrsim \nu^{-1} \left(\left(1 - \frac{\varepsilon}{2} C_{\operatorname{div}} \sigma\right) \|\pi_0(q_h)\|_\Omega^2 - \frac{C_{\operatorname{div}}}{2\varepsilon} \sigma \|\tilde{q}_{1,h}\|_\Omega^2 \right). \quad (21)$$

Let $\mathbf{v}_{T,h} := \mathbf{v}_{T,h}(q_{1,h})$ be like in Lemma 3. Using (9) and (10), we get:

$$b_h(\mathbf{v}_{T,h}, q_{1,h}) = (\nabla q_{1,h}, \mathbf{v}_{T,h})_\Omega \gtrsim \nu^{-1} \sigma^{-2(d-2)} \|\tilde{q}_{1,h}\|_\Omega^2 \quad (22)$$

Last, we consider $\mathbf{v}_h^* = \mu \mathbf{v}_{N,h} + \mathbf{v}_{T,h}$, where $\mu > 0$. Using (21) and (22), we have:

$$\begin{aligned} b_h(\mathbf{v}_h^*, q_h) &= \mu b_h(\mathbf{v}_{N,h}, q_h) + b_h(\mathbf{v}_{T,h}, q_h), \\ &\gtrsim \nu^{-1} \left(\mu \left(1 - \frac{\varepsilon}{2} C_{\text{div}} \sigma\right) \|\pi_0(q_h)\|_{\Omega}^2 + \left(\sigma^{-2(d-2)} - \frac{\mu C_{\text{div}}}{2\varepsilon} \sigma\right) \|\tilde{q}_{1,h}\|_{\Omega}^2 \right). \end{aligned}$$

Let us choose $\varepsilon = (C_{\text{div}} \sigma)^{-1} < 1$ and $\mu = \sigma^{-2(d-2)} \varepsilon^2$. We obtain that:

$$b_h(\mathbf{v}_h^*, q_h) \gtrsim C_{\min} \nu^{-1} \|q_h\|_{\Omega}^2 \text{ with } C_{\min} = \varepsilon^2 \sigma^{-2(d-2)} = C_{\text{div}}^{-2} \sigma^{-2(d-1)}. \quad (23)$$

We will now bound $\|\mathbf{v}_h^*\|_h$ by $\|q_h\|_{\Omega}$. We have: $\|\mathbf{v}_h^*\|_h^2 \leq 2(\mu^2 \|\mathbf{v}_{N,h}\|_h^2 + \|\mathbf{v}_{T,h}\|_h^2)$. Using (8) and (11) $\|\mathbf{v}_h^*\|_h^2 \leq 2\nu^{-2} (\mu^2 C_{\text{div}}^2 \|\pi_0(q_h)\|_{\Omega}^2 + \sigma^4 \|\tilde{q}_{1,h}\|_{\Omega}^2)$, hence:

$$\|\mathbf{v}_h^*\|_h^2 \leq C_{\max}^2 \nu^{-2} \|q_h\|_{\Omega}^2 \text{ with } C_{\max} = \sqrt{2} \sigma^2. \quad (24)$$

Using (24) in (23), we get (17) with $C_{\min} C_{\max}^{-1} = \frac{1}{\sqrt{2}} C_{\text{div}}^{-2} \sigma^{-2d}$. \square

Let $\mathbf{V}_h := \{\mathbf{v}_h \in \mathbf{X}_{0,h} \mid \forall q_h \in Q_h, b_h(\mathbf{v}_h, q_h) = 0\}$. Let $(\mathbf{v}_h, q_{1,h}) \in \mathbf{V}_h \times Q_{1,h}$. Integrating by parts, we have $0 = b_h(\mathbf{v}_h, q_{1,h}) = -(\text{div}_h \mathbf{v}, \pi_0(q_{1,h}))_{\Omega} + \sum_{F \in \mathcal{F}_h} \int_F q_{1,h} [\mathbf{v}_h] \cdot \mathbf{n}_F = 0 + \sum_{F \in \mathcal{F}_h} \int_F q_{1,h} [\mathbf{v}_h] \cdot \mathbf{n}_F$. Hence:

Lemma 4 For all $(\mathbf{v}_h, q_{1,h}) \in \mathbf{V}_h \times Q_{1,h}$, it holds: $\sum_{F \in \mathcal{F}_h} \int_F q_{1,h} [\mathbf{v}_h] \cdot \mathbf{n}_F = 0$.

Let $\ell_{\mathbf{f}} \in \mathcal{L}(\mathbf{X}_h, \mathbb{R})$ be such that $\forall \mathbf{v}_h \in \mathbf{X}_h, \ell_{\mathbf{f}}(\mathbf{v}_h) = (\mathbf{f}, \mathbf{v}_h)_{\Omega}$ if $\mathbf{f} \in \mathbf{L}^2(\Omega)$, $\ell_{\mathbf{f}}(\mathbf{v}_h) = \langle \mathbf{f}, \mathcal{I}_h(\mathbf{v}_h) \rangle_{\mathbf{H}_0^1(\Omega)}$ if $\mathbf{f} \notin \mathbf{L}^2(\Omega)$, where $\mathcal{I}_h : \mathbf{X}_{0,h} \rightarrow \mathbf{Y}_{0,h}$ is for instance an averaging operator [10, §22.4.1], with $\mathbf{Y}_{0,h} = \{\mathbf{v}_h \in \mathbf{H}_0^1(\Omega) \mid \forall K \in \mathcal{T}_h, \mathbf{v}_h|_K \in \mathbf{P}^k(K)\}$. The discretization of Problem (1) with the $\mathbf{P}_{nc}^1 - (P^0 + P^1)$ element reads: Find $(\mathbf{u}_h, p_h) \in \mathbf{X}_{0,h} \times Q_h$ such that $\forall (\mathbf{u}_h, p_h) \in \mathbf{X}_{0,h} \times Q_h$

$$\nu(\mathbf{u}_h, \mathbf{v}_h)_h + b_h(\mathbf{v}_h, p_h) + b_h(\mathbf{u}_h, q_h) = \ell_{\mathbf{f}}(\mathbf{v}_h). \quad (25)$$

Due to Proposition 3 and Theorem 1, Problem (25) is well posed. We can derive error estimates similar to those given by [1, Theorems 3, 4, 6].

3 Convergence Estimates

When $\Omega \subset \mathbb{R}^2$, we have the two Lemmas below.

Lemma 5 For all $(\mathbf{v}_h, q_h) \in \mathbf{V}_h \times Q_{2,h}$, $(\nabla q_h, \mathbf{v}_h)_{\Omega} = 0$.

Proof. Let $(\mathbf{v}_h, q_h) \in \mathbf{V}_h \times Q_{2,h}$. Integrating by parts and using Lemma 4, we get that $\forall q_{1,h} \in Q_{1,h}$: $(\nabla q_h, \mathbf{v}_h)_{\Omega} = -\sum_{F \in \mathcal{F}_h} \int_F q_h [\mathbf{v}_h] \cdot \mathbf{n}_F = -\sum_{F \in \mathcal{F}_h} \int_F (q_h - q_{1,h}) [\mathbf{v}_h] \cdot \mathbf{n}_F = \sum_{F \in \mathcal{F}_h} \int_F (q_h - q_{1,h}) [\mathbf{v}_h - \mathbf{v}_h(M_F)] \cdot \mathbf{n}_F$. Hence, we obtain: $(\nabla q_h, \mathbf{v}_h)_{\Omega} = -\sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} \int_F (q_h - q_{1,h}) (\mathbf{v}_h - \mathbf{v}_h(M_F)) \cdot \mathbf{n}_{F,K}$. Choose $q_{1,h}$ such that $\forall S \in \mathcal{V}_h, q_{1,h}(S) = q_{2,h}(S)$. Then, $\forall F \in \mathcal{F}_h$, the degree 3 polynomial $(q_h - q_{1,h})(\mathbf{v}_h - \mathbf{v}_h(M_F))$ vanishes at the quadrature points of Simpson's rule. Since

it is exact for degree 3 polynomials, $\forall F \in \mathcal{F}_h$, $\int_F (q_h - q_{1,h})(\mathbf{v}_h - \mathbf{v}_h(M_F)) \cdot \mathbf{n}_{F,K} = 0$ and $(\nabla q_h, \mathbf{v}_h)_\Omega = 0$. \square

Lemma 6 *Let $\phi \in H^3(\Omega) \cap L^2_{zmv}(\Omega)$ such that $|\phi|_{H^3(\Omega)} \neq 0$.*

Then, $\forall \mathbf{v}_h \in \mathbf{V}_h$, $(\nabla \phi, \mathbf{v}_h)_\Omega \lesssim \sigma h^2 |\phi|_{H^3(\Omega)} \|\mathbf{v}_h\|_\Omega$.

Proof. Let $\mathbf{v}_h \in \mathbf{V}_h$. Using Lemma 5 and Cauchy-Schwarz, it holds: $|(\nabla \phi, \mathbf{v}_h)_\Omega| = |(\nabla(\phi - q_h), \mathbf{v}_h)_\Omega|$ for all $q_h \in \mathcal{Q}_{2,h}$. We then use [10, Lemma 11.9] to conclude. \square

Theorem 2 *Suppose that Ω is convex. Let $\phi \in H^3(\Omega) \cap L^2_{zmv}(\Omega)$ such that $|\phi|_{H^3(\Omega)} \neq 0$. Let (\mathbf{u}_h, p_h) be the solution of Problem (25) with $\mathbf{f} = \nabla \phi$. The solution of Problem (1) is then $(\mathbf{u}, p) = (0, \phi)$. It holds:*

$$\nu \|\mathbf{u}_h\|_h \lesssim \sigma^2 h^3 |\phi|_{H^3(\Omega)}; \quad \nu \|\mathbf{u}_h\|_\Omega \lesssim \sigma^3 h^4 |\phi|_{H^3(\Omega)}.$$

Proof. Setting $\mathbf{v}_h = \mathbf{u}_h$ as test-function in Problem (25) and using Lemma 6, it holds: $\nu \|\mathbf{u}_h\|_h^2 = (\nabla \phi, \mathbf{u}_h)_\Omega \lesssim \sigma h^2 |\phi|_{H^3(\Omega)} \|\mathbf{u}_h\|_\Omega$. From [1, Theorem 4], have: $\|\mathbf{u}_h\|_\Omega \lesssim \sigma h \|\mathbf{u}_h\|_h$. We deduce that $\nu \|\mathbf{u}_h\|_h \lesssim \sigma^2 h^3 |\phi|_{H^3(\Omega)}$. Using this estimate, we get that $\nu \|\mathbf{u}_h\|_\Omega \lesssim \sigma^3 h^4 |\phi|_{H^3(\Omega)}$. \square

When $\Omega \subset \mathbb{R}^3$, let \mathcal{E}_h be the set of edges of \mathcal{T}_h . Let $E := S_i S_j \in \mathcal{E}_h$. We set $\phi_E := 4\lambda_i \lambda_j$ where λ_i (resp. λ_j) is the barycentric coordinate related to vertex S_i (resp. S_j). Let $\tilde{P}^2(\mathcal{T}_h) = \text{vect}((\tilde{\phi}_E)_{E \in \mathcal{E}_h})$, where for every edge E , $\tilde{\phi}_E$ is the canonical extension of ϕ_E to all simplices intersecting with $\text{int}(E)$. In order to obtain optimal results in 3D similar to those obtained in 2D, the discrete pressure space can be $Q_h = Q_{0,h} + Q_{1,h} + \tilde{Q}_{2,h}$, where $\tilde{Q}_{2,h}$ is a subspace of $\tilde{P}^2(\mathcal{T}_h) \cap L^2_{zmv}(\Omega)$ [9]. One can then prove Lemma 5 in 3D using a 7-point quadrature rule on a triangle, exact for polynomials of degree 3. The quadrature points are the barycentre of the triangle, its vertices and the midpoints of its edges. Lemma 6 follows. Finally, in [9, Theorem 4.4.6, page 92], it is claimed that one obtains a convergence result identical to the one of Theorem 2, now in 3D.

4 Numerical Results For The Stationary Navier-Stokes Model

Let us consider stationary Navier-Stokes equations set in $\Omega = (0, 1)^d$:

$$\text{Find } (\mathbf{u}, p) \text{ such that } -\nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = \mathbf{f} \text{ and } \text{div } \mathbf{u} = 0 \text{ in } \Omega, \quad (26)$$

with $\mathbf{u}|_{\partial\Omega=0}$ and $\int_\Omega p = 0$. Computations are done with the TrioCFD code [6, 7] with the $\mathbf{P}_{nc}^1 - (P^0 + P^1)$ finite element. The convection term is discretized using the MUSCL scheme [6]. Let us set: $\|(\mathbf{u}, p)\|_\nu^2 := \|\nabla \mathbf{u}\|_\Omega^2 + \nu^{-2} \|p\|_\Omega^2$. The discrete errors values are $\varepsilon_0^\nu(\mathbf{u}_h) := \|\mathbf{u} - \mathbf{u}_h\|_\Omega / \|(\mathbf{u}, p)\|_\nu$, and $\varepsilon_0^\nu(p_h) := \nu^{-1} \|p - p_h\|_\Omega / \|(\mathbf{u}, p)\|_\nu$. We let N be the number of degrees of freedom ($h \approx N^{1/d}$).

We first consider the 2D case and the prescribed solution $\mathbf{u} = (y, -x)^T$, $p = (x^2 +$

$y^2)/2 - 1/3$ so that $\mathbf{f} = 0$. Fig. 1 shows $\varepsilon_0^\nu(\mathbf{u}_h)$ (left) and $\varepsilon_0^\nu(p_h)$ (right) against N , for $\nu = 10^{-1}$, $\nu = 10^{-2}$ and $\nu = 10^{-3}$.

Fig. 1 2D case. Plots of $\varepsilon_0^\nu(\mathbf{u}_h)$ and $\varepsilon_0^\nu(p_h)$ for $\nu = 10^{-1}$, $\nu = 10^{-2}$ or $\nu = 10^{-3}$.




Table 1 shows the convergence rates for \mathbf{u} , $\tau_{\mathbf{u}}$ (left), and for p , τ_p (right).

Table 1 2D case, convergence rates.

ν	10^{-1}	10^{-2}	10^{-3}	ν	10^{-1}	10^{-2}	10^{-3}
$\tau_{\mathbf{u}}$	3.9	3.3	2.3	τ_p	3.4	3.4	2.5

We then consider the 3D case and the prescribed solution $\mathbf{u} = (y - z, z - x, x - y)^T$, $p = \frac{1}{2}(x^2 + y^2 + z^2) - xy - xz - yz - 1/4$ so that $\mathbf{f} = 0$. Fig. 2 shows $\varepsilon_0^\nu(\mathbf{u}_h)$ (left) and $\varepsilon_0^\nu(p_h)$ (right) against the number of degrees of freedom N , for $\nu = 10^{-1}$, $\nu = 10^{-2}$ and $\nu = 10^{-3}$.

Fig. 2 3D case. Plots of $\varepsilon_0^\nu(\mathbf{u}_h)$ and $\varepsilon_0^\nu(p_h)$ for $\nu = 10^{-1}$, $\nu = 10^{-2}$ or $\nu = 10^{-3}$.




Table 2 shows the convergence rates for \mathbf{u} , $\tau_{\mathbf{u}}$ (left), and p , τ_p (right).

Table 2 3D case, convergence rates.

ν 10^{-1} 10^{-2} 10^{-3}	ν 10^{-1} 10^{-2} 10^{-3}
$\tau_{\mathbf{u}}$ 5.6 4.7 3.3	τ_p 5.2 4.8 3.7

More numerical results are available in [8, 9, 6, 21]. The $\mathbf{P}_{nc}^1 - (P^0 + P^1)$ scheme has good approximation properties and can therefore be used to solve the Navier-Stokes equations in the context of industrial simulation. We plan to study discrete pressure enrichment to higher order nonconforming Crouzeix-Raviart mixed FEM [2, 3].

Acknowledgements We would like to thank P.-E. Angeli for his contribution to §4.

References

1. Crouzeix M., Raviart, P.-A.: *RAIRO, Sér. Anal. Numer.*, **7**, 33, 33–75 (1973).
2. Sauter, S.: *Computers & Mathematics with Applications*, **149**, 49–70 (2023).
3. Sauter, S., Torres, C.: *Mathematics of Computation*, **92**, 341, 1033–1059 (2023).
4. Hecht, F.: *RAIRO, Sér. Anal. Numér.* **15**, 2, 119-150 (1981).
5. Linke, A.: *Comput. Methods Appl. Mech. Engrg.* **268**, 782-800 (2014).
6. Angeli, P.-E. et al: FVCA8 benchmark for the Stokes and Navier-Stokes equations with the TrioCFD code, in: Cancès, C., Omnes, P. (eds.) *Finite Volumes for Complex Applications VIII - Methods and Theoretical Aspects*, pp. 181-302, Springer (2017).
7. CEA/STMF: <https://trio CFD.cea.fr/>.
8. Heib, S.: *Nouvelles discrétisations non structurées pour des écoulements de fluides à incompressibilité renforcée*, PhD thesis, Univ. Pierre et Marie Curie - Paris VI, France (2003).
9. Fortin, T.: *Une méthode éléments finis à décomposition L^2 d'ordre élevé motivée par la simulation d'écoulement diphasique bas Mach*, Univ. Pierre et Marie Curie - Paris VI (2006).
10. Ern, A., Guermond, J.-L.: *Finite elements I*, Springer (2021).
11. Amrouche, C., Ciarlet, P. G., Mardare, C.: *Journal de Mathématiques Pures et Appliquées*, **104**, 2, 207-226 (2015).
12. Bonnet-Ben Dhia, A.-S., Ciarlet Jr., P.: Méthodes variationnelles pour l'analyse de problèmes non coercifs, *M.Sc. AMS lecture notes (ENSTA-IPP)* (2022).
13. Barré, M., Ciarlet Jr., P.: The T-coercivity approach for mixed problems, *C. R. Acad. Sci. Paris, Ser. I* (to appear).
14. Jamelot, E.: Improved stability estimates for solving Stokes problem with Fortin-Soulie finite elements, submitted.
15. Warburton, T., Hesthaven, J.S.: *Comput. Methods Appl. Mech. Engrg.* **192**, 25, 2765–2773 (2003).
16. Apel, T., Nicaise, S., Schöberl, J.: *Numerische Mathematik* **89**, 2, 193–223 (2001).
17. Emonot, P.: *Méthode de volumes éléments finis : applications aux équations de Navier-Stokes et résultats de convergence*, PhD thesis, Université Claude Bernard - Lyon I, France (1992).
18. John, V., Linke, A., Merdon, C., Neilan M., Rebholz, L.G.: *SIAM REVIEW* **59**, 3, 492–544 (2017).
19. Bernardi, B., Hecht F.: *ESAIM: M2AN*, **34**, 5, 953-980 (2000).
20. Assous, F., Ciarlet Jr., P., Labrunie, S. : *Mathematical Foundations of Computational Electromagnetism*, Springer (2018).
21. Gounand, S.: <https://cea.hal.science/cea-02489502/document> (2015).