



## Stability of the P1nc-(P0+P1) element

Erell Jamelot, Patrick Ciarlet, Stefan Sauter

### ► To cite this version:

Erell Jamelot, Patrick Ciarlet, Stefan Sauter. Stability of the P1nc-(P0+P1) element. ENUMATH 2023 - The European Conference on Numerical Mathematics and Advanced Applications, Sep 2023, Lisbonne, Portugal. hal-04414894v4

HAL Id: hal-04414894

<https://hal.science/hal-04414894v4>

Submitted on 15 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Stability Of The $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$ Element

Erell Jamelot<sup>[0000-0002-9919-0793]</sup> and  
Patrick Ciarlet, Jr<sup>[0000-0003-1198-5063]</sup> and  
Stefan Sauter<sup>[0000-0002-1403-1874]</sup>

**Abstract** We solve the Stokes problem numerically. We analyse the  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  mixed finite element method which exhibits interesting numerical features. However, only an incomplete proof of the inf-sup condition is available. We prove here this condition and the stability of the method.

## 1 Stokes Problem

A popular low-order method for solving the Stokes problem is the nonconforming Crouzeix-Raviart mixed finite element method [1, Example 4] (cf. [2, 3] for higher orders). The discrete velocity is piecewise affine, continuous in the barycentre of the interfaces between elements, but not globally continuous,  $\mathbf{P}_{nc}^1$ ; and the discrete pressure is piecewise constant,  $P^0$ . This method introduces some consistency error on the discrete velocity, which may lead to an unphysical solution when solving the Navier-Stokes equations, especially when the source field is a strong gradient [4, 5]. To recover consistency, the discrete pressure space can be generated by  $P^1$ -continuous plus  $P^0$  basis functions, cf. the TrioCFD code [6, 7]. This leads to the  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  mixed finite element method, which exhibits interesting numerical features. However, only an incomplete proof of the stability is available in [8, 9]. Below, we prove the inf-sup condition. Consider the Stokes problem set in an open,

---

Erell Jamelot  
Université Paris-Saclay, CEA, Service de Thermo-hydraulique et de Mécanique des Fluides, 91191 Gif-sur-Yvette cedex, France, e-mail: [rell.jamelot@cea.fr](mailto:rell.jamelot@cea.fr)

Patrick Ciarlet, Jr  
POEMS, CNRS, INRIA, ENSTA Paris, Institut Polytechnique de Paris, 91120 Palaiseau, France  
e-mail: [patrick.ciarlet@ensta-paris.fr](mailto:patrick.ciarlet@ensta-paris.fr)

Stefan Sauter  
Institut für Mathematik, Universität Zürich, Winterthurerstr 190, CH-8057 Zürich, Switzerland  
e-mail: [stas@math.uzh.ch](mailto:stas@math.uzh.ch)

bounded, connected subset  $\Omega$  of  $\mathbb{R}^d$ ,  $d = 2, 3$ , with a Lipschitz boundary  $\partial\Omega$ :

$$\text{Find } (\mathbf{u}, p) \text{ such that } -\nu \Delta \mathbf{u} + \nabla p = \mathbf{f} \text{ and } \operatorname{div} \mathbf{u} = 0 \text{ in } \Omega, \quad (1)$$

with homogeneous boundary conditions  $\mathbf{u} = 0$  on  $\partial\Omega$  and a normalisation condition:  $\int_{\Omega} p = 0$ . The case of inhomogeneous Dirichlet boundary conditions, that is  $\mathbf{u} = \mathbf{g}$  on  $\partial\Omega$  with  $\mathbf{g} \in \mathbf{H}^{\frac{1}{2}}(\partial\Omega)$  such that  $\int_{\partial\Omega} \mathbf{g} \cdot \mathbf{n}_{|\partial\Omega} = 0$  can be studied similarly. Given  $D$  an open subset of  $\mathbb{R}^d$ , let us set  $\mathbf{L}^2(D) = (L^2(D))^d$ , and  $L_{zmv}^2(D) = \{q \in L^2(D) \mid \int_D q = 0\}$ . We denote by  $(\cdot, \cdot)_D$  the  $L^2(D)$  or  $\mathbf{L}^2(D)$  inner product, and  $\|\cdot\|_D$  the associated norm. Due to Poincaré-Steklov inequality [10, Lemma 3.24], the semi-norm in  $H_0^1(\Omega)$  is equivalent to the natural norm, so that  $(v, w)_{H_0^1(\Omega)} = (\nabla v, \nabla w)_{\Omega}$  and  $\|v\|_{H_0^1(\Omega)} = \|\nabla v\|_{\Omega}$ . Let us set  $\mathbf{V} = \{\mathbf{v} \in \mathbf{H}_0^1(\Omega) \mid \operatorname{div} \mathbf{v} = 0\}$  which is a closed subset of  $\mathbf{H}_0^1(\Omega)$ . We denote by  $\mathbf{V}^\perp$  the orthogonal of  $\mathbf{V}$  in  $\mathbf{H}_0^1(\Omega)$ .

**Proposition 1** [13, Theorem 3.1 (e)]. *The operator  $\operatorname{div} : \mathbf{H}_0^1(\Omega) \rightarrow L^2(\Omega)$  is an isomorphism of  $\mathbf{V}^\perp$  onto  $L_{zmv}^2(\Omega)$ . We call  $C_{\operatorname{div}} > 1$  the constant such that:  $\forall p \in L_{zmv}^2(\Omega), \exists! \mathbf{v}_p \in \mathbf{V}^\perp \mid \operatorname{div} \mathbf{v}_p = p \text{ and } \|\mathbf{v}_p\|_{\mathbf{H}_0^1(\Omega)} \leq C_{\operatorname{div}} \|p\|_{\Omega}$ .*

Let  $\mathbf{f} \in (\mathbf{H}_0^1(\Omega))'$ . The variational formulation of Problem (1) reads:  
Find  $(\mathbf{u}, p) \in \mathbf{H}_0^1(\Omega) \times L_{zmv}^2(\Omega)$  such that  $\forall (\mathbf{v}, q) \in \mathbf{H}_0^1(\Omega) \times L_{zmv}^2(\Omega)$ :

$$\nu(\mathbf{u}, \mathbf{v})_{\mathbf{H}_0^1(\Omega)} - (p, \operatorname{div} \mathbf{v})_{\Omega} - (q, \operatorname{div} \mathbf{u})_{\Omega} = \langle \mathbf{f}, \mathbf{v} \rangle_{\mathbf{H}_0^1(\Omega)}. \quad (2)$$

One can prove that Problem (2) is well-posed with the T-coercivity theory [14, 15, 16], using Poincaré-Steklov inequality [10, Lemma 3.24] and Proposition 1.

## 2 Discretization And Stability Of The $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$ Scheme

Consider  $(\mathcal{T}_h)_h$  a family of simplicial triangulations of  $\Omega$ . For a given  $\mathcal{T}_h := \{K\}$ , we call  $\mathcal{F}_h := \{F\}$  its set of  $(d-1)$ -simplices, called facets (faces if  $d = 3$ , edges if  $d = 2$ ), and  $\mathcal{V}_h := \{S\}$  its set of vertices. In the bounds, we will use  $\lesssim$  and  $\gtrsim$  for (hidden) constants independent of  $\mathcal{T}_h$ .

- For  $K \in \mathcal{T}_h$ , we call  $h_K$  and  $\rho_K$  the diameters of  $K$  and its inscribed sphere respectively, and we let:  $\sigma_K = \frac{h_K}{\rho_K}$ . When the  $(\mathcal{T}_h)_h$  is a shape-regular triangulation family (see e.g. [10, def. 11.2]), there exists a constant  $\sigma > 1$ , called the shape regularity parameter, such that for all  $h$ , for all  $K \in \mathcal{T}_h$ ,  $\sigma_K \leq \sigma$ .
- Let  $\mathcal{F}_h = \mathcal{F}_h^i \cup \mathcal{F}_h^b$ , where  $\mathcal{F}_h^b = \{F \in \mathcal{F}_h \mid F \subset \partial\Omega\}$ ,  $\mathcal{F}_h^i = \mathcal{F}_h \setminus \mathcal{F}_h^b$ . For  $K \in \mathcal{T}_h$ , we set  $\mathcal{F}_K^i = \{F \in \mathcal{F}_h^i \mid F \subset \partial K\}$ .

For  $F \in \mathcal{F}_h$ ,  $M_F$  denotes the barycentre of  $F$ , and  $\mathbf{n}_F$  its unit normal (outward oriented if  $F \subset \partial\Omega$ ). For  $K \in \mathcal{T}_h$  and  $F \subset \partial K$ , we denote by  $\mathbf{n}_{F,K}$  the normal vector of  $F$  outgoing from  $K$ . Let  $\hat{K}$  be the reference simplex. For  $K \in \mathcal{T}_h$ , we denote by  $T_K : \hat{K} \rightarrow K$  the geometric mapping such that  $\forall \hat{\mathbf{x}} \in \hat{K}$ ,  $\mathbf{x}|_K = T_K(\hat{\mathbf{x}}) = \mathbb{B}_K \hat{\mathbf{x}} + \mathbf{b}_K$ ,

where  $\mathbb{B}_K \in \mathbb{R}^{d \times d}$  and  $\mathbf{b}_K \in \mathbb{R}^d$ . Let us set  $J_K = \det(\mathbb{B}_K)$ . Recall that [10, Lemma 11.1]:  $|J_K| = |K|/|\hat{K}|$ ,  $\|\mathbb{B}_K\| \leq h_K/\rho_{\hat{K}}$ ,  $\|\mathbb{B}_K^{-1}\| \leq h_{\hat{K}}/\rho_K$ . For  $v \in L^2(\Omega)$ , we set:  $v_K = v|_K$ ,  $\hat{v}_K = v \circ T_K$ . By changing the variable, we get:  $\|v\|_K^2 = |J_K| \|\hat{v}_K\|_{\hat{K}}^2$ . Let  $v \in H^1(K)$ , then  $\nabla v|_K = (\mathbb{B}_K^{-1})^T \nabla_{\hat{x}} \hat{v}$ , and  $\|\nabla v\|_K^2 \leq \|\mathbb{B}_K^{-1}\|^2 |J_K| \|\nabla_{\hat{x}} \hat{v}\|_{\hat{K}}^2$ ,  $\|\nabla_{\hat{x}} \hat{v}\|_{\hat{K}}^2 \leq \|\mathbb{B}_K\|^2 |J_K|^{-1} \|\nabla v\|_K^2$ . Let  $K \in \mathcal{T}_h$  and  $F \subset \partial K$ , we denote by  $T_{F,K}$  the geometric mapping such that  $\forall \mathbf{x} \in F$ ,  $T_{F,K}(T_K^{-1}(\mathbf{x})) = \mathbf{x}$ . We let  $\hat{T}_{F,K} = T_K^{-1}(F)$ . According to [1, Eq. (3.17)]:

$$\forall K \in \mathcal{T}_h, \forall F \subset \partial K, \quad |F| |K|^{-1} \lesssim \rho_K^{-1}. \quad (3)$$

We set  $\mathcal{P}_h H^1 = \{v \in L^2(\Omega) \mid \forall K \in \mathcal{T}_h, v|_K \in H^1(K)\}$ , endowed with:  $(v, w)_h := \sum_{K \in \mathcal{T}_h} (\nabla v, \nabla w)_K$ ,  $\|v\|_h^2 = \sum_{K \in \mathcal{T}_h} \|\nabla v\|_K^2$ . Let  $F \in \mathcal{F}_h^i$  such that  $F = \partial K_L \cap \partial K_R$  and  $\mathbf{n}_F = \mathbf{n}_{F,K_L}$ . The jump of  $v \in \mathcal{P}_h H^1$  across  $F$  is defined by:  $[v]_F := v|_{K_L} - v|_{K_R}$ . For  $F \in \mathcal{F}_h^b$ , we set:  $[v]_F := v|_F$ .

For a subset  $D$  of  $\mathbb{R}^d$ , for  $k \in \mathbb{N}$ , we call  $P^k(D)$  the set of degree  $k$  polynomials on  $D$ ,  $\mathbf{P}^k(D) = (P^k(D))^d$ , and  $\mathcal{P}_{disc}^k(\mathcal{T}_h) = \{q \in L^2(\Omega) \mid \forall K \in \mathcal{T}_h, q|_K \in P^k(K)\}$ .

For  $k > 0$  we define  $\mathcal{P}^k(\mathcal{T}_h) = C^0(\bar{\Omega}) \cap \mathcal{P}_{disc}^k(\mathcal{T}_h)$ . We call  $\pi_0 : L^2(\Omega) \mapsto \mathcal{P}_{disc}^0(\mathcal{T}_h)$  the  $L^2$ -orthogonal projection onto  $\mathcal{P}_{disc}^0(\mathcal{T}_h)$ , such that  $\forall q \in L^2(\Omega)$ ,  $\pi_0(q) = \sum_{K \in \mathcal{T}_h} q_K \mathbb{1}_K$  where  $\forall K \in \mathcal{T}_h$ ,  $q_K = \int_K q / |K|$ .

For  $K \in \mathcal{T}_h$  and a vertex  $S$  of  $K$ ,  $\lambda_{S,K}$  denotes the barycentric coordinate related to  $S$ . We can endow  $\mathcal{P}^1(\mathcal{T}_h)$  with the basis  $(\phi_S)_{S \in \mathcal{V}_h}$  such that:  $\forall K \in \mathcal{T}_h$ ,  $\phi_{S|K} = \lambda_{S,K}$  if  $S \in K$ , zero otherwise, so that  $\mathcal{P}^1(\mathcal{T}_h) = \text{vect}((\phi_S)_{S \in \mathcal{V}_h})$ .

Using [12, Thm. 5] and Eq. (3), one can prove that:

$$\forall w \in P^k(K), \forall F \subset \partial K, \quad \|w\|_F \lesssim (|F| |K|^{-1})^{\frac{1}{2}} \|w\|_K \lesssim (\rho_K)^{-\frac{1}{2}} \|w\|_K. \quad (4)$$

Using Eq. (4) with  $w = v - \underline{v}_K$  and Poincaré-Steklov inequality [10, Lemma 12.11] in  $K$ , it holds (with below,  $\underline{v}_F = \int_F v / |F|$ ):

$$\forall v \in P^k(K), \forall F \subset \partial K, \quad \|v - \underline{v}_F\|_F \lesssim \|v - \underline{v}_K\|_F \lesssim (\sigma_K h_K)^{1/2} \|\nabla v\|_K. \quad (5)$$

Using the proof of [17, Lemma 2] and [12, Thm. 5], we can prove that:

$$\forall v \in P^1(K), \quad \|\nabla v\|_K \lesssim (\rho_K)^{-1} \|v - \underline{v}_K\|_K. \quad (6)$$

Let  $\mathbf{X}_{0,h}$  be the space of the discrete velocity of the nonconforming Crouzeix-Raviart mixed finite element method, defined by:

$$\mathbf{X}_h = (X_h)^d, \quad X_h = \{v_h \in \mathcal{P}_{disc}^1(\mathcal{T}_h) \mid \forall F \in \mathcal{F}_h^i, \int_F [v_h] = 0\}; \quad (7)$$

$$\mathbf{X}_{0,h} = (X_{0,h})^d, \quad X_{0,h} = \{v_h \in X_h \mid \forall F \in \mathcal{F}_h^b, \int_F v_h = 0\}. \quad (8)$$

The condition on the jumps of  $v_h$  on the inner facets is often called the patch-test condition. Due to this condition, one can prove that :

**Proposition 2** [1, Lemma 2]. *The broken norm  $v_h \rightarrow \|v_h\|_h$  is a norm over  $X_{0,h}$ .*

For  $K \in \mathcal{T}_h$  and  $F \subset \partial K$ ,  $\lambda_{F,K}$  denotes the barycentric coordinate related to the vertex  $S_{F,K}$  opposite to facet  $F$  in  $K$ . We can endow  $X_h$  with the basis  $(\psi_F)_{F \in \mathcal{F}_h}$  such that for all  $K \in \mathcal{T}_h$ ,  $\psi_F|_K = 1 - d\lambda_{F,K}$  if  $F \subset \partial K$  and zero otherwise. We then have  $\psi_F|_F = 1$ , so that  $[\psi_F]_F = 0$  if  $F \in \mathcal{F}_h^i$ , and for all  $F' \neq F$ ,  $\int_{F'} \psi_F = 0$ . We have:  $X_h = \text{vect}((\psi_F)_{F \in \mathcal{F}_h})$  and  $X_{0,h} = \text{vect}((\psi_F)_{F \in \mathcal{F}_h^i})$ .

Let  $\Pi_h : \mathbf{H}^1(\Omega) \mapsto \mathbf{X}_h$  be the Crouzeix-Raviart interpolation operator, such that for all  $\mathbf{v} \in \mathbf{H}^1(\Omega)$ ,  $\Pi_h(\mathbf{v}) = \sum_{F \in \mathcal{F}_h} \underline{\mathbf{v}}_F \psi_F$ ,  $\underline{\mathbf{v}}_F = \int_F \mathbf{v} / |F|$ .

**Proposition 3** [17, Lemma 2]. *The interpolation operator  $\Pi_h$  is such that for all  $\mathbf{v}_h \in \mathbf{H}_0^1(\Omega)$ ,  $\|\Pi_h(\mathbf{v}_h)\|_h \leq \|\mathbf{v}_h\|_{\mathbf{H}_0^1(\Omega)}$ .*

However, due to the consistency error on the discrete velocity, it is well-known [18, 19] that choosing the space of the discrete pressures equal to  $\mathcal{P}_{disc}^0(\mathcal{T}_h) \cap L_{zmv}^2(\Omega)$  yields inaccurate results when solving Navier-Stokes equations. Starting from [20], a new element is proposed in [8]. The space of the discrete pressures is then:  $Q_h = Q_{0,h} + Q_{1,h}$ , where  $Q_{0,h} := \mathcal{P}_{disc}^0(\mathcal{T}_h) \cap L_{zmv}^2$  and for  $k > 0$ ,  $Q_{k,h} := \mathcal{P}^k(\mathcal{T}_h) \cap L_{zmv}^2(\Omega)$ . One can build an  $L^2$ -orthogonal sum letting  $\tilde{Q}_{1,h} = \{q_h \in Q_h \mid \forall K \in \mathcal{T}_h, \int_K q_h = 0\}$ .

**Proposition 4** *We have:  $Q_h = Q_{0,h} \overset{\perp}{\oplus} \tilde{Q}_{1,h}$ . It holds for all  $q_h \in Q_h$ :  $\|q_h\|_\Omega^2 = \|\pi_0(q_h)\|_\Omega^2 + \|q_h - \pi_0(q_h)\|_\Omega^2$ .*

To prove the discrete inf-sup condition of  $\mathbf{X}_{0,h} \times Q_h$ , we need some Lemmas.

**Lemma 1** *Let  $k > 0$ ,  $(\mathbf{v}, q) \in \mathbf{X}_{0,h} \times Q_{k,h}$ . It holds:*

$$|\sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\mathbf{v} \cdot \mathbf{n}_{F,K}, q)_F| \lesssim \sigma \|\mathbf{v}\|_h \|q - \pi_0(q)\|_\Omega. \quad (9)$$

*Proof.* We start with the proof of [20, Lemma 3.1]. Using the patch-test condition, one obtains:

$$\sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\mathbf{v} \cdot \mathbf{n}_{F,K}, q)_F = \sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} ((\mathbf{v}_K - \underline{\mathbf{v}}_F) \cdot \mathbf{n}_{F,K}, q_K - \underline{q}_K)_F.$$

We have, using Eq. (4) with  $w = q_K - \underline{q}_K$  and Eq. (5) for the components of  $\mathbf{v}$ :

$$\left| ((\mathbf{v}_K - \underline{\mathbf{v}}_F) \cdot \mathbf{n}_{F,K}, q_K - \underline{q}_K)_F \right| \lesssim \sigma_K \|\underline{\mathbf{v}}_K\|_K \|q_K - \underline{q}_K\|_K.$$

Summing and using the discrete Cauchy-Schwarz inequality, we obtain (9).  $\square$

Introducing the element-wise divergence operator  $\text{div}_h$ , we prove next that one can build an explicit right inverse of  $\text{div}_h$  from  $Q_h$  to  $\mathbf{X}_{0,h}$ .

**Lemma 2** *For all  $q_h \in Q_h$ , there exists  $\mathbf{v}_{0,h} \in \mathbf{X}_{0,h}$  such that:*

$$-(\text{div}_h \mathbf{v}_{0,h}, \pi_0(q_h))_\Omega = \nu^{-1} \|\pi_0(q_h)\|_\Omega^2, \quad (10)$$

$$\|\mathbf{v}_{0,h}\|_h \leq C_{\text{div}} \nu^{-1} \|\pi_0(q_h)\|_\Omega. \quad (11)$$

*Proof.* Let  $q_h \in Q_h$ . From Proposition 1,  $\exists \mathbf{v}_{\pi_0(q_h)} \in \mathbf{H}_0^1(\Omega) \mid \operatorname{div} \mathbf{v}_{\pi_0(q_h)} = \pi_0(q_h)$  and  $\|\mathbf{v}_{\pi_0(q_h)}\|_{\mathbf{H}_0^1(\Omega)} \leq C_{\operatorname{div}} \|\pi_0(q_h)\|_\Omega$ . Let  $\mathbf{v}_{0,h} = -\nu^{-1} \Pi_h(\mathbf{v}_{\pi_0(q_h)})$ . Integrating by part twice, we get:  $\forall K \in \mathcal{T}_h, \int_K \operatorname{div} \mathbf{v}_{0,h} = -\nu^{-1} \int_K \operatorname{div} \mathbf{v}_{\pi_0(q_h)}$ , hence (10). Using Proposition 3, we get (11).  $\square$

We can also build an explicit right inverse of the discrete divergence operator from  $Q_{1,h}$  to  $\mathbf{X}_{T,h} = \{\mathbf{v}_h \in \mathbf{X}_{0,h} \mid \forall F \in \mathcal{F}_h^i, \mathbf{v}_h \cdot \mathbf{n}_F = 0\}$ , if [20, Hyp. 4.1] holds:

**Hypothesis 1.** We suppose that  $\mathcal{T}_h$  is such that each element  $K \in \mathcal{T}_h$  has at most  $d-1$  facets that lie on  $\partial\Omega$ .

**Lemma 3** Assuming Hypothesis 1,  $\forall q_{1,h} \in Q_{1,h}, \exists \mathbf{v}_{1,h} \in \mathbf{X}_{T,h}$  such that:

$$(\nabla q_{1,h}, \mathbf{v}_{1,h})_\Omega \gtrsim \nu^{-1} \sigma^{-2(d-2)} \|q_{1,h} - \pi_0(q_{1,h})\|_\Omega^2, \quad (12)$$

$$(\operatorname{div} \mathbf{v}_{1,h}, q_{0,h})_K = 0 \quad \forall q_{0,h} \in Q_{0,h}, \quad (13)$$

$$\|\mathbf{v}_{1,h}\|_h \lesssim \nu^{-1} \sigma^2 \|q_{1,h} - \pi_0(q_{1,h})\|_\Omega. \quad (14)$$

*Proof.* Let us consider  $q_{1,h} \in Q_{1,h}$ . For all  $K \in \mathcal{T}_h$ , we let  $q_K := q_{1,h|K}$  and  $\hat{q}_K = q_K \circ T_K$ . Since  $q_{1,h} \in H^1(\Omega)$  we have (see e.g. [21, Prop. 2.2.10]):  $\forall F \in \mathcal{F}_h^i \mid F = K \cap K', \nabla q_K \times \mathbf{n}_F = \nabla q_{K'} \times \mathbf{n}_F$ . We construct  $\mathbf{v}_{1,h} \in \mathbf{X}_{T,h} \mid \forall F \in \mathcal{F}_h^i, \mathbf{v}_{1,h}(M_F) = \nu^{-1} h_F^2 \mathbf{n}_F \times (\nabla q_{1,h} \times \mathbf{n}_F)$ , where  $h_F$  is the diameter of  $F$ . We have:

$$\forall K \in \mathcal{T}_h \quad (\nabla q_{1,h}, \mathbf{v}_{1,h})_K = \nu^{-1} (d+1)^{-1} |K| \sum_{F \in \mathcal{F}_K^i} h_F^2 |\nabla q_K \times \mathbf{n}_F|^2. \quad (15)$$

Let  $\mathbf{n}_{\hat{F}_{F,K}}$  be the unit normal vector of the facet  $\hat{F}_{F,K}$  outgoing from  $\hat{K}$ . We have:

$$\mathbf{n}_{F,K} = |K| (\mathbb{B}_K^{-1})^T |\hat{F}_{F,K}| |F|^{-1} \mathbf{n}_{\hat{F}_{F,K}}.$$

Consider  $d=2$ . Changing the variable, assuming Hypothesis 1, we get:

$$\sum_{F \in \mathcal{F}_K^i} h_F^2 |\nabla q_K \times \mathbf{n}_F|^2 = \sum_{F \in \mathcal{F}_K^i} |\hat{F}_{F,K}|^2 |\nabla_{\hat{x}} \hat{q}_K \times \mathbf{n}_{\hat{F}_{F,K}}|^2 \gtrsim |\nabla_{\hat{x}} \hat{q}_K|^2. \quad (16)$$

Consider  $d=3$ . Changing the variable, one now finds that  $|\nabla q_K \times \mathbf{n}_F| = |\hat{F}_{F,K}| |F|^{-1} |\mathbb{B}_K \nabla_{\hat{x}} \hat{q}_K \times \mathbf{n}_{\hat{F}_{F,K}}|$ . Hence:  $|\nabla q_K \times \mathbf{n}_F| \gtrsim |F|^{-1} \|\mathbb{B}_K^{-1}\|^{-1} |\nabla_{\hat{x}} \hat{q}_K \times \mathbf{n}_{\hat{F}_{F,K}}|$ . Notice that  $h_F |F|^{-1} \|\mathbb{B}_K^{-1}\|^{-1} \gtrsim \sigma_K^{-1}$ . Assuming Hypothesis 1, we then have:

$$\sum_{F \in \mathcal{F}_K^i} h_F^2 |\nabla q_K \times \mathbf{n}_F|^2 \gtrsim \sigma_K^{-2} |\nabla_{\hat{x}} \hat{q}_K|^2. \quad (17)$$

Using Poincaré-Steklov inequality [10, Lemma 12.11] in  $\hat{K}$ , we have:  $|\hat{K}| |\nabla_{\hat{x}} \hat{q}_K|^2 = \|\nabla_{\hat{x}} \hat{q}_K\|_{\hat{K}}^2 \gtrsim \|\hat{q}_K - \underline{q}_K\|_{\hat{K}}^2 = |K|^{-1} \|q_K - \underline{q}_K\|_K^2$  with  $\underline{q}_K = \int_K q_K / |K|$ . Using (16) and (17) in (15), it holds:

$$(\nabla q_{1,h}, \mathbf{v}_{1,h})_K \gtrsim \nu^{-1} \sigma_K^{-2(d-2)} |K| |\nabla_{\hat{x}} \hat{q}_K|^2 \gtrsim \nu^{-1} \sigma_K^{-2(d-2)} \|q_K - \underline{q}_K\|_K^2. \quad (18)$$

We obtain (12) by summation. Remark that  $\nabla \psi_{F|K} = |K|^{-1} |F| \mathbf{n}_{F,K}$ .

Let us prove (13). Recall the orthogonality  $\nabla \psi_{F|K}(M_F) \perp \mathbf{v}_{1,h}(M_F)$ , so that  $(\operatorname{div} \mathbf{v}_{1,h}, q_{0,h})_K = |K| q_{0,h}(K) \sum_{F \in \mathcal{F}_K^i} \nabla \psi_{F|K}(M_F) \cdot \mathbf{v}_{1,h}(M_F) = 0$ .

Let us prove (14). For  $K \in \mathcal{T}_h$ ,  $\underline{\underline{\mathbf{v}}}_{1,h|K} = \sum_{F \in \mathcal{F}_K^i} \mathbf{v}_{1,h}(M_F) \otimes \nabla \psi_{F|K}$ .

Hence:  $\|\underline{\nabla} \mathbf{v}_{1,h}\|_K^2 \lesssim \nu^{-2} \sum_{F \in \mathcal{F}_K^i} |F|^2 |K|^{-1} h_F^4 |\nabla q_K \times \mathbf{n}_F|^2$ . Using (3), and assuming Hypothesis 1, we get:  $\|\underline{\nabla} \mathbf{v}_{1,h}\|_K^2 \lesssim \nu^{-2} \sigma_K^2 h_K^2 |K| \sum_{F \in \mathcal{F}_K^i} |\nabla q_K \times \mathbf{n}_F|^2 \lesssim \nu^{-2} \sigma_K^2 h_K^2 \|\nabla q_K\|_K^2$ . From Eq. (6):  $\|\underline{\nabla} \mathbf{v}_{1,h}\|_K \lesssim \nu^{-1} \sigma_K^2 \|q_K - \underline{q}_K\|_K$ .  $\square$

Let  $b_h : \mathbf{X}_h \times Q_h \rightarrow \mathbb{R}$  be defined by  $b_h(\mathbf{v}_h, q_h) = -(\operatorname{div}_h \mathbf{v}_h, q_{0,h})_\Omega + (\mathbf{v}_h, \nabla q_{1,h})_\Omega$  with  $q_h = q_{0,h} + q_{1,h}$  such that  $(q_{0,h}, q_{1,h}) \in Q_{0,h} \times Q_{1,h}$ .

**Theorem 1 (Stability)** *The following continuity property holds:*

$$\forall (\mathbf{v}_h, q_h) \in \mathbf{X}_{0,h} \times Q_h, \quad |b_h(\mathbf{v}_h, q_h)| \lesssim \sigma \|\mathbf{v}_h\|_h \|q_h\|_\Omega. \quad (19)$$

Assuming Hypothesis 1, the following inf-sup condition holds:

$$\forall q_h \in Q_h, \exists \mathbf{v}_h \in \mathbf{X}_{0,h}, \quad b_h(\mathbf{v}_h, q_h) \gtrsim \frac{1}{\sqrt{2}} C_{\operatorname{div}}^{-2} \sigma^{-2d} \|\mathbf{v}_h\|_h \|q_h\|_\Omega. \quad (20)$$

*Proof.* Let us prove (19). Let  $(\mathbf{v}_h, q_h) \in \mathbf{X}_{0,h} \times Q_h$ . Integrating by parts, we have:

$$b_h(\mathbf{v}_h, q_h) = -(\operatorname{div}_h \mathbf{v}_h, \pi_0(q_h))_\Omega + \sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} (\mathbf{v}_h \cdot \mathbf{n}_{F,K}, q_{1,h})_F. \quad (21)$$

Using Cauchy-Schwarz, we have:  $|(\operatorname{div}_h \mathbf{v}_h, \pi_0(q_h))_\Omega| \leq \sqrt{d} \|\mathbf{v}_h\|_h \|\pi_0(q_h)\|_\Omega$ .

Using (9) in (21), we obtain (19) from Proposition 4. Let us prove (20), starting from the proof of [20, Lemma 4.2]. Let  $q_h \in Q_h \setminus \{0\}$ , where  $q_h = q_{0,h} + q_{1,h}$  is such that  $(q_{0,h}, q_{1,h}) \in Q_{0,h} \times Q_{1,h}$ . Let  $\mathbf{v}_h := \mathbf{v}_{0,h}$  be like in Lemma 2. Using (21) with  $\mathbf{v}_h = \mathbf{v}_{0,h}$ , (9) and (10), letting  $\tilde{q}_{1,h} = q_h - \pi_0(q_h)$ , we have:

$$b_h(\mathbf{v}_{0,h}, q_h) \gtrsim \nu^{-1} \|\pi_0(q_h)\|_\Omega^2 - \sigma \|\mathbf{v}_{0,h}\|_h \|\tilde{q}_{1,h}\|_\Omega. \quad (22)$$

Using (11) and Young inequality, we have for all  $\varepsilon > 0$ :

$$-\|\mathbf{v}_{0,h}\|_h \|\tilde{q}_{1,h}\|_\Omega \geq -\frac{1}{2} C_{\operatorname{div}} \nu^{-1} (\varepsilon \|\pi_0(q_h)\|_\Omega^2 + \varepsilon^{-1} \|\tilde{q}_{1,h}\|_\Omega^2). \quad (23)$$

We now insert the bound (23) in (22) to get:

$$b_h(\mathbf{v}_{0,h}, q_h) \gtrsim \nu^{-1} \left( (1 - \frac{\varepsilon}{2} C_{\operatorname{div}} \sigma) \|\pi_0(q_h)\|_\Omega^2 - \frac{C_{\operatorname{div}}}{2\varepsilon} \sigma \|\tilde{q}_{1,h}\|_\Omega^2 \right). \quad (24)$$

Let  $\mathbf{v}_{1,h}$  be like in Lemma 3. Using (12) and (13), we get:

$$b_h(\mathbf{v}_{1,h}, q_h) = (\nabla q_{1,h}, \mathbf{v}_{1,h})_\Omega \gtrsim \nu^{-1} \sigma^{-2(d-2)} \|\tilde{q}_{1,h}\|_\Omega^2 \quad (25)$$

Last, we set  $\mathbf{v}_h^\star = \mu \mathbf{v}_{0,h} + \mathbf{v}_{1,h}$ , where  $\mu > 0$ . Using (24) and (25), we have:

$$\begin{aligned} b_h(\mathbf{v}_h^\star, q_h) &= \mu b_h(\mathbf{v}_{0,h}, q_h) + b_h(\mathbf{v}_{1,h}, q_h), \\ &\gtrsim \nu^{-1} \left( \mu (1 - \frac{\varepsilon}{2} C_{\operatorname{div}} \sigma) \|\pi_0(q_h)\|_\Omega^2 + (\sigma^{-2(d-2)} - \frac{\mu C_{\operatorname{div}}}{2\varepsilon} \sigma) \|\tilde{q}_{1,h}\|_\Omega^2 \right). \end{aligned}$$

Let us choose  $\varepsilon = (C_{\operatorname{div}} \sigma)^{-1} < 1$  and  $\mu = \sigma^{-2(d-2)} \varepsilon^2$ . We obtain that:

$$b_h(\mathbf{v}_h^\star, q_h) \gtrsim C_{\min} \nu^{-1} \|q_h\|_\Omega^2 \text{ with } C_{\min} = \varepsilon^2 \sigma^{-2(d-2)} = C_{\text{div}}^{-2} \sigma^{-2(d-1)}. \quad (26)$$

Let us bound  $\|\mathbf{v}_h^\star\|_h$  by  $\|q_h\|_\Omega$ . We have:  $\|\mathbf{v}_h^\star\|_h^2 \lesssim (\mu^2 \|\mathbf{v}_{0,h}\|_h^2 + \|\mathbf{v}_{1,h}\|_h^2)$ . Using (11) and (14)  $\|\mathbf{v}_h^\star\|_h^2 \lesssim \nu^{-2} (\mu^2 C_{\text{div}}^2 \|\pi_0(q_h)\|_\Omega^2 + \sigma^4 \|\tilde{q}_{1,h}\|_\Omega^2)$ , hence:

$$\|\mathbf{v}_h^\star\|_h^2 \leq C_{\max}^2 \nu^{-2} \|q_h\|_\Omega^2 \text{ with } C_{\max} = \sqrt{2} \sigma^2. \quad (27)$$

Using (27) in (26), we get (20) since  $C_{\min} C_{\max}^{-1} = \frac{1}{\sqrt{2}} C_{\text{div}}^{-2} \sigma^{-2d}$ .  $\square$

Let  $\mathbf{V}_h = \{\mathbf{v}_h \in \mathbf{X}_{0,h} \mid \forall q_h \in Q_h, b_h(\mathbf{v}_h, q_h) = 0\}$ . We know from [22, Lemma 3.1] that  $\mathbf{V}_h \subset \{\mathbf{v}_h \in \mathbf{X}_{0,h} \mid \forall K \in \mathcal{T}_h, \text{div } \mathbf{v}_h|_K = 0\}$ . Using  $P^0 + P^1$  discrete pressures improves consistency. Indeed, integrating by parts, we have that for all  $(\mathbf{v}_h, q_{1,h}) \in \mathbf{V}_h \times Q_{1,h}$ ,  $0 = b_h(\mathbf{v}_h, q_{1,h}) = -(\text{div}_h \mathbf{v}, \pi_0(q_{1,h}))_\Omega + \sum_{F \in \mathcal{F}_h} \int_F q_{1,h} [\mathbf{v}_h] \cdot \mathbf{n}_F = 0 + \sum_{F \in \mathcal{F}_h} \int_F q_{1,h} [\mathbf{v}_h] \cdot \mathbf{n}_F$ . Hence:

**Lemma 4** For all  $(\mathbf{v}_h, q_{1,h}) \in \mathbf{V}_h \times Q_{1,h}$ , it holds:  $\sum_{F \in \mathcal{F}_h} \int_F q_{1,h} [\mathbf{v}_h] \cdot \mathbf{n}_F = 0$ .

Let  $\ell_f \in \mathcal{L}(\mathbf{X}_h, \mathbb{R})$  be such that  $\forall \mathbf{v}_h \in \mathbf{X}_h$ ,  $\ell_f(\mathbf{v}_h) = (\mathbf{f}, \mathbf{v}_h)_\Omega$  if  $\mathbf{f} \in \mathbf{L}^2(\Omega)$ ,  $\ell_f(\mathbf{v}_h) = \langle \mathbf{f}, I_h(\mathbf{v}_h) \rangle_{\mathbf{H}_0^1(\Omega)}$  if  $\mathbf{f} \notin \mathbf{L}^2(\Omega)$ , where  $I_h : \mathbf{X}_{0,h} \rightarrow \mathbf{Y}_{0,h}$  is for instance an averaging operator [10, §22.4.1], with  $\mathbf{Y}_{0,h} = \{\mathbf{v}_h \in \mathbf{H}_0^1(\Omega) \mid \forall K \in \mathcal{T}_h, \mathbf{v}_h|_K \in \mathbf{P}^k(K)\}$ . The discretization of Problem (1) with the  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  element reads: Find  $(\mathbf{u}_h, p_h) \in \mathbf{X}_{0,h} \times Q_h$  such that  $\forall (\mathbf{u}_h, p_h) \in \mathbf{X}_{0,h} \times Q_h$

$$\nu(\mathbf{u}_h, \mathbf{v}_h)_h + b_h(\mathbf{v}_h, p_h) + b_h(\mathbf{u}_h, q_h) = \ell_f(\mathbf{v}_h). \quad (28)$$

Due to Proposition 2 and Theorem 1, Problem (28) is well posed.

### 3 Convergence Of The $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$ Element

Following the proofs of [1, Theorems 3, 4, 6], we can prove

**Theorem 2 (Convergence)** Suppose that Hypothesis 1 holds and that the solution of Problem (1) is such that  $(\mathbf{u}, p) \in (\mathbf{V} \cap \mathbf{H}^2(\Omega)) \times (H^1(\Omega) \cap L_{zmv}^2(\Omega))$ . It holds:

$$\|\mathbf{u} - \mathbf{u}_h\|_h \lesssim \sigma h \nu^{-1} \|\mathbf{f}\|_\Omega. \quad (29)$$

$$\text{If } \Omega \text{ is convex} \quad \|\mathbf{u} - \mathbf{u}_h\|_\Omega \lesssim \sigma^2 h^2 \nu^{-1} \|\mathbf{f}\|_\Omega. \quad (30)$$

$$\|(p - p_h) - \pi_0(p - p_h)\|_\Omega \lesssim \nu \sigma^{(2d-1)} h \|\mathbf{f}\|_\Omega. \quad (31)$$

$$\|\pi_0(p - p_h)\|_\Omega \lesssim \nu C_{\text{div}} \sigma^{(2d+1)} h \|\mathbf{f}\|_\Omega. \quad (32)$$

In the case of inhomogeneous Dirichlet boundary conditions, one can also recover (29) and (30), following [1, §7], where a lifting of  $\mathbf{g}$  is used.

When  $\Omega \subset \mathbb{R}^2$ , we have moreover the two Lemmas below.

**Lemma 5** For all  $(\mathbf{v}_h, q_h) \in \mathbf{V}_h \times Q_{2,h}$ ,  $(\nabla q_h, \mathbf{v}_h)_\Omega = 0$ .

*Proof.* Let  $(\mathbf{v}_h, q_h) \in \mathbf{V}_h \times Q_{2,h}$ . Integrating by parts and using Lemma 4, we get that  $\forall q_{1,h} \in Q_{1,h}$ :  $(\nabla q_h, \mathbf{v}_h)_\Omega = -\sum_{F \in \mathcal{F}_h} \int_F q_h [\mathbf{v}_h] \cdot \mathbf{n}_F = -\sum_{F \in \mathcal{F}_h} \int_F (q_h - q_{1,h}) [\mathbf{v}_h] \cdot \mathbf{n}_F = \sum_{F \in \mathcal{F}_h} \int_F (q_h - q_{1,h}) [\mathbf{v}_h - \mathbf{v}_h(M_F)] \cdot \mathbf{n}_F$ . Hence, we obtain:  $(\nabla q_h, \mathbf{v}_h)_\Omega = -\sum_{K \in \mathcal{T}_h} \sum_{F \subset \partial K} \int_F (q_h - q_{1,h})(\mathbf{v}_h - \mathbf{v}_h(M_F)) \cdot \mathbf{n}_{F,K}$ . Choose  $q_{1,h}$  such that  $\forall S \in \mathcal{V}_h$ ,  $q_{1,h}(S) = q_{2,h}(S)$ . Then,  $\forall F \in \mathcal{F}_h$ , the degree 3 polynomial  $(q_h - q_{1,h})(\mathbf{v}_h - \mathbf{v}_h(M_F))$  vanishes at the quadrature points of Simpson's rule. Since it is exact for degree 3 polynomials,  $\forall F \in \mathcal{F}_h$ ,  $\int_F (q_h - q_{1,h})(\mathbf{v}_h - \mathbf{v}_h(M_F)) \cdot \mathbf{n}_{F,K} = 0$  and  $(\nabla q_h, \mathbf{v}_h)_\Omega = 0$ .  $\square$

**Lemma 6** *Let  $\phi \in H^3(\Omega) \cap L^2_{zmv}(\Omega)$  such that  $|\phi|_{H^3(\Omega)} \neq 0$ .*

*Then,  $\forall \mathbf{v}_h \in \mathbf{V}_h$ ,  $(\nabla \phi, \mathbf{v}_h)_\Omega \lesssim \sigma h^2 |\phi|_{H^3(\Omega)} \|\mathbf{v}_h\|_\Omega$ .*

*Proof.* Let  $\mathbf{v}_h \in \mathbf{V}_h$ . Using Lemma 5 and Cauchy-Schwarz, it holds:  $|(\nabla \phi, \mathbf{v}_h)_\Omega| = |(\nabla(\phi - q_h), \mathbf{v}_h)_\Omega|$  for all  $q_h \in Q_{2,h}$ . We then use [10, Lemma 11.9] to conclude.  $\square$

**Theorem 3 (Pressure Robustness)** *Suppose that  $\Omega$  is convex. Let  $p \in H^3(\Omega) \cap L^2_{zmv}(\Omega)$  such that  $|p|_{H^3(\Omega)} \neq 0$ . Consider Problem (1) with  $\mathbf{f} = \nabla p$ . Then  $(\mathbf{u}_h, p_h)$ , the solution of Problem (28) is such that (where  $\delta p = p - p_h$ ):*

$$\nu \|\mathbf{u}_h\|_h \lesssim \sigma^2 h^3 |p|_{H^3(\Omega)}, \quad \nu \|\mathbf{u}_h\|_\Omega \lesssim \sigma^3 h^4 |p|_{H^3(\Omega)}. \quad (33)$$


$$\|\delta p - \pi_0(\delta p)\|_\Omega \lesssim \sigma^4 h^2 |p|_{H^2(\Omega)}, \quad \|\pi_0(\delta p)\|_\Omega \lesssim C_{\text{div}} \sigma^6 h^2 |p|_{H^2(\Omega)}. \quad (34)$$

*Proof.* Setting  $\mathbf{v}_h = \mathbf{u}_h$  as test-function in Problem (28) and using Lemma 6, it holds:  $\nu \|\mathbf{u}_h\|_h^2 = (\nabla \phi, \mathbf{u}_h)_\Omega \lesssim \sigma h^2 |\phi|_{H^3(\Omega)} \|\mathbf{u}_h\|_\Omega$ . From [1, Theorem 4], we have:  $\|\mathbf{u}_h\|_\Omega \lesssim \sigma h \|\mathbf{u}_h\|_h$ . We deduce that  $\nu \|\mathbf{u}_h\|_h \lesssim \sigma^2 h^3 |\phi|_{H^3(\Omega)}$ . Using this estimate, we get that  $\nu \|\mathbf{u}_h\|_\Omega \lesssim \sigma^3 h^4 |\phi|_{H^3(\Omega)}$ . The proof of (34) is similar to that of Eqs (31)-(32), using (33) and noticing that  $\forall K \in \mathcal{T}_h$ ,  $\|\nabla(\phi - \pi_1 \phi)\|_K \lesssim \sigma_K h_K |\phi|_{H^2(K)}$ .  $\square$

## 4 Numerical Results

To get  $p_h \in L^2_{zmv}(\Omega)$ , we eliminate a degree of freedom in  $\mathcal{P}_{disc}^0(\mathcal{T}_h)$  and in  $\mathcal{P}^1(\mathcal{T}_h)$  and post-process by subtracting the mean-value. Let us set:  $\|(\mathbf{u}, p)\|_\nu^2 := \|\nabla \mathbf{u}\|_\Omega^2 + \nu^{-2} \|p\|_\Omega^2$ . Let  $\delta_h \mathbf{u} = \Pi_h \mathbf{u} - \mathbf{u}_h$ ,  $\delta p = p - p_h$  and  $\pi_{disc}^1$  be the  $L^2$ -orthogonal projection onto  $\mathcal{P}_{disc}^1(\mathcal{T}_h)$ . We compute the discrete errors values:  $\varepsilon_0^\nu(\mathbf{u}_h) := \|\delta_h \mathbf{u}\|_\Omega / \|(\mathbf{u}, p)\|_\nu$ ,  $\varepsilon_1^\nu(\mathbf{u}_h) := \|\delta_h \mathbf{u}\|_h / \|(\mathbf{u}, p)\|_\nu$  and  $\varepsilon_0^\nu(p_h) := \nu^{-1} (\|\pi_0(\delta p)\|_\Omega^2 + \|\pi_{disc}^1(\delta p) - \pi_0(\delta p)\|_\Omega^2)^{\frac{1}{2}} / \|(\mathbf{u}, p)\|_\nu$ . Let  $\tau_{0,\mathbf{u}}$ ,  $\tau_{1,\mathbf{u}}$  and  $\tau_p$  be the averaged convergence rates of  $\varepsilon_0^\nu(\mathbf{u})$ ,  $\varepsilon_1^\nu(\mathbf{u})$  and  $\varepsilon_0^\nu(p)$ .

We first consider a 2D case illustrating Thm. 3, with the prescribed solution  $\mathbf{u} = 0$ ,  $\mathbf{f} = \nabla \psi$  with  $\psi(x, y) = \exp(-10(1 - x + 2y))$ . Fig. 1 shows  $\varepsilon_0^\nu(\mathbf{u}_h)$ ,  $\varepsilon_1^\nu(\mathbf{u}_h)$  and  $\varepsilon_0^\nu(p_h)$  against  $h$ , for  $\nu = 10^{-2}$ , with  $P^0$  (left) or  $P^0 + P^1$  (right) discrete pressures, comparing the  $\mathbf{P}_{nc}^1 \times P^0$  [1, Example 4] and  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  schemes. Tab. 1 reports observed convergence rates, which are better for the  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  scheme.

**Fig. 1** 2D case. Plots of  $\varepsilon_0^\nu(\mathbf{u}_h)$ ,  $\varepsilon_1^\nu(\mathbf{u}_h)$  and  $\varepsilon_0^\nu(p_h)$  for  $\nu = 10^{-2}$ .**Table 1** 2D case, averaged convergence rates on the last five meshes.

$P^0   P^0 + P^1  $	$P^0   P^0 + P^1  $	$P^0   P^0 + P^1  $			
$\tau_{0,\mathbf{u}} 2.02$	$4.03$	$\tau_{1,\mathbf{u}} 1.01$	$3.07$	$\tau_p 1.60$	$2.10$

Let us consider stationary Navier-Stokes equations set in  $\Omega = (0, 1)^3$ : Find  $(\mathbf{u}, p)$  such that  $-\nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = \mathbf{f}$  and  $\operatorname{div} \mathbf{u} = 0$  in  $\Omega$ , with inhomogeneous Dirichlet boundary conditions. Computations are done with the TrioCFD code [6, 7] with the  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  scheme. The convection term is discretized using the MUSCL scheme [6]. We study the prescribed solution  $\mathbf{u} = (y - z, z - x, x - y)^T$ ,  $p = \frac{1}{2}(x^2 + y^2 + z^2) - xy - xz - yz - 1/4$  so that  $\mathbf{f} = 0$ . Fig. 2 shows  $\varepsilon_0^\nu(\mathbf{u}_h)$  (left) and  $\varepsilon_0^\nu(p_h)$  (right) against  $h$ , for  $\nu = 10^{-1}$ ;  $10^{-2}$ ;  $10^{-3}$ .


**Fig. 2** 3D case. Plots of  $\varepsilon_0^\nu(\mathbf{u}_h)$  and  $\varepsilon_0^\nu(p_h)$  for  $\nu = 10^{-1}$ ,  $\nu = 10^{-2}$  or  $\nu = 10^{-3}$ .

Table 2 reports the convergence rates  $\tau_{0,\mathbf{u}}$  (left), and  $\tau_p$  (right). More numerical results are available in [8, 9, 6, 23]. The  $\mathbf{P}_{nc}^1 \times (P^0 + P^1)$  scheme shows good approximation properties and should therefore be suitable for solving the Navier-Stokes

**Table 2** 3D case, averaged convergence rates on the last three meshes.

$\frac{\nu}{\tau_{0,u}}$	$10^{-1}$	$ 10^{-2} $	$ 10^{-3} $	$\frac{\nu}{\tau_p}$	$10^{-1}$	$ 10^{-2} $	$ 10^{-3} $
	1.68	1.88	1.41		0.70	1.72	1.35

equations in the context of industrial simulation. More generally, it seems interesting to design and study discrete pressure enrichment for higher order nonconforming Crouzeix-Raviart mixed FEM [2, 3].

**Acknowledgements** We would like to thank P.-E. Angeli for his contribution to §4.

## References

1. Crouzeix M., Raviart, P.-A.: *RAIRO, Sér. Anal. Numer.*, **7**, 33, 33–75 (1973).
2. Sauter, S.: *Computers & Mathematics with Applications*, **149**, 49–70 (2023).
3. Sauter, S., Torres, C.: *Mathematics of Computation*, **92**, 341, 1033–1059 (2023).
4. Hecht, F.: *RAIRO, Sér. Anal. Numér.* **15**, 2, 119–150 (1981).
5. Linke, A.: *Comput. Methods Appl. Mech. Engrg.* **268**, 782–800 (2014).
6. Angeli, P.-E. et al: FVCA8 benchmark for the Stokes and Navier-Stokes equations with the TrioCFD code, in: Cancès, C., Omnes, P. (eds.) *Finite Volumes for Complex Applications VIII - Methods and Theoretical Aspects*, pp. 181–302, Springer (2017).
7. CEA/DM2S, TRUST-TrioCFD code, <https://trioefd.cea.fr/>.
8. Heib, S.: *Nouvelles discréétisations non structurées pour des écoulements de fluides à incompressibilité renforcée*, PhD thesis, Univ. Pierre et Marie Curie - Paris VI, France (2003).
9. Fortin, T.: *Une méthode éléments finis à décomposition  $L^2$  d'ordre élevé motivée par la simulation d'écoulement diphasique bas Mach*, Univ. Pierre et Marie Curie - Paris VI (2006).
10. Ern, A., Guermond, J.-L.: *Finite elements I*, Springer (2021).
11. Ern, A., Guermond, J.-L.: *Finite elements II*, Springer (2021).
12. Warburton, T., Hesthaven, J.S.: *Comput. Methods Appl. Mech. Engrg.*, **192**, 25, 2765–2773 (2003).
13. Amrouche, C., Ciarlet, P. G., Mardare, C.: *Journal de Mathématiques Pures et Appliquées*, **104**, 2, 207–226 (2015).
14. Bonnet-Ben Dhia, A.-S., Ciarlet Jr., P.: Méthodes variationnelles pour l’analyse de problèmes non coercifs, *M.Sc. AMS lecture notes (ENSTA-IPP)* (2023).
15. Barré, M., Ciarlet Jr., P.: The T-coercivity approach for mixed problems, *C. R. Acad. Sci. Paris, Ser. I*, to appear, <https://univ-rennes2.hal.science/INRIA/hal-03820910>.
16. Jamelot, E.: Improved stability estimates for solving Stokes problem with Fortin-Soulie finite elements, submitted, <https://cea.hal.science/cea-03833616v3>.
17. Apel, T., Nicaise, S., Schöberl, J.: *Numerische Mathematik* **89**, 2, 193–223 (2001).
18. Emonot, P.: *Méthode de volumes éléments finis : applications aux équations de Navier-Stokes et résultats de convergence*, PhD thesis, Université Claude Bernard - Lyon I, France (1992).
19. John, V., Linke, A., Merdon, C., Neilan M., Rebholz, L.G.: *SIAM REVIEW*, **59**, 3, 492–544 (2017).
20. Bernardi, C., Hecht F.: *ESAIM: M2AN*, **34**, 5, 953–980 (2000).
21. Assous, F., Ciarlet Jr., P., Labrunie, S.: Mathematical Foundations of Computational Electromagnetism, Springer (2018).
22. Burman, E., Hansbo, P.: *Numer. Methods Partial Differ. Equ.*, **21**, 5, 986–997 (2005).
23. Gounand, S.: <https://cea.hal.science/cea-02489502/document> (2015).