



HAL
open science

Cyclic Diaryl λ 3 -Chloranes: Reagents and Their C–C and C–O Couplings with Phenols via Aryne Intermediates

Matteo Lanzi, Torben Rogge, Tan Sang Truong, K N Houk, Joanna Wencel-Delord

► **To cite this version:**

Matteo Lanzi, Torben Rogge, Tan Sang Truong, K N Houk, Joanna Wencel-Delord. Cyclic Diaryl λ 3 -Chloranes: Reagents and Their C–C and C–O Couplings with Phenols via Aryne Intermediates. *Journal of the American Chemical Society*, 2022, 145 (1), pp.345-358. 10.1021/jacs.2c10090 . hal-04415921

HAL Id: hal-04415921

<https://hal.science/hal-04415921>

Submitted on 24 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cyclic Diaryl λ^3 -chloranes: reagents and their C-C and C-O couplings with phenols *via* aryne intermediates

Matteo Lanzi,^a Torben Rogge,^b Tan Sang Truong,^a K. N. Houk,^{*b} and Joanna Wencel-Delord^{*a}

^a Laboratoire d'Innovation Moléculaire et Applications (UMR CNRS 7042), Université de Strasbourg/Université de Haute Alsace, ECPM, 67087 Strasbourg, France

^b Department of Chemistry and Biochemistry, University of California, Los Angeles, CA 90095, USA

KEYWORDS: Hypervalent chlorine, cross-coupling, direct arylation, metal-free coupling

ABSTRACT: Hypervalent chloranes are a class of rare and poorly explored reagents. Their unique electronic properties confer them reactivity that is complementary to that of the common iodanes and emerging bromanes. Highly chemo- and regio-selective, metal-free and mild C-C and C-O couplings are reported here. Experimental and computational mechanistic studies elucidate the unprecedented reactivities and selectivities of these systems and the intermediacy of aryne intermediates. The synthetic potential of these transformations is further demonstrated *via* post-functionalization of C-C and C-O coupling products obtained from reactions of chloranes with phenols under different conditions.

Introduction. Hypervalent compounds are nowadays fundamental tools in organic chemistry,^{1,2} both in academia and industry, serving as oxidants, electrophiles, and radical initiators.³⁻⁷ Widely investigated, the λ^3 -iodanes represent the most common hypervalent reagents. On the contrary, their isoelectronic bromine⁸⁻¹¹ and chlorine¹²⁻¹⁴ congeners have remained unexplored for decades as a consequence of a lack of a general and efficient synthesis.¹⁵ We recently reported a safe and practical method for the preparation of cyclic diaryl λ^3 -bromanes.¹⁶ The enhanced nucleofugality and strong electron-withdrawing property of λ^3 -bromanes arising from the higher ionization potential of Br confer to these compounds a unique reactivity (**Figure 1a**). Therefore, cyclic diaryl bromanes serve as original aryne precursors (**Figure 1b**)^{16,17} and halogen-bond organo-catalysts.^{18,19} In clear contrast, cyclic diaryl iodines undergo transition metal catalyzed *ortho*-functionalization,²⁰ clearly illustrating the differences between the cyclic diaryl λ^3 -bromanes and the corresponding iodanes.

Considering the higher ionization potential and electronegativity of Cl, translating into a decreased positive charge in hypervalent λ^3 -chloranes, an amplified reactivity of diaryl λ^3 -chloranes as aryne precursors could be anticipated.^{12,21} Such a superior reactivity of Cl(III)-compounds, in comparison with Br(III)-species, is expected to translate into unprecedented reactivity under metal-free conditions. Challenging synthesis of such compounds remains, however, a major impediment in uncovering this reactivity. A general synthetic route furnishing such compounds followed by demonstration of their chemical behavior would though open new vistas in the chemistry of emerging hypervalent compounds.




Figure 1: a) Electronic properties of iodine, bromine and chlorine; b) Reactivity of cyclic diaryl λ^3 -iodonium and bromonium salts; c) Common C-O and C-C couplings of phenols; d) Selected examples of biologically active terphenyls; e) This work


Due to the prevalence of the phenol motif in medicinal chemistry, agrochemistry, and natural products,^{22–24} the development of site-selective and transition metal-free methodologies for their late-stage functionalization is highly desirable.²⁵ In particular, cross-dehydrogenative coupling (CDC)-type reactions rapidly expanding molecular complexity on the phenol scaffolds have attracted considerable attention over the last years (**Figure 1c**).^{26,27} Various approaches towards O-arylation *via* cross-couplings of phenols²⁸ have been disclosed, including C-O coupling between phenols and arynes,^{29–31} and direct arylation with diaryl iodonium salts.⁵ In clear contrast, *ortho*-C-functionalization of phenols *via* cross-couplings are less developed and mainly limited to electrochemical transformations,³² oxidative couplings, and metal-catalyzed reactions.²⁷ On the contrary, the direct C-arylation of phenols with arynes under transition metal-free and mild conditions remains unprecedented.^{33,34} Consequently, considering the expected unusual reactivity of λ^3 -chloranes, combined with the interest in the designing chemoselective C-O vs. C-C cross-couplings of phenols, and diversity of the terphenyl compound in natural and biologically active products (**Figure 1d**),³⁵ we have embarked on exploring the reactivity of chloranes with phenols.

We report herein: 1) the synthesis and characterization of a large panel of cyclic diaryl λ^3 -chloranes; 2) their amplified reactivity as aryne precursors as illustrated in metal-free chemoselective C-O coupling; 3) previously unachievable site-selective, metal-free C-C couplings (**Figure 1e**). Noteworthy, the selectivity issues associated with this transformation are multifaceted and go far beyond the C-O vs. C-C chemoselectivity. Aiming synthetically useful protocols, control of *ortho*/*meta*-functionalization should be reached, together with the site-selectivity, relating to the selective decoration of only one aromatic ring in case of unsymmetrical hypervalent chloranes. Moreover, mechanistic investigations and density functional theory (DFT) calculations rationalized the superior reactivity of chloranes, the solvent-dependent chemoselectivity of C-O vs. C-C couplings, as well as the preferential *meta*-functionalization.

RESULTS AND DISCUSSION

Synthesis of cyclic diaryl λ^3 -chloranes. We began our investigation by developing a simple and efficient approach for the preparation of cyclic diaryl λ^3 -chloranes **2**. Encouraged by our previously described synthetic route to cyclic diaryl λ^3 -bromanes, we hypothesized that similar mild reaction conditions, involving *t*BuONO as organic oxidant and a Brønsted acid could enable the preparation of **2** starting from a simple 1,1'-chloro(amino)biphenyl **1**³⁶ (**Scheme 1**). Remarkably, an aqueous solution of HBF₄ assented to isolate **2a-BF₄** in 79% yield through simple precipitation with diethyl ether. In analogy, the HPF₆, MsOH and TfOH acids enabled the preparation of **2a** in moderate to good yields (52 – 63%). Subsequently, the generality of this method was evidenced while engaging various 1,1'-chloro(amino)biphenyls **1a-z**. A large variety of cyclic diaryl λ^3 -chloranes **2a-z** was thus furnished in excellent to moderate yields. Electron-donating groups such as methyl, *tert*-butyl and methoxy were well tolerated at the 1-2-, 3- and 4-positions, respectively, delivering the desired products **2b-k** in good to excellent yields (46 – 81%). Substrates bearing various electron-withdrawing substituents, including chlorine, methyl and benzyl ester as well as methyl-ketone were also compatible, further illustrating an interesting functional group tolerance. Notably,

Scheme 1: Scope of cyclic diaryl λ^3 -chloranes



even a radical sensitive -CN moiety could be installed (**2l**), furnishing **2l** in synthetically useful 47% yield. Selected unsymmetric substrates (**1q – 1z**) were successfully submitted to the standard reaction conditions furnishing highly decorated cyclic diaryl λ^3 -chloranes. Particularly interesting are the “push-pull” λ^3 -chloranes, **2q**, **2t** and **2x**, bearing an electron-donating (Me- and MeO-) and electron withdrawing groups (MeOOC- and Cl-) which were obtained in excellent yields. The robustness and usefulness of the procedure were furthermore demonstrated by performing the synthesis of **2d**, **2w**, **2x** and **2z** at gram-scale level (1.2 or 2.2 mmol scale).

Single crystal X-ray analysis of **2a-BF₄** and **2w** confirmed the tricyclic structure of the hypervalent cyclic diaryl λ^3 -chloranes. Considering the scarcity of the hypervalent chlorine-type compounds and targeting their comparison with the corresponding bromanes and iodanes congeners, the crystallographic structures of these hypervalent compounds have been examined (**Figure 2**). The crystallographic data for the compounds **2a-BF₄**, **3a-OMs** and **4a-OMs** revealed a peculiar structural trend. The bond length between the carbon and the hypervalent halogen atom increases in the order Cl(III) < Br(III) < I(III). Accordingly, a significant enlargement of the C-X-C angle is observed, translating into an almost perfect T-shape structure for the chlorine compound (**4a-OMs**, 81.9°; **3a**, 86.9°; **2a-BF₄**, 92.0°), establishing the λ^3 -hypervalent compounds three-center-four-electron bond (3c-4e) feature.




Figure 2: X-ray structure analysis of **2a-BF₄** (CCDC 2176956), **3a-OMs** (CCDC 2063898) and **4a-OMs** (CCDC 2063903)

Reactivity. With the simple and scalable protocol delivering a large family of cyclic diaryl λ^3 -chloranes **2** in hand, we embarked on exploring their reactivity in sustainable, transition metal-free couplings. Aiming at the development of straightforward and site-selective diversification strategies of phenols we began our study by investigating the reaction of hypervalent λ^3 -iodane (**4a-OTf**), λ^3 -bromane (**3a-OTf**), and λ^3 -chlorane (**2a-OTf**) with 4-methylphenol **5a** in the presence of a weak base, such as cesium or potassium carbonate (Table 1).

Table 1: Reactivity of cyclic diaryl hypervalent λ^3 -reagents

X(III)-OTf	Conditions ^[a]	Yields ^[b]	
		C-C	C-O
1. 4a-OTf		No conversion	
2. 3a-OTf	Cs ₂ CO ₃ ; CHCl ₃	15%	70%
3. 2a-OTf		35%	35%
4. 4a-OTf		No conversion	
5. 3a-OTf	K ₂ CO ₃ ; CHCl ₃	26%	9%
6. 2a-OTf		49%	Traces
7. 4a-OTf		No conversion	
8. 3a-OTf	K ₂ CO ₃ ; H ₂ O	No conversion	
9. 2a-OTf		13%	60%

^[a] Conditions: 0.1 mmol of X(III)-OTf reagent (**4a-OTf**, **3a-OTf**, **2a-OTf**), 0.15 mmol of **5a**, and 0.3 mmol of Cs₂CO₃ or K₂CO₃ in 1 mL of CHCl₃ or H₂O at room temperature for 16h. ^[b] Isolated yields.


While no reaction occurred using the hypervalent λ^3 -iodane compounds (Table 1, entries 1, 4 and 7), the λ^3 -bromane and λ^3 -

chlorane substrates reacted smoothly. Encouragingly, in addition to the expected *meta*-selective C-O coupling, metal-free dehydrogenative *meta*-selective C-C bond formation was also observed, delivering product **6a** in a significant amount while using **2a-OTf**. Remarkably, the nature of the base and the solvent appeared to have a crucial impact on the C-O vs. C-C chemoselectivity. Indeed, the hypervalent bromide **3a-OTf** in the presence of cesium carbonate provided the C-O coupling product **9a** in 70% yield (Table 1, entry 2). In clear contrast, under the same reaction conditions **2a-OTf** showed a complementary reactivity, favoring the formation of the challenging C-C coupling product (**6**), together with the C-O product (**7**) (Table 1, entry 3). The simple replacement of the Cs₂CO₃ base by the potassium salt almost completely inhibited the formation of the ether-derived product, furnishing selectively the direct arylation product, **6a**, in 49% yield (Table 1, entry 6). The use of water turned out to be an additional handle to control the reaction outcome, as in the aqueous medium the reaction between **2a-OTf** and **5a** provided **7a** in 60% isolated yield (Table 1, entry 9), while no coupling occurred when using the hypervalent iodine **4a-OTf** and bromane **3a-OTf** (Table 1, entries 7 and 8), further highlighting the complementary and potent reactivity of these reagents. Additional control experiments were carried out to elucidate the nature and the role of the base and the solvent (see the SI, Table S-3) identifying potassium carbonate as crucial to avoid the concomitant production of both C-C and C-O coupling products whereas the choice of the solvent gives a unique handle to reach high chemoselectivity. Noteworthy, in the case of both transformations the generation of the *meta*-substituted products is largely favored and only a trace amount of the *ortho*-coupling product is observed (see the SI, Table S-2).

Scope of C-C and C-O couplings. With the optimal reaction conditions in hand, we first investigated the generality of the metal-free dehydrogenative *ortho*-arylation of phenols. (Scheme 2). In all cases, very high *meta*-selectivity has been observed with regard to the biaryl unit of the hypervalent chlorine precursor. Alkyl substituents (Me, *t*Bu and Et) at the *para*-positions were well tolerated providing **6a**, **6b**, and **6c** in moderate to good yields. Functionalization of the natural product chavicol (**5d**) afforded **6d** in 64% yield, illustrating compatibility with a synthetically useful allyl moiety. Functional groups such as iodo (**5e**) and methyl ester (**5f**) derivatives were tolerated under these mild reaction conditions, albeit **6e** was obtained only in 30% yield. Considering the special role of biosteres in medicinal chemistry,³⁷ the 4-adamantyl phenol **5g** was also investigated providing the desired **6g** in 72% yield whereas **5h** leads to **6h** in moderate amount. Protected 1,4-dihydroxyquinones **5i** and **5j** conveyed respectively **6i** and **6j** in synthetically useful yields. Couplings with sensitive moieties such as propargyl **5k**, allyl **5l** and cyclopropyl **5m** derivatives occurred smoothly, yielding highly valuable products. Disubstituted phenol **5n** furnished a 1:1 mixture of regioisomers of **6n**. Interestingly, the introduction of an *ortho*-substituent on the phenol (**5o**) does not affect the reaction yield. Furthermore, a remarkable *ortho*-regioselectivity (with respect to the OH group of the phenol) has been achieved while investigating unsymmetric and sterically hindered substrates such as **5p**, **5q**, **5r** and **5s**, delivering the desired products as single regioisomers in high yields (up to 76% yield). Finally, the arylation of the phenol **5t** and 2-naphthol **5u** supplied regioisomerically pure **6t** and **6u**.

Intrigued by the discovery of a solvent-dependent chemoselectivity, we then focused our attention on the development of a water mediated metal-free C-O coupling (**Scheme 3a**). The versatility and the robustness of this approach were validated regarding the compatibility of this reaction with a large diversity of *ortho*-, *meta*- and *para*-functionalized substrates **5**. Synthetically useful functional groups, including nitro **5v**, iodo **5z**, methyl ketone **5x** as well as cyanide **5c**, were well tolerated furnishing the corresponding products in good yields. Phenols bearing electron withdrawing-groups, such as, **5w** and **5y**, out-competed the electron rich coupling partners, due to the acidity. Remarkably, the C-O arylation of the challenging sterically hindered phenol **5h** was still possible, albeit **7h** was isolated in moderate yield. Moreover, L-tyrosine derivative **5i** was submitted to the standard reaction conditions and a good yield of the arylated product **7i** was obtained, further illustrating the high versatility of the method.


Scheme 2: Scope of metal-free C-C bond formation with cyclic diaryl λ^3 -chloranes



Cyclic diaryl hypervalent λ^3 -bromane **3a-OTf** was also briefly tested for the metal-free carbon-oxygen bond formation (**Scheme 3b**). Due to the lower reactivity of Br(III)-species under the standard procedure, an alternative protocol involving

$CHCl_3$ as solvent and cesium carbonate as base, was identified as suitable for various electron-rich and electron-poor phenols which delivered products **9** in good to excellent yields.


Scheme 3: Scope of metal free C-O coupling; a) with cyclic diaryl λ^3 -chloranes; b) with cyclic diaryl λ^3 -bromanes



The complexity of the coupling between diaryl λ^3 -chloranes and phenols does not only rely on the chemoselectivity between C-O and C-C coupling, but an additional degree of selectivity arises from 1) a possible functionalization of the biaryl unit of the λ^3 -chloranes either in *ortho*- or *meta*-positions and 2) selective functionalization of one ring over the other in case of unsymmetrical hypervalent substrates. While using the simple **2a-OTf** substrate, *meta*-selective coupling is highly favored (*m/o* selectivity from 99:1 to 88:12). To further investigate the *meta/ortho*-regioselectivity trends, the reactivity of unsymmetric cyclic diaryl λ^3 -chloranes were tested (**Scheme 4**). Under the C-O coupling protocol, the 2-methyl substituted chloranes **2d** and **2w** were efficiently reacted with the phenol **5w**, delivering selectively *meta*-substituted products **7dw** and **7ww** respectively in 74% and 71% isolated yields. The presence of an electron-withdrawing group at the 3-position, such as methyl ester group, (**2z**), provided **7zw** in good yield but a regioselectivity-switch occurred, translating into formation of the *ortho*-regioisomer as the major product (*m/o* 31:69). On the other hand, the impact of the ester substituent, when introduced at the 2-position, is less pronounced, thus generating the *meta*-product **7yw** as the major regioisomer (*m/o* 68:32). In addition, the unsymmetric λ^3 -chloranes prone to undergo the functionalization at

both, different biaryl aromatic units, tend to be functionalized at the less electron-rich aryl motif.^{38,39} Indeed, OMe- substituted chlorane **2k** delivered **7kw** as a mixture of two products in *a:b* regioselectivity ratio 73:27. Impressively, a perfect regioselectivity was reached by designing a “push-pull”-type substrate. In case of **2t** bearing one electron-rich and one electron-withdrawing substituent a complete regioselective-control was achieved while reacting with **5w**. Finally, we extended the regioselectivity studies taking into consideration the effect of different phenols on the transformation with **2w** and **2b** into the C-C bond formation. Sterically hindered phenols **5u** and **5b** provided **6wu** and **6wb** with a similar regioselectivity ratio of 87:13 *m:o*, while a methyl group **5a** furnished **6wa** in 59% yield, albeit with a marginally reduced regioselectivity.

Scheme 4: Unsymmetric cyclic diaryl λ^3 -chloranes in C-O and C-C bond formations



In order to demonstrate even further the sustainability of this methodology, the C-O coupling was performed under flow protocol. Reaction of **2a**-OTf and the phenol **5a** delivered the desired product **7a** in 72% isolated yield after a simple liquid-liquid extraction and filtration through silica gel. Remarkably, a drastic reduction of the reaction time (13.3 min of residence time vs. 16 hours) and an improved chemoselectivity were reached under flow chemistry conditions (see the SI, **Scheme S-12**).

Mechanistic Studies. Captivated by the singular reactivity of this class of cyclic diaryl λ^3 -chloranes, mechanistic studies were undertaken, combining both, experimental and DFT approaches.


Experimental studies: Generation of aryne intermediate: Firstly, competition tests between λ^3 -chloranes and λ^3 -bromanes, in both C-C and C-O coupling, were performed (see the SI, **Scheme S-14** and **Scheme S-15**). Remarkably, C-C coupling is twice as efficient for λ^3 -chlorane **2a** than for λ^3 -bromane **3a**. Even more impressive is the almost 5 times increased reactivity observed for **2a** in the C-O coupling reaction. Following this

trend, in case of the parallel reactions conducted with **2a**-OTf and **3a**-OTf, the bromine substrate provided the desired products **8b** and **9w** in only low yields, with 2-bromo-biphenyl being a major decomposition product. On the contrary, **2a**-OTf furnished the desired product in good yields, proving the higher and complementary reactivity of the latter.

Subsequently, deuteration studies were conducted (**Scheme 5a**). C-O coupling between **2a**-OTf and **5w**, conducted in deuterated solvents, furnished **7w**-[D₁] in good yield with high deuterium incorporation (up to 95%) at the *ortho*-position. This result perfectly validates the expected formation of the aryne intermediate and the *meta*-functionalization. Upon a nucleophilic attack, a carbanion intermediate is formed which can be easily trapped in the presence of [D⁺]-sources.⁴⁰

Afterward, aiming at the determination of the kinetic isotope effect characterizing these metal-free C-C and C-O couplings, deuterated substrate **2a**-[D₄]-OTf was reacted with **5w** and **5b**, respectively (**Scheme 5b**). Both reactions delivered **6a**-[D] and **7w**-[D] as a mixture of -[D₄] and -[D₃] adducts with an identical [D₄] : [D₃] ratio of 77:23, suggesting that the aryne generation features a KIE of 3.3 and might be expected to be the rate-determining step of these transformations.

Scheme 5: Mechanistic studies




Generation of the aryne intermediate was also proven by trapping such a highly reactive species (**Scheme 6**). Taking advantage of the highly versatile cycloaddition reactions for the construction of new carbon-carbon and carbon-nitrogen bonds, **2a**-OTf was reacted with 2,5-dimethylfuran and benzyl azide, thus delivering **10** in excellent yield and **11** as a separable mixture of product in 58% yield.

To better understand the aryne formation process the initial rates of both, C-C and C-O coupling were measured (see the SI, **Scheme S-20** and **Scheme S-21**). Similar rates have been observed within 30 min of the reactions leading to 50% of conversions. These results support a limited influence of the reaction

media on the formation of the aryne intermediate step from **2a**-OTf, which can further rapidly react with the phenol.

Scheme 6: Cyclic diaryl λ^3 -chloranes in cycloaddition reactions



DFT calculations: To gain further insights into the formation of arynes from hypervalent compounds **2a-4a** and their subsequent reaction with phenols **5**, density functional theory (DFT) calculations at the ω B97X-D/def2-QZVPP+SMD(CHCl₃)/ ω B97X-D/def2-TZVP level of theory including corrections for solvation by chloroform were performed (see the SI for full computational details).⁴¹⁻⁴³

Aryne generation: The formation of highly reactive aryne intermediates from **2a-4a** is proposed to take place *via* deprotonation with carbonate and concurrent C-X bond cleavage. Coordination of carbonate to intermediate **A** was found to be thermodynamically favorable (**Figure 3**), and the larger stabilization of **B-I** compared to **B-Br** and of **B-Cl** can be rationalized by the greater electrostatic stabilization due to the significantly larger positive charge of iodine in **A-OTf** (+1.20 for **I** compared to +0.87 for **Br**, and +0.67 for **Cl**; see the SI, **Figure S-3**).




Figure 3: Calculated energy diagram (in kcal mol⁻¹) for the formation of arynes **C** from hypervalent halonium **2a** (red color), **3a** (black) and **4a** (blue).

Subsequent deprotonation and C-X bond cleavage occur in a concerted process *via* seven-membered cyclic transition state **TS1**, with an activation free energy of 16.2 kcal mol⁻¹ for **I**, 9.4 kcal mol⁻¹ for **Br** and only 4.3 kcal mol⁻¹ for **Cl** with respect to the corresponding intermediate **B**. The substantially smaller activation free energy as well as the considerably higher stabiliza-

tion of **C** agrees well with the experimentally observed preferred aryne formation with hypervalent λ^3 -bromane and λ^3 -chlorane compounds compared to well-established λ^3 -iodane reagents. Furthermore, the experimentally determined increased reactivity of hypervalent λ^3 -chloranes is consistent with the surprisingly low energy barrier required for aryne formation. The geometries and electronic properties of 3-(*o*-halophenyl)benzynes **C** were found to be largely independent of the halide substituent (**Figure 4**).




Figure 4: Calculated geometries of aryne intermediates **C**. Distances are given in Å; values in red correspond to NBO charges.

meta- vs. ortho-selectivity: While the slightly larger distortion towards linearity at C2, *i.e.* the aryne **C** atom in *ortho*-position with respect to the ArX substituent, indicates a marginally preferred nucleophilic attack at C2 due to the more pronounced p-orbital character, partial positive charge, and lesser transition state distortion energy,³⁸ we hypothesize that the significant steric bulk of the ArX substituent overrides the predicted selectivity, thus leading to the observed attack at *meta*-position.⁴⁴

Chemoselectivity: The C-O vs. C-C, condition-controlled chemoselectivity was subsequently investigated. The formation of a solvent-separated potassium and phenolate ion pair is likely to occur in water, whereas in less polar chloroform, the potassium phenolate ion pair is formed, presumably rendering an attack of the phenolate oxygen onto the benzyne less facile. In contrast, cesium phenolate is expected to form loose ion-pairs in chloroform, due to the more complete solvation in part caused by the significantly lower charge density compared to potassium. To shed light on the *meta-ortho*-selectivity of the transformation as well as the solvent-dependent switch from C-O to C-C bond formation, we performed a series of calculations at different C-O and C-C bond distances, respectively. These computations were performed because there is no electronic energy barrier to the reaction, although there will be a free energy barrier due to the decrease of entropy for this bimolecular reaction. The barrier will be low (less than the 8-12 kcal/mol $-T\Delta S$ of a bimolecular reaction). For the reaction of intermediate **C-Cl** with phenolate, a clear preference for C-O over C-C bond formation can be observed for all relevant bond distances (**Figure 5a**).⁴⁵

In addition, C-O as well as C-C bond formation was found to preferentially occur in the *meta*-position, which is in good agreement with the experimental observation. In contrast, replacing phenolate with potassium phenolate led to a distinct change in reactivity, with C-C bond formation in the *meta*-position now preferred over C-O bond formation for bond distances $d > 1.8$ Å (**Figure 5b**).

Based on these mechanistic evidence, a plausible reaction mechanism for the formation of C-C and C-O coupling is proposed (Scheme 7). Initially, an anion exchange leads to the energetically more stable 2a-carbonate intermediate **I**, which through a low barrier process, evolves into a seven-membered ring TS. Concerted base-mediated deprotonation and carbon-halogen cleavage generate the aryne intermediate **II**. The *meta*-selective nucleophilic attack onto the benzyne intermediate **II** can provide the formation of the new C-C and C-O bond respectively yielding intermediate **IIIa** and **IIIb**. The rearomatization of the intermediate **IIIa** and the subsequent protonation of the carbanion generate the products **6a** and **7a**.





Figure 5: Calculated energy profile (in kcal mol⁻¹) for C-O bond (black and red squares) and C-C bond (blue and green triangles) formation with (a) phenolate and (b) potassium phenolate.

Post-functionalization. The transition metal-free C-C and C-O couplings described here provide a sustainable and rapid access to highly modular building blocks, straightforwardly diversifiable into various interesting compounds (Scheme 8). To demonstrate several possible post-modifications, large-scale couplings were first performed. The 15 mmol scale synthesis of the cyclic diaryl λ^3 -chlorane was performed, yielding 3 grams of 2a-OTf in 60% yield within 90 minutes, ensuring a useful and simple approach to these hypervalent reagents. Subsequently, a vast diversification study was carried out to further display the utility of our products. The 2 mmol scale reactions of 2a-OTf with 5a and 5d under the C-C coupling conditions provided 6a and 6d in synthetically useful amounts. Interestingly, exploiting the allyl fragment of 6d, a Grubbs catalyzed metathesis furnished the highly functionalized dichloro-diphenol compound 12 in 61% yield. A straightforward conversion


of the hydroxy group to the corresponding triflate 13 enabled the synthesis of the *m*-quinquephenyl 14 via a double Suzuki reaction in 58% yield, while a Cl-selective palladium catalyzed Suzuki coupling furnished the product 15 in 59%. The peculiar presence of a hydroxyl residue was also exploited for the synthesis of a new phosphine 16 which was prepared in 55% yield. Subsequently, the relevance of C-O coupling product was evaluated. The scale-up reactions up to 3 mmol scales, produced the desired products 7y, 7z and 7J in good to excellent yields (55%, 78% and 68% respectively). Taking advantages of the selective reactivity of the iodine with respect to the chlorine atom, a Suzuki reaction and a lithium base mediated exchange were performed starting from 7z, supplying respectively 17 and 18 in 40% and 50% yields. On the other hand, a palladium catalyzed C-H activation delivered the dibenzofuran 19 from 7J in 64% yield. Notably, the resulting products contain the residual chlorine substituent as a handle for further post-modifications. Accordingly, a palladium catalyzed C-N Buchwald-Hartwig and a Suzuki reaction were performed starting from 7y granting respectively access to 20 and 21 in 64% and 83% yields. Finally, a palladium-catalyzed dechlorination process yielded the benzofuran 22.

Scheme 7: Proposed mechanism



Conclusion. We have here reported a new, simple, and robust synthetic method for the preparation of novel cyclic diaryl λ^3 -chloranes 2. The higher reactivity of these compounds proved to be complementary to the well-established chemistry of cyclic diaryl λ^3 -iodane and more recent λ^3 -bromane compounds. The increased reactivity of cyclic diaryl λ^3 -chloranes as aryne precursors enhances the unparalleled metal-free *ortho*-arylation of phenols under mild base-mediated reaction conditions. Furthermore, experimental and DFT mechanistic investigations elucidated the origins of the reactivities, as well as the chemo- and regio-selectivity observed in this work. A variety of synthetic applications show the considerable potential of these new reagents in synthetic organic chemistry

Scheme 8: Post-Functionalization



ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

General procedure, product characterizations, computational methods, energies and geometries of calculated structures (PDF)

AUTHOR INFORMATION

Corresponding Author

*Joanna Wencel-Delord - wenceldelord@unistra.fr

*K. N. Houk - houk@chem.ucla.edu

ACKNOWLEDGMENT

We thank the Centre National de la Recherche Scientifique (CNRS), the “Ministère de l’Éducation Nationale et de la Recherche” (France) for financial support. J.W.D., M.L., and T. S. T. are very grateful to the European Commission for the ERC-Starting Grant “AICHIMIE” no. 949804. Generous support by the Alexander von Humboldt-foundation (Feodor Lynen fellowship, T.R.) and the National Science Foundation (CHE-1764328 to K.N.H.) is gratefully acknowledged. Calculations were performed on the Hoffman2 cluster at the University of California, Los Angeles. We are grateful to Dr. Lydia Karmazin, Dr. Corinne Bailly and Dr. Nathalie Gruber for X-ray diffraction analysis (Service de Radiocristallographie, CNRS-Université de Strasbourg, Strasbourg, France). The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

REFERENCES

- Yoshimura, A.; Zhdankin, V. V. Advances in Synthetic Applications of Hypervalent Iodine Compounds. *Chem. Rev.* **2016**, *116* (5), 3328–3435. <https://doi.org/10.1021/acs.chemrev.5b00547>.
- Zhdankin, V. V.; Stang, P. J. Chemistry of Polyvalent Iodine. *Chem. Rev.* **2008**, *108* (12), 5299–5358. <https://doi.org/10.1021/cr800332c>.
- Silva, Jr., L. F.; Olofsson, B. Hypervalent Iodine Reagents in the Total Synthesis of Natural Products. *Nat. Prod. Rep.* **2011**, *28* (10), 1722. <https://doi.org/10.1039/c1np00028d>
- Morimoto, K.; Sakamoto, K.; Ohnishi, Y.; Miyamoto, T.; Ito, M.; Dohi, T.; Kita, Y. Metal-Free Oxidative Para Cross-Coupling of Phenols. *Chem. Eur. J.* **2013**, *19* (27), 8726–8731. <https://doi.org/10.1002/chem.201301028>.
- Stridfeldt, E.; Lindstedt, E.; Reitti, M.; Blid, J.; Norrby, P.-O.; Olofsson, B. Competing Pathways in O-Arylations with Diaryliodonium Salts: Mechanistic Insights. *Chem. Eur. J.* **2017**, *23* (53), 13249–13258. <https://doi.org/10.1002/chem.201703057>.
- Merritt, E.; Olofsson, B. Diaryliodonium Salts: A Journey from Obscurity to Fame. *Angew. Chem. Int. Ed.* **2009**, *48* (48), 9052–9070. <https://doi.org/10.1002/anie.200904689>.
- Wang, X.; Studer, A. Iodine(III) Reagents in Radical Chemistry. *Acc. Chem. Res.* **2017**, *50* (7), 1712–1724. <https://doi.org/10.1021/acs.accounts.7b00148>.
- Winterson, B.; Patra, T.; Wirth, T. Hypervalent Bromine(III) Compounds: Synthesis, Applications, Prospects. *Synthesis* **2022**, *54* (05), 1261–1271. <https://doi.org/10.1055/a-1675-8404>.
- Ochiai, M. Hypervalent Aryl-, Alkynyl-, and Alkenyl- λ^3 -Bromanes. *Synlett* **2009**, *2009* (2), 159–173. <https://doi.org/10.1055/s-0028-1087355>.

- (10) Farooq, U.; Shah, A.-H. A.; Wirth, T. Hypervalent Bromine Compounds: Smaller, More Reactive Analogues of Hypervalent Iodine Compounds. *Angew. Chem. Int. Ed.* **2009**, *48* (6), 1018–1020. <https://doi.org/10.1002/anie.200805027>.
- (11) Miyamoto, K.; Saito, M.; Tsuji, S.; Takagi, T.; Shiro, M.; Uchiyama, M.; Ochiai, M. Benchtop-Stable Hypervalent Bromine(III) Compounds: Versatile Strategy and Platform for Air- and Moisture-Stable λ^3 -Bromanes. *J. Am. Chem. Soc.* **2021**, *143* (25), 9327–9331. <https://doi.org/10.1021/jacs.1c04536>.
- (12) Miyamoto, K.; Uchiyama, M. Hypervalent Organo- λ^3 -Chloranes. *Chem. Lett.* **2021**, *50* (4), 832–838. <https://doi.org/10.1246/cl.200849>.
- (13) Nakajima, M.; Miyamoto, K.; Hirano, K.; Uchiyama, M. Diaryl- λ^3 -Chloranes: Versatile Synthesis and Unique Reactivity as Aryl Cation Equivalent. *J. Am. Chem. Soc.* **2019**, *141* (16), 6499–6503. <https://doi.org/10.1021/jacs.9b02436>.
- (14) Watanabe, Y.; Takagi, T.; Miyamoto, K.; Kanazawa, J.; Uchiyama, M. Shelf-Stable (E)- and (Z)-Vinyl- λ^3 -Chlorane: A Stereospecific Hyper-Vinylating Agent. *Org. Lett.* **2020**, *22* (9), 3469–3473. <https://doi.org/10.1021/acs.orglett.0c00924>.
- (15) Sandin, R. B.; Hay, A. S. Stable Bromonium and Chloronium Salts. *J. Am. Chem. Soc.* **1952**, *74* (1), 274–275. <https://doi.org/10.1021/ja01121a524>.
- (16) Lanzi, M.; Dherbassy, Q.; Wencel-Delord, J. Cyclic Diaryl λ^3 -Bromanes as Original Aryne Precursors. *Angew. Chem. Int. Ed.* **2021**, *60* (27), 14852–14857. <https://doi.org/10.1002/anie.202103625>.
- (17) Lanzi, M.; Ali Abdine, R. A.; De Abreu, M.; Wencel-Delord, J. Cyclic Diaryl λ^3 -Bromanes: A Rapid Access to Molecular Complexity via Cycloaddition Reactions. *Org. Lett.* **2021**, *23* (23), 9047–9052. <https://doi.org/10.1021/acs.orglett.1c03278>.
- (18) Yoshida, Y.; Ishikawa, S.; Mino, T.; Sakamoto, M. Bromonium Salts: Diaryl- λ^3 -Bromanes as Halogen-Bonding Organocatalysts. *Chem. Commun.* **2021**, *57* (20), 2519–2522. <https://doi.org/10.1039/D0CC07733J>.
- (19) Yoshida, Y.; Mino, T.; Sakamoto, M. Chiral Hypervalent Bromine(III) (Bromonium Salt): Hydrogen- and Halogen-Bonding Bifunctional Asymmetric Catalysis by Diaryl- λ^3 -Bromanes. *ACS Catal.* **2021**, *11* (12), 13028–13033. <https://doi.org/10.1021/acscatal.1c04070>.
- (20) (a) Zhang, X.; Zhao, K.; Li, N.; Yu, J.; Gong, L.; Gu, Z. Atroposelective Ring Opening of Cyclic Diaryliodonium Salts with Bulky Anilines Controlled by a Chiral Cobalt(III) Anion. *Angew. Chem. Int. Ed.* **2020**, *59*, 19899–19904. <https://doi.org/10.1002/anie.202008431>; (b) Li, Q.; Zhang, M.; Zhan, S.; Gu, Z. Copper-Catalyzed Enantioselective Ring-Opening of Cyclic Diaryliodoniums and O-Alkylhydroxylamines. *Org. Lett.* **2019**, *21* (16), 6374–6377. <https://doi.org/10.1021/acs.orglett.9b02267>.
- (21) Karandikar, S. S.; Bhattacharjee, A.; Metzke, B. E.; Javaly, N.; Valente, E. J.; McCormick, T. M.; Stuart, D. R. Orbital Analysis of Bonding in Diarylhalonium Salts and Relevance to Periodic Trends in Structure and Reactivity. *Chem. Sci.* **2022**, *13* (22), 6532–6540. <https://doi.org/10.1039/D2SC02332F>.
- (22) Quideau, S.; Deffieux, D.; Douat-Casassus, C.; Pouységu, L. Plant Polyphenols: Chemical Properties, Biological Activities, and Synthesis. *Angew. Chem. Int. Ed.* **2011**, *50* (3), 586–621. <https://doi.org/10.1002/anie.201000044>.
- (23) Scott, K. A.; Cox, P. B.; Njardarson, J. T. Phenols in Pharmaceuticals: Analysis of a Recurring Motif. *J. Med. Chem.* **2022**, *65* (10), 7044–7072. <https://doi.org/10.1021/acs.jmedchem.2c00223>.
- (24) *The Chemistry of Phenols*; Rappoport, Z., Ed.; The chemistry of functional groups; Wiley: Hoboken, NJ, 2003.
- (25) Huang, Z.; Lumb, J.-P. Phenol-Directed C–H Functionalization. *ACS Catal.* **2019**, *9* (1), 521–555. <https://doi.org/10.1021/acscatal.8b04098>.
- (26) Eisenhofer, A.; Hioe, J.; Gschwind, R. M.; König, B. Photocatalytic Phenol-Arene C–C and C–O Cross-Dehydrogenative Coupling: Photocatalytic Phenol-Arene C–C and C–O Cross-Dehydrogenative Coupling. *Eur. J. Org. Chem.* **2017**, *2017* (15), 2194–2204. <https://doi.org/10.1002/ejoc.201700211>.
- (27) Tian, T.; Li, Z.; Li, C.-J. Cross-Dehydrogenative Coupling: A Sustainable Reaction for C–C Bond Formations. *Green Chem.* **2021**, *23* (18), 6789–6862. <https://doi.org/10.1039/D1GC01871J>.
- (28) Krylov, I. B.; Vil', V. A.; Terent'ev, A. O. Cross-Dehydrogenative Coupling for the Intermolecular C–O Bond Formation. *Beilstein J. Org. Chem.* **2015**, *11*, 92–146. <https://doi.org/10.3762/bjoc.11.13>.
- (29) Liu, Z.; Larock, R. C. Facile O-Arylation of Phenols and Carboxylic Acids. *Org. Lett.* **2004**, *6* (1), 99–102. <https://doi.org/10.1021/ol0361406>.
- (30) Gebara, K. S.; Casagrande, G. A.; Raminelli, C. An Efficient Fluoride-Mediated O-Arylation of Sterically Hindered Halophenols with Silylaryl Triflates under Mild Reaction Conditions. *Tetrahedron Letters* **2011**, *52* (22), 2849–2852. <https://doi.org/10.1016/j.tetlet.2011.03.124>.
- (31) Castillo, J.-C.; Quiroga, J.; Rodriguez, J.; Coquerel, Y. Time-Efficient Synthesis of Pyrido-[2,3-*d*]Pyrimidinones via α -Oxoket-enes. *Eur. J. Org. Chem.* **2016**, *2016* (11), 1994–1999. <https://doi.org/10.1002/ejoc.201600171>.
- (32) Röckl, J. L.; Pollok, D.; Franke, R.; Waldvogel, S. R. A Decade of Electrochemical Dehydrogenative C,C-Coupling of Aryls. *Acc. Chem. Res.* **2020**, *53* (1), 45–61. <https://doi.org/10.1021/acs.accounts.9b00511>.
- (33) Truong, T.; Daugulis, O. Divergent Reaction Pathways for Phenol Arylation by Arynes: Synthesis of Helicenes and 2-Arylphenols. *Chem. Sci.* **2013**, *4* (1), 531–535. <https://doi.org/10.1039/C2SC21288A>.
- (34) Yoshimura, A.; Fuchs, J. M.; Middleton, K. R.; Maskae, V.; Rohde, G. T.; Saito, A.; Postnikov, P. S.; Yusubov, M. S.; Nemykin, V. N.; Zhdankin, V. V. Pseudocyclic Arylbenzodioxaboroles: Efficient Benzyne Precursors Triggered by Water at Room Temperature. *Chem. Eur. J.* **2017**, *23* (66), 16738–16742. <https://doi.org/10.1002/chem.201704393>.
- (35) (a) Kitamura, S.; Hvorecny, K. L.; Niu, J.; Hammock, B. D.; Madden, D. R.; Morisseau, C. Rational Design of Potent and Selective Inhibitors of an Epoxide Hydrolase Virulence Factor from *Pseudomonas aeruginosa*. *J. Med. Chem.* **2016**, *59* (10), 4790–4799. <https://doi.org/10.1021/acs.jmedchem.6b00173>; (b) Xu, Y.; Wang, Y.; Wu, D.; He, W.; Wang, L.; Zhu, W. P-Terphenyls From *Aspergillus* Sp. GZWMJZ-055: Identification, Derivation, Antioxidant and α -Glycosidase Inhibitory Activities. *Front. Microbiol.* **2021**, *12*, 654963. <https://doi.org/10.3389/fmicb.2021.654963>; (c) Guo, R.-T.; Cao, R.; Liang, P.-H.; Ko, T.-P.; Chang, T.-H.; Hudock, M. P.; Jeng, W.-Y.; Chen, C. K.-M.; Zhang, Y.; Song, Y.; Kuo, C.-J.; Yin, F.; Oldfield, E.; Wang, A. H.-J. Bisphosphonates Target Multiple Sites in Both Cis- and Trans-Prenyltransferases. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104* (24), 10022–10027. <https://doi.org/10.1073/pnas.0702254104>. (d) Kovács, A.; Vasas, A.; Hohmann, J. Natural Phenanthrenes and Their Biological Activity. *Phytochemistry* **2008**, *69* (5), 1084–1110. <https://doi.org/10.1016/j.phytochem.2007.12.005>; (e) Hangeland, J. J.; Doweiko, A. M.; Dejneka, T.; Friends, T. J.; Devasthale, P.; Mellström, K.; Sandberg, J.; Grynfarb, M.; Sack, J. S.; Einspahr, H.; Färnegårdh, M.; Husman, B.; Ljunggren, J.; Koehler, K.; Sheppard, C.; Malm, J.; Ryono, D. E. Thyroid Receptor Ligands. Part 2: Thyromimetics with Improved Selectivity for the Thyroid Hormone Receptor Beta. *Bioorg. Med. Chem. Lett.* **2004**, *14* (13), 3549–3553. <https://doi.org/10.1016/j.bmcl.2004.04.032>.
- (36) Synthesis of **1a** was conducted under transition metal free protocol (see the SI, page 14)
- (37) Tse, E. G.; Houston, S. D.; Williams, C. M.; Savage, G. P.; Rendina, L. M.; Hallyburton, I.; Anderson, M.; Sharma, R.;

- Walker, G. S.; Obach, R. S.; Todd, M. H. Nonclassical Phenyl Bioisosteres as Effective Replacements in a Series of Novel Open-Source Antimalarials. *J. Med. Chem.* **2020**, *63* (20), 11585–11601. <https://doi.org/10.1021/acs.jmedchem.0c00746>.
- (38) Medina, J. M.; Mackey, J. L.; Garg, N. K.; Houk, K. N. The Role of Aryne Distortions, Steric Effects, and Charges in Regioselectivities of Aryne Reactions. *J. Am. Chem. Soc.* **2014**, *136* (44), 15798–15805. <https://doi.org/10.1021/ja5099935>.
- (39) Bronner, S. M.; Mackey, J. L.; Houk, K. N.; Garg, N. K. Steric Effects Compete with Aryne Distortion To Control Regioselectivities of Nucleophilic Additions to 3-Silylarynes. *J. Am. Chem. Soc.* **2012**, *134* (34), 13966–13969. <https://doi.org/10.1021/ja306723r>.
- (40) Shi, J.; Li, L.; Li, Y. *O*-Silylaryl Triflates: A Journey of Kobayashi Aryne Precursors. *Chem. Rev.* **2021**, *121* (7), 3892–4044. <https://doi.org/10.1021/acs.chemrev.0c01011>.
- (41) Chai, J.-D.; Head-Gordon, M. Long-Range Corrected Hybrid Density Functionals with Damped Atom–Atom Dispersion Corrections. *Phys. Chem. Chem. Phys.* **2008**, *10* (44), 6615. <https://doi.org/10.1039/b810189b>.
- (42) Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7* (18), 3297. <https://doi.org/10.1039/b508541a>.
- (43) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* **2009**, *113* (18), 6378–6396. <https://doi.org/10.1021/jp810292n>.
- (44) Cheong, P. H.-Y.; Paton, R. S.; Bronner, S. M.; Im, G.-Y. J.; Garg, N. K.; Houk, K. N. Indolyne and Aryne Distortions and Nucleophilic Regioselectivities. *J. Am. Chem. Soc.* **2010**, *132* (4), 1267–1269. <https://doi.org/10.1021/ja9098643>
- (45) Previous studies already showed that additions of nucleophiles to arynes often do not display a maximum (*i.e.* transition state) on the electronic energy surface, especially with anionic nucleophiles, thereby rendering the geometry optimization of a transition state structure for the comparison of activation free energies unfeasible (Refs. 31 and 37).