



HAL
open science

**Aramazdospirifer orbelianus (Abich, 1858) n. comb., a
new cyrtospiriferid brachiopod genus and a
biostratigraphically important species from the lower
Famennian (Upper Devonian) of Armenia**

Vahram Serobyany, Taniel Danelian, Catherine Crônier, Araik Grigoryan,
Bernard Mottequin

► **To cite this version:**

Vahram Serobyany, Taniel Danelian, Catherine Crônier, Araik Grigoryan, Bernard Mottequin. Aramazdospirifer orbelianus (Abich, 1858) n. comb., a new cyrtospiriferid brachiopod genus and a biostratigraphically important species from the lower Famennian (Upper Devonian) of Armenia. Comptes Rendus. Palevol, 2022, 21 (6), pp.145-156. 10.5852/cr-palevol2022v21a6 . hal-04418776

HAL Id: hal-04418776

<https://hal.science/hal-04418776v1>

Submitted on 26 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Aramazdospirifer orbelianus (Abich, 1858) n.comb., a new cyrtospiriferid brachiopod genus and a biostratigraphically important species from the lower Famennian (Upper Devonian) of Armenia

VAHRAM SEROBYAN ^{a, b, *}, TANIEL DANELIAN ^b, CATHERINE CRÔNIER ^b,
ARAIK GRIGORYAN ^a, BERNARD MOTTEQUIN ^c

^a Institute of Geological Sciences of the National Academy of Sciences of the Republic of Armenia–Yerevan, Armenia, 24A, Marshal Baghramyan Avenue, Yerevan 0019, Republic of Armenia

^b Univ. Lille, CNRS, UMR 8198–Evo–Eco–Paleo, F–59000 Lille, France

^c O.D. Earth and History of Life, Royal Belgian Institute of Natural Sciences, rue Vautier 29, B 1000 Brussels, Belgium

Abstract

The new genus *Aramazdospirifer* (Cyrtospiriferidae) is erected to include *Spirifer orbelianus* Abich, 1858 from the lower Famennian of Central Armenia as its type species and to resolve long standing issues related to the generic affinity of the latter. The micro–ornament and internal structure of this species are investigated and documented for the first time, on the basis of recently collected material from the Ertych horizon of three different sections. Additionally, a neotype is selected for Abich’s species as the type material is lost. *Aramazdospirifer orbelianus* (Abich, 1858) is a biostratigraphically important species for the lower Famennian strata of the Lesser Caucasus. It appears to be restricted to the South–Armenian Block; reports outside this Gondwanan area are discussed and discarded.

Keywords

Brachiopoda; Spiriferida; lower Famennian; Gondwana; Armenia.

* Corresponding author. vahram.serobyian@univ-lille.fr, vahramserobyian@gmail.com (V. Serobyian)

Aramazdospirifer orbelianus (Abich, 1858) *n. comb.*, un nouveau genre de brachiopode cyrtospiriféride et une espèce biostratigraphique importante du Famennien inférieur (Dévonien supérieur) d'Arménie

Le nouveau genre *Aramazdospirifer* (Cyrtospiriferidae) est ici érigé, afin d'inclure *Spirifer orbelianus* Abich, 1858 du Famennien inférieur d'Arménie centrale comme son espèce type et résoudre ainsi des discussions de longue date sur l'affinité générique de cette espèce. La micro-ornementation et la structure interne de celle-ci sont étudiées et documentées pour la première fois, sur la base de matériel récemment récolté dans trois coupes différentes au sein de l'horizon Ertych. En outre, un néotype est sélectionné pour cette espèce décrite initialement par Abich étant donné que le matériel type est perdu. *Aramazdospirifer orbelianus* (Abich, 1858) est une espèce biostratigraphiquement importante du Famennien inférieur du Petit Caucase. Cette espèce semble être restreinte au bloc Sud-Arménien ; des rapports la mentionnant à l'extérieur du domaine gondwanien sont discutés et écartés.

Mots clés

Brachiopodes; Spiriferida; Famennien inférieur; Gondwana; Arménie.

1. Introduction

The Upper Devonian sedimentary sequences of Armenia are highly fossiliferous and contain a diverse and well-preserved brachiopod fauna (Abrahamyan, 1957, 1964, 1974; Arakelyan, 1964), which remain largely undocumented from a taxonomic and biostratigraphic point of view. Re-investigation of this fauna is crucial to better understand Late Devonian changes in brachiopod diversity and palaeobiogeographic distribution. In Armenia, Devonian outcrops were first examined by the famous German geologist Hermann Abich, considered as the 'Father of Caucasian Geology' (Milanovsky, 2007). Abich (1858) described a number of new brachiopod species from the Lesser Caucasus, notably the spiriferide species *Spirifer orbelianus*. The latter was erroneously reported afterwards in the Franco-Belgian Basin by Gosselet (1874), and later in different parts of the world (see Sartenaer, 1974 for an up-to-date review, and

references below). Due to its great potential for biostratigraphic correlations of lower Famennian successions throughout the Lesser Caucasus, re–investigation of *S. orbelianus* is a necessary task as its affinities at the genus level still remain unclear, in spite of the extensive taxonomic studies led by Abrahamyan (1957, 1974) and Afanasjeva (in Alekseeva et al., 2018a). The purpose of the present study is to reassess the taxonomy of *S. orbelianus* Abich, 1858 on the basis of recently collected material from the lower Famennian of Central Armenia, with implications for its palaeobiogeographic distribution.

2. *Stratigraphic and palaeogeographic settings*

In the southern part of Central Armenia crops out a ca. 1,500 m–thick Middle Devonian–Lower Carboniferous sequence of platform carbonate deposits (Fig. 1). They record the earliest depositional history of Palaeozoic sediments in the area; they were accumulated on a Gondwanan passive margin that was facing the Palaeotethys, situated to the north; this part of Gondwana was later individualized as the South–Armenian Block, following its northward migration and opening of Neotethys further to the South (Sosson et al., 2010). The Middle Devonian–Lower Carboniferous sequences of Armenia constitute a continuous succession of mixed carbonate–siliciclastic deposits, developed in a shallow water environment. The Upper Devonian–Lower Carboniferous sequences were subdivided by Abrahamyan (1964) and Arakelyan (1964) into twelve ‘formations’. However, in practice, most of them have very similar lithological characteristics and they cannot be recognized in the field without knowledge of their brachiopod assemblages. Thus, they have a biostratigraphic rather than lithostratigraphic significance (see Serobyán et al., 2019). It is worth noting that Abrahamyan et al. (1975) described these stratigraphic units as ‘horizons’ and no longer used the term ‘formation’. In Armenia, Famennian deposits are more widespread than the Frasnian ones. The latter consist mainly of marly and sandy limestones, biogenic limestones rich in brachiopods and corals (a few gastropods and cephalopods were also found) with interbedded sandstones and quartzites, whereas the Famennian is chiefly represented by biogenic and sandy limestones, quartzites and shales. The first Devonian biostratigraphical zonal scheme based on brachiopods was established by Rzhonsnitskaya (1948). Afterwards, in her groundbreaking monograph, Abrahamyan (1957) discussed the occurrence and stratigraphic distribution of the brachiopod species she dealt with;

she thus proposed a new continuous biostratigraphic scheme characterized by marker species or species assemblages (Fig. 2). This was a major step forward, as the existing Devonian brachiopod biostratigraphic scheme available at the time for the region was very rudimentary. As the Upper Palaeozoic sedimentary sequences of Armenia continue into Nakhichevan, the stratigraphic and faunal similarities in terms of brachiopods allowed Mamedov and Rzhonsnitskaya (1985) to use and refine Abrahamyan's zonal scheme for the entire region of the Lesser Caucasus (or Transcaucasia). Subsequently, their zonal scheme was updated by Rzhonsnitskaya and Mamedov (2000) and correlated with the international conodont biozonation based on the conodont study of Aristov (1994) carried out in Nakhichevan.

3. *Material and methods*

The examined material was collected from the Noravank, Ertych and Djravank sections (Fig. 1) during several field seasons organized in 2018 and 2019. It comes from marly limestones and shales of the lower Famennian Ertych horizon (Fig. 2), which can be correlated with the '*Cyrtospirifer orbelianus* Zone of Abrahamyan (1957) and the '*Cyrtiopsis orbelianus*–*Cyrtiopsis armenicus* Zone of Rzhonsnitskaya and Mamedov (2000). In total 80 articulated brachiopod shells and 15 dissociated valves were collected, the main part of which is derived from a soft, weathered surface that provided well-preserved, nearly sediment-free specimens. The bulk of the specimens illustrated and investigated herein is housed at the Geological Museum of the Institute of Geological Sciences of the National Academy of Sciences of Armenia, Yerevan (IGSNASRAGM). Some specimens are deposited at the Royal Belgian Institute of natural Sciences, Brussels (RBINS) and others at the University of Lille (USTL).

The recently collected material was compared with Abrahamyan's specimens, collected in Armenia during the 1940s to 1980s, most of which are now stored at the IGSNASRAGM. The internal structure of the newly collected articulated specimens identified as *A. orbelianus* (Abich, 1858) found in the three sections (Fig. 1) was investigated by using the standard technique of serial sections and acetate peels. The latter were assembled between microscope slides and photographed under a binocular microscope Olympus SZX 12. Afterwards, the photographs were transferred to CorelDRAW X7 software and internal details were drawn using a digital drawing tablet. Furthermore, in order to capture the fine details of the internal structure, the

ground specimens were photographed directly under a Canon EOS 700D camera that was attached on a Zeiss SteREO Discovery V20 Microscope. Intact brachiopod specimens were coated with magnesium oxide or ammonium chloride sublimate before being photographed. All images have been further processed using Adobe Photoshop CS6. Additionally, the IGSNASRAGM 3895/PS 3001 specimen was coated with gold and the digitization of the micro-ornament of the latter was performed by ZEISS EVO Scanning Electron Microscope. RBINS specimens selected for scanning electron microscopy were observed using a low vacuum SEM, an ESEM FEI Quanta 200, but not coated with gold.

4. *Systematic palaeontology*

The supraspecific classification adopted herein follows Carter *et al.* (2006) for the Order Spiriferida. The synonymy list only concerns the report of the species in the South–Armenian Block.

Order Spiriferida Waagen, 1883

Suborder Spiriferidina Waagen, 1883

Superfamily Cyrtospiriferoidea Termier and Termier, 1949

Family Cyrtospiriferidae Termier and Termier, 1949

Subfamily Cyrtospiriferinae Termier and Termier, 1949

Genus *Aramazdospirifer* gen. nov.

Type species: *Spirifer orbelianus* Abich, 1858.

Other species: It is currently difficult to assign other species to *Aramazdospirifer* gen. nov. with certainty in the absence of taxonomical revisions of the diverse cyrtospiriferide fauna from the Famennian of the ex–USSR. *Spirifer (Cyrtospirifer) aperturatus* von Schlotheim *sensu* Nalivkin (1930) described in the Central Kara–Tau and along the headwaters of the Naryn river (Kazakhstan) likely belongs to the new genus, although the latter has a less globular shell, cubcircular tongue and lacks median and longitudinal elevations in sulcus and on fold. It is worth

nothing that the real *aperturatus* from Germany and the Franco–Belgian basin differs from Nalivkin’s species and does not belong to *Aramazdospirifer* gen. nov. (see Paeckelmann, 1942, Vandercammen, 1959). It is also probable that *Cyrtospirifer* (*Cyrtospirifer*) *pentagonalis* Sidjachenko (1962) described from the Famennian of Central Kara–Tau should be assigned to *Aramazdospirifer* gen. nov. as it shares many common features with the new genus. More precisely its inflated and subpentagonal shell that is ornamented with closely spaced and flattened ribs, acute to mucronate cardinal extremities and apsacline triangular ventral interarea that fit well with the diagnosis of the new genus. Nevertheless, the micro–ornament, the internal morphology and the type of pseudodeltidium of the latter species is unknown. Finally, *Cyrtospirifer aquilinus* Romanowski *sensu* Nalivkin (1930) described from the Central Kara–Tau and along the headwaters of the Naryn river (Kazakhstan) might also be assigned to the new genus, although it lacks median and longitudinal elevations in sulcus and on fold. However, its micro–ornament and internal morphology is unknown. All these uncertainties preclude the possibility of a definite assignment to *Aramadzospirifer*.

Etymology: In honor of Aramazd, who was the chief god in pre–Christian Armenian mythology.

Diagnosis: Shell of medium–size, ventribiconvex, subpentagonal, wider than long to longer than wide, with mucronate cardinal extremities; generally widest at hinge line; ventral interarea moderately high, apsacline; delthyrium wide and most of its height covered by a pseudodeltidium, the latter is composed of several distinct plates, with submesothyrid foramen; fold and sulcus moderately wide, well–defined; median longitudinal elevation developed in sulcus and frequently on fold; ribs numerous, usually simple on flanks, increasing by bifurcation in sulcus and on fold; micro–ornament capillate both on ribs and in interspaces; dental plates long, strong and intrasinal; delthyrial plate well–developed; unsupported ctenophoridium with numerous vertical lamellae; crural plates short; dorsal myophragm present.

Remarks: *Aramazdospirifer* gen. nov. is included in the Superfamily Cyrtospiriferoidea based on the presence of dental plates, a ctenophoridium, a well–developed delthyrial plate, and capillate ornamentation. Although a thorough revision of this superfamily, recommended by Ma and Day (2000), is still pending, *Aramazdospirifer* gen. nov. is assigned to the Family Cyrtospiriferidae, as defined by Johnson (2006), due to the development of ribs in the sulcus and on fold. Additionally, its wide hinge line argues for its assignment to the Subfamily Cyrtospiriferinae

rather than to the Cyrtiopsinae.

Afanasjeva in Alekseeva *et al.* (2018a) examined some specimens collected from the Armenian sections and reassigned Abich's species to the late Givetian?–early Frasnian genus *Uchtospirifer* Liashenko, 1957 known from South Timan (Russia). *Aramazdospirifer* gen. nov. shares indeed several external and internal characters with *Uchtospirifer* Ljaschenko, 1957 *sensu* Sokiran (2006). More particularly, both genera display a subpentagonal shell that is ornamented with closely spaced and flattened ribs (also in sulcus and on fold), an apsacline and triangular ventral interarea, divergent dental plates, and an unsupported ctenophoridium. However, *Aramazdospirifer* gen. nov. differs from *Uchtospirifer* by its much more inflated shell, acute to mucronate cardinal extremities, wider hinge line, clearly defined longitudinal and median elevation in sulcus and similar elevation often developed on fold, and the type of pseudodeltidium formed by several distinct plates. Furthermore, it is worth noting that *Aramazdospirifer* gen. nov. has narrower ribs perceptible along the whole length of its sulcus and fold, whereas the median ribs of *Uchtospirifer* are variably developed, sometimes flattened to imperceptible or absent. Moreover, the micro-ornament of the new genus lacks the tubercles observed on *Uchtospirifer*.

Externally *Aramazdospirifer* gen. nov. resembles also to the Famennian genus *Wenjukovispirifer* Oleneva, 2016 mainly in terms of its rounded subpentagonal outline, ventribiconvex lateral profile, acute to mucronate cardinal extremities, wide hinge line, well-defined and triangular ventral interarea, wide sulcus and fold bearing bifurcating ribs that are narrower than those present on the flanks. However, *Aramazdospirifer* gen. nov. differs by its median and longitudinal elevation in sulcus and similar elevation frequently displayed on fold, and its pseudodeltidium that is formed by several distinct plates with minute submesothyrid foramen. Additionally, *Aramazdospirifer* gen. nov. lacks pustules that are observed in *Wenjukovispirifer*. It is difficult to compare the internal morphology of these genera as Oleneva (2016) did not illustrate any serial sections. Nevertheless, she mentioned the presence of a median septum situated in the ventral interior of *Wenjukovispirifer*, a character that is not observed in *Aramazdospirifer* gen. nov.

The new genus differs strongly from *Cyrtospirifer* Nalivkin in Frederiks, 1924, by its more globular, longitudinally elongated and inflated shell, well-developed pseudodeltidium, longitudinal and median elevation in sulcus and similar elevation often developed on fold and the

type of micro-ornament. Internally, these genera appear to be similar, though *Aramazdospirifer* gen. nov. possesses an unsupported ctenophoridium, while the latter is supported by an apical callosity in many *Cyrtospirifer* representatives.

Aramazdospirifer gen. nov. is distinguished externally from *Lamarckispirifer* Gatinaud, 1949 *sensu* Ma and Day (2007), known from the lower Famennian of South China, by its more rounded and inflated shell, longitudinal and median elevation in sulcus and analogous elevation often developed on fold and its high tongue. Additionally, *Aramazdospirifer* gen. nov. lacks a median furrow on fold of the dorsal valve and the *hayasakai*-type micro-ornament. Internally, the most substantial difference is the presence of a dorsal median septum in *Lamarckispirifer*, while this character is absent in *Aramazdospirifer* gen. nov.

Aramazdospirifer gen. nov. can be distinguished from the genus *Pseudocyrtiopsis* Ma and Day, 1999, known from the lower Famennian of South China, by its shell outline, longitudinal and median elevation in sulcus and analogous elevation often developed on fold, simple and flattened ribs, lack of well-developed pustules in ribs and interspaces. Internally, *Aramazdospirifer* gen. nov. differs by its unsupported ctenophoridium.

Aramazdospirifer orbelianus (Abich, 1858)

Figures 3–7, Table 1

1858 *Spirifer Orbelianus*, nov. sp. Abich, p. 438, 440, 524–525, pl. 1, figs. 2–3; pl. 2, figs. 4–5.

1948 *Cyrtospirifer orbelianus* (Abich); Rzhonsnitskaya, p. 1480.

1952 *Cyrtospirifer orbelianus* (Abich); Arakelyan, p. 40, 42.

1957 *Cyrtospirifer orbelianus* (Abich, 1858); Abrahamyan, p. 70, pl. 8, fig. 3.

1964 *Cyrtospirifer orbelianus* (Abich, 1858); Arakelyan, p.75, 77, 82, 94.

1973 *Cyrtospirifer orbelianus* (Abich, 1858); Abrahamyan *et al.*, p. 218.

1974 *Cyrtiopsis orbelianus* (Abich, 1858); Abrahamyan, p. 60, pl. 22, figs. 2–3.

1974 *Spirifer orbelianus* Abich, H., 1858; Sartenaer, p. 10 (only the Armenian specimens).

1975 *Cyrtiopsis orbelianus* (Abich, 1858); Arakelyan *et al.*, p. 24.

2000 *Cyrtiopsis orbelianus* (Abich, 1858); Rzhonsnitskaya and Mamedov, p. 331, table 1.

2018a *Uchtospirifer orbelianus* (Abich, 1858); Afanasjeva in Alekseeva *et al.*, pl. 30, fig. 4;

text–fig. 103.

2018b *Uchtospirifer orbelianus* (Abich, 1858); Grechishnikova in Alekseeva *et al.*, p. 832, 855.

Neotype.— Abich (1858) did not designate a type specimen for his new species among the four ones he illustrated. Abich's collection is stored in the Saint Petersburg Mining Institute Museum, but all the spiriferides collected by Abich are lost (D. Bezgodova, personal communication, 2019). Therefore, the single specimen illustrated by Abrahamyan (1957, pl. 8, fig. 3) is hereby designated as the neotype and figured in Figure 3.9–14. It is stored at the Geological Museum in Yerevan under the collection number IGSNASRAGM 3897/AB97/48.

Type locality and horizon.— Marly/sandy limestone layers (Abrahamyan, 1957, 10th limestone layer of the Amaghu section, appendix 1) of the Ertych horizon, Noravank section (formerly Amaghu), Central Armenia.

Occurrence and age.— This species is one of the most biostratigraphically valuable species in Armenia for the recognition of the lower Famennian and is used for the definition of a brachiopod zone, namely the '*Cyrtospirifer orbelianus* Zone of Abrahamyan (1957), as well as the '*Cyrtiopsis orbelianus*–'*Cyrtiopsis armenicus* Zone of Rzhonsnitskaya and Mamedov (2000) (Fig. 2). Initially, Abrahamyan (1957) reported this species from sequences considered as constituting the lower part of the upper Famennian. Following the extensive stratigraphic study of Abrahamyan (1964) and Arakelyan (1964), it appeared that the previously reported *orbelianus* bearing sequences are actually early Famennian in age. Therefore, Abrahamyan (1974) specified that *A. orbelianus* is restricted to the lower Famennian (possibly corresponding to the equivalent of the *Palmatolepis crepida*–*P. glabra pectinata* conodont zones of Spalleta *et al.* (2017)). Previously this species has been observed in Armenia in the Argichi, Lanjanist (Kadrlu), Chanakchi (Zangakatun), Ertych and Noravank sections (Fig. 1; Abrahamyan, 1957; Arakelyan, 1964) and doubtfully in the Gyumushlug section of Nakhichevan (Fig. 1; Abrahamyan, 1957). It is worth noting that only two poorly preserved specimens of possible *A. orbelianus* have been found so far in Nakhichevan by Abrahamyan (1957) (IGSNASRAGM collections). Nevertheless, both specimens lack longitudinal and median elevations, and display an erect beak, which are not characters present in *A. orbelianus*. Moreover, further sampling in several sections exposing the lower Famennian strata in Nakhichevan, including the Gyumushlug section, did not reveal the

presence of Abich's species (e.g. Arakelyan, 1964; Aristov *et al.*, 1979; Feliks *et al.*, 1980; Grechishnikova, 1986).

Material examined.— Eighty articulated specimens, ten ventral and five dorsal valves from the marly/sandy limestones of the Ertych horizon of the Djravank (sixty articulated specimens and ten ventral valves), Noravank (type locality; twelve articulated specimens and four dorsal valves) and Ertych (eight articulated specimens and one dorsal valve) sections. Three articulated specimens in Abrahamyan's collection (IGSNASRAGM) from the Noravank section and two ventral valves from the Gyumushlug section (Nakhichevan).

Description.— Shell medium-sized (up to 39 mm in width and 39.1 mm in length), wider than long to longer than wide, generally ventribiconvex, rounded subpentagonal in outline, widest at hinge line, highest at about midlength; cardinal extremities acute to mucronate, but strongly extended in some specimens; anterior margin strongly excavated by sulcus, emarginate; anterior commissure uniplicate.

Ventral valve strongly inflated, rounded pentagonal in outline, with convex flanks sloping steeply towards lateral commissures; highest at about midlength or slightly posteriorly; umbo strongly inflated, large and prominent; beak erect (85–90°), interarea apsacline, triangular, moderately high, well-defined, concave; delthyrium wide, with most of its height covered by pseudodeltidium, the latter formed by several distinct plates; foramen minute, rounded to ovate; sulcus wide, shallow to moderately deep, originating from beak, widening and becoming deeper anteriorly, sulcus margins gentle. Sulcus bears a longitudinal and median elevation which inconspicuously originates in the posterior part of the valve; it widens, thickens and becomes more conspicuous anteriorly (highest and widest at the anterior margin); tongue high, perpendicular to commissural plane with its distal part sometimes bent dorsally, subogival in outline.

Dorsal valve wider than long, inflated with flanks sloping moderately to strongly towards lateral commissures, subquadrangular to subtrapezoidal in outline; highest in the anterior third of the valve, but gradually decreasing towards the anterior margin; interarea linear (up to 3 mm high), slightly concave, orthocline; fold well-defined, wide, moderately high, inconspicuously originating from beak, widening and becoming higher anteriorly, often bearing a longitudinal and median elevation starting from dorsal beak.

Ornamentation of up to 35 rounded (generally 28–30, 5–6 ribs per 5 mm at anterior margin near sulcus and fold), simple, flattened, low ribs on each flank, becoming fainter towards posterolateral margins; in sulcus and on fold, up to 25 ribs, increasing by bifurcations, much narrower than those present on flanks; interspaces as wide as ribs on flanks, but wider than ribs in sulcus and on fold; micro-ornament of capillae both on ribs and in interspaces with concentric growth lines sometimes thickened as growth varices.

Ventral valve interior with thin, long, intrasinal and divergent dental plates, becoming much less divergent to almost parallel more anteriorly, converging dorsally in umbonal region (as seen in transverse section); delthyrial plate well-developed, thick (particularly in large specimens); umbonal callus well-developed, central and lateral apical cavities large and filled in by callus; teeth relatively small, subrectangular.

Dorsal valve interior with unsupported ctenophoridium composed of up to 33 well developed, relatively long lamellae; hinge plate divided; outer hinge plates slightly concave, crural bases dorsally convergent; spiral cones not preserved in the sectioned specimens.

Variability.— The shell shape varies from almost globular forms with thickness exceeding as width and length, having narrower, shallow sulcus to transversely elongated wider forms with broad sulcus and relatively less inflated valves. The beak angle, the height of the longitudinal elevation and the number of ribs are also shifting.

Ontogeny.— Juvenile forms differ from adults in having less inflated valves, straight beaks, shallower sulcus, weak fold, fainter ribs, inconspicuous elevations developed only anteriorly and the less convex delthyrial plate. The size distribution during growth represented by the length/width, thickness/width, width of sulcus/width and length of dorsal valve/width plots (Fig. 6) shows a continuous and progressive growth with no distinct grouping. The relative proportions of *Aramazdospirifer orbelianus* represented by sufficient material remain constant (linear regression: $y = ax + b$; significant probability value: $p < 0.001$ *** whatever the degree of development of individuals (Fig. 6)). Moreover, the correlation is positive with width varying proportionally with length, thickness, width of sulcus and the length of dorsal valve. To complete the scatter plots, the measurements (in mm) of numerous individuals of *Aramazdospirifer orbelianus* are also presented in table 1 and figure 7. The length of the dorsal valve shows less dispersed values (Fig. 7).

5. Discussion

Aramazdospirifer orbelianus was named by Abich (1858) in honor of the Orbelian family, one of the strongest dynasties in medieval Armenia, characterized by a long history of political influence documented in inscriptions throughout the provinces of Vayots Dzor and Syunik. Abich found this species in the lower Famennian sequence exposed at Noravank although he did not explicitly mention this locality. Abich (1858, p. 524) indicated that ‘*Sp. Orbelianus* ist bis jetzt nur in dem Baranco des Erhebungsthalles von Gyneschik [= Gnishik], in den Schichten h and k Profils pag. 440, und zwar in sehr grossen Mengenverhältnissen gefunden worden’, i.e., the author found this species in large numbers only in the h (impure limestones) and k (limestones with glauconitic sand) horizons of the profile exposed on the slopes of the valley of the Gnishik river (Fig. 1). These are apparently the marly/sandy limestone layers constituting the lower Famennian sequence exposed below the Noravank monastery, which is one of the most notable monasteries built by the Orbelian dynasty.

Abich’s species was later reported by Gosselet (e.g. 1874, 1880, 1894) in northern France and southern Belgium, from sequences accumulated close to the Givetian–Frasnian boundary. However, according to Sartenaer (1974) and Mottequin (2019), the Franco–Belgian material was erroneously identified as *Spirifer orbelianus* and needs further investigation. Reed (1922) reported a single specimen from Pamir, identified as *S. orbelianus* and stated that the species displays all the superficial features typical for Abich’s species. Nonetheless, his single illustrated specimen has rounded cardinal extremities, coarser ribs with very narrow interspaces and fewer ribs in sulcus and on fold; it also lacks a median longitudinal elevation. We thus consider this specimen as distinct from Abich’s species and in need of re–examination. Bonnet (1947, p. 32, p. 39) noted the presence of *Spirifer orbelianus* in Frasnian strata of Armenia, but this occurrence remains highly questionable as he did not illustrate his material (Sartenaer, 1974). Gatinaud (1949) erected the genus *Lamarckispirifer* and assigned many species to it, including Abich’s species, but as noted by Ma and Day (2007), the latter species differs externally by its inflated shell and cannot be attributed to that genus. During the ‘golden era’ of Palaeozoic studies in Armenia (1950s to 70s) *A. orbelianus* was reported extensively by Abrahamyan (1957, 1974), Abrahamyan *et al.* (1973) and Arakelyan (1952, 1964). Abrahamyan (1957) assumed that *A. orbelianus* could be a sister species of ‘*Cyrtiopsis*’ *armenicus* as the latter appeared in a slightly younger horizon. Therefore, Abrahamyan (1974) incorrectly assigned both species to the lower

Famennian genus *Cyrtiopsis* Grabau, 1923. Sartenaer (1974) discussed several reports of *A. orbelianus* from Africa, the Franco–Belgian basin and Iran. He rightly concluded that none of them corresponds to Abich’s species. In sum, *A. orbelianus* is known only from the South–Armenian Block (Central Armenia and possibly Nakhichevan).

6. Conclusions

The taxonomic revision of *Spirifer orbelianus* Abich, 1858, based on material collected from its type locality in Central Armenia and complemented by the specimens described by Abrahamyan (1957), led us to define the new cyrtospiriferine genus *Aramazdospirifer*. *A. orbelianus* (Abich, 1858) is one of the most useful species for the biostratigraphy of the lower Famennian in the Lesser Caucasus due to its short stratigraphic range and abundance. In addition to its type species, which appears to be endemic in the South–Armenian Block, several other Famennian species described from Central Kazakhstan and the East European Platform may be also considered for assignment to the genus *Aramazdospirifer*, although they are in need of a modern taxonomic reassessment. Further studies on Armenian material are still needed in order to complement the recent revisions of the Famennian brachiopod faunas from the area of northern Gondwana that now corresponds to the Lesser Caucasus, Afghanistan and Iran (Alekseeva *et al.*, 2018a, b; Mottequin and Brice, 2019).

Acknowledgments

The Ph.D. studies of V. Serobyanyan in France are funded by the French Embassy in Yerevan and the MOBILLEX International Mobility Grant Programme of the University of Lille. T. Danelian and A. Grigoryan are grateful to the Erasmus Programme for funding their staff mobility in 2018 and 2019, respectively. Fieldwork was also facilitated by the logistic support of the Institute of Geological Sciences (Armenian Academy of Sciences). The authors thank the Région Hauts–de–France, and the Ministère de l’Enseignement Supérieur et de la Recherche (CPER Climibio), and the European Fund for Regional Economic Development for their financial support. Sylvie Régner (UMR Evo–Evo–Paleo, ULille) and Julien Cillis (RBINS, Brussels) are gratefully

acknowledged for the high-quality SEM pictures obtained on specimens deposited at IGNASRAGM and at the RBINS, respectively. Daria Bezgodova is thanked for kindly providing information about Abich's collection. Constructive remarks by Jed Day, Elena Sokiran and an anonymous reviewer improved the initial manuscript. The authors are also grateful to E. Sokiran for her advice on literature related to brachiopod genera described from the ex-USSR.

References

Abich, H., 1858. Vergleichende geologische Grundzüge der Kaukasischen, Armenischen und Nordpersischen Gebirge. Prodrômus einer Geologie der Kaukasischen Länder. Mémoires de l'Académie impériale des Sciences de Saint-Pétersbourg, 6^{ème} série, Sciences mathématiques et physiques, 7, 359–534.

Abrahamyan, M.S., 1957. Brakhiopody verkhnefamenskikh i etrenskikh otlozhenii yugozapadnoi Armenii (Brachiopods of the Upper Famennian and Etroeungt deposits of the South-Western Armenia). Akademiya Nauk Armyanskoy SSR, Institut Geologicheskikh Nauk, Yerevan, 142 p. (in Russian).

Abrahamyan, M.S., 1964. Karbon (The Carboniferous). In: Mkrtchian, S.S., Vardaniants, L.A., Gabrielian, A.A., Maghakian, I.G., Paffenholz, C.N (Eds.). Geologiya Armyanskoy SSR (Geology of the Armenian USSR). Akademiya Nauk Armyanskoy SSR, Yerevan, 2, pp. 96–118 (in Russian).

Abrahamyan, M.S., 1974. Opisanie fauny, Devonskaya sistema, Tip Brachiopoda, Brakhiopody (Description of fauna, Devonian System, Phylum Brachiopoda, brachiopods). In: Akopian, V.T. (Ed.), Atlas iskopaemoy fauny Armyanskoy SSR (Atlas of the fossil fauna of the Armenian SSR). Akademiya Nauk Armyanskoy SSR, Institut Geologicheskikh Nauk, Yerevan, pp. 48–64 (in Russian).

Abrahamyan, M.S., Arakelyan R.A., Azizbekov Sh.A., 1973. Malyy Kavkaz (Yuzhnoye Zakavkaz'ye) (Lesser Caucasus (South Transcaucasia)). In: Rzhonsnitskaya, M.A. (Ed.), Kavkazskaya geosinklinal'naya oblast' i Predkavkaz'e (Caucasus geosynclinal region and Ciscaucasia). In Nalivkin, D.V., Rzhonsnitskaya, M.A., Markovski, B.P. (eds.): Stratigrafiya SSSR, Devonskaya sistema (Stratigraphy of the USSR, the Devonian system). Nedra, Moscow, 1, pp. 210–219 (in Russian).

Abrahamyan, M.S., Arakelyan, R.A., Papoyan, A.S., 1975. Granitsa Devona i Nizhnego Karbona (The Devonian and Lower Carboniferous boundary). In: Stratigrafiya Karbona i Geologiya Uglenosnykh Formatsii SSSR (Stratigraphy of the Carboniferous and Geology of Coal-bearing Formations of the USSR). Nedra, Moscow, pp. 49–55 (in Russian).

Alekseeva, R.E., Afanasjeva, G.A., Grechishnikova, I.A., Oleneva, N.V., Pakhnevich, A.V., 2018a. Devonian and Carboniferous Brachiopods and Biostratigraphy of Transcaucasia (Ending). *Paleontological Journal*, 52/9, 969–1085, doi.org/10.1134/S0031030118090010

Alekseeva, R.E., Afanasjeva, G.A., Grechishnikova, I.A., Oleneva, N.V., Pakhnevich, A.V., 2018b. Devonian and Carboniferous Brachiopods and Biostratigraphy of Transcaucasia. *Paleontological Journal*, 52/8, 829–967, doi.org/10.1134/S0031030118080014

Arakelyan, R.A., 1952. Stratigrafiya paleozoyskikh otlozheniy yugo–zapadnoy Armenii i privileyushchikh chastey Nakhichevanskoy ASSR (Stratigraphy of the Palaeozoic deposits of southwestern Armenia and the adjacent parts of the Nakhichevan ASSR). *Akademiya Nauk Armyanskoy SSR, Institut Geologicheskikh Nauk, Yerevan*, 142 p. (in Russian).

Arakelyan, R.A., 1964. Devon (The Devonian). In: Mkrtchian S.S., Vardaniants L.A., Gabrielian A.A., Maghakian I.G., Paffenholz C.N. (Eds.). *Geologiya Armyanskoy SSR (Geology of the Armenian USSR)*. *Akademiya Nauk Armyanskoy SSR, Yerevan*, 2, pp. 46–96 (in Russian).

Arakelyan, R.A., Malxasyan, E.G., Mkrtchyan, C.C., Paffenholz, C.N., 1975. *Geologicheskii ocherk Armyanskoy CCP (Geological sketch of the Armenian SSR)*. *Akademiya Nauk Armyanskoy SSR, Institut Geologicheskikh Nauk, Yerevan*, 175 p. (in Russian).

Aristov, V.A., 1994. Konodonty devona–nizhnego karbona Evrazii: soobshchestva, zonalnoye raschleneniye, korrelyatsiya raznofatsialnykh otlozheniy (Conodonts from the Devonian and Lower Carboniferous of Eurasia: communities, zonation and correlation of heterofacial deposits). *Proceedings of the Geological Institute of the Russian Academy of Sciences* 484, 1–193 (in Russian).

Aristov, V.A., Gretschischnikova, I.A., Tschigova, V.A., Felix, V.P., 1979. Subdivision and correlation of the Famennian and the lower Tournaisian deposits of Transcaucasia (on brachiopods, conodonts and ostracods). *Geological Survey of Belgium, Professional Paper* 161, 91–95.

Bonnet, P., 1947. Description géologique de la Transcaucasie méridionale (chaînes de l'Araxe moyen). *Mémoires de la Société géologique de France, nouvelle série* 25/53, 1–62.

Carter, J.L., Johnson, J.G., Gourvenec, R., Hou, H.F., 2006. Spiriferida. In: Kaesler, R.L. (Ed.), Treatise on invertebrate paleontology: part H, Brachiopoda, revised, 5. Geological Society of America, and University of Kansas, Boulder, Colorado, and Lawrence, Kansas, pp. 1689–1870.

Feliks, V.P., Grechishnikova, I.A., Levitskii, E.S., Nagiev, V.N., 1980. Novyye dannyye o geologii rudnika Gyumyushlung (New data on the geology of the Gyumushlug mine). Izvestiya vysshikh uchebnykh zavedeniy. Geologiya i razvedka 2, 69–73 (in Russian).

Frederiks, G.N., 1924. O verkhnekamennougol'nykh spiriferidakh Urala (On the Upper Carboniferous spiriferids of the Urals). Izvestiya Geologicheskogo Komiteta 38 (1919), 295–324 (in Russian).

Gatinaud, G., 1949. Contribution à l'étude des Brachiopodes Spiriferidae. I.—Exposé d'une nouvelle méthode d'étude de la morphologie externe des Spiriferidae à sinus plissé. Bulletin du Museum National d'Histoire Naturelle, série 2/4, 487–492.

Gosselet, J., 1874. Carte géologique de la bande méridionale des calcaires dévoniens de l'Entre Sambre—et—Meuse. Bulletin de l'Académie royale des Sciences, des Lettres et des Beaux—Arts de Belgique, 2ème série 37, 81–114.

Gosselet, J., 1880. Esquisse géologique du Nord de la France et des contrées voisines. 1^{er} fascicule. Terrains primaires, Imprimerie Six—Horemans, Lille, 167 p.

Gosselet, J., 1894. Étude sur les variations du *Spirifer Verneuli* et sur quelques espèces voisines. Mémoires de la Société géologique du Nord 4, 1–61.

Grechishnikova, I.A., 1986. Novyye devonskiye spiriferidy Zakavkaz'ye (New Devonian spiriferides of Transcaucasia). Zapiski Gornogo Instituta 107, 52–60 (in Russian).

Johnson, J.G., 2006. Cyrtospiriferoidea. In: Kaesler, R.L. (Ed.), Treatise on Invertebrate Paleontology: Part H, Brachiopoda, Revised, 5. Geological Society of America, and University of Kansas, Boulder, Colorado, and Lawrence, Kansas, pp. 1722–1732.

Liashenko, A. I., 1957. Novyi rod Devonskikh brachiopod *Uchtospirifer* (A new genus of Devonian brachiopods: *Uchtospirifer*). Akademiia Nauk SSSR, Doklady 117/ 5, 885–888 (in Russian).

Ma, X. –P., Day, J., 1999. The late Devonian brachiopod *Cyrtiopsis davidsoni* Grabau 1923, and related forms from central Hunan of South China. *Journal of Paleontology* 73, 608–624, doi.org/10.1017/S0022336000032431

Ma, X. –P., Day, J., 2000. Revision of *Tenticospirifer* Tien, 1938, and morphologically similar spiriferid brachiopod genera from the Late Devonian (Frasnian) of Eurasia, North America, and Australia. *Journal of Paleontology* 74, 444–463, doi.org/10.1017/S0022336000031711

Ma, X. –P., Day, J., 2007. Morphology and revision of Late Devonian (Early Famennian) *Cyrtospirifer* (Brachiopoda) and related genera from South China and North America. *Journal of Paleontology* 81, 286–311, doi.org/10.1666/0022-3360(2007)81[286:MAROLD]2.0.CO;2

Mamedov, A.B., Rzhonsnitskaya, M.A., 1985. Devonian of the south Transcaucasus: Zonal subdivision, boundaries of series and stages, correlation. *Courier Forschungsinstitut Senckenberg* 75, 135–156.

Milanovsky, E.E., 2007. Hermann Abich (1806–1886): the “Father of Caucasian Geology” and his travels in the Caucasus and Armenian Highlands. *Geological Society, London, Special Publications* 287, 177–181, doi.org/10.1144/SP287.14

Mottequin, B., 2019. An annotated catalogue of types of Silurian–Devonian brachiopod species from southern Belgium and northern France in the Royal Belgian Institute of Natural Sciences (1870–1945), with notes on those curated in other Belgian and foreign institutions. *Geologica Belgica* 22 (1–2), 47–89, doi.org/10.20341/gb.2019.005

Mottequin, B., Brice, D., 2019. Reappraisal of some Upper Devonian (Famennian) spiriferide brachiopods from the Band-e Bayan Domain (Afghanistan). *Geobios*, 52, 47–65, doi.org/10.1016/j.geobios.2018.11.003

Nalivkin, D.V., 1930. Brachiopody verkhnego i srednego devona Turkestana (Brachiopods from the Upper and Middle Devonian of Turkestan), *Trudy Geologicheskogo Komiteta. Novaya seriya*, 180, 1–221 (in Russian).

- Oleneva, N.V., 2016. Devonian Brachiopods of the Orders Spiriferida and Spiriferinida of the European Russia and Transcaucasia: Systematics, Shell Microstructure. *Paleontological Journal*, 50 (11), 1207–1296, doi:10.1134/S0031030116110010
- Reed, F.R.C., 1922. Devonian fossils from Chitral and the Pamirs. *Memoirs of the Geological Survey of India. Palaeontologia Indica*, 6/2, 1–134.
- Rzonsnitskaya, M.A., 1948. Devonskiye otlozheniya Zakavkaz'ye (Devonian deposits of Transcaucasia). *Doklady Akademii Nauk SSSR*, 59/ 8, 1477–1480 (in Russian).
- Rzhonsnitskaya, M.A., Mamedov, A.B., 2000. Devonian stage boundaries in the southern Transcaucasus. *Courier Forschungsinstitut Senckenberg*, 225, 329–333.
- Sartenaer, P., 1974. Signification stratigraphique du 'niveau des monstres' du Frasnien franco-belge. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, 50/7, 1–19.
- Serobyanyan, V., Grigoryan, A., Mottequin, B., Mayilyan, R., Crônier, C., Danelian, T., 2019. Biostratigraphy of the Upper Devonian trigonirhynchiid brachiopods (Rhynchonellida) from Armenia. *Proceedings of National Academy of Sciences of the Republic of Armenia. Earth Sciences*, 72/2, 3–18.
- Sidjachenko, A.I., 1962. Spiriferidy i stratigrafiya famenskikh otlozheniy tsentral'nogo i yugo-vostochnogo Karatau (Spiriferides and stratigraphy of the Famennian deposits of the Central and Southeastern Kara-Tau). *Akademiya Nauk SSSR, Sibirskoje otdelenie, Moskva*, 147 p. (in Russian).
- Sokiran, E. V., 2006. Early–Middle Frasnian cyrtospiriferid brachiopods from the East European Platform. *Acta Palaeontologica Polonica*, 51/4, 759–772.
- Sosson, M., Rolland, Y., Müller, C., Danelian, T., Melkonyan, R., Kekelia, S., Adamia, S., Babazadeh, V., Kangarli, T., Avagyan, A., Galoyan, G., Mosar, J., 2010. Subductions, obduction and collision in the Lesser Caucasus (Armenia, Azerbaijan, Georgia), new insights. In: M. Sosson, N. Kaymakci, R. A. Stephenson, F. Bergerat, V. Starostenko, (Eds), *Sedimentary Basin*

Tectonics from the Black Sea and Caucasus to the Arabian Platform. Geological Society, London, Special Publications, 340, pp. 329–352, doi.org/10.1144/SP340.14

Spaletta, C., Perri, M.C., Over, D.J., Corradini, C., 2017. Famennian (Upper Devonian) conodont zonation: revised global standard. *Bulletin of Geosciences*, 92, 31–57.

Termier, H., Termier, G., 1949. Essai sur l'évolution des Spiriféridés. *Notes et Mémoires, Service géologique, Division des Mines et de la Géologie (Maroc)*, 74, 85–112.

Waagen, W.H., 1883. Salt Range fossils, I. *Productus*–Limestone Fossils. Brachiopoda. *Palaeontologia Indica, Series*, 13–14, 391–546.

Figure 1. Schematic geological map of the South–Armenian Block; distribution of the Upper Devonian–Lower Carboniferous deposits with localization of the sections containing *Aramazdospirifer orbelianus* (Abich, 1858).

Figure 1. Carte géologique schématique du Bloc Sud–Arménien; distribution des dépôts du Dévonien supérieur–Carbonifère inférieur avec localisation des coupes contenant *Aramazdospirifer orbelianus* (Abich, 1858).

Figure 2. Biostratigraphic scheme of the Lower Famennian of the Lesser Caucasus (Transcaucasus) correlated with the major lithostratigraphic units.

Figure 2. Schéma biostratigraphique du Famennien inférieur de la partie méridionale du Petit Caucase (Transcaucasie) corrélé avec les principales unités lithostratigraphiques.

Figure 3. *Aramazdospirifer orbelianus* (Abich, 1858). 1–6 and 18–20. Almost complete specimen (IGSNASRAGM 3895/PS 3001) from the Djravank section in ventral, dorsal, posterior, lateral, posterodorsal and anterior views (1–6), close–up of micro–ornament (capillae and growth lines) on dorsal valve (close to anterior margin; 18–20). 7–8. Incomplete specimen (IGSNASRAGM 3896/PS 3002) from the Noravank section, close–up of the ventral interarea (delthyrium; 7). 9–14. Almost complete specimen (IGSNASRAGM 3897/AB97/48; neotype, Noravank section) in ventral, dorsal, posterodorsal, posterior, anterior and lateral views. 15–22. Almost complete specimen (IGSNASRAGM 3898/PS 3004) from the Ertych section in ventral, dorsal, lateral, posterior and anterior views. 23–27. Almost complete juvenile specimen (IGSNASRAGM 3899/PS 3005) from the Djravank section in ventral, dorsal, posterior, anterior and lateral views. Scale bars: 10 mm (1–6, 9–14, 15–17, 21–27), 2 mm (7), 5 mm (8), 200 μm (18), 400 μm (19), 500 μm (20).

Figure 3. *Aramazdospirifer orbelianus* (Abich, 1858). 1–6 et 18–20. Spécimen quasi–complètement (IGSNASRAGM 3895/PS 3001) de la coupe de Djravank en vues ventrale, dorsale, postérieure, latérale, postérodorsale et antérieure (1–6), vue de détail de la micro–ornementation (capillae et lignes de croissance) sur la valve dorsale (proche de la marge antérieure 18–20). 7–8. Spécimen incomplet (IGSNASRAGM 3896/PS 3002) de la coupe de Noravank, vue de détail de l’interarea ventrale (delthyrium; 7). 9–14. Spécimen articulé quasi complet (IGSNASRAGM 3897/AB97/48; néotype, coupe de Noravank) en vues ventrale,

dorsale, postérodorsale, postérieure, antérieure et latérale. 15–22. Spécimen quasi complet (IGSNASRAGM 3898/PS 3004) de la coupe d'Ertych en vues ventrale, dorsale, latérale, postérieure et antérieure. 23–27. Spécimen juvénile quasi complet (IGSNASRAGM 3899/PS 3005) de la coupe de Djravank en vues ventrale, dorsale, postérieure, antérieure et latérale. Echelles: 10 mm (1–6, 9–14, 15–17, 21–27), 2 mm (7), 5 mm (8), 200 µm (18), 400µm (19), 500 µm (20).

Figure 4. *Aramazdospirifer orbelianus* (Abich, 1858) from the Ertych section. The arrows indicate the anterior margin. 1–7. Almost complete specimen (RBINS a13459) in ventral (muscle field and long intrasinal dental plates are visible), dorsal and anterolateral views, close-up of the pseudodeltidium, and detail of the capillate micro-ornament observed in sulcus near the anterior margin. 8–11. Incomplete specimen (RBINS a13460) in ventral (slightly inclined showing the intrasinal dental plates and the median fold in sulcus) and dorsal views, close-up of the myophragm, and detail of the capillate micro-ornament observed on the left flank of the dorsal valve. 12–13. Poorly preserved articulated specimen (RBINS a13461), view of the ventral area showing the pseudodeltidium and close-up of the latter. Scale bars: 10 mm (1–3, 8–9), 5 mm (4–5, 10, 13), 5 mm (12), 2 mm (11), 1 mm (6, 7).

Figure 4. *Aramazdospirifer orbelianus* (Abich, 1858) de la coupe d'Ertych. Les flèches indiquent le bord antérieure. 1–7. Spécimen quasi complet (RBINS a13459) en vue ventrale (les régions du muscle et des longues plaques dentaires intrabasinales sont visibles), vues dorsale et antérolatérale, vue de près du pseudodeltidium et détail de la micro-ornementation capillate observée au niveau du sulcus près de la marge antérieure. 8–11. Spécimen incomplet (RBINS a13460) en vues ventrale (légèrement inclinée montrant les plaques dentaires intrasinales et le plissement médian au niveau du sulcus) et dorsale, vue de détail du myophragme, et détail de la micro-ornementation capillate observée sur le côté gauche de la valve dorsale. 12–13. Spécimen articulé mal préservé (RBINS a13461), vue de la région ventrale montrant le pseudodeltidium et vue de près de ce dernier. Echelles: 10 mm (1–3, 8–9), 5 mm (4–5, 10, 13), 5 mm (12), 2 mm (11), 1 mm (6, 7).

Figure 5. Transverse serial sections of *Aramazdospirifer orbelianus* (Abich, 1858) from the Djravank section. Numbers refer to distances in mm from the tip of the ventral umbo. Scale bars: 5 mm.

Figure 5. Coupes s eries transversales de *Aramazdospirifer orbelianus* (Abich, 1858) de la , coupe de Djravank. Les chiffres se r f rent   des distances en mm de l'extr mit  de l'umbo ventral. Echelle: 5 mm.

Figure 6. Scatter diagrams of *Aramazdospirifer orbelianus* (Abich, 1858). Abbreviations: N, number of specimens measured; p: probability value; r: coefficient of correlation. 1, Relation between shell width and length. 2, Relation between shell width and thickness. 3, Relation between shell width and width of sulcus. 4, Relation between shell width and length of dorsal valve.

Figure 6. Diagrammes de dispersion d'*Aramazdospirifer orbelianus* (Abich, 1858). Abr viations: N, nombre de sp cimens mesur s; p: valeur de probabilit ; r: coefficient de corr lation. 1, Relation entre la largeur et la longueur de la coquille. 2, Relation entre la largeur et l' paisseur de la coquille. 3, Relation entre la largeur de la coquille et la largeur du sulcus. 4, Relation entre la largeur et la longueur de la valve dorsale.

Figure 7. Measurements and ratios in mm of *Aramazdospirifer orbelianus* (Abich, 1858) Abbreviations: W–width of the shell, L–length of the shell, T–thickness of the shell, Ws–width of the sulcus, dL–length of the dorsal valve.

Figure 7. Mesures et rapports en mm de *Aramazdospirifer orbelianus* (Abich, 1858) Abr viations: W–largeur de la coquille, L–longueur de la coquille, T– paisseur de la coquille, Ws–largeur du sulcus, dL–longueur de la valve dorsale.

Table 1. Measurements and ratios in mm of *Aramazdospirifer orbelianus* (Abich, 1858) Abbreviations: W–width of the shell, L–length of the shell, T–thickness of the shell, Ws–width of the sulcus, dL–length of the dorsal valve.

Tableau 1. Mesures et rapports en mm de *Aramazdospirifer orbelianus* (Abich, 1858) Abr viations: W–largeur de la coquille, L–longueur de la coquille, T– paisseur de la coquille, Ws–largeur du sulcus, dL–longueur de la valve dorsale.