



HAL
open science

Cancer's vulnerability to food availability is evolutionarily conserved: diet modulates tumorigenesis in both Hydra and zebrafish

Sophie Tissot, Lena Guimard, Jordan Meliani, Justine Boutry, Antoine M Dujon, Jean-pascal Capp, Jácint Tökölyi, Peter A. Biro, Christa Beckmann, Laura Fontenille, et al.

► To cite this version:

Sophie Tissot, Lena Guimard, Jordan Meliani, Justine Boutry, Antoine M Dujon, et al.. Cancer's vulnerability to food availability is evolutionarily conserved: diet modulates tumorigenesis in both Hydra and zebrafish. 2023. hal-04419834

HAL Id: hal-04419834

<https://hal.science/hal-04419834>

Preprint submitted on 26 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Cancer's vulnerability to food availability is evolutionarily conserved: diet modulates tumorigenesis in both Hydra and zebrafish

Sophie TISSOT (✉ sophie.tissot@ird.fr)

Université de Montpellier, CNRS, IRD

Lena GUIMARD

Université de Montpellier, CNRS, IRD

Jordan MELIANI

Université de Montpellier, CNRS, IRD

Justine BOUTRY

Université de Montpellier, CNRS, IRD

Antoine M DUJON

Deakin University

Jean-Pascal CAPP

University of Toulouse, INSA, CNRS, INRAE

Jácint TOKOLYI

University of Debrecen

Peter A. BIRO

Deakin University

Christa BECKMANN

Western Sydney University

Laura FONTENILLE

AZELEAD

Nam DO KHOA

AZELEAD

Rodrigo HAMEDE

University of Tasmania

Benjamin ROCHE

Université de Montpellier, CNRS, IRD

Beata UJVARI

Deakin University

Aurora M NEDELCO

University of New Brunswick

Frédéric THOMAS

Article

Keywords:

Posted Date: September 5th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3255124/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on November 14th, 2023. See the published version at <https://doi.org/10.1038/s41598-023-46896-1>.

Abstract

Cancer is a disease that affects the control of cell proliferation in many multicellular lineages. Nonetheless, little is known about the extent of conservation of the biological traits and ecological factors that promote or inhibit tumorigenesis across the metazoan tree. Particularly, changes in food quantity and quality have been linked to increased cancer incidence in humans, an outcome of evolutionary mismatch. Here, we apply evolutionary ecology principles to test the hypothesis whether food availability, regardless of the multicellular lineage considered, should govern tumorigenesis. We used two phylogenetically unrelated model systems, namely the cnidarian *Hydra oligactis* and the fish *Danio rerio* to investigate the impact of resource availability on tumor emergence and progression. Individuals from healthy and tumor-prone lines were placed on four diets that differed in feeding frequency and quantity. For both models, frequent overfeeding favored tumor emergence, while lean diets appear more protective. When investigating tumor progression, high food availability promoted it, whereas low resources controlled it, but without curing the animals. We discuss our results in light of current ideas about the possible conservation of basic processes governing cancer in metazoans (including ancestral life history trade-offs at the cell level) and in the framework of evolutionary medicine.

Introduction

One of the leading hypotheses of evolutionary oncology is that the increase in human cancer incidence in industrialized countries reflects the mismatch between our adaptations to ancestral environments/lifestyles and the fast-changing existence that humans in modern societies are currently experiencing [1]. Among the long list of lifestyle factors that have been proposed to promote oncogenesis in industrialized populations (e.g., lack of exercise, sun exposure, chronic stress, lack of sleep), one that is often cited is diet. For instance, the westernization of the diet in Asian countries, characterized by an overabundance of readily available high caloric foods, has been accompanied by an increase in cancer rates and/or associated mortality (e.g. [2], [3]). The exploration of the oncogenic consequences of diet-related evolutionary mismatches is more than ever a timely topic given that many human populations have recently (i.e. since the industrial revolution), or are in the process, of changing their dietary habits [4]. Further, as it is common for wildlife to feed on human waste and consume food that they have not evolved with, the problem of food mismatches generating cancers is also relevant to several wildlife species [5], [6].

An important aspect of dietary evolutionary mismatch, at least for humans in modern societies, is not only the quality but also the quantity of food. Indeed, compared to ancestral times, a major dietary change in industrialized populations concerns the food intake per meal and the frequency of meals, which have both increased in many geographical areas [7]–[9]. The generally accepted evolutionary explanation behind this propensity to overeat in these populations is that contemporary humans have remained adapted to past environments, in which it was advantageous in terms of selective value to constantly seek and consume food, because food was scarce and/or of variable availability over time

[10]. The current maintenance of these habits is now maladaptive in most developed countries due to the abundance and regularity of food resources [11].

Many health problems (e.g., obesity, cardiovascular disease, diabetes, cancers) are associated with the resulting excessive dietary intakes, especially when combined with reduced physical activity relative to our ancestors [12]–[14]. Not surprisingly, several studies have shown that calorie restriction has a protective effect against cancer, especially when applied at an early stage ([15], [16]; for review [17], [18]). While previous data indicate that intermittent fasting might be more important than caloric restriction in preventing prostate cancer growth in SCID mice [19], recent work on a murine breast cancer model revealed that daily calorie restriction is more potent against primary tumor growth and metastatic burden than periods of fasting and diet composition [20]. In a related vein, Simone et al. [21] proposed that there are actually three types of diets that can not only prevent cancer but also reduce its progression: calorie restriction without malnutrition, intermittent fasting and the ketogenic diet. All have the advantage of limiting, among other things, the carbohydrate intake necessary for tumor growth with limited, if any, detriment to healthy cells.

Mechanistically, cancer cells' vulnerability to food availability is thought to be the result of their inability to switch to the maintenance program under nutrient-limited conditions [22]. Instead, cancer cells continue to grow and proliferate and become susceptible to stress-induced damage [23], [24]. The ability to switch from a reproductive to a survival mode in stressful environments is a fundamental property of all cells and reflects a basic life history trade-off [25]. If the cancer cells' vulnerability to food availability, well-known in human and rodent models, reflects this fundamental trade-off between reproduction/proliferation and survival/maintenance, we would expect to observe similar outcomes in other animal systems that develop tumors. Note that although not all tumors are, or have the potential to become, cancer (i.e., malignant), tumor development is a pre-requisite for any oncogenic process. Thus, any negative effect of food availability on tumorigenesis will directly affect the potential for cancer and its progression.

To address the possibility that food availability can also affect tumorigenesis in non-human systems, we used a freshwater invertebrate (*Hydra oligactis*) and a vertebrate (the zebrafish *Danio rerio*) model and investigated if/how food levels and frequency of availability can affect the initiation and progression of spontaneous, transmitted, and experimentally induced tumors. Specifically, the experiments in this study were designed to (i) investigate whether the reported increased sensitivity to starvation of cancer cells in humans is also characteristic of tumoral cells in other animal lineages (and might be partially responsible for the reduced incidence of tumors/cancers in the wild) and (ii) address the potential role of evolutionary mismatch (in terms of increased food availability relative to natural/ancestral systems) in promoting tumorigenesis in non-human animal lineages.

Both model systems used in this study possess strains that can develop tumors with known dynamics. In hydras, tumor-like abnormal growths (resulting from the uncontrolled proliferation of the germ stem cells) have been reported in several species and strains in laboratory settings [26], [27]. Such tumors can appear

spontaneously, but in some strains they can also be transmitted vertically during asexual reproduction. Specifically, in one *Hydra oligactis* strain maintained for more than 15 years in the lab, the tumors can be transmitted from the parental polyp to offspring (through the bud), which will develop tumors in about four weeks (see Fig. 1) [26]. In addition, individuals collected from the wild, which are tumor-free, can sporadically develop tumors when they are kept in the lab [28]. [28]. The mechanisms underlying the development and the transmission of these tumors as well as their impact on the biology and the ecology of the hosts are not well understood [29], [30]. Notably, at least in one *H. oligactis* strain bearing transmissible tumors, the tumor-associated microbiome might also play a role in the maintenance of tumors [31]. In zebrafish, a genetically modified strain has been developed that spontaneously develops a pigmentation abnormality (i.e. nevi) that can then progress into skin tumors in almost 100% of cases (see Fig. 2) [32]. We used both healthy individuals as well as tumor-bearing individuals at early and late stages of tumorigenesis (i.e., before and after the appearance of nevi or tumors).

We formulated three hypotheses and specific predictions. (i) Under nutrient deprivation, pre-cancerous cells that are defective in controlling their proliferation are unable to activate their maintenance program and will continue to divide and sustain stress-induced damage and death; thus, low resource levels or fluctuations in resource levels (such as during calorie restriction or intermittent fasting; and also likely to characterize natural systems) should have a preventive or even curative/purging role during early stages of tumorigenesis. (ii) Abundant and/or frequent food availability (reflecting artificial settings; i.e., evolutionary mismatch) allows pre-cancerous cells to avoid the consequences of the proliferation/maintenance trade-off (and its purging role) induced by nutrient limitation and take advantage of available resources to promote proliferation; thus, such diets facilitate tumorigenesis and/or tumor progression. (iii) As tumor cells accumulate genetic and epigenetic disorders, they are less affected by ancestral trade-offs (as they acquire additional ways to withstand stress in response to changes in the tumor microenvironment); thus, fully developed tumors should be less impacted by the scarcity of resources. Additionally, a lack of resources at this late stage may in fact promote tumor progression by lowering the efficiency of host anticancer defenses.

Materials and Methods

The general experimental design used for this experiment is summarized in Fig. 3 and detailed in the next sections.

Hydra

We used two different sets of hydras. One set originated from the so-called St. Petersburg lineage (*H. oligactis*), which include both healthy (tumor-free) and tumoral clonal strains that have been kept in culture for more than 15 years (described in [26]) (see Fig. 1). The other set was composed of wild hydras collected 02/05/2022 from the field (Montaud lake in France (43°44'52"N; 3°59'23"E) on May 2nd, 2022); under our lab conditions, these wild hydras can spontaneously develop tumors [28]. The specific lab

culture conditions are detailed in [30]; briefly, polyps are grown at 18°C and fed *ad libitum* 3 times a week with *artemia nauplii*.

For hydras from the St. Petersburg lineage, to test the effect of nutrition on tumor emergence, four groups (corresponding to the feeding regimes described below) of 60 juveniles (i.e. freshly detached from the parental polyp) from both the tumor-free strain and the tumoral strain (note that tumors are not visible at this stage) were isolated and placed individually in wells of standard cell culture plates (12-well plates, Thermo Scientific, 1.5 ml/well; the plates were changed every month).

Then, to evaluate the effect of nutrition on tumor progression, the experiment was started with individuals that already developed tumors. To obtain those individuals, four groups (corresponding to the feeding regimes described below) of 60 juveniles from the tumor-free strain and 60 juveniles from the tumoral strain were placed in similar plates for five weeks, as in the previous experiment. They were fed *ad libitum* 3 times a week with *artemia nauplii* (*Artemia salina*, Planktovie S.A.S., Marseille, France) hatched according to the procedure described in [21], then 8 hours after feeding the remaining *artemia* were removed from the wells and the water level was restored. The experiment was initiated when the hydras were five weeks old – corresponding to the time when the tumor phenotype emerges in the tumoral strain (i.e. tumors are visible size and the number of tentacles increases; [26]). At that time, the tumoral strain presented heterogeneous intensities of expression of the tumor phenotype depending on the individuals (discussed later). Details on the number of juveniles and aged hydras at the start of the experiment (i.e. considering the mortality that occurs between birth and the start of the experiment) for each batch, diet and status are available in the supplementary material (Table S1).

Concerning the wild hydras from Montaud, directly after sampling, four groups of 48 individuals were placed individually under the four different feeding regimes in the same plates as described above. Details on the numbers of individuals for each diet are available in the supplementary material (Table S2).

All the polyps were exposed to four diets: (i) high-abundance and frequent, with *ad libitum* feeding five times a week, (ii) high-abundance and intermittent with *ad libitum* feeding once a week, (iii) low-abundance and frequent feeding with 5 *artemia* individuals five times a week, and (iv) low-abundance and intermittent feeding with 5 *artemia* individuals once a week. For juvenile hydras, the appearance of tumors was monitored for two months, and the intensity of the tumor phenotype expression was visually quantified under a dissecting microscope using a 6-level ordinal scale of tumor phenotype development. This scale was validated using the ratio of large to small germinal cells (the cell type associated with tumors in hydras, [26]) ensuring that visual tumor progression matched the accumulation of abnormal cells. For 5-week-old tumoral hydras, the tumor phenotype expression was measured at the beginning of the experiment and one month later. Full detail on the scale used for the measurement is provided in the supplementary material (Figure S1 and Figure S2).

Zebrafish

The kita-GFP-RAS transgenic zebrafish line, which is known to spontaneously develop premalignant lesions (nevi) and tumours/melanomas (see Fig. 2) [33], was used in this study to non-invasively measure the prevalence and the timing of nevi and tumors. The strain AB was used as control. Larval and adult zebrafish were maintained on a light/dark cycle of 12/12h in a system with a partial renewal of the water up to 30%. Embryos were obtained by natural spawning of breeding pairs and were kept at 28.5°C in the tank water. At 9 days post fertilization (dpf) the larvae were introduced into the rearing system, with 50 individuals in a tank in order to obtain at least 20 fish at the age of three months. All experimental procedures on zebrafish were carried out in accordance with European directives and French Ministry of Health animal protection regulations (approval number: F341725) by the company Azelead based in Montpellier, France. The experiments were approved by the ethics committee for animal experimentation (Direction départementale de la protection des populations - Hérault). The study was conducted in accordance with ARRIVE guidelines.

To test the effect of nutrition on tumor emergence (i.e., nevi onset), the number of larvae was fixed at 200 larvae from the tumoral strain (i.e. kita-GFP-RAS strain) and 200 larvae from the control strain (i.e. AB strain) to consider the high natural mortality expected at the larval stage affecting about 50% of the individuals (L. Fontenille, *personal communication*). For each strain the larvae were separated into four groups and placed in identical aquariums to those used for rearing. These zebrafish were fed with four diets from 5 dpf.

To evaluate the effect of nutrition on tumor progression, 400 larvae were placed in identical aquariums and fed in a conventional manner (described in supplementary, Table S3) for three months. Then, for the tumoral strain, 80 individuals with nevi were selected and separated into the four experimental groups; for the control strain 80 individuals were also randomly selected and subjected to the same treatment as the tumoral strain.

All zebrafish were fed with four different diets, using the same categories as for the hydra (see above), but this time on a dry food basis and with artemia as enrichment. For juvenile zebrafish (i.e. 5dpf), the diets have been adapted to follow the change in their food needs over time and respected the differences between each diet. The constitution of the different diets used for zebrafish is available in the supplementary material (Table S3). In juvenile zebrafish, the appearance of nevi was followed for six and a half months, and in three-month-old zebrafish that already had nevi, the appearance of tumors was followed for seven months.

Data analysis

The analyses presented here were performed with the R software (version 4.2.2) [34] and the graphical representations were realized with the “GGplot2” package [35]. The prevalence of tumors across time for hydras from the tumor-free and the tumoral strains, and of nevi and tumors for zebrafish from the tumoral strain were analyzed according to the diet with survival regressions. These specific regressions allow to treat censure data whether or not the instantaneous risk of observing the phenomenon of interest (in this case, the appearance of a tumor or nevi) is constant, by changing the distribution law in the model

[36]. The intensity of tumor phenotype expression of hydras from the tumoral strain and its progression over one month were analyzed according to diet with respectively cumulative link models (CLM) and cumulative link mixed models (CLMM), a random effect was included for the latter to account for the lack of independence between repeated measurement of each individual. The number of replicates was too small to quantify potential batch effects in hydra experiments while the zebrafish experimental design only used a single batch. The Table 1 summarizes the distribution, function, and associated packages used to explore each trait. Then, the procedure detailed by [37] was followed to operate the model selection based on the weight of the Akaike information criterion (AIC) or on this corrected version (AICc) for models with a random effect. The formulas of all the different models constructed, as well as the value of AIC or AICc's weight were obtained with the "MuMIn" package [38], and are detailed in the supplementary material (Table S4). The prevalence of tumors for wild hydras was compared between diets with a test of given proportion.

Table 1

Details of distributions, functions, and R packages used for each trait measured for juvenile hydras (*H. oligactis*), juvenile zebrafish (*D. rerio*), five-week-old hydras, and three month old zebrafish. Functions are presented in italics and the associated R packages in parenthesis.

Biological model	Trait	Distributions, functions, and R packages used	
		Juvenile individuals	Aged individuals
Hydras from tumor-free strain	Tumor prevalence	Weibull <i>survreg</i> (survival [39])	
	Tumor timing		
Hydra from tumoral strain	Tumor prevalence	Weibull <i>survreg</i> (survival [39])	
	Tumor timing		
	Tumor size	Logistic <i>clm</i> (ordinal [76])	Logistic <i>clmm</i> (ordinal [76])
Zebrafish from tumoral strain	Nevi prevalence	Weibull <i>survreg</i> (survival [39])	
	Nevi timing		
	Tumor prevalence	Weibull <i>survreg</i> (survival [39])	
	Tumor timing		
Wild hydras	Tumor prevalence	Binomial <i>prop.test</i> (stat [77])	

During the experiments with hydras, the appearance of spontaneous tumors in individuals from the tumor-free strain was monitored. This special type of tumor, which differs from those observed in the St. Petersburg tumoral strain, is known to appear sporadically and with a low prevalence in hydra kept in

laboratory (a more detailed description is available in [28]). Therefore, the prevalence of these spontaneous tumors was analyzed according to the diet treatment using a survival regression as previously described in [39]. The formulas of all the different models constructed, as well as the value of AIC or conditional AIC's weight are detailed in the supplementary material (Table S4) and details of the model selected are summarized in Table 1.

Results

The effect of food availability on tumorigenesis in juvenile hydras from the St. Petersburg lineage

To assess the effect of food quantity and frequency on tumor emergence and/or progression, juvenile hydras from both tumor-free and tumoral strains (before tumors are visible) were subjected to four diets varying in food quantity and frequency. For hydras from the tumor-free strain we recorded the rate of appearance of spontaneous tumors. For hydras from the tumoral strain, tumor development was monitored for two months and, at the end, the intensity of tumor phenotype expression was measured to assess the effect of feeding on tumor progression.

We found that diet significantly affected the emergence of spontaneous tumors in juvenile hydras from the tumor-free strain. The proportion of individuals developing tumors increased as the frequency and/or quantity of food increased. The high-abundance and frequent diet led to a significantly higher rate (90% vs 35%) of tumor occurrence (Fig. 4A; survival regression; estimate = 0.07, SE = 0.04, p-value = 1.5e-06) compared to the low-abundance and frequent diet (Fig. 4A; survival regression; estimate = 0.19, SE = 0.53, p-value = 2e-03).

For juvenile hydras from the tumoral strain, the models selected to explain the tumor prevalence across time indicated a significant effect of the diet. The high-abundance and frequent diet was associated with the highest tumor development rate (Fig. 4B; survival regression; estimate = 0.11, SE = 0.04, p-value = 3.4e-10), followed by the high-abundance and intermittent diet (Fig. 4B; survival regression; estimate = 0.27, SE = 0.10, p-value = 3.9e-04), the low-abundance and frequent diet (Fig. 4B; survival regression; estimate = 0.46, SE = 0.17, p-value = 3.3e-02), and finally the low-abundance and intermittent diet with the lowest tumor development rate.

Interestingly, in the tumor-free strain, the low-abundance and frequent diet has a stronger effect compared to the high-abundance and intermittent diet whereas we observed the reverse situation in the tumoral strain. Thus, the frequency seems to be more important in the former case whereas it is the abundance in the latter, suggesting different underlying mechanisms in the two strains.

Lastly, for hydras in the tumoral strain that developed tumors, the model selected to explain the intensity of tumor phenotype expression includes the effect of the diet, by revealing that hydras on a high-

abundance and frequent diet developed more intense tumor phenotypes than those on the other three diets (Fig. 4C. CLM; OR = 10.32, SE = 9.66, p value = 1.3e-02).

The effect of food availability on tumorigenesis in five-week-old hydras from the St. Petersburg lineages

Five-week-old hydra from both the tumor-free and the tumoral strains were subjected, as above, to four diets varying in food quantity and frequency. The rate of spontaneous tumor development in hydra from the tumor-free strain was recorded. For hydra from the tumoral strain (with visible tumors), the intensity of tumor phenotype expression was measured at the beginning of the experiment and one month later.

To explain the proportion of hydras from the tumor-free strain developing spontaneous tumors when the different diets are applied at five weeks of age, the model with the best AIC contained a significant effect of diet. The proportion of hydras that developed spontaneous tumors increased with the frequency and quantity of provided food. The low-abundance and intermittent diet led to the lowest spontaneous tumor development rate with less than 5% of individuals showing tumors. The high-abundance and intermittent diet presented a higher rate with 25% of individuals developing tumors (Fig. 4D; survival regression; estimate = 0.20 SE = 0.12, p-value = 5.05e-03), the low-abundance and frequent diet has a rate of 50% of individuals (Fig. 4D; survival regression; estimate = 0.10, SE = 0.06, p-value = 8e-05), and finally the high-abundance and frequent diet with shows the highest rate of 85% of individuals developing spontaneous tumors (Fig. 4D; survival regression; estimate = 0.04, SE = 0.02, p-value = 3.6e-08).

In the tumoral hydra strain, we detected a significant effect of the interaction between diet and time on tumor progression. The intensity of tumor phenotype expression increased over time in hydras in the high-abundance and frequent diet group (Fig. 4E; CLMM; OR = 5.00, SE = 3.44, p-value = 1.9e-02) while it decreased in hydras in the low-abundance and intermittent diet group (Fig. 4E; CLMM; OR = 0.18, SE = 0.15, p-value = 3.8e-02). In addition, we observed a high mortality rate (half-life of less than 30 days) in hydras in the low-abundance and intermittent diet group. For those individuals, the intensity of tumor phenotype expression was measured on moribund animals, whose body was altered, leading to an underestimation of this intensity (our personal observations).

The effect of food availability on tumorigenesis in wild hydras

We found that out of the four diet treatments, wild hydra only developed spontaneous tumors in the high-abundance and frequent group and at a rate of 33% of individuals (Fig. 5; test of given proportion; X-squared = 16.87, p-value = 3.99e-05).

The effect of food availability on tumorigenesis in juvenile zebrafish

Juvenile zebrafish from both control and tumoral strains were subjected to four diets varying in food quantity and frequency, and tumor onset was surveyed for six months and half. Zebrafish from the control strain did not show nevi or tumor development, regardless of diet. In the fish from the tumoral strain, the proportion of individuals developing nevi is well explained by the effect of the diet; an increase in food availability was associated with an increased development of tumors. Zebrafish exposed to the high-abundance and frequent diet had the highest nevi development rate with 100% of individuals affected (Fig. 6A; survival regression; estimate = 0.40, SE = 0.05, p-value = 3.6e-12), followed by the low-abundance and frequent diet with 88% of individuals affected (Fig. 6A; survival regression; estimate = 0.54, SE = 0.07, p-value = 4.7e-06), the high-abundance and intermittent diet with 68% of individuals affected (Fig. 6A; survival regression; estimate = 0.69, SE = 0.10, p-value = 9.1e-03), and finally the low-abundance and intermittent diet with 32% of individuals developing nevi.

The effect of food availability on tumor progression in three-month-old zebrafish

Three-month-old zebrafish from the control strain and tumoral strain (with visible nevi) were also fed the four diets. Tumor development and/or progression (i.e. from nevi to tumors) was studied for two to seven months. Neither nevi nor tumors were observed in the zebrafish control strain in any of the four diet treatments. Concerning the proportion of zebrafish from the tumoral strain developing tumor across time, the relevant model contained the significant effect of diet. Zebrafish following the high-abundance and frequent diet had higher tumor development rates than those in the three other diets, with 55% of individuals affected (Fig. 6B; survival regression; estimate = 0.15, SE = 0.12, p-value = 1.8e-02).

Discussion

As a result of extensive research on the roles of diet on human cancer dynamics, important advances have been made over the past decades [40]–[44]. For example, it is now clearly established that many cancers can be prevented by adopting appropriate dietary measures [45]–[47]. There is also growing evidence that certain diets are more favorable than others in terms of the likelihood of recovery in cancer patients [48]. However, the subject remains topical, particularly because numerous human populations, especially in industrialized countries, have recently altered, or are in the process of, changing their dietary habits [4]. This will likely generate some evolutionary mismatches with still underestimated oncological consequences [1]. The objectives of this study were to (i) test whether food quantity and frequency affect tumor initiation and progression in non-human systems, and (ii) identify whether tumors' vulnerability to nutrient availability is a conserved trait in evolutionarily distant animal lineages (i.e., mammals, fish and hydra).

The effect of food intake on tumor emergence

In hydra, we explored the link between tumor emergence and diet by considering 3 categories of polyps. They included two types of healthy polyps and one polyp type presumably tumoral but asymptomatic.

The healthy polyps included (i) polyps from the tumor-free laboratory strain, and (ii) polyps from the wild, which are likely healthy since tumor polyps have never been found in nature [28]). The presumed tumoral polyps were juvenile polyps that just detached from tumoral parents, which in standard laboratory conditions usually develop tumors in about four weeks as a result of vertical transmission (see [49]). Our results confirm that diet is a key variable in the likelihood of tumor emergence, with frequent overeating being a factor favoring tumor emergence, whereas lean diets (restricted diet and/or intermittent fasting) appear to be protective.

For the two categories of healthy hydras, it is unclear if high food availability is directly responsible for the initiation of tumors, and/or if it favors the early accumulation of abnormal cells when tumoral cells appear due to other reasons. The food we used (artemia) is not itself mutagenic, but its high availability is likely to boost cell proliferation (since the budding rate of hydras is positively correlated with food intake [50]–[52]) and hence tumoral risks linked to cell divisions. This impact of food abundance is likely due to metabolic effects because reduced calorie intake produces metabolic reprogramming [53] that can prevent or slow down cancer progression [54]. Of note, recent studies showed that reduced caloric intake and periodic fasting do not have the same metabolic effects. The different effects of these modes of caloric restriction might explain their differential impact on tumor growth and metastatic burden [20], suggesting a possible explanation for the effects of the different diets in this study. Furthermore, because tumor development in *H. oligactis* is mediated by bacteria [31], we cannot rule out that food availability alters the composition of the microbiome, which in turn influences tumorigenesis. In any case, our findings support the hypothesis that low food availability has a protective effect, perhaps by limiting the energy supply that first tumoral cells need to grow. Since this observation also applies to hydras from the tumoral strain, it suggests that the protective role of lean diets is strong and durable even when the first tumoral cells are already present inside the body.

Another major lesson from the hydra experiments comes from wild individuals, since this experiment in fact simulated a dietary evolutionary mismatch. Indeed, hydras in the wild consume freshwater crustaceans [55], but not artemia which live in brackish ecosystems. Moreover, it is unlikely that a similar high-abundance and frequent diet as the one we used would be found in the wild. We therefore simulated a substantial qualitative and quantitative dietary change, and our results, showing that only individuals overfed with this new diet develop tumors, strongly supports the idea that dietary evolutionary mismatches can promote the dynamics of tumoral processes.

Finally, our experiments on hydra indicate that a high-abundance and frequent diet favors the emergence of tumors, irrespective of whether the individuals have had this diet from early in life or from five weeks of age. Thus, a lean diet early in life does not provide a lasting protective effect if a switch to food-rich lifestyles occurs later in life. This result is consistent with those of Pomatto-Watson et al. [20] who showed that daily calorie restriction limits tumor processes better than calorie cycling. These findings support the idea that tumor processes are continuous, and that it is the availability of resources at the time they are initiated that explains whether tumors will succeed in progressing. This observation also suggests that a lean diet is protective mainly through non-genetic processes. If it was protective against

mutagenic processes compared to the high-abundance and frequent diet, it would reduce or delay the appearance of tumors after the switch to food-rich lifestyle because less mutated cells would be present to fuel tumor development.

In zebrafish, the link between the emergence of tumoral processes and diet was explored in both healthy (control) individuals and individuals that are expected to develop tumors (i.e. the equivalent of the third category of hydra). In the control group, there was no tumoral development regardless of the diet used, while in the tumoral group there was a higher prevalence of nevi among the overfed individuals, in accordance with previous works showing with a different feeding system that increased feeding enhances melanoma formation in a *p53/BRAF*-dependent model [16]. These findings confirm that a high availability of food is not in itself directly oncogenic, but it boosts the proliferation of (pre)malignant cells that may appear. Stated differently, these findings support the theoretical expectation that low resource levels are detrimental to emerging tumors [56] because malignant cells are highly dependent on energy to proliferate, and/or are more sensitive to nutrient limitation and stress-induced damage. Of note, spontaneous tumorigenesis does not occur in the zebrafish non-tumoral strain regardless of the diet whereas the hydra tumor-free strain developed tumors at a different rate depending on the diet. This result indicates a different sensitivity to diet for tumor development in these different organisms, vertebrates being less sensitive possibly due to stronger defense systems against tumorigenesis.

Results on the impact of food availability quantity and frequency on tumor emergence in different hydra lines and zebrafish also highlight differential sensitivities according to tumor type. Indeed, spontaneous tumors developed by hydra of the tumor-free strain, as well as those developed by zebrafish, show greater sensitivity to feeding frequency than to food quantity, in contrast to the transmissible tumors present in the hydra tumor strain. Although further research is needed to confirm this link, these preliminary results suggest that transmissible tumor cells, thanks to their longer evolutionary time than non-transmissible tumor cells, may have acquired adaptations to better withstand a momentary cut in food availability. Among other things, these adaptations could involve exploiting other metabolic pathways [57] to ensure their maintenance while waiting for a sufficiently large energy supply to develop. Non-transmissible tumor cells, having just occurred in the host organism, are confronted for the first time with this selection pressure which will thus eliminate most of these premalignant cells whereas a low but constant food availability would be sufficient to maintain a greater quantity of tumor cells in the organism and allow some to develop by diverting the energy resources of the host, probably to its detriment.

The effect of food intake on tumor progression

In hydra, overfeeding juvenile polyps results in larger tumors relative to individuals fed lean diets. These results clearly indicate that tumor progression in tumoral polyps is positively influenced by the amount and the frequency of food intake by the host. Consistent with this finding, when the different diets were applied on older hydras with already developed tumors, overfeeding favored tumor progression while the lean diets were associated with a stabilization of tumor size, but not with regression. This supports the idea that lean diets can help control tumor size in tumor-bearing individuals, but is not curative in itself.

In zebrafish, the measure of tumor progression is the proportion of individuals with precancerous lesions (i.e., nevi) that develop tumors following the various diets. The overfed individuals showed higher tumor prevalence while those on lean diets were less likely to develop tumors, even if they kept their premalignant lesions (nevi). These data confirm, as for hydra, that high availability of food boosts the proliferation of tumor cells that are already present and that lower resource levels help to control tumor progression but without being curative. These findings are thus consistent with the conceptual explanation that in advanced tumors, cells may have already evolved ways to survive stress such as that induced by low food availability in the tumor microenvironment [56].

It is interesting to speculate on the reasons why the most food rich treatment, high-abundance and frequent diet, was most detrimental for cancer initiation and progression. One possibility is that this very substantial food ration has used up a considerable proportion of available energy to the body in digestion itself, forcing an energetic tradeoff which might leave less energy available to fight cancer. Indeed, high feeding rations can use upwards up to 70% of fish aerobic scope [58] leaving little energy for other purposes, including energy which is known to be important in fueling anti-cancer processes in the body [59], [60]. If one were to use a human analogy, this situation might be described as a low level ‘food coma’ akin to what we experience after eating excessively, leaving us lethargic due to energy diversion to digestion. Another possibility is that reduced calorie intake reduces the incidence and progression of spontaneous tumors in both hydra and zebrafish through common mechanisms relying on evolutionary conserved intracellular pathways like the Insulin/Insulin-like growth factor (IGF) signaling pathway that connects nutrient levels to metabolism, growth, development, longevity in vertebrates and invertebrates. Of note, *Hydra* utilizes the insulin signaling to regulate its own size [61]. This evolutionary conserved control of cell proliferation and growth promoting factors by insulin signaling might be of particular relevance to explain the fact that cancer’s vulnerability to food availability is a conserved trait.

Cancer’s vulnerability to food availability is evolutionarily conserved and reflects ancestral life history trade-offs

Our results argue that tumor/cancer’s vulnerability to food availability is conserved among three evolutionarily distant lineages. Such deep conservation might reflect ancestral processes associated with the control of cell proliferation. Specifically, although unicellular organisms are often considered “immortal” and in a constant proliferative mode (hence the common view of cancer cells as a reversal to unicellularity; e.g., [62]), they also need to be able to control cell proliferation in response to signals from environment or neighboring cells. For instance, under stress (including nutrient deprivation), single-celled organisms repress proliferation to ensure long-term survival (i.e., a basic life history trade-off) (e.g., [63]). Mutants that are unable to control proliferation do have an immediate reproductive advantage but incur long-term costs in their ability to withstand stress-induced damage (e.g., see discussion and examples in [64]–[71]). It has been suggested that during the transition to multicellularity, the genes involved in the expression of such trade-offs were co-opted into the evolution of the two main specialized cell types – somatic and reproductive [72], [73]. Consequently, mutations in such genes can result in uncontrolled proliferation but also increased sensitivity to stress. Such antagonistic effects are thought to contribute

to the stability of cooperative behaviors, including during the evolution of multicellularity, by eliminating “cheater” cells [74], [75]. This inability to trade-off reproduction for survival in response to stress can still play a role in purging cheaters – including pre-cancerous cells, in extant multicellular lineages [75]. Our findings, together with the proposed differential stress resistance of cancer cells (e.g., [23], [24]) are consistent with this possibility.

Concluding remarks

Although further functional studies would be needed to explain our findings, this study supports the theoretical assumptions that low food availability has a preventive role against tumoral processes when applied at first stages of tumorigenesis, in both human and non-human systems. From an evolutionary perspective, these findings might reflect ancestral life history trade-offs at the cellular level that are expected to be conserved in all multicellular lineages. However, when the tumoral processes are advanced, the scarcity of resources alone is not enough to be curative. Increases in the quantity and/or the frequency of food may facilitate tumorigenesis and in all cases tumor progression. From an evolutionary medicine perspective, these results support the idea that fluctuations in food availability may have acted in the past (and still be acting in species in the wild) as a purging mechanism against premalignant cells that can occur spontaneously and frequently in the host body. The recent increase in food availability by-passes this evolutionarily conserved anti-cancer mechanism and generates evolutionary mismatches leading to higher cancer risk.

Declarations

Acknowledgments

We are grateful to Marina Mione for providing us with the pictures of zebrafish.

Author contributions

The original idea for the article came from discussions between A.M.N., F.T., and S.T.; S.T., J.B., and F.T. designed the study. S.T., L.G., and J.M. collected the data on hydra, whereas L.F., and N.D.K. collected the data on zebrafish. S.T. supported by J.B., and A.M.D. performed the analyses. S.T. wrote the manuscript with the support of F.T., B.U., A.M.N., J.T., P.A.B., C.B., R.H., J-P.C, and B.R., and all authors significantly contributed to revisions.

Data availability statement

Scripts and data are provided in supplementary information.

Competing interests statement

Authors declare no competing interests.

Funding

This work was funded by the MAVA Foundation, André Hoffmann, and by the following grants: ANR EVOSEXCAN.

References

1. C. A. Aktipis and R. M. Nesse, "Evolutionary foundations for cancer biology," *Evolutionary Applications*, vol. 6, no. 1, pp. 144–159, 2013, doi: 10.1111/eva.12034.
2. Y. Kagawa, "Impact of westernization on the nutrition of Japanese: Changes in physique, cancer, longevity and centenarians," *Preventive Medicine*, vol. 7, no. 2, pp. 205–217, Jun. 1978, doi: 10.1016/0091-7435(78)90246-3.
3. C. Heidemann, M. B. Schulze, O. H. Franco, R. M. van Dam, C. S. Mantzoros, and F. B. Hu, "Dietary Patterns and Risk of Mortality From Cardiovascular Disease, Cancer, and All Causes in a Prospective Cohort of Women," *Circulation*, vol. 118, no. 3, pp. 230–237, Jul. 2008, doi: 10.1161/CIRCULATIONAHA.108.771881.
4. J. E. Singh, A. K. Illner, K. Dokova, N. Usheva, T. Kostadinova, and K. Aleksandrova, "Mapping the global evidence on nutrition transition: a scoping review protocol," *BMJ open*, vol. 10, no. 6, Art. no. 6, 2020, doi: 10.1136/bmjopen-2019-034730.
5. M. Giraudeau, T. Sepp, B. Ujvari, P. W. Ewald, and F. Thomas, "Human activities might influence oncogenic processes in wild animal populations," *Nat Ecol Evol*, vol. 2, no. 7, Art. no. 7, Jul. 2018, doi: 10.1038/s41559-018-0558-7.
6. P. A. Pesavento, D. Agnew, M. K. Keel, and K. D. Woolard, "Cancer in wildlife: patterns of emergence," *Nat Rev Cancer*, vol. 18, no. 10, pp. 646–661, Oct. 2018, doi: 10.1038/s41568-018-0045-0.
7. J. M. Brunstrom, A. C. L. Drake, C. G. Forde, and P. J. Rogers, "Undervalued and ignored: Are humans poorly adapted to energy-dense foods?," *Appetite*, vol. 120, pp. 589–595, Jan. 2018, doi: 10.1016/j.appet.2017.10.015.
8. A. Drewnowski and N. Darmon, "The economics of obesity: dietary energy density and energy cost," *Am J Clin Nutr*, vol. 82, no. 1 Suppl, pp. 265S-273S, Jul. 2005, doi: 10.1093/ajcn/82.1.265S.
9. C. Peter Herman, J. Polivy, P. Pliner, and L. R. Vartanian, "Mechanisms underlying the portion-size effect," *Physiol Behav*, vol. 144, pp. 129–136, May 2015, doi: 10.1016/j.physbeh.2015.03.025.
10. K. D. Brownell and K. B. Horgen, *Food fight: the inside story of the food industry, America's obesity crisis, and what we can do about it*. New York: McGraw-Hill, 2004.
11. J. M. Brunstrom and B. K. Cheon, "Do humans still forage in an obesogenic environment? Mechanisms and implications for weight maintenance," *Physiol Behav*, vol. 193, no. Pt B, pp. 261–267, Sep. 2018, doi: 10.1016/j.physbeh.2018.02.038.
12. P. Carrera-Bastos, M. Fontes-Villalba, J. H. O'Keefe, S. Lindeberg, and L. Cordain, "The western diet and lifestyle and diseases of civilization," *RRCC*, vol. 2, pp. 15–35, Mar. 2011, doi: 10.2147/RRCC.S16919.

13. P. Gluckman and M. Hanson, *Mismatch: why our world no longer fits our bodies*. Oxford University Press, 2006. Accessed: Mar. 01, 2023. [Online]. Available: <https://eprints.soton.ac.uk/61151/>
14. P. Gluckman *et al.*, *Principles of Evolutionary Medicine*, Second Edition, Second Edition. Oxford, New York: Oxford University Press, 2016.
15. S. D. Hursting, S. M. Dunlap, N. A. Ford, M. J. Hursting, and L. M. Lashinger, "Calorie restriction and cancer prevention: a mechanistic perspective," *Cancer & Metabolism*, vol. 1, no. 1, p. 10, Mar. 2013, doi: 10.1186/2049-3002-1-10.
16. V. Grigura, M. Barbier, A. P. Zarov, and C. K. Kaufman, "Feeding amount significantly alters overt tumor onset rate in a zebrafish melanoma model," *Biology Open*, vol. 7, no. 1, p. bio030726, Jan. 2018, doi: 10.1242/bio.030726.
17. M. Alidadi, M. Banach, P. C. Guest, S. Bo, T. Jamialahmadi, and A. Sahebkar, "The effect of caloric restriction and fasting on cancer," *Seminars in Cancer Biology*, vol. 73, pp. 30–44, Aug. 2021, doi: 10.1016/j.semcancer.2020.09.010.
18. M. Lv, X. Zhu, H. Wang, F. Wang, and W. Guan, "Roles of Caloric Restriction, Ketogenic Diet and Intermittent Fasting during Initiation, Progression and Metastasis of Cancer in Animal Models: A Systematic Review and Meta-Analysis," *PLOS ONE*, vol. 9, no. 12, p. e115147, Dec. 2014, doi: 10.1371/journal.pone.0115147.
19. W. C. Buschemeyer III *et al.*, "Effect of intermittent fasting with or without caloric restriction on prostate cancer growth and survival in SCID mice," *The Prostate*, vol. 70, no. 10, pp. 1037–1043, 2010, doi: 10.1002/pros.21136.
20. L. C. D. Pomatto-Watson *et al.*, "Daily caloric restriction limits tumor growth more effectively than caloric cycling regardless of dietary composition," *Nat Commun*, vol. 12, no. 1, Art. no. 1, Oct. 2021, doi: 10.1038/s41467-021-26431-4.
21. B. A. Simone *et al.*, "Selectively starving cancer cells through dietary manipulation: methods and clinical implications," *Future Oncology*, vol. 9, no. 7, pp. 959–976, Jul. 2013, doi: 10.2217/fon.13.31.
22. R. Buono and V. D. Longo, "Starvation, Stress Resistance, and Cancer," *Trends in Endocrinology & Metabolism*, vol. 29, no. 4, pp. 271–280, Apr. 2018, doi: 10.1016/j.tem.2018.01.008.
23. C. Lee *et al.*, "Fasting cycles retard growth of tumors and sensitize a range of cancer cell types to chemotherapy," *Sci Transl Med*, vol. 4, no. 124, p. 124ra27, Mar. 2012, doi: 10.1126/scitranslmed.3003293.
24. L. Raffaghello *et al.*, "Starvation-dependent differential stress resistance protects normal but not cancer cells against high-dose chemotherapy," *Proceedings of the National Academy of Sciences*, vol. 105, no. 24, pp. 8215–8220, Jun. 2008, doi: 10.1073/pnas.0708100105.
25. S. C. Stearns, "Trade-Offs in Life-History Evolution," *Functional Ecology*, vol. 3, no. 3, pp. 259–268, 1989, doi: 10.2307/2389364.
26. T. Domazet-Lošo *et al.*, "Naturally occurring tumours in the basal metazoan Hydra," *Nat Commun*, vol. 5, no. 1, Art. no. 1, Jun. 2014, doi: 10.1038/ncomms5222.

27. J. Boutry *et al.*, "Spontaneously occurring tumors in different wild-derived strains of hydra," *Sci Rep*, vol. 13, no. 1, Art. no. 1, May 2023, doi: 10.1038/s41598-023-34656-0.
28. J. Boutry *et al.*, "Spontaneously occurring germline tumors in different wild-derived strains of hydra," in press.
29. J. Boutry *et al.*, "Tumors (re)shape biotic interactions within ecosystems: Experimental evidence from the freshwater cnidarian Hydra," *Science of The Total Environment*, vol. 803, p. 149923, Jan. 2022, doi: 10.1016/j.scitotenv.2021.149923.
30. J. Boutry *et al.*, "Tumors alter life history traits in the freshwater cnidarian, Hydra oligactis," *iScience*, vol. 25, no. 10, p. 105034, Oct. 2022, doi: 10.1016/j.isci.2022.105034.
31. K. Rathje, B. Mortzfeld, M. P. Hoepfner, J. Taubenheim, T. C. G. Bosch, and A. Klimovich, "Dynamic interactions within the host-associated microbiota cause tumor formation in the basal metazoan Hydra," *PLOS Pathogens*, vol. 16, no. 3, p. e1008375, Mar. 2020, doi: 10.1371/journal.ppat.1008375.
32. C. Bevona, W. Goggins, T. Quinn, J. Fullerton, and H. Tsao, "Cutaneous Melanomas Associated With Nevi," *Archives of Dermatology*, vol. 139, no. 12, pp. 1620–1624, Dec. 2003, doi: 10.1001/archderm.139.12.1620.
33. C. Santoriello *et al.*, "Kita driven expression of oncogenic HRAS leads to early onset and highly penetrant melanoma in zebrafish," *PLoS One*, vol. 5, no. 12, p. e15170, Dec. 2010, doi: 10.1371/journal.pone.0015170.
34. R Development Core Team, "R: a language and environment for statistical computing." R Foundation for Statistical Computing, Vienna, Austria, 2020.
35. H. Wickham *et al.*, "ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics." Feb. 10, 2023. Accessed: Feb. 13, 2023. [Online]. Available: <https://CRAN.R-project.org/package=ggplot2>
36. "Maxime HERVÉ - R et statistiques." <https://www.maximeherve.com/r-et-statistiques> (accessed Jul. 04, 2023).
37. A. F. Zuur, E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith, *Mixed Effects Models and Extensions in Ecology With R*. 2009.
38. K. Bartoń, "{MuMIn}: {Multi}-{Model} {Inference}." 2020.
39. T. M. Therneau, T. L. (original S.->R port and R. maintainer until 2009), A. Elizabeth, and C. Cynthia, *survival: Survival Analysis*. 2022. Accessed: Jun. 15, 2022. [Online]. Available: <https://CRAN.R-project.org/package=survival>
40. S. T. Mayne, M. C. Playdon, and C. L. Rock, "Diet, nutrition, and cancer: past, present and future," *Nat Rev Clin Oncol*, vol. 13, no. 8, Art. no. 8, Aug. 2016, doi: 10.1038/nrclinonc.2016.24.
41. G. Grosso *et al.*, "Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk," *Nutr Rev*, vol. 75, no. 6, pp. 405–419, Jun. 2017, doi: 10.1093/nutrit/nux012.

42. R. Murphy, "An Integrative Approach to Assessing Diet–Cancer Relationships," *Metabolites*, vol. 10, no. 4, p. 123, Mar. 2020, doi: 10.3390/metabo10040123.
43. N. Papadimitriou *et al.*, "An umbrella review of the evidence associating diet and cancer risk at 11 anatomical sites," *Nat Commun*, vol. 12, no. 1, Art. no. 1, Jul. 2021, doi: 10.1038/s41467-021-24861-8.
44. S. K. Veettil *et al.*, "Role of Diet in Colorectal Cancer Incidence: Umbrella Review of Meta-analyses of Prospective Observational Studies," *JAMA Netw Open*, vol. 4, no. 2, p. e2037341, Feb. 2021, doi: 10.1001/jamanetworkopen.2020.37341.
45. M. J. Glade, "Food, nutrition, and the prevention of cancer: a global perspective. American Institute for Cancer Research/World Cancer Research Fund, American Institute for Cancer Research, 1997," *Nutrition*, vol. 15, no. 6, pp. 523–526, Jun. 1999, doi: 10.1016/s0899-9007(99)00021-0.
46. K. F. Brown *et al.*, "The fraction of cancer attributable to modifiable risk factors in England, Wales, Scotland, Northern Ireland, and the United Kingdom in 2015," *Br J Cancer*, vol. 118, no. 8, pp. 1130–1141, Apr. 2018, doi: 10.1038/s41416-018-0029-6.
47. D. M. Parkin, L. Boyd, and L. C. Walker, "16. The fraction of cancer attributable to lifestyle and environmental factors in the UK in 2010," *Br J Cancer*, vol. 105 Suppl 2, no. Suppl 2, pp. S77-81, Dec. 2011, doi: 10.1038/bjc.2011.489.
48. C. Castro-Espin and A. Agudo, "The Role of Diet in Prognosis among Cancer Survivors: A Systematic Review and Meta-Analysis of Dietary Patterns and Diet Interventions," *Nutrients*, vol. 14, no. 2, p. 348, Jan. 2022, doi: 10.3390/nu14020348.
49. T. Domazet-Lošo *et al.*, "Naturally occurring tumours in the basal metazoan Hydra," *Nat Commun*, vol. 5, no. 1, Art. no. 1, Jun. 2014, doi: 10.1038/ncomms5222.
50. J. J. Otto and R. D. Campbell, "Tissue economics of hydra: regulation of cell cycle, animal size and development by controlled feeding rates," *Journal of Cell Science*, vol. 28, no. 1, pp. 117–132, Dec. 1977, doi: 10.1242/jcs.28.1.117.
51. S. Gurkewitz, M. Chow, and R. D. Campbell, "Hydra size and budding rate: influence of feeding," *International Journal of Invertebrate Reproduction*, vol. 2, no. 3, pp. 199–201, Jan. 1980, doi: 10.1080/01651269.1980.10553355.
52. B. Hecker and L. B. Slobodkin, "Responses of Hydra Oligactis to Temperature and Feeding Rate," in *Coelenterate Ecology and Behavior*, G. O. Mackie, Ed., Boston, MA: Springer US, 1976, pp. 175–183. doi: 10.1007/978-1-4757-9724-4_19.
53. R. M. Anderson and R. Weindruch, "Metabolic reprogramming, caloric restriction and aging," *Trends Endocrinol Metab*, vol. 21, no. 3, pp. 134–141, Mar. 2010, doi: 10.1016/j.tem.2009.11.005.
54. C. Vidoni *et al.*, "Calorie Restriction for Cancer Prevention and Therapy: Mechanisms, Expectations, and Efficacy," *J Cancer Prev*, vol. 26, no. 4, pp. 224–236, Dec. 2021, doi: 10.15430/JCP.2021.26.4.224.
55. M. I. Deserti, K. S. Esquiús, A. H. Escalante, and F. H. Acuña, "Trophic ecology and diet of Hydra vulgaris (Cnidaria; Hydrozoa)," *Animal Biology*, vol. 67, no. 3–4, pp. 287–300, Jan. 2017, doi:

10.1163/15707563-00002537.

56. C. A. Aktipis, A. M. Boddy, R. A. Gatenby, J. S. Brown, and C. C. Maley, "Life history trade-offs in cancer evolution," *Nat Rev Cancer*, vol. 13, no. 12, Art. no. 12, Dec. 2013, doi: 10.1038/nrc3606.
57. L. K. Borouhgs and R. J. DeBerardinis, "Metabolic pathways promoting cancer cell survival and growth," *Nat Cell Biol*, vol. 17, no. 4, Art. no. 4, Apr. 2015, doi: 10.1038/ncb3124.
58. T. Norin and T. D. Clark, "Fish face a trade-off between 'eating big' for growth efficiency and 'eating small' to retain aerobic capacity," *Biology Letters*, vol. 13, no. 9, p. 20170298, Sep. 2017, doi: 10.1098/rsbl.2017.0298.
59. P. A. Biro, F. Thomas, B. Ujvari, and C. Beckmann, "Can Energetic Capacity Help Explain Why Physical Activity Reduces Cancer Risk?," *Trends in Cancer*, vol. 6, no. 10, pp. 829–837, Oct. 2020, doi: 10.1016/j.trecan.2020.06.001.
60. P. A. Biro, F. Thomas, B. Ujvari, and C. Beckmann, "A novel perspective suggesting high sustained energy expenditure may be net protective against cancer," *Evol Med Public Health*, vol. 10, no. 1, pp. 170–176, 2022, doi: 10.1093/emph/eoac012.
61. B. M. Mortzfeld, J. Taubenheim, A. V. Klimovich, S. Fraune, P. Rosenstiel, and T. C. G. Bosch, "Temperature and insulin signaling regulate body size in Hydra by the Wnt and TGF-beta pathways," *Nat Commun*, vol. 10, no. 1, Art. no. 1, Jul. 2019, doi: 10.1038/s41467-019-11136-6.
62. H. Chen, F. Lin, K. Xing, and X. He, "The reverse evolution from multicellularity to unicellularity during carcinogenesis," *Nat Commun*, vol. 6, no. 1, Art. no. 1, Mar. 2015, doi: 10.1038/ncomms7367.
63. S. A. West, S. P. Diggle, A. Buckling, A. Gardner, and A. S. Griffin, "The Social Lives of Microbes," *Annual Review of Ecology, Evolution, and Systematics*, vol. 38, no. 1, pp. 53–77, 2007, doi: 10.1146/annurev.ecolsys.38.091206.095740.
64. S. E. Finkel, "Long-term survival during stationary phase: evolution and the GASP phenotype," *Nat Rev Microbiol*, vol. 4, no. 2, Art. no. 2, Feb. 2006, doi: 10.1038/nrmicro1340.
65. P. Eraso and J. M. Gancedo, "Use of glucose analogues to study the mechanism of glucose-mediated cAMP increase in yeast," *FEBS Letters*, vol. 191, no. 1, pp. 51–54, Oct. 1985, doi: 10.1016/0014-5793(85)80991-1.
66. R. B. Wilson and K. Tatchell, "SRA5 encodes the low-Km cyclic AMP phosphodiesterase of *Saccharomyces cerevisiae*," *Mol Cell Biol*, vol. 8, no. 1, pp. 505–510, Jan. 1988, doi: 10.1128/mcb.8.1.505-510.1988.
67. C.-W. Chang, J. L. Moseley, D. Wykoff, and A. R. Grossman, "The LPB1 Gene Is Important for Acclimation of *Chlamydomonas reinhardtii* to Phosphorus and Sulfur Deprivation," *Plant Physiol*, vol. 138, no. 1, pp. 319–329, May 2005, doi: 10.1104/pp.105.059550.
68. J. L. Moseley, C.-W. Chang, and A. R. Grossman, "Genome-based approaches to understanding phosphorus deprivation responses and PSR1 control in *Chlamydomonas reinhardtii*," *Eukaryot Cell*, vol. 5, no. 1, pp. 26–44, Jan. 2006, doi: 10.1128/EC.5.1.26-44.2006.
69. J. P. Davies, F. H. Yildiz, and A. Grossman, "Sac1, a putative regulator that is critical for survival of *Chlamydomonas reinhardtii* during sulfur deprivation.," *EMBO J*, vol. 15, no. 9, pp. 2150–2159, May

1996.

70. L. Váchová and Z. Palková, "Physiological regulation of yeast cell death in multicellular colonies is triggered by ammonia," *J Cell Biol*, vol. 169, no. 5, pp. 711–717, Jun. 2005, doi: 10.1083/jcb.200410064.
71. J. E. Strassmann and D. C. Queller, "Evolution of cooperation and control of cheating in a social microbe," *Proceedings of the National Academy of Sciences*, vol. 108, no. supplement_2, pp. 10855–10862, Jun. 2011, doi: 10.1073/pnas.1102451108.
72. A. M. Nedelcu and R. E. Michod, "The evolutionary origin of an altruistic gene," *Mol Biol Evol*, vol. 23, no. 8, pp. 1460–1464, Aug. 2006, doi: 10.1093/molbev/msl016.
73. R. M. S. Saggere, C. W. J. Lee, I. C. W. Chan, D. G. Durnford, and A. M. Nedelcu, "A life-history trade-off gene with antagonistic pleiotropic effects on reproduction and survival in limiting environments," *Proceedings of the Royal Society B: Biological Sciences*, vol. 289, no. 1967, p. 20212669, Jan. 2022, doi: 10.1098/rspb.2021.2669.
74. K. R. Foster, G. Shaulsky, J. E. Strassmann, D. C. Queller, and C. R. L. Thompson, "Pleiotropy as a mechanism to stabilize cooperation," *Nature*, vol. 431, no. 7009, Art. no. 7009, Oct. 2004, doi: 10.1038/nature02894.
75. M. E. Cameron-Pack, S. G. König, A. Reyes-Guevara, A. Reyes-Prieto, and A. M. Nedelcu, "A personal cost of cheating can stabilize reproductive altruism during the early evolution of clonal multicellularity," *Biology Letters*, vol. 18, no. 6, p. 20220059, Jun. 2022, doi: 10.1098/rsbl.2022.0059.
76. R. H. B. Christensen, "ordinal: Regression Models for Ordinal Data." Nov. 16, 2022. Accessed: Feb. 09, 2023. [Online]. Available: <https://CRAN.R-project.org/package=ordinal>
77. K. Bolar, "STAT: Interactive Document for Working with Basic Statistical Analysis." Apr. 01, 2019. Accessed: Oct. 29, 2022. [Online]. Available: <https://CRAN.R-project.org/package=STAT>

Figures




Figure 1

Phenotype of healthy and tumoral hydras from the clonal strains of St. Petersburg. (A) Healthy hydra from the tumor-free strain: the body is long and thin, and the number of tentacles does not exceed 7. (B) Tumoral hydra from the tumoral strain: many tumefactions thickening the body, and the tentacles are supernumerary (i.e. equal or superior to 8). The pictures are taken with a trinocular magnifier, scale bar: 1 mm.




Figure 2

Phenotype of healthy and tumoral zebrafish. (A) Healthy zebrafish from the AB strain. (B) Zebrafish with nevi (i.e. premalignant lesions) from the kita-GFP-RAS transgenic strain: the pigmentation pattern is altered. (C) Zebrafish with nevi and tumor from the kita-GFP-RAS transgenic strain: in addition to the modified pattern, a melanoma has developed on the fish's back.




Figure 3

Graphical summary of the general experimental design.




Figure 4

The effects of four diets on tumorigenesis in hydras from tumoral and tumor-free St. Petersburg strains.

(A) The proportion of juvenile hydras from the tumor-free strain developing spontaneous tumors across time (days) according to the diet followed.

(B) The proportion of juvenile hydras from the tumoral strain developing tumors across time (days) according to the diet followed.

(C) The frequency of the intensity of tumor phenotype expression in juvenile hydras from the tumoral strain according to the diet followed.

(D) The proportion of five-week-old hydras from the tumor-free strain developing spontaneous tumors across time (days) according to the diet followed since their fifth week.

(E) Changes in the frequency of the intensity of tumor phenotype expression in hydras from the tumoral strain (five-week-old) according to the diet followed over one month (i.e. after tumor development).

Wild hydras




Figure 5

The proportion of wild hydras developing tumors across time (days) according to diet.




Figure 6

The effects of four diets on tumorigenesis in zebrafish from the tumoral strain

(A) The proportion of juvenile zebrafish from the tumoral strain developing nevi across time (days) according to the diet followed. Given the high natural mortality (see Material and Methods), the analysis was performed only on individuals still alive at the end of the study or having developed nevi before dying.

(B) The proportion of zebrafish with nevi from the tumoral strain developing tumors across time (days) according to the diet followed since their third month (i.e. after nevi development).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Sldataandscripts.zip](#)
- [Supplementaryinformation.docx](#)