



HAL
open science

Antimicrobial resistance and geographical distribution of *Staphylococcus* sp. isolated from whiting (*Merlangius merlangus*) and seawater in the English Channel and the North sea

Thomas Brauge, Erwan Bourdonnais, Sylvain Trigueros, Pierre Cresson, Sabine Debuiche, Sophie Granier, Graziella Midelet

► To cite this version:

Thomas Brauge, Erwan Bourdonnais, Sylvain Trigueros, Pierre Cresson, Sabine Debuiche, et al.. Antimicrobial resistance and geographical distribution of *Staphylococcus* sp. isolated from whiting (*Merlangius merlangus*) and seawater in the English Channel and the North sea. *Environmental Pollution*, 2024, 345, pp.123434. 10.1016/j.envpol.2024.123434 . hal-04423681

HAL Id: hal-04423681

<https://hal.science/hal-04423681v1>

Submitted on 29 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 **Antimicrobial resistance and geographical distribution of**
2 ***Staphylococcus* sp. isolated from whiting (*Merlangius***
3 ***merlangus*) and seawater in the English Channel and the**
4 **North Sea**

5

6 **Thomas Brauge^{1*}, Erwan Bourdonnais¹, Sylvain Trigueros¹, Pierre Cresson², Sabine**
7 **Debuiche¹, Sophie A. Granier³, Graziella Midelet¹**

8

9 ¹ French Agency for Food, Environmental and Occupational Health & Safety, Laboratory for Food
10 Safety, Bacteriology and Parasitology of Fishery and Aquaculture Products Unit, Boulogne-sur-Mer,
11 France.

12 ² Ifremer, Channel/North Sea Fisheries Laboratory, Boulogne-sur-Mer, France

13 ³ French Agency for Food, Environmental and Occupational Health & Safety, Fougères Laboratory,
14 Fougères, France

15

16 *** Correspondence:**

17 Thomas Brauge

18 thomas.brauge@anses.fr

19

20 **Keywords: Antimicrobial resistance, marine environment, *Staphylococcus*,**
21 **environmental parameters.**

22

23 **Abstract**

24

25 *Staphylococcus* is a significant food safety hazard. The marine environment serves as a source
26 of food for humans and is subject to various human-induced discharges, which may contain
27 *Staphylococcus* strains associated with antimicrobial resistance (AMR). The aim of this study
28 was to assess the occurrence and geographical distribution of AMR *Staphylococcus* isolates in
29 seawater and whiting (*Merlangius merlangus*) samples collected from the English Channel
30 and the North Sea. We isolated and identified 238 *Staphylococcus* strains, including 12
31 coagulase-positive (CoPs) and 226 coagulase-negative (CoNs) strains. All CoPs isolates

32 exhibited resistance to at least one of the 16 antibiotics tested. Among the CoNs strains, 52%
33 demonstrated resistance to at least one antibiotic, and 7 isolates were classified as multi-drug
34 resistant (MDR). In these MDR strains, we identified AMR genes that confirmed the
35 resistance phenotype, as well as other AMR genes, such as quaternary ammonium resistance.
36 One CoNS strain carried 9 AMR genes, including both antibiotic and biocide resistance
37 genes. By mapping the AMR phenotypes, we demonstrated that rivers had a local influence,
38 particularly near the English coast, on the occurrence of AMR *Staphylococcus*. The analysis
39 of marine environmental parameters revealed that turbidity and phosphate concentration were
40 implicated in the occurrence of AMR *Staphylococcus*. Our findings underscore the crucial
41 role of wild whiting and seawater in the dissemination of AMR *Staphylococcus* within the
42 marine environment, thereby posing a risk to human health.

43

44 **1 Introduction**

45

46 Antimicrobial resistance (AMR) is a significant public health issue that can increase the
47 lethality of pathogenic bacteria and lead to economic losses. In 2019, it was estimated that
48 antibiotic-resistant bacteria were responsible for 1.27 million deaths in Europe alone (Norrby
49 et al., 2009; Mestrovic et al., 2022). While the accuracy of this estimate is questioned by some
50 authors, the clinical, economic, and public health burden associated with antimicrobial
51 resistance is undeniable (O'Neill, 2016). Humans can be exposed to antibiotic-resistant
52 bacteria through their diet or environment, which aligns with the One Health concept.
53 *Staphylococcus* sp. is a bacterial agent that is particularly problematic in terms of AMR and is
54 found in humans, animals, and the environment. Among the *Staphylococcus* genus,
55 coagulase-positive (CoPS) *S. aureus* is the main pathogen responsible for a variety of clinical
56 infections in humans, including bacteremia, endocarditis, and various invasive medical device
57 infections (Tong et al., 2015). Coagulase-negative *Staphylococcus* (CoNS), especially
58 *Staphylococcus epidermidis* and *Staphylococcus haemolyticus*, have also emerged as recurrent
59 causative agents of nosocomial infections, particularly those associated with indwelling
60 devices (Becker et al., 2014). The study of CoNS in the context of antimicrobial resistance
61 and the safety of seafood products is relevant. Indeed, CoNS can act as reservoirs of
62 antimicrobial resistance genes. Even though they are not the primary pathogens, they can
63 carry resistance genes that may be transferred to other bacteria, including more virulent
64 pathogens. This contributes to the dissemination of antimicrobial resistance (Fišarová et al.,

65 2019). Additionally, CoNS can share ecological niches with other bacteria, including
66 foodborne pathogens. Investigating their presence and antimicrobial resistance can help
67 understand the complex dynamics of microbial communities in aquatic environments, which
68 is crucial for assessing potential risks to human health associated with the consumption of
69 seafood products (Regecova et al., 2014; Chajęcka-Wierzchowska et al., 2015). Seafood can
70 be contaminated by *Staphylococcus* in two ways: post-capture contamination through human
71 contact or contaminated surfaces, and pre-capture contamination due to the presence of
72 *Staphylococcus* in the marine environment. Several studies have highlighted the
73 contamination of the marine environment with *Staphylococcus*, which is not indigenous to
74 this environment but rather a consequence of anthropogenic discharges such as wastewater
75 treatment plants, run-off water, and other discharges from terrestrial environments (Gabutti et
76 al., 2000; Soge et al., 2009; Goodwin et al., 2012; Akanbi et al., 2017). These anthropogenic
77 discharges also transport residues of drugs used in human or veterinary medicine, as well as
78 antimicrobial resistance genes (ARGs), into the marine environment (Marti et al., 2014;
79 Bourdonnais et al., 2022; Bourdonnais et al., 2023). Additionally, the genome of aquatic
80 bacteria contains a high abundance of mobile genetic elements that frequently contribute to
81 the selection and dissemination of ARGs (Stalder et al., 2012). Aquatic bacteria serve as
82 potential intermediate vectors for the transfer of naturally occurring ARGs from
83 environmental bacteria to human pathogenic or commensal bacteria, such as *Staphylococcus*.
84 The presence of numerous AMR genes and the omnipresence of antibiotic residues in the
85 marine environment, due to anthropization, could significantly contribute to the emergence of
86 AMR *Staphylococcus*, which poses a threat to human health. Only a few studies have reported
87 the presence of methicillin-resistant *Staphylococcus aureus* (MRSA) in fish and fishery
88 products in various regions of the world (Vaiyapuri et al., 2019).

89 Therefore, questions remain regarding the presence of AMR *Staphylococcus* in the marine
90 environment, the influence of the environment on its occurrence, and the associated risk
91 posed by AMR *Staphylococcus* in wild fish intended for human consumption, such as
92 whiting. Whiting is a keystone fish species in the area, known for its high trophic position and
93 predominant biomass in the fish assemblage (Cresson et al., 2020; Timmerman et al., 2020;
94 Timmerman et al., 2021). To our knowledge, no data on this subject is available in the
95 literature. To address this information gap, we investigated the occurrence and geographical
96 distribution of AMR *Staphylococcus* in areas subject to heterogeneous human impacts and
97 various environmental conditions. We collected seawater and whiting (*Merlangius*
98 *merlangus*) samples from the English Channel and the North Sea, which are ideal locations

99 due to their numerous areas with high anthropogenic influences, particularly near the coasts
100 and in regions with intense fishing activities and maritime traffic.

101

102 **2 Material and methods**

103

104 *2.1 Marine samples and preparation*

105

106 Forty-six sampling stations were defined in the English Channel and the North Sea area
107 (Figure 1). At each sampling station, 10 whiting (*Merlangius merlangus*) and 300 mL of
108 seawater were collected. These samples were collected during the International Bottom Trawl
109 Survey (IBTS) in February 2017 on the R/V Thalassa (Verin, 2017). The IBTS is an
110 ecosystemic survey, initially designed to estimate fish abundance to inform fisheries
111 management indicators but which includes sampling of all ecosystem components. Protocols
112 of the survey are currently being evaluated by Ifremer and are validated by the ICES IBTS
113 Working Group (<https://www.ices.dk/community/groups/pages/ibtswg.aspx>). In addition,
114 survey's PIs received training about animal well-being and ethics. The physico-chemical
115 parameters at the water surface, including water temperature, pH, salinity, turbidity, oxygen
116 saturation, as well as the concentrations of phosphate, nitrite, nitrate, silicate, ammonium,
117 chlorophyll a, total phaeopigments, total suspended matter, organic, and inorganic suspended
118 matter, were measured at each sampling station during sample collection.

119 The forty-six seawater samples were collected at a depth of about 30 meters using Niskin
120 bottles and transferred in sterile containers. All seawater samples were filtered on 0.45 µm
121 pore-size mixed cellulose ester membranes (Millipore, Burlington, USA) with a Nalgene
122 individual vacuum filtration system (Thermo Fisher Scientific, Waltham, USA). The 0.45 µm
123 filter membranes were placed in a tube containing 30 mL of alkaline peptone saline water
124 (APSW, Oxoid, Basingstoke Hampshire, UK) with 20% (v/v) of glycerol and stored at -20°C
125 on the ship. Back to the laboratory, the tubes containing the filters were thawed at room
126 temperature and vortexed for 1 minute. The filter was removed using sterilized forceps. The
127 tubes were centrifuged for 10 minutes at 5000 ×g and the pellet was remained in 4 mL of the
128 supernatant. Two milliliters were analyzed to isolate the *Staphylococcus* strains.

129 A total of 460 fish were analyzed. After thawing the fish at room temperature, we swabbed
130 both surfaces of each fish using a sterile sponge soaked with buffered peptone water (BPW)
131 (3M, Cergy-Pontoise, France) to study fish skin. The homogenate was recovered and we

132 added 20% (v/v) of glycerol. The fish gills were extracted, diluted by half with BPW (v/m),
133 mixed and supplemented with 20% (v/v) of glycerol. The fish skin and gill samples were
134 pooled for each sampling station (10 individuals per sampling station) except for the V0116,
135 V0117, V0127 and V0133 sample stations (the fish samples were analyzed individually).

136

137 2.2 Isolation and identification of bacterial isolates

138

139

140 Each sample was thawed and diluted at 1/10 in BPW and APSW enrichment broths. The
141 broths were incubated at 30°C overnight and plated on total 3 selective and 2 non-selective
142 media. The ASPW enrichment broths were plated on non-selective saline nutritive agar (SNA,
143 2% NaCl) (Biorad, Marne-la-Coquette, France, Oxoid) then on the following selective agars:
144 ChromID[®] MRSA (Biomérieux, Lyon, France), Chapman (Oxoid) and Thiosulfate Citrate
145 Bile Saccharose (TCBS, Oxoid). Although the SNA is a non-selective medium, its salt
146 content (2% NaCl) still influences the preferential growth of certain bacterial groups, thereby
147 promoting the development of halophilic bacteria. This approach has allowed us to increase
148 our chances of isolating halophilic bacteria such as staphylococci. The use of a non-selective
149 saline nutrient agar has facilitated the growth of a diverse range of bacteria without
150 discrimination, thus optimizing the recovery of microorganisms present in the sample,
151 including those in low abundance. The BPW enrichment broths were plated on non-selective
152 Trypticase Soy Agar with 0.6% Yeast extract (TSAYe, Oxoid) and on selective Polymyxin
153 pyruvate Egg yolk Mannitol Bromothymol blue Agar (PEMBA, Oxoid). PEMBA were
154 incubated at 30°C for 48h and TSAYE, TCBS, Chapman and ChromID[®] MRSA agars at
155 37°C for 24h. After incubation, one colony of different morphologies of each medium was
156 plated on a non-selective agar (TSAYe for the BPW enrichments and SNA for the ASPW
157 enrichments) and incubated for 24 hours at the same temperature of the bacterial isolation
158 agars. The bacterial isolates obtained were stored in brain heart infusion (BHI) (Biomérieux)
159 supplemented with 20% (v/v) of glycerol at -80°C. The strains were then identified by 16
160 rRNA gene sequencing and MALDI TOF spectroscopy according to the protocols described
161 by (Brauge et al., 2021).

162

163 2.3 Antimicrobial susceptibility test of the Staphylococcus isolates

164

165 *Staphylococcus* isolates from BHI supplemented with 20% (v/v) of glycerol at -80°C were
166 grown on SNA or TSA Ye for 24h at 37°C. Antimicrobial susceptibility was tested by disk
167 diffusion following the EUCAST standard (EUCAST, 2012). A 0,5Mc Farland
168 staphylococcal suspension was plated with a cotton swab on a Muller Hinton Agar (MHA,
169 Biorad, Marnes La Coquette, France) and the plates were incubated at 35°C for 16-18h. The
170 antibiotic disks (Biorad) tested were: penicillin G (1 U), cefoxitin (30 µg), ciprofloxacin (5
171 µg), kanamycin (30 µg), tobramycin (10 µg), gentamycin (10 µg), erythromycin (15 µg),
172 clindamycin (2 µg), vancomycin (5 µg), linezolid (10 µg), sulfamide (200 µg), trimethoprim
173 (5 µg), tetracycline (30 µg), chloramphenicol (30 µg), rifampicin (5 µg) and fusidic acid (10
174 µg). Cefoxitin was used to detect methicillin resistance and the clindamycin disk was
175 positioned next to the erythromycin disk to visualize MLS_B mechanism. Zone diameters were
176 measured with a Biomic V3 (Giles Scientific, Santa Barbara, USA). The *Staphylococcus*
177 ATCC 25923 strain was used as a quality control.

178

179 2.4 Whole genome sequencing of multi-drug resistant strains

180

181

182 Among all the strains identified as resistant, we specifically targeted and sequenced in wole
183 genome the strains of *Staphylococcus* classified as multi-drug resistant (MDR). The
184 designation of MDR among the *Staphylococcus* strains was applied when these strains
185 exhibited phenotypic resistance to at least three drugs belonging to three different antibiotic
186 classes (Gandolfi-Decristophoris et al., 2013). This categorization denoted a comprehensive
187 resistance profile, indicating the ability to withstand the effects of multiple antimicrobial
188 agents. Following the classification as MDR, the *Staphylococcus* strains were cultured on
189 Mueller-Hinton Agar (MHA) media. The genomic DNA was then extracted from few
190 colonies using the DNeasy[®] Blood & Tissue Kit (Qiagen, Hilden, Germany) following
191 manufacturer's protocol for Gram positive strains. DNA quality control and sequencing was
192 carried out by Genoscreen society (Lille, France). The ARGs were identified with the
193 Resfinder database from the CGE server (<http://genomicepidemiology.org/>, last access March
194 22nd 2023). The presence of the ARGs was validated when it had a 100% percent identity.

195

196 2.5 Data analysis and statistics

197

198 Antibiotic susceptibility test results were analyzed to define the phenotypic resistances
199 according to the clinical thresholds established in the CLSI (CLSI, 2015, 2016) and EUCAST
200 (EUCAST, 2018) guidelines and according to (Briet, 2018). All data analysis were performed
201 as described in our previous study (Bourdonnais et al., 2022), using RStudio Software version
202 1.4.1717 (RStudio, Inc, Boston, USA). The occurrence of the antimicrobial resistance
203 phenotypes was plotted on a map, using the leaflet and ggplot2 packages of the software.
204 Correlations between the environmental factors and the number of *Staphylococcus* strains
205 with phenotypic resistances were determined with Pearson's rank correlation coefficient and
206 plotted using the corrplot package. Statistical analysis of correlations (*p*-values) was
207 performed with a significance test. Only the Pearson correlation coefficients with a significant
208 *p*-value (*p* < 0.05) were considered.

209

210 **3 Results**

211

212 Among the 1,742 bacteria isolated from seawater and whiting samples, we identified
213 238 isolates as belonging to the *Staphylococcus* genus at the species level. Twelve strains of
214 *Staphylococcus* were coagulase positive (CoPS) and were identified as *S. aureus* (Table 1).
215 Four of these strains were isolated from seawater, four from fish skin and four from fish gills
216 samples. We observed that 100% of the CoPS were resistant to penicillin G. We identified
217 three CoPS strains from fish gill samples that exhibited resistance to both erythromycin and
218 fusidic acid, representing a resistance prevalence of 25%. In comparison, erythromycin
219 resistance has been observed at a prevalence of approximately 30% in *S. aureus* strains
220 isolated from fish imported from Egypt, India, Yemen, and Turkey (Ertas Onmaz et al., 2015;
221 Obaidat et al., 2015). Fusidic acid resistance was also noted in 22% of *S. aureus* strains
222 isolated from raw shrimp in China (Dai et al., 2023). Additionally, the induction of
223 clindamycin resistance by erythromycin was observed in one CoPS strain (B3PA-RCPE17-
224 62à71PEPE1), indicating a MLSB phenotype. This particular strain was isolated from a fish
225 skin sample.

226

227 **Table 1: Antibiotic resistance phenotypes of coagulase-positive *Staphylococcus* (CoPS, n**
228 **= 12).** R: resistant; S: susceptible. CFOX: cefoxitin; CLND: clindamycin; CLOR:
229 chloramphenicol; CPRX: ciprofloxacin; EARTH: erythromycin; FSAD: fusidic acid; GENT:
230 gentamycin; KANA: kanamycin; LNZD: linezolid; PENG: penicillin G; RFPN: rifampicin;

231 SULF: sulfamide; TETR: tetracycline; TOBR: tobramycin; TRIM: trimethoprim; VNCO:
232 vancomycin.

233

234 Two hundred and twenty six coagulase-negative *Staphylococcus* (CoNS) were isolated
235 (Table 2) with 46 strains isolated from seawater, 63 strains from fish skin and 117 strains
236 from fish gills. The predominantly identified CoNS *Staphylococcus* species were *S. pasteurii*
237 (n = 89) and *S. warneri* (n = 85) followed by *S. epidermidis* (n = 19), *S. equorum* (n = 13), *S.*
238 *haemolyticus* (n = 9), *S. saprophyticus* (n = 5), *S. hominis* (n = 4) and *S. cohnii* (n = 2). We
239 noticed that 52.2% of these strains had phenotypic resistance to at least one antibiotic. All
240 samples combined (fish skin, gills and seawater), the highest number of resistance was
241 associated with erythromycin (40.7% of the CoNS), followed by fusidic acid (12.4% of the
242 CoNS) and tetracycline (10.2% of the CoNS). Resistances were detected to tobramycin in
243 3.5%, to cefoxitin in 2.2% and to kanamycin in 0.4% of the CoNS. No phenotypic resistance
244 to ten other antibiotics tested was observed for these strains and there was no specific
245 resistance for the type of samples analyzed. We identified 7 MDR strains among the CoNS.
246 Two strains isolated from seawater had three phenotypic resistances, one (*S. hominis* B3PA-
247 RCPE-6-FE-AL1) to fusidic acid, cefoxitin, tobramycin and the other (*S. pasteurii* B3PA-
248 RCPE-24FEPE1) to erythromycin, fusidic acid, tobramycin. Another CoNS strain (*S. warneri*
249 B3PA-RCPE-17-7280OUE(2)) with all three resistances to erythromycin, fusidic acid and
250 tetracycline was isolated from fish gills. We also observed resistance to 4 antibiotics
251 (erythromycin, fusidic acid, tobramycin and tetracycline) for 2 strains of *S. hominis* (B3PA-
252 RCPE-17-418OUE(3) and B3PA-RCPE-17-424OUE(1)) isolated from fish gills and one
253 strain of *S. warneri* (B3PA-RCPE-14FEPE1) isolated from seawater. Finally, one strain of *S.*
254 *haemolyticus* B3PA-RCPE2FEMRSA1 resistant to 5 antibiotics (erythromycin, fusidic acid,
255 cefoxitin, tobramycin and kanamycin) was also isolated from seawater. In order to assess the
256 genetic basis of these phenotypic resistances, we sequenced the total genome of the seven
257 MDR strains and identified the resistance genes present in their genomes with 100% identity
258 (Table 3). Ten antibiotic resistance genes confirming phenotypic resistance results were
259 identified in the MDR *Staphylococcus* strains. These genes were consistent with the observed
260 phenotypic resistances: the *fusC* (4 strains) and *fusB* (3 strains) genes coding for fusidic acid
261 resistance, *mecA* (2 strains) for cefoxitin resistance, *aadD* (5 strains) for tobramycin
262 resistance, *mphC* (3 strains), *msrA* (3 strains) and *ermC* (1 strain) for erythromycin resistance,
263 *tet(K)* (4 strains) for tetracycline resistance and *aph(3')-III* (1 strain) for kanamycin resistance.
264 Moreover other resistance genes were also identified: *blaZ* gene encoding penicillin resistance

265 through the production of a β -lactamase in all the MDR strains, *bleO* gene (bleomycin
266 resistance) in 6 MDR strains and the *fosD* gene (fosfomycin resistance) in the *S. warneri*
267 B3PA-RCPE-14FEPE1 strain isolated from seawater. Other resistance genes were also
268 identified in the *S. haemolyticus* B3PA-RCPE2FEMRSA1 strain isolated from seawater such
269 as the *vgaA* gene coding for resistance to streptogramin A and relative compounds as well as
270 the *qacA* gene involved in resistance to quaternary ammoniums. This latest strain carried 9
271 antimicrobial genes. And, the induction of resistance to clindamycin by erythromycin was
272 observed for three CoNS strain (*S. haemolyticus* B3PA-RCPE2FEMRSA1 isolated in
273 seawater sample at the V0003 station, *S. epidermidis* B3PA-RCPE17-72à80OUTSA(2)
274 isolated in seawater sample at the V00039 station and *S. warneri* B3PA-RCPE17-
275 134à144PECH(1)) isolated in seawater sample at the V0060 station), which is a marker of a
276 phenotype MLS_B. For the MDR *S. haemolyticus* B3PA-RCPE2FEMRSA1 strain, we
277 identified the *ermC* gene.

278

279 **Table 2: Antibiotic resistance phenotypes of coagulase-negative *Staphylococcus* (CoNS, n**
280 **= 226).** R: resistant; S: susceptible. CFOX: cefoxitin; CLND: clindamycin; CLOR:
281 chloramphenicol; CPRX: ciprofloxacin; EARTH: erythromycin; FSAD: fusidic acid; GENT:
282 gentamycin; KANA: kanamycin; LNZD: linezolid; PENG: penicillin G; RFPN: rifampicin;
283 SULF: sulfamide; TETR: tetracycline; TOBR: tobramycin; TRIM: trimethoprim; VNCO:
284 vancomycin.

285

286 **Table 3: Phenotypic and genotypic resistances of multidrug resistant strains of**
287 ***Staphylococcus*.** FSAD: fusidic acid; CFOX: cefoxitin; TOBR: tobramycin; EARTH:
288 erythromycin; TETR: tetracycline; KANA: kanamycin.

289

290 The geographical distribution of all antimicrobial resistant CoPS and CoNS strains
291 was investigated (Figure 2). Regarding the number of isolated strains, a significantly higher
292 number of AMR *Staphylococcus* strains were located at the sampling stations near the
293 Thames, Humber and Rhine estuaries. Erythromycin-resistant strains were widely distributed
294 in the English Channel and the North Sea extending from coastal to offshore stations without
295 distinction. Similar observations have been made for fusidic acid resistance in these seas.
296 Penicillin G and cefoxitin-resistant strains were isolated from the samples collected near the
297 French and English coasts, especially off the Humber estuary and the mouth of the Thames.
298 The kanamycin-resistant *Staphylococcus* strain was located between the French and English

299 coasts in the English Channel. Tobramycin and tetracycline resistant-strains were mostly
300 isolated from the samples collected along the English coast, near the Thames and Humber
301 estuaries, but also in the open sea in the Northern North Sea. We identified a tetracycline-
302 resistant strain along the Dutch coast close to the port of Rotterdam and the Rhine estuary. It
303 is clear that the diversity of antimicrobial resistance was more important at stations located
304 near the French and English coasts, more precisely in estuaries of the Thames, Humber and
305 Seine. In contrast, few antimicrobial resistance diversity was observed along the German and
306 Dutch coasts. Moreover, the MDR strains of *Staphylococcus* were isolated from the samples
307 collected in the middle of the English Channel (sampling station V0003), at the mouth of the
308 Thames (V0029 and V0133), off the Humbert estuary (V0039 and V0052) but also from the
309 northern coast of the Netherlands (V0082).

310 In order to evaluate the potential relationships between the occurrence of the
311 phenotypic resistance of *Staphylococcus* and the environmental parameters, we performed a
312 correlation analysis (Figure 3). For all the nature of samples (seawater and fish), we observed
313 that the number of tobramycin-resistant strains was moderately and positively correlated with
314 the number of strains with a resistance to erythromycin, fusidic acid and ceftiofur. The
315 number of tetracycline-resistant strains was positively correlated with the number of strains
316 resistant to erythromycin and fusidic acid as well as the turbidity, the phosphate concentration
317 and the inorganic suspended matter (ISMP). Regarding the number of kanamycin-resistant
318 strains, we observed a positive correlation with the number of ceftiofur and gentamicin-
319 resistant strains. Finally, a positive correlation was also observed between the number of
320 strains resistant to penicillin G and fusidic acid.

321

322 **4 Discussion**

323

324 In this study, we investigated the occurrence and geographical distribution of AMR
325 *Staphylococcus* in whiting (*Merlangius merlangus*), a widely harvested and consumed wild
326 fish, as well as in seawater samples from the English Channel and the North Sea. These areas
327 experience significant anthropogenic pressures, both directly and through river discharges
328 (Tappin and Millward, 2015; Schöneich-Argent et al., 2020; Bourdonnais et al., 2022;
329 Bourdonnais et al., 2023; Mauffret et al., 2023). Initially, we isolated and identified 238
330 *Staphylococcus* strains before assessing their resistance to 16 antibiotics crucial for human
331 health, using the disk diffusion method following the EUCAST guidelines. Methicillin-

332 resistant *Staphylococcus* represents a major concern, as these strains are typically resistant not
333 only to all β -lactams but also to a wide range of other antibiotics. The prevalence of
334 methicillin-resistant *Staphylococcus aureus* (MRSA) varies significantly across institutions
335 and geographical regions. In Europe, a North-South gradient is generally observed, with
336 MRSA strains being less common in Scandinavian hospitals (<2%) compared to
337 Mediterranean ones (>40%) (Stefani and Varaldo, 2003). Our study conducted in northern
338 waters seems to support this trend, as no MRSA was detected. Specifically, the cefoxitin
339 phenotypic resistance was not observed in the 12 *S. aureus* (CoPS) isolates in our samples,
340 and all isolates were resistant to penicillin G, which aligns with previous observations in
341 beach sand and bathing seawater samples Akanbi et al. (2017). It has been reported by Daurel
342 and Leclercq (2008) that over 80% of *S. aureus* produce a penicillinase enzyme. We observed
343 the MLS_B phenotype in one CoPS strain and three CoNS strains. In the MDR CoNS strains,
344 we detected the presence of the *ermC* gene, which confirmed the underlying phenotypic
345 mechanism in the *S. haemolyticus* B3PA-RCPE2FEMRSA1 strain. The *ermC* gene that we
346 identified in the MDR *S. haemolyticus* strain has previously been detected in marine
347 sediments collected in the Arctic region, but its presence has been associated with human
348 activities (Tan et al., 2018).

349
350 Furthermore, only 2.2% of the CoNS strains (2 *S. cohnii*, 1 *S. haemolyticus*, 1 *S. hominis*
351 isolated from seawater samples, and 1 *S. saprophyticus* isolated from fish gills) showed
352 resistance to cefoxitin, indicating methicillin resistance. These values are lower than those
353 reported in other studies. The prevalence of MRSA in fish and fishery products varied
354 between 1% and 60%, depending on the sample type and origin (Vaiyapuri et al., 2019) and
355 1.0% of wild freshwater fish samples in Germany were found to be positive for MRSA
356 (EFSA and ECDC, 2023). In the case of the MDR *S. haemolyticus* B3PA-RCPE2FEMRSA1
357 and *S. hominis* B3PA-RCPE-6-FE-AL1 strains isolated from seawater samples, we identified
358 the *mecA* gene, which is responsible for methicillin resistance and can be carried by the
359 mobile chromosomal cassette *SCCmec*, as described by Gomez et al. (2017). *S.*
360 *hamemolyticus* is associated with nosocomial infections such as bacteremia and infective
361 endocarditis. Methicillin-resistant *Staphylococcus* strains have previously been isolated from
362 mackerel, salmon, tuna, flounder, and yellowtail sashimi (Hammad et al., 2012). We also
363 identified one *S. saprophyticus* strain showing phenotypic resistance to cefoxitin. This species
364 is commonly implicated in certain types of CoNS clinical infections, such as skin and urinary
365 tract infections (Silva et al., 2020). Similar to previous studies that identified the *mecA* and

366 *aph(3')-III* genes in marine *S. aureus* strains and river waters (Levin-Edens et al., 2012;
367 Paulus et al., 2020), we observed that the *S. haemolyticus* B3PA-RCPE2FEMRSA1 strain
368 also carried these two antimicrobial resistance genes. However, we did not observe
369 phenotypic penicillin resistance in any of the 7 MDR CoNS strains, but we identified the
370 presence of the *blaZ* gene for further whole-genome sequencing analysis. Similarly, Pimenta
371 et al. (2023) demonstrated that a species of *S. haemolyticus* isolated from a clinical sample
372 harbored the *blaZ* genes and *femA* plasmid but showed a negative result for phenotypic
373 resistance to cefoxitin and penicillin G. Two hypotheses could explain this result: the first is
374 that *blaZ* might not be expressed due to genetic mutations in the regulatory regions of the
375 gene. Indeed, the *blaZ* gene is regulated by two other genes, the *blaR1* antirepressor and the
376 *blal* repressor. Following exposure to beta-lactams, *blaR1*, a transmembrane sensor-
377 transducer, undergoes autocatalytic cleavage, promoting the cleavage of the repressor gene,
378 *blal*, thereby allowing transcription of *blaZ*. A mutation in these regions could affect the
379 expression of *blaZ* and thus the expression of phenotypic resistance (El Feghaly et al., 2012;
380 Pimenta et al., 2023). Another hypothesis could be that penicillin resistance might not be
381 consistently detected using commonly employed susceptibility testing methods. In support of
382 this, El Feghaly et al. (2012) demonstrated that 9.5% of clinically susceptible *S. aureus*
383 isolates to penicillin G carried the *blaZ* gene. They proposed that enhancing interpretation
384 criteria for the disc diffusion zone size from 29 to 35 mm could improve the sensitivity of
385 phenotypic penicillin resistance testing. Furthermore, our investigation revealed the presence
386 of the *bleO* gene in 6 out of the 7 MDR CoNS strains, indicating resistance to bleomycin.
387 Numerous studies have shown that over 80% of CoPS strains resistant to aminoglycosides
388 also exhibit resistance to bleomycin.

389

390 Regarding the CoNS isolates, we identified resistance to erythromycin (40.7%), fusidic
391 acid (12.4%), tetracycline (10.2%), tobramycin (3.5%), cefoxitin (2.2%) and kanamycin
392 (0.4%) which were lower than those reported in clinical settings (Archer and Climo, 1994;
393 Schmitz et al., 1999; Schreckenberger et al., 2004; Koksai et al., 2009). It is widely accepted
394 that higher levels of antibiotic resistance in clinical isolates are due to constant exposure to
395 antibiotics compared to the marine environment. The open marine environment allows for
396 dilution and dispersion of antibiotic residues and antimicrobial resistance (Stefani and
397 Varaldo, 2003). However, the environment can also contribute to the development of
398 antibiotic resistance in microorganisms due to human/animal therapeutics, wastewater,
399 agriculture, and industrial use of antibiotics (Larsson and Flach, 2021). Bourdonnais et al.

400 (2022) highlighted the presence of the *sul1* and *int11* genes in seawater from the English
401 Channel and the North Sea with concentrations of up to 3.5 log gene copies / L of seawater.
402 Marine environment is thus a reservoir of antimicrobial resistance, notably as it is subjected to
403 anthropogenic discharges. Therefore, the wide distribution of AMR *Staphylococcus* in the
404 marine environment can be considered as a source of concern. Two hundred and thirty eight
405 strains of *Staphylococcus* were isolated from the samples collected in the English Channel
406 and the North Sea, and 54.6% showed at least one antibiotic resistance with 3% of multi-
407 resistant strains. We also revealed that the number of resistant *Staphylococcus* and the
408 diversity of resistance associated with them varied between the different sampling stations in
409 the English Channel and the North Sea. The highest prevalence of antibiotic resistance was
410 found along the English coast. Furthermore, the majority of MDR *Staphylococcus* was
411 isolated in these areas (6/7 strains). Several hypotheses can be put forward to explain this
412 occurrence. The first would be the anthropogenic impact associated to river discharges from
413 the Thames and the Humber effluents. These two rivers are the largest sources of freshwater
414 that England discharges into the North Sea. Previous observations have shown the presence of
415 numerous antimicrobial residues (erythromycin, clarithromycin and azithromycin)
416 downstream of the Thames source (White et al., 2019). In addition, Xu et al. (2019) quantified
417 the antimicrobial resistance genes (ARGs) *bla_{TEM}*, *tetA*, *sul1*, *sul2*, *ermB*, *tetQ*, *tet(X)*, *drfA1*,
418 *drfA2* and *int11* in these waters with abundances ranging from 10⁵ to 10⁸ copies / L. We did
419 not identify these genes in the genomic DNA of the MDR *Staphylococcus* strains isolated
420 from the fish and seawater samples collected near these two areas. To our knowledge, this
421 was the first time that the *mphC*, *aadD*, *fusC* and *fusB* genes were identified in the marine
422 environment. We identified the *tetK* gene in MDR CoNS 4 strains isolated in fish gills. In
423 Chile, the *tetK* resistance gene was identified in *Pseudoalteromonas*, *Shewanella*, *Cobetia* and
424 *Vibrio* strains isolated from sediment and water samples from an aquaculture site but also
425 from the marine environment (Tomova et al., 2015). The prevalence of resistance to fusidic
426 acid varies between 4% and 30% depending on the country (Briet, 2018). Resistance to this
427 antibiotic is acquired either by mutation of the *fusA* target or by acquisition of a *fusB* plasmid
428 determinant (Briet, 2018). In our study, the MDR *S. hominis* and *S. haemolyticus* strains
429 carried the *fusC* gene and the MDR *S. warneri* and *S. pasteurii* strains carried the *fusB* gene
430 that conferred the resistance at fusidic acid confirming by phenotypic and genotypic
431 resistance. For one *S. warneri* strain, the *fusD* gene was described in more by genotypic
432 method. In addition, we have identified the *qacA* gene in a *S. haemolyticus* B3PA-
433 RCPE2FEMRSA1 strain isolated from the English Channel seawater. This gene codes for a

434 subunit of the QacA multidrug efflux pump conferring resistance to antimicrobial compounds
435 such as quaternary ammoniums. This gene has been identified in several MDR *S.*
436 *haemolyticus* strains isolated from dairy cows, a human with septicemia and also in a
437 persistent strain from a veterinary clinic floor despite disinfection (Anthonisen et al., 2002).
438 The *qacA* gene was located on a plasmid with the *blaZ* gene, coding for a β -lactamase and
439 identified in the 7 MDR *Staphylococcus* strains in our study. Disinfectant resistant bacteria
440 surviving disinfection thus represent a threat to the food industry and to the public health.
441 This *S. haemolyticus* B3PA-RCPE2FEMRSA1 strain, isolated in seawater sample, carried 9
442 ARGs (*ermC* (ERTH), *fusC* (FSAD), *mecA* (CFOX), *aadD* (TOBR), *aph(3')-III* (KANA),
443 *blaZ* (penicillin), *bleO* (bleomycin), *vga(A)* (streptogramin A), *qacA* (quaternary
444 ammoniums)) with resistance to antibiotics but also to disinfectant in food industry. The
445 transfer of AMR genes between CoNS and *S. aureus* has been reported that CoNS may act as
446 a AMR gene reservoir for CoPS due to their ability to exchange genetic material inter/intra
447 species (Rossi et al., 2020).

448

449 Another study conducted in the estuarine waters of the Humber revealed that *Vibrio*
450 *parahaemolyticus* strains were phenotypically resistant to various antibiotics such as
451 kanamycin, gentamycin, cefazolin and tetracycline (Daramola et al., 2009), indicating a
452 potential source of AMR in the Humber estuary. We can also postulate that this
453 contamination by AMR could come from man-made structures in the English Channel and the
454 North Sea. Indeed, various ports (Port of Hull, Green Port Hull, Immingham) and offshore
455 installations (oil and gas platforms, wind farms) are present in these areas. These different
456 areas can be a source of AMR contamination due to the human activities or to the marine
457 animals concentration such as birds which are recognized as a vector of ARGs in the North
458 sea (Vanermen et al., 2015). These could have an additional impact on the occurrence of
459 resistance in the marine environment, but to our knowledge, no studies have been carried out
460 on the impact of these artificial structures on the occurrence of antimicrobial resistance.

461

462 A high abundance of AMR *Staphylococcus* (especially to erythromycin) was also observed
463 near the mouths of the Rhine and the Meuse, two large rivers flowing into the North Sea. This
464 is in agreement with the observations of Blaak et al. (2012) who found several *Enterococcus*
465 *faecium* strains resistant to erythromycin, tetracycline and ampicillin in the Meuse, Rhine and
466 New Meuse rivers. These *E. faecium* strains may have been responsible for horizontal transfer
467 of ARGs to *Staphylococcus* strains. More recently, Paulus et al. (2020) have demonstrated the

468 presence of resistance genes to erythromycin (*ermB* and *ermF*), sulfonamide (*sulI*),
469 tetracycline (*tetM*), aminoglycosides (*aph(III)*) and carbapenem (*bla_{OXA}*) along the entire
470 length of the Rhine in Switzerland, Germany and the Netherlands due to potential pollution
471 by pharmaceutical industries in the Lower Rhine. The impact of these river effluents on the
472 pollution of the North Sea is cumulative with that of the city and harbor of Rotterdam, the
473 largest harbor in Europe and therefore the source of much anthropogenic pollution. However,
474 there is no information on the contamination of effluents from port activities with ARGs or
475 antibiotic residues.

476

477 Few antibiotic resistances has been observed in the North Sea environment. This area, in
478 contrast to the others, is quite far from the coast and thus from its influence, but has a
479 concentration of offshore platforms for oil and gas exploitation (Fujii, 2015). This
480 dissemination of AMR *Staphylococcus* within the marine environment may raise questions
481 about the involvement of heavy maritime traffic and their various discharges. Indeed,
482 previous studies have shown that ballast water from ships can contain antibiotic resistant
483 bacteria and ARGs (Brinkmeyer, 2016; Lv et al., 2020).

484

485 A further consideration is the impact of environmental factors that may favor the spread
486 and persistence of AMR *Staphylococcus*. Thus, in this study we looked at environmental
487 factors related to phenotypic resistance of *Staphylococcus*. We showed that resistance to
488 tobramycin and tetracycline was strongly correlated with turbidity and suspended matter.
489 Both descriptors are proxies of organic matter and of aggregation of microorganisms.
490 Similarly, Bourdonnais et al. (2022) observed that turbidity values were positively correlated
491 with the abundance of the *sulI* and *intI1* genes, and with the abundance of the bacterial
492 population in the North Sea and the English Channel seawaters. Chen et al. (2020) showed
493 that the occurrence of ARGs in an estuary in China was correlated with turbidity with a higher
494 prevalence of the ARGs in areas with high turbidity. Although not carried by the same strains,
495 co-occurrence of penicillin G and cefoxitin resistance was observed in several strains isolated
496 from samples along the English coast. A plausible explanation lies in the fact that
497 *Staphylococcus* strains are exposed to similar environmental conditions or medical treatments
498 along the English coast, potentially creating a common selective pressure that favors the
499 development of resistance mechanisms to both penicillin G and cefoxitin. This situation may
500 result from the frequent use of these antibiotics in the region. Indeed, the study by Curtis et al.
501 (2019) clearly indicated that antibiotic prescription trends in the community in England from

1998 to 2016 were predominantly represented by penicillins, cephalosporins, and other beta-lactams. The potential presence of antibiotic residues in English effluents could exert selective pressure on *Staphylococcus* strains, prompting them to evolve and adapt to their local environment. If resistance to penicillin G and cefoxitin confers a selective advantage in this particular environment, these resistance characteristics can coexist in different strains. Several studies have identified the *blaZ* gene, encoding β -lactamase, as responsible for resistance to penicillin G and/or cefoxitin in *Staphylococcus* strains (Mok et al., 2023; Pimenta et al., 2023). Despite the absence of phenotypic resistance, genetic analysis of the 7 multidrug-resistant *Staphylococcus* strains in our study confirmed the significant presence of this gene. The lack of phenotypic results for beta-lactams despite the presence of the *blaZ* gene suggests that other, currently unknown, resistance genes may be responsible for resistance to cefoxitin and penicillin G. This aligns with the observations of Lienen et al. (2021), who demonstrated that 14 out of 15 non-golden *Staphylococcus* isolates lacking the *blaZ*, *mecA*, and *mecC* genes exhibited reduced sensitivity to several beta-lactam antibiotics, including cefoxitin (MIC 4 to 8 mg/L) and penicillin (MIC 0.25 to 0.5 mg/L). Two kanamycin-resistant strains were also resistant to gentamycin and tobramycin. Gentamycin is the most commonly active aminoglycoside on *Staphylococcus*. This resistance is mainly due to the acquisition of aminoglycoside inactivating enzymes which can be encoded by the *aadD* or *aph(3')-III* genes, two genetic determinants identified in this study by WGS approach and which also allow resistance to kanamycin and tobramycin (KTG phenotype).

522

523 **5 Conclusion**

524

This study provided, for the first time, information on the presence of AMR *Staphylococcus* in wild fish destined for human consumption and in seawater from the North Sea and the English Channel. Two hundred and thirty eight *Staphylococcus* strains were isolated from seawater and fish samples, of which 52.2% showed at least one antibiotic resistance and 2.9% were multi-resistant. Only 12 *Staphylococcus aureus* strains were identified. All of them were resistant to penicillin G. In contrast, 226 CoNS were isolated and 3% of these strains were resistant to cefoxitin. The next most common resistances found were to erythromycin (40.7%), fusidic acid (12.4%) and tetracycline (10.2%). We had observed the phenotype MLS_B for 1 CoPS and 3 CoNS and the KTG phenotype for two CoNS. One strain, *S. haemolyticus* B3PA-RCPE2FEMRSA1 isolated in seawater sample, carried 9 ARGs (*ermC*

535 (ERTH), *fusC* (FSAD), *mecA* (CFOX), *aadD* (TOBR), *aph(3')-III* (KANA), *blaZ* (penicillin),
536 *bleO* (bleomycin), *vga(A)* (streptogramin A), *qacA* (quaternary ammoniums)) with resistance
537 to antibiotics but also to disinfectant in food industry. This isolate was a potential vector of
538 ARGs in clinic and in food industry. The geographical distribution of AMR *Staphylococcus*
539 clearly showed an influence of river and port discharges on the occurrence of these strains and
540 the diversity of resistances, especially near the English coasts. This study showed for the first
541 time the occurrence of AMR *Staphylococcus* in the North Sea and the English Channel, and
542 the carriage by wild fish thus representing a health risk for consumers.

543

544 **6 Conflict of interests**

545

546 The authors declare that the research was conducted in the absence of any commercial or
547 financial relationships that could be construed as a potential conflict of interest.

548

549 **7 Funding**

550

551 This study was supported by CPER MARCO and ECOANTIBIO

552

553 **8 Acknowledgments**

554

555 We would like to thank IFREMER, and the crew of R/V Thalassa, the PI and scientific staff
556 of IBTS survey for their help for fish and seawater sampling, and for the provision of
557 environmental parameters.

558

559 **9 References**

560

561 Akanbi, O.E., Njom, H.A., Fri, J., Otigbu, A.C., and Clarke, A.M. (2017) Antimicrobial
562 susceptibility of *Staphylococcus aureus* isolated from recreational waters and beach sand in
563 Eastern Cape Province of South Africa. *Int J Environ Res Public Health* **14**: 1001. Doi:
564 <https://doi.org/10.3390/ijerph14091001>.

565 Anthonisen, I.L., Sunde, M., Steinum, T.M., Sidhu, M.S., and Sørum, H. (2002) Organization
566 of the antiseptic resistance gene *qacA* and Tn552-related beta-lactamase genes in multidrug-
567 resistant *Staphylococcus haemolyticus* strains of animal and human origins. *Antimicrob*
568 *Agents Chemother* **46**: 3606-3612. Doi: <https://doi.org/10.1128/aac.46.11.3606-3612.2002>.

569 Archer, G.L., and Climo, M.W. (1994) Antimicrobial susceptibility of coagulase-negative
570 staphylococci. *Antimicrob Agents Chemother* **38**: 2231-2237. Doi:
571 <https://doi.org/10.1128/aac.38.10.2231>.

572 Becker, K., Heilmann, C., and Peters, G. (2014) Coagulase-negative staphylococci. *Clin*
573 *Microbiol Rev* **27**: 870-926. Doi: <https://doi.org/10.1128/cmr.00109-13>.

574 Blaak, H., Van Rooijen, S., Schuijt, M., Docters van Leeuwen, A., Italiaander, R., van den
575 Berg, H. et al. (2012) Prevalence of antibiotic resistant bacteria in the rivers Meuse, Rhine
576 and New Meuse. In *Bilthoven, Netherlands: National Institute for Public Health and the*
577 *Environment*.

578 Bourdonnais, E., Le Bris, C., Brauge, T., and Midelet, G. (2023) Tracking antimicrobial
579 resistance indicator genes in wild flatfish from the English Channel and the North Sea area: A
580 one health concern. *Environ Pollut*: 123274. Doi:
581 <https://doi.org/10.1016/j.envpol.2023.123274>.

582 Bourdonnais, E., Colcanap, D., Le Bris, C., Brauge, T., and Midelet, G. (2022) Occurrence of
583 Indicator Genes of Antimicrobial Resistance Contamination in the English Channel and North
584 Sea Sectors and Interactions With Environmental Variables. *Front Microbiol*: 1750. Doi:
585 <https://doi.org/10.3389/fmicb.2022.883081>.

586 Brauge, T., Trigueros, S., Briet, A., Debuiche, S., Leleu, G., Gassilloud, B. et al. (2021)
587 MALDI-TOF Mass Spectrometry Fingerprinting Performance Versus 16S rDNA Sequencing
588 to Identify Bacterial Microflora From Seafood Products and Sea Water Samples. *Front Mar*
589 *Sci* **8**. Doi: <https://doi.org/10.3389/fmars.2021.650116>.

590 Briet, A. (2018) Étude de la flore bactérienne et de sa résistance aux antibiotiques des
591 produits de la pêche et de l'aquaculture. In: Université du Littoral Côte d'Opale.

592 Brinkmeyer, R. (2016) Diversity of bacteria in ships ballast water as revealed by next
593 generation DNA sequencing. *Mar Pollut* **107**: 277-285. Doi:
594 <https://doi.org/10.1016/j.marpolbul.2016.03.058>.

595 Chajęcka-Wierzchowska, W., Zadernowska, A., Nalepa, B., Sierpińska, M., and Łaniewska-
596 Trokenheim, Ł. (2015) Coagulase-negative staphylococci (CoNS) isolated from ready-to-eat
597 food of animal origin—phenotypic and genotypic antibiotic resistance. *Food Microbiol* **46**:
598 222-226. Doi: <https://doi.org/10.1016/j.fm.2014.08.001>.

599 Chen, Y.-r., Guo, X.-p., Niu, Z.-s., Lu, D.-p., Sun, X.-l., Zhao, S. et al. (2020) Antibiotic
600 resistance genes (ARGs) and their associated environmental factors in the Yangtze Estuary,
601 China: From inlet to outlet. *Mar Pollut* **158**: 111360. Doi:
602 <https://doi.org/10.1016/j.marpolbul.2020.111360>.

603 CLSI (2015) Methods for Antimicrobial Dilution and Disk Susceptibility Testing of
604 Infrequently Isolated or Fastidious Bacteria. In.

605 CLSI (2016) Methods for antimicrobial disk susceptibility testing of bacteria isolated from
606 aquatic animals; approved guideline. In.

607 Cresson, P., Chouvelon, T., Bustamante, P., Bănar, D., Baudrier, J., Le Loc'h, F. et al. (2020)
608 Primary production and depth drive different trophic structure and functioning of fish
609 assemblages in French marine ecosystems. *Prog Oceanogr* **186**: 102343. Doi:
610 <https://doi.org/10.1016/j.pocean.2020.102343>.

611 Curtis, H.J., Walker, A.J., Mahtani, K.R., and Goldacre, B. (2019) Time trends and
612 geographical variation in prescribing of antibiotics in England 1998-2017. *J Antimicrob*
613 *Chemother* **74**: 242-250. Doi: <https://doi.org/10.1093/jac/dky377>.

614 Dai, J., Huang, J., Wu, S., Zhang, F., Li, Y., Rong, D. et al. (2023) Occurrence, Antibiotic
615 Susceptibility, Biofilm Formation and Molecular Characterization of *Staphylococcus aureus*
616 Isolated from Raw Shrimp in China. *Foods* **12**. Doi: <https://doi.org/10.3390/foods12142651>.

617 Daramola, B.A., Williams, R., and Dixon, R.A. (2009) In vitro antibiotic susceptibility of
618 *Vibrio parahaemolyticus* from environmental sources in northern England. *Int J Antimicrob*
619 *Agents* **34**: 499-500. Doi: <https://doi.org/10.1016/j.ijantimicag.2009.06.015>.

620 Daurel, C., and Leclercq, R. (2008) L'antibiogramme de *Staphylococcus aureus*. *Revue*
621 *Francophone des laboratoires* **2008**: 81-90.

622 EFSA, and ECDC (2023) The European Union Summary Report on Antimicrobial Resistance
623 in zoonotic and indicator bacteria from humans, animals and food in 2020/2021. *EFSA*
624 *Journal* **21**: e07867. Doi: <https://doi.org/10.2903/j.efsa.2022.7209>.

625 El Feghaly, R.E., Stamm, J.E., Fritz, S.A., and Burnham, C.A. (2012) Presence of the bla(Z)
626 beta-lactamase gene in isolates of *Staphylococcus aureus* that appear penicillin susceptible by
627 conventional phenotypic methods. *Diagn Microbiol Infect Dis* **74**: 388-393. Doi:
628 <https://doi.org/10.1016/j.diagmicrobio.2012.07.013>.

629 Ertas Onmaz, N., Abay, S., Karadal, F., Hizlisoy, H., Telli, N., and Al, S. (2015) Occurrence
630 and antimicrobial resistance of *Staphylococcus aureus* and *Salmonella* spp. in retail fish
631 samples in Turkey. *Mar Pollut Bull* **90**: 242-246. Doi:
632 <https://doi.org/10.1016/j.marpolbul.2014.10.046>.

633 EUCAST (2012) Antimicrobial susceptibility testing EUCAST disk diffusion method. In.
634 EUCAST (2018) Breakpoint tables for interpretation of MICs and zone diameters Version
635 8.0. In.

636 Fišarová, L., Pantůček, R., Botka, T., and Doškař, J. (2019) Variability of resistance plasmids
637 in coagulase-negative staphylococci and their importance as a reservoir of antimicrobial
638 resistance. *Res Microbiol* **170**: 105-111. Doi: <https://doi.org/10.1016/j.resmic.2018.11.004>.

639 Fujii, T. (2015) Temporal variation in environmental conditions and the structure of fish
640 assemblages around an offshore oil platform in the North Sea. *Mar Environ Res* **108**: 69-
641 82. Doi: <https://doi.org/10.1016/j.marenvres.2015.03.013>.

642 Gabutti, G., De Donno, A., Bagordo, F., and Montagna, M. (2000) Comparative survival of
643 faecal and human contaminants and use of *Staphylococcus aureus* as an effective indicator of
644 human pollution. *Mar Pollut* **40**: 697-700. Doi: [https://doi.org/10.1016/S0025-326X\(00\)00007-2](https://doi.org/10.1016/S0025-326X(00)00007-2).

645

646 Gandolfi-Decristophoris, P., Regula, G., Petrini, O., Zinsstag, J., and Schelling, E. (2013)
647 Prevalence and risk factors for carriage of multi-drug resistant Staphylococci in healthy cats
648 and dogs. *J Vet Sci* **14**: 449-456. Doi: <https://doi.org/10.4142/jvs.2013.14.4.449>.

649 Gomez, P., Casado, C., Saenz, Y., Ruiz-Ripa, L., Estepa, V., Zarazaga, M., and Torres, C.
650 (2017) Diversity of species and antimicrobial resistance determinants of staphylococci in
651 superficial waters in Spain. *FEMS Microbiol Ecol* **93**. Doi:
652 <https://doi.org/10.1093/femsec/fiw208>.

653 Goodwin, K.D., McNay, M., Cao, Y., Ebentier, D., Madison, M., and Griffith, J.F. (2012) A
654 multi-beach study of *Staphylococcus aureus*, MRSA, and enterococci in seawater and beach
655 sand. *Water Res* **46**: 4195-4207. Doi: <https://doi.org/10.1016/j.watres.2012.04.001>.

656 Hammad, A.M., Watanabe, W., Fujii, T., and Shimamoto, T. (2012) Occurrence and
657 characteristics of methicillin-resistant and-susceptible *Staphylococcus aureus* and methicillin-
658 resistant coagulase-negative staphylococci from Japanese retail ready-to-eat raw fish. *Int J*
659 *Food Microbiol* **156**: 286-289. Doi: <https://doi.org/10.1016/j.ijfoodmicro.2012.03.022>.

660 Koksall, F., Yasar, H., and Samasti, M. (2009) Antibiotic resistance patterns of coagulase-
661 negative staphylococcus strains isolated from blood cultures of septicemic patients in Turkey.
662 *Microbiol Res* **164**: 404-410. Doi: <https://doi.org/10.1016/j.micres.2007.03.004>.

663 Larsson, D., and Flach, C.-F. (2021) Antibiotic resistance in the environment. *Nat Rev*
664 *Microbiol*: 1-13. Doi: <https://doi.org/10.1038/s41579-021-00649-x>.

665 Levin-Edens, E., Soge, O.O., No, D., Stiffarm, A., Meschke, J.S., and Roberts, M.C. (2012)
666 Methicillin-resistant *Staphylococcus aureus* from Northwest marine and freshwater

667 recreational beaches. *FEMS Microbiol Ecol* **79**: 412-420. Doi: [https://doi.org/10.1111/j.1574-](https://doi.org/10.1111/j.1574-6941.2011.01229.x)

668 [6941.2011.01229.x](https://doi.org/10.1111/j.1574-6941.2011.01229.x).

669 Lienen, T., Schnitt, A., Hammerl, J.A., Marino, S.F., Maurischat, S., and Tenhagen, B.A.

670 (2021) Multidrug-resistant *Staphylococcus cohnii* and *Staphylococcus urealyticus* isolates

671 from German dairy farms exhibit resistance to beta-lactam antibiotics and divergent

672 penicillin-binding proteins. *Sci Rep* **11**: 6075. Doi: [https://doi.org/10.1038/s41598-021-85461-](https://doi.org/10.1038/s41598-021-85461-6)

673 [6](https://doi.org/10.1038/s41598-021-85461-6).

674 Lv, B., Cui, Y., Tian, W., Wei, H., Chen, Q., Liu, B. et al. (2020) Vessel transport of

675 antibiotic resistance genes across oceans and its implications for ballast water management.

676 *Chemosphere* **253**: 126697. Doi: <https://doi.org/10.1016/j.chemosphere.2020.126697>.

677 Marti, E., Variatza, E., and Balcazar, J.L. (2014) The role of aquatic ecosystems as reservoirs

678 of antibiotic resistance. *Trends Microbiol* **22**: 36-41. Doi:

679 <https://doi.org/10.1016/j.tim.2013.11.001>.

680 Mauffret, A., Chouvelon, T., Wessel, N., Cresson, P., Bănanaru, D., Baudrier, J. et al. (2023)

681 Trace elements, dioxins and PCBs in different fish species and marine regions: Importance of

682 the taxon and regional features. *Environ Res* **216**: 114624. Doi:

683 <https://doi.org/10.1016/j.envres.2022.114624>.

684 Mestrovic, T., Aguilar, G.R., Swetschinski, L.R., Ikuta, K.S., Gray, A.P., Weaver, N.D. et al.

685 (2022) The burden of bacterial antimicrobial resistance in the WHO European region in 2019:

686 A cross-country systematic analysis. *Lancet Public Health* **7**: e897-e913. Doi:

687 [https://doi.org/10.1016/S2468-2667\(22\)00225-0](https://doi.org/10.1016/S2468-2667(22)00225-0).

688 Mok, H.T., Teng, C.B., Bergin, S., Hon, P.Y., Lye, D.C., De, P.P., and Vasoo, S. (2023)

689 Treatment outcomes with benzylpenicillin and non-benzylpenicillin antibiotics, and the

690 performance of the penicillin zone-edge test versus molecular detection of *blaZ* in penicillin-

691 susceptible *Staphylococcus aureus* (PSSA) bacteraemia. *J Antimicrob Chemother* **78**: 2515-

692 2523. Doi: <https://doi.org/10.1093/jac/dkad263>.

693 Norrby, R., M. Powell, B., Aronsson, D., Monnet, I., Lutsar, I., Bocsan, O. et al. (2009) The

694 bacterial challenge: time to react. *Stockholm: European Centre for Disease Prevention and*

695 *Control*. Doi: doi 10.2900/2518.

696 O'Neill, J. (2016) Tackling drug-resistant infections globally: final report and

697 recommendations. In.

698 Obaidat, M.M., Salman, A.E., and Lafi, S.Q. (2015) Prevalence of *Staphylococcus aureus* in

699 Imported Fish and Correlations between Antibiotic Resistance and Enterotoxigenicity. *J Food*

700 *Prot* **78**: 1999-2005. Doi: <https://doi.org/10.4315/0362-028x.jfp-15-104>.

701 Paulus, G.K., Hornstra, L.M., and Medema, G. (2020) International tempo-spatial study of

702 antibiotic resistance genes across the Rhine river using newly developed multiplex qPCR

703 assays. *Sci Total Environ* **706**: 135733. Doi: <https://doi.org/10.1016/j.scitotenv.2019.135733>.

704 Pimenta, L.K.L., Rodrigues, C.A., Filho, A.R.G., Coelho, C.J., Goes, V., Estrela, M. et al.

705 (2023) *Staphylococcus* spp. Causatives of Infections and Carrier of *blaZ*, *femA*, and *mecA*

706 Genes Associated with Resistance. *Antibiotics* **12**: 671. Doi:

707 <https://doi.org/10.3390/antibiotics12040671>.

708 Regecova, I., Pipova, M., Jevinova, P., Maruskova, K., Kmet, V., and Popelka, P. (2014)

709 Species identification and antimicrobial resistance of coagulase-negative staphylococci

710 isolated from the meat of sea fish. *J Food Sci* **79**: M898-902. Doi:

711 <https://doi.org/10.1111/1750-3841.12429>.

712 Rossi, C.C., Pereira, M.F., and Giambiagi-deMarval, M. (2020) Underrated *Staphylococcus*

713 species and their role in antimicrobial resistance spreading. *Genet Mol Biol* **43**. Doi:

714 <https://doi.org/10.1590/1678-4685-GMB-2019-0065>.

715 Schmitz, F.-J., Fluit, A.C., Gondolf, M., Beyrau, R., Lindenlauf, E., Verhoef, J. et al. (1999)

716 The prevalence of aminoglycoside resistance and corresponding resistance genes in clinical

717 isolates of staphylococci from 19 European hospitals. *J Antimicrob Chemother* **43**: 253-
718 259. Doi: <https://doi.org/10.1093/jac/43.2.253>.

719 Schöneich-Argent, R.I., Dau, K., and Freund, H. (2020) Wasting the North Sea?—A field-
720 based assessment of anthropogenic macrolitter loads and emission rates of three German
721 tributaries. *Environ Pollut* **263**: 114367. Doi: <https://doi.org/10.1016/j.envpol.2020.114367>.

722 Schreckenberger, P.C., Ilendo, E., and Ristow, K.L. (2004) Incidence of constitutive and
723 inducible clindamycin resistance in *Staphylococcus aureus* and coagulase-negative
724 staphylococci in a community and a tertiary care hospital. *J Clin Microbiol* **42**: 2777-
725 2779. Doi: <https://doi.org/10.1128/jcm.42.6.2777-2779.2004>.

726 Silva, V., Caniça, M., Capelo, J.L., Igrejas, G., and Poeta, P. (2020) Diversity and genetic
727 lineages of environmental staphylococci: A surface water overview. *FEMS Microbiol Ecol*
728 **96**: fiae191. Doi: <https://doi.org/10.1093/femsec/fiae191>.

729 Soge, O.O., Meschke, J.S., No, D.B., and Roberts, M.C. (2009) Characterization of
730 methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative
731 *Staphylococcus* spp. isolated from US West Coast public marine beaches. *J Antimicrob*
732 *Chemother* **64**: 1148-1155. Doi: <https://doi.org/10.1093/jac/dkp368>.

733 Stalder, T., Barraud, O., Casellas, M., Dagot, C., and Ploy, M.-C. (2012) Integron
734 involvement in environmental spread of antibiotic resistance. *Front Microbiol* **3**: 119. Doi:
735 <https://doi.org/10.3389/fmicb.2012.00119>.

736 Stefani, S., and Varaldo, P. (2003) Epidemiology of methicillin-resistant staphylococci in
737 Europe. *Clin Microbiol Infect* **9**: 1179-1186. Doi: <https://doi.org/10.1111/j.1469-0691.2003.00698.x>.

738 Tan, L., Li, L., Ashbolt, N., Wang, X., Cui, Y., Zhu, X. et al. (2018) Arctic antibiotic
739 resistance gene contamination, a result of anthropogenic activities and natural origin. *Sci*
740 *Total Environ* **621**: 1176-1184. Doi: <https://doi.org/10.1016/j.scitotenv.2017.10.110>.

741 Tappin, A., and Millward, G. (2015) The English Channel: contamination status of its
742 transitional and coastal waters. *Mar Pollut* **95**: 529-550. Doi:
743 <https://doi.org/10.1016/j.marpolbul.2014.12.012>.

744 Timmerman, C.-A., Marchal, P., Denamiel, M., Couvreur, C., and Cresson, P. (2020)
745 Seasonal and ontogenetic variation of whiting diet in the Eastern English Channel and the
746 Southern North Sea. *Plos one* **15**: e0239436. Doi:
747 <https://doi.org/10.1371/journal.pone.0239436>.

748 Timmerman, C.-A., Giraldo, C., Cresson, P., Ernande, B., Travers-Trolet, M., Rouquette, M.
749 et al. (2021) Plasticity of trophic interactions in fish assemblages results in temporal stability
750 of benthic-pelagic couplings. *Mar Environ Res* **170**: 105412. Doi:
751 <https://doi.org/10.1016/j.marenvres.2021.105412>.

752 Tong, S.Y., Davis, J.S., Eichenberger, E., Holland, T.L., and Fowler Jr, V.G. (2015)
753 *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations,
754 and management. *Clin Microbiol Rev* **28**: 603-661. Doi: <https://doi.org/10.1128/cmr.00134-14>.

755 Vaiyapuri, M., Joseph, T.C., Rao, B.M., Lalitha, K.V., and Prasad, M.M. (2019)
756 Methicillin- Resistant *Staphylococcus aureus* in Seafood: prevalence, laboratory detection,
757 clonal nature, and control in seafood chain. *J Food Sci* **84**: 3341-3351. Doi:
758 <https://doi.org/10.1111/1750-3841.14915>.

759 Vanermen, N., Onkelinx, T., Courtens, W., Van De Walle, M., Verstraete, H., and Stienen,
760 E.W. (2015) Seabird avoidance and attraction at an offshore wind farm in the Belgian part of
761 the North Sea. *Hydrobiologia* **756**: 51-61. Doi: <https://doi.org/10.1007/s10750-014-2088-x>.


762 Verin, Y. (2017) IBTS 2017 cruise. *RV Thalassa*. Doi: <https://doi.org/10.17600/17001700>.

763 White, D., Lapworth, D.J., Civil, W., and Williams, P. (2019) Tracking changes in the
764 occurrence and source of pharmaceuticals within the River Thames, UK; from source to sea.
765 *Environ Pollut* **249**: 257-266. Doi: <https://doi.org/10.1016/j.envpol.2019.03.015>.

766


767 Xu, L., Chen, H., Canales, M., and Ciric, L. (2019) Use of synthesized double-stranded gene
768 fragments as qPCR standards for the quantification of antibiotic resistance genes. *J Microbiol*
769 *Methods* **164**: 105670. Doi: <https://doi.org/10.1016/j.mimet.2019.105670>.

770



771


772 Figure 1: Sampling stations in the English Channel and the North Sea.



773

774 Figure 2: Geographical distribution of antimicrobial resistant *Staphylococcus* strains (CoNS and
 775 CoPS).

776



777

778 Figure 3: Pearson correlation matrix between the phenotypic resistances of coagulase-negative and
 779 coagulase-positive *Staphylococcus* strains and the environmental parameter values. ERTH:
 780 erythromycin; FSAD: fusidic acid; CFOX: ceftiofur; TOBR: tobramycin; TETR: tetracycline; KANA:
 781 kanamycin; PENG: penicillin G. Temp: temperature; Sal: salinity; Oxy: oxygen saturation; Turb:
 782 turbidity; PHOS phosphate; NTRA: nitrate; NTRI: nitrite; SLCA: silicate; CPHL: chlorophyll a;
 783 TPHP: total phaeopigments; AMON: ammonium; OSMP: organic suspended matter; ISMP:
 784 inorganic suspended matter.

785