



A *Bos taurus* sequencing methods benchmark for assembly, haplotyping, and variant calling

Camille Ech , Carole Iampietro, Cl ment Birbes, Andreea Dr au, Claire Kuchly, Arnaud Di Franco, Christophe Klopp, Thomas Faraut, Sarah Djebali, Adrien Castinel, et al.

► To cite this version:

Camille Ech , Carole Iampietro, Cl ment Birbes, Andreea Dr au, Claire Kuchly, et al.. A *Bos taurus* sequencing methods benchmark for assembly, haplotyping, and variant calling. Plant and Genome San Diego 2024, Jan 2024, San Diego (California), United States. 10, 2024. hal-04432803

HAL Id: hal-04432803

<https://hal.science/hal-04432803>

Submitted on 26 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin e au d p t et   la diffusion de documents scientifiques de niveau recherche, publi s ou non,  manant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv s.



Distributed under a Creative Commons Attribution 4.0 International License


Inspired by the production of reference datasets in the Genome in a Bottle project, we sequenced a Charolais heifer using different technologies: Illumina paired-end, Oxford Nanopore, Pacific Biosciences (HiFi and CLR), 10X Genomics linked-reads, and Hi-C. To generate haplotypic assemblies, we also sequenced both parents with short reads. From these data, we constructed two haplotyped trio high quality reference genomes and a consensus assembly, using current software packages. **The assemblies generated using PacBio HiFi reach a size of 3.2 Gb**, which is significantly larger than the 2.7 Gb ARS-UCD1.2 reference. The BUSCO score of the consensus assembly reaches a completeness of 95.8% completeness among highly conserved mammalian genes. **We also identified 35,866 structural variants greater than 50 base pairs**. This assembly is a contribution to the bovine pangenome for the Charolais breed. These datasets will prove to be useful, allowing the community to gain additional insight into sequencing technologies for applications such as SNP, indel or structural variant calling, and de novo assembly.

More information in our data paper → <https://doi.org/10.1038/s41597-023-02249-1>

More information in our data paper → <https://doi.org/10.1038/s41597-023-02249-1>

1	2	3	4
Sample	Sequencing technology	Library type	Dataset
Father	Oxford Nanopore Illumina	Ligation sequencing gDNA 10X Genomics Chromium Hi-C	a,b,c
Mother	Oxford Nanopore Illumina PacBio	Ligation sequencing gDNA 10X Genomics Chromium Hi-C Circular Long Read	a,b,c
Heifer	Oxford Nanopore Illumina PacBio	Ligation sequencing gDNA 10X Genomics Chromium PCR Free 2x250bp Hi-C Circular Long Read Consensus Long Read	a, b, c, d

The parents and the heifer were sequenced using Oxford Nanopore Technologies on GridION and PromethION, Chromium 10X and the Hi-C method on MiSeq, HiSeq or NovaSeq6000. The heifer was additionally sequenced using Illumina 2x250pb on NovaSeq6000 and PacBio Sequel II (CLR and CCS (i.e. HiFi reads) mode). For the Trio approach, parent reads (2x150pb) from 10X Genomics Chromium data were used.




a-Long reads assemblies from Oxford Nanopore Technologies and Pacific Biosciences followed by a polishing step for erroneous assemblies and scaffolding step. **b**-10XChromium assembly and scaffolding with Supernova. **c**-Phased assembly with HiFi and parental Illumina reads. **d**-Phased assembly with HiFi and Hi-C data.

Comparison and evolution of the heifer assemblies size metrics after different steps in different assemblies pipelines.

	ARS-UCD1.2	Heifer w/ Parents short read data		Heifer w/ Heifer Hi-C data	
		Hap1	Hap2	Hap1	Hap2
Pipeline used		c		d	
Data type	CLR	CLR + Trio	CLR + Trio	CCS + Hi-C	CCS + Hi-C
Number of contigs	3 077	2 871	2 300	2 658	2 136
Total size (Gb)	2.7	3.16	3.11	3.08	3.18
N50 contigs length	12 000 000	71 619 842	69 165 538	80 106 842	71 644 334
BUSCO	95.7%	95.8%	95.3%	95.8%	95.7%
Phasing ratio	**	97.3%	96.7%	62.6%	60.5%
Contigs phasing ratio	**	97.5%	96.9%	84.6%	85.6%

Details of pipeline used in this study are shown in Figure 2. CLR (80X), CCS (40X) and Hi-C (28X) Illumina (30X)* BUSCO analysis was performed on polished contigs. ** Reference is not haplotyped

Length distribution of DEL (left) and INS (right) variants, detected by the different sequencing technologies on the Charolais heifer. Sharp peaks correspond to the structural variability generated by transposable elements (SINE, LTR and LINE). All the long reads technologies exhibit roughly the same distribution underlining the fact that they are comparable in their ability to detect DEL and INS variants (see also Fig 5). This holds also with short reads for DEL variants. In contrast, there is a significant difference between long and short read technologies in their ability to detect INS variants, short reads failing to detect a large proportion of INS variants, the larger the size the more pronounced the trend. This is an expected consequence of the limited size of reads that prevents to recover long insertions. **Long read technology seems imperative for an accurate detection of the full spectrum of structural variants.**



The majority of variants are identified by all the long read technologies as well as for the assembly comparison based variant detection. This upset plot again highlights the added value of long reads for structural variant detection.

High-quality phased and consensus genome assemblies for Charolais heifer with PacBio, Hi-C and parents data !
Very promising SV detection with long reads data !