

AvrLm10A/AvrLm10B a part of a multigene family of cooperating effectors largely conserved in Dothideomycetes and Sordariomycetes

Nacera Talbi, Yohann Petit-Houdenot, Claire-Line Marais, Bénédicte Ollivier, Francoise Blaise, Thierry Rouxel, Marie-hélène Balesdent, Isabelle Fudal

▶ To cite this version:

Nacera Talbi, Yohann Petit-Houdenot, Claire-Line Marais, Bénédicte Ollivier, Francoise Blaise, et al.. AvrLm10A/AvrLm10B a part of a multigene family of cooperating effectors largely conserved in Dothideomycetes and Sordariomycetes. Fungal Genetics Symposium IDF, Nov 2021, Paris, France. hal-04433125

HAL Id: hal-04433125 https://hal.science/hal-04433125

Submitted on 1 Feb2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AvrLm10A/AvrLm10B a part of a multigene family of cooperating effectors largely conserved in Dothideomycetes and Sordariomycetes.

<u>Nacera Talbi</u>¹, Like Fokkens², Corinne Audran³, Yohann Petit-Houdenot¹, Françoise Blaise¹,, Angélique Gautier¹, Thierry Rouxel¹, Marie-Hélène Balesdent¹, Martijn Rep² and Isabelle Fudal¹

¹UMR BIOGER, INRAE, AgroParisTech, Université Paris-Saclay, Thiverval-Grignon, France ²Molecular Plant Pathology, University of Amsterdam, Amsterdam, Netherlands ³UMR LIPME, Université de Toulouse, INRAE, CNRS, Castanet-Tolosan, France.

With only a few exceptions, fungal effectors (corresponding to small secreted proteins) have long been considered as species- or even isolate-specific. With the increasing availability of high-quality fungal genomes and annotations, including improved strategies to predict fungal effector repertoires and decipher their 3-D structure, trans-species or trans-genera families of effectors are unveiled.

Two avirulence effectors of *Leptosphaeria maculans*, the fungus responsible for stem canker of oilseed rape, AvrLm10A and AvrLm10B, are members of such a large family of effectors. AvrLm10A and AvrLm10B are two neighbor genes organized in opposite transcriptional orientation. Both are necessary to trigger recognition by the Rlm10 resistance gene, and the two corresponding proteins physically interact *in vitro* and *in planta* [1]. Sequence searches within the L. maculans genome indicated that AvrLm10A/AvrLm10B belong to a multigene family comprising five pairs of genes with a similar tail-to-tail organization, a very unusual feature for this species in which RIP is very active to prevent duplications of genes. A search for homologues in fungal protein databases found that AvrLm10A was conserved in more than 30 Dothideomycetes and Sordariomycetes plant pathogenic fungi, with other cases of variable copy numbers within a genome. One of the AvrLm10A homologues, SIX5, is an avirulence effector from *Fusarium oxysporum* f.sp. *lycopersici* functioning as a cooperating pair with Avr2 [2]. Compared to AvrLm10A, less AvrLm10B homologues were identified and we found that AvrLm10A homologues could, in fact, be associated with at least five distinct effector families, suggesting an ability of AvrLm10A / SIX5 to cooperate with several effectors. A detailed characterization of the AvrLm10 family was performed in L. maculans, including transcriptomic and functional analyses. The two genes in a pair always had the same expression pattern and, depending on the pair of genes, two expression profiles were distinguished, associated with the biotrophic colonization of cotyledons or petioles and stems. One of the five pairs had no detectable expression and could correspond to pseudogenes. Of the three pairs functionally investigated, two had the ability to physically interact. Since SIX5 and Avr2 also physically interact, these data point to a general role of the AvrLm10A protein as a transporter able to interact with a limited number of effector families whose encoding gene is co-regulated with the neighboring AvrLm10A homologue.

[1] Petit-Houdenot, Y., Degrave, A., Meyer, M., Blaise, F., Ollivier, B., Marais, C. L. *et al.* 2019. A two genesfor-one gene interaction between Leptosphaeria maculans and Brassica napus. New Phytologist, 223, 397-411.

[2] Ma L, Houterman PM, Gawehns F, Cao L, Sillo F, Richter H, Clavijo-Ortiz MJ, Schmidt SM, Boeren S, Vervoort J et al. 2015. The AVR2–SIX5 gene pair is required to activate I-2-mediated immunity in tomato. New Phytologist 208: 507–518.

Keywords: Leptosphaeria maculans, oilseed rape, avirulence, effectors, conservation, multigene family