



**HAL**  
open science

# Multi-scale modelling of nano-sensors based on carbon nanotubes (cnts) and conjugated polymers for water quality monitoring

Robert Benda, Eric Cancès, Bérengère Lebental

## ► To cite this version:

Robert Benda, Eric Cancès, Bérengère Lebental. Multi-scale modelling of nano-sensors based on carbon nanotubes (cnts) and conjugated polymers for water quality monitoring. Graphene & Co Annual Meeting 2019, Oct 2019, Bad Herrenalb, Germany. hal-04433924

**HAL Id: hal-04433924**

**<https://hal.science/hal-04433924v1>**

Submitted on 2 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# MULTI-SCALE MODELLING OF NANO-SENSORS BASED ON CARBON NANOTUBES (CNTS) AND CONJUGATED POLYMERS FOR WATER QUALITY MONITORING

R. Benda<sup>1,2,3</sup>, Eric Cancès<sup>2,3</sup> and B. Lebental<sup>1,4</sup>

<sup>1</sup>Laboratoire de physique des interfaces et des couches minces – Ecole Polytechnique, Centre National de la Recherche Scientifique : UMR7647, Route de Saclay 91128 PALAISEAU CEDEX, France

<sup>2</sup>Centre d'Enseignement et de Recherche en Mathématiques et Calcul Scientifique (CERMICS) – Université Paris Est (UPE), École des Ponts ParisTech (ENPC) – 6 et 8 avenue Blaise Pascal Cité Descartes - Champs sur Marne 77455 Marne la Vallée Cedex 2, France

<sup>3</sup>MATERIALS (INRIA Paris) – INRIA, École des Ponts ParisTech (ENPC) – France

<sup>4</sup>Laboratoire Instrumentation, Simulation et Informatique Scientifique (IFSTTAR/COSYS/LISIS) – IFSTTAR, PRES Université Paris-Est – 14-20 Boulevard Newton - Cité Descartes, Champs sur Marne

We study resistive chemical sensors based on percolating networks of CNTs functionalized non-covalently by conjugated polymers. A change of resistance is expected in water upon complexation of a target ion by a specifically designed polymer probe. Multi-scale modelling aims at understanding and predicting the response of the sensor.

First, we derive the functional dependence of the effective resistance of random percolating networks of stick nanowires on elementary physical parameters (nanowire linear resistance per unit length, nanowire/nanowire contact resistance, metallic electrode/nanowire contact resistance) [1]. This understanding of the three main contributions, of different physical origins, to the whole, experimentally measured sensor resistances, is useful for the design of optimal sensor dimensions and sensible element densities.

Then, we study by means of variable-charge molecular dynamics, based on ReaxFF force field (within LAMMPS), the geometrical features of non-covalent functionalization of CNTs by conjugated polymers [2]. We benchmark the performance of ReaxFF force field for pi-pi stacking interacting compounds against other high-level electronic (DFT) methods, with a focus on the level of theory needed to model accurately London dispersion interactions (wave-function based theory vs. density functional theory) and to capture charge transfer from the organic molecule (or conjugated polymer) to the CNT -- this charge transfer is suspected to act as a doping and to modulate the sensor response.

Finally, we study the complexation of target ions (e.g. chlorine, copper, calcium) by the polymer probes -- designed specifically for their affinity towards given ions -- by means of molecular dynamics simulations in solvent (water) with GROMOS classical force field, within GROMACS code. We test numerically the expected selectivity of the polymer probes to different ions, finding the possible binding sites and their corresponding binding energies. We confirm the binding energies obtained with this classical force field by ab-initio (DFT) calculations -- with CP2K code -- and derive association constants for the different ions and binding sites, with free energy calculation techniques using GROMACS.

## References

[1] R. Benda, E. Cancès and B. Lebental, arXiv:1904.04369, to be published in *Journal of Applied Physics*.

[2] R. Benda and al., *in preparation*.




Figure 1 : CNT / conjugated polymer interaction.




Figure 2: Random 2D percolating network of stick nanowires (e.g. CNTs) whose effective resistance is decomposed into three different contributions.