

**Mobility and transport of nalidixic acid and niflumic acid in saturated soil
columns**

Tao Luo^{a,b*}, Tao Chen^{a,b}, Jean-François Boily^b, Khalil Hanna^{a,c,*}

^a Univ Rennes, Ecole Nationale Supérieure de Chimie de Rennes, CNRS, ISCR-UMR 6226,
F-35000, Rennes, France.

^b Department of Chemistry, Umeå University, Umeå, SE-901 87, Sweden

^c Institut Universitaire de France (IUF), MESRI, 1 rue Descartes, 75231 Paris, France.

*Corresponding authors: tao.luo@ensc-rennes.fr; khalil.hanna@ensc-rennes.fr




Fig. S1 The breakthrough curves of NA in F soil. Initial condition: 100 mM NaCl, 10 μ M NA, pH 7.1.




Fig. S2 The breakthrough curves of NFA in S soil with or without NaN_3 . Initial condition: 100 mM NaCl, 10 μ M NFA, 100 mg/L NaN_3 , pH 6.5.




Fig. S3 The breakthrough curves of non-reactive tracer and modeling results. Initial condition: 200 mM NaCl, injection volume = 4 mL.




Fig. S4 Sorption kinetics of NA and NFA on two soils. Initial conditions: 2 g/L soil, 20 μ M NA, 20 μ M NFA, 100 mM NaCl, pH 7.




Fig. S5 pH edges of NA or NFA sorption on two soils using NaCl as electrolyte in single system. Initial condition: 2 g/L soil, 100 mM NaCl, 20 μ M NA, 20 μ M NFA.




Fig. S6 The breakthrough curves of NFA in F soil. Initial condition: 100 mM NaCl, 10 μ M NFA, pH 7.1. The subscript 's' and 'b' indicated NFA existed in the single and binary system, respectively. The vertical dash lines indicated the period of 24 h interruption.




Fig. S7 pH edges of NA and NFA co-sorption on two soils using NaCl as electrolyte in binary system. Initial condition: 2 g/L soil, 100 mM NaCl, 20 μ M NA, 20 μ M NFA.




Fig. S8 pH edges of NA and NFA sorption on two soils using CaCl_2 as electrolyte in a, b) single and c, d) binary system. Initial condition: 2 g/L soil, 35 mM CaCl_2 , 20 μM NA, 20 μM NFA.




Fig. S9 pH edges of CIP sorption in soil. Initial condition: 2 g/L soil, 20 μM CIP.

Table S1 soil properties

	S soil	F soil
pH	6.5	7.1
Humidity, %	2.0	1.4
CEC, m _{eq} /100g	13.4	13.2
OC, %	2.9	2.2
Sand, wt %	10.8	20.5
Silt, wt %	66.9	59.8
Clay, wt %	22.3	19.7
TN, g/kg N	2.2	2.3
P ₂ O ₅	16	241
Exchangeable elements, mg/kg	CaO	2364
	MgO	61
	K ₂ O	31
	Cd	<0.5
	Al	26000
	Co	13
Total elements, mg/kg	Cu	35
	Fe	34829
	Mn	688
	Zn	76
		128

Table S2 Parameters of NA and NFA transport under flow condition

	F soil								S soil											
	NA		NFA		NFA+NA				NFA+NA+Ca ²⁺				NA		NFA		NFA+NA			
					NA	NFA	NA	NFA	NA	NFA	NA	NFA					NA	NFA		
pH					7.1										6.5					
^a θ					0.68										0.645					
^b q					0.24868										0.24868					
^c λ					0.2639										0.398					
^d Stage	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2		
^e f	0.76	0.23	0.03	0.13	0.76	0.23	0.01	0.17	0.76	0.23	0.03	0.05	0.48	0.40	0.04	0.07	0.48	0.40		
^f K	115	115	64	381	115	115	137	889	115	115	93	135	329	135	82	95	329	135		
^g β	1.2	1.2	1.47	0.12	1.2	1.2	1.3	0.1	1.2	1.2	1	0.1	1.2	0.69	1.3	0.36	1.2	0.69		
^h α	7.2E ⁻⁴	1.3E ⁻³	2.4E ⁻⁴	3.7E ⁻⁴	7.2E ⁻⁴	1.3E ⁻³	9.8E ⁻⁴	3.1E ⁻⁴	7.2E ⁻⁴	1.3E ⁻³	1E ⁻⁴	1.6E ⁻⁴	8.6E ⁻⁷	7E ⁻⁴	2.4E ⁻⁴	2.4E ⁻⁴	8.6E ⁻⁷	7E ⁻⁴		
R ²	0.99	0.97	0.99	0.94	0.99	0.97	0.99	0.95	0.99	0.97	0.99	0.99	0.99	0.95	0.99	0.99	0.95	0.99		

^aθ is water content.

^bq is Darcy velocity.

^cλ is the dispersivity.

^dStage 1, 2 are the two stages of injection solution. Stage 1 is the injection of reactive solute from 0-110 PV. Stage 2 is the injection of background solution from 110 PV-end.

^ef is the fraction of equilibrium sites.

^fK is the empirical distribution coefficient in freundlich isotherm. Unit: g⁻¹ cm³.

^gβ is the index of freundlich isotherm.

^hα is the adsorption rate.

Table S3 Freundlich fitting results of NA and NFA sorption isotherm.

	Equation	k_f^a	n	R^2
S soil, NA	Freundlich	0.647	0.59	0.99
S soil, NFA	Freundlich	0.703	0.337	0.98
F soil, NA	Freundlich	0.345	0.678	0.99
F soil, NFA	Freundlich	0.609	0.234	0.87

^a unit: g⁻¹ L.