



HAL
open science

Effects of cosmic rays on hydrocarbon interstellar dust

Marie Godard, Géraldine Féraud, Marin Chabot, Yvain Carpentier, Thomas Pino, Rosario Brunetto, J. Duprat, Cécile Engrand, Philippe Brechignac, Louis Le Sergeant D'hendecourt, et al.

► **To cite this version:**

Marie Godard, Géraldine Féraud, Marin Chabot, Yvain Carpentier, Thomas Pino, et al.. Effects of cosmic rays on hydrocarbon interstellar dust. ECLA - European Conference on Laboratory Astrophysics, Sep 2011, Paris, France. pp.395 - 398, 10.1051/eas/1258067 . hal-04435178

HAL Id: hal-04435178

<https://hal.science/hal-04435178v1>

Submitted on 2 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EFFECTS OF COSMIC RAYS ON HYDROCARBON INTERSTELLAR DUST

M. Godard^{1,2}, G. Féraud³, M. Chabot⁴, Y. Carpentier^{3,5}, T. Pino³,
R. Brunetto¹, J. Duprat⁶, C. Engrand⁶, P. Bréchnac³,
L. d'Hendecourt¹ and E. Dartois¹

Abstract. Hydrogenated amorphous carbons, an important component of the interstellar carbonaceous dust, possess infrared spectral signatures (at 3.4, 6.9 and 7.3 μm) that are ubiquitous in the diffuse interstellar medium of galaxies, but not observed in dense clouds. To better understand the role played by cosmic rays in the disappearance of these absorption bands, irradiation experiments of hydrocarbon dust analogues have been performed with different swift ions. The results obtained through the *in situ* infrared monitoring of the samples during the irradiations allow to infer the dehydrogenation effect of the cosmic ray distribution on the interstellar hydrogenated amorphous carbons. The importance of this interstellar dust destruction by cosmic rays is discussed in comparison to other energetic processes in different interstellar environments.

1 Introduction

Dust is a major component of the interstellar medium that continuously evolves with the different energetic processes active in such environments (UV photons, cosmic rays, shocks, heating, ...). Observations of the interstellar spectral signatures, together with laboratory simulations of dust processing, allow to get insights into the composition and evolution of the different types of interstellar grains.

¹ Institut d'Astrophysique Spatiale, Univ. Paris-Sud, Bât. 121, 91405 Orsay, France

² Astrochemistry Laboratory (Code 691), NASA Goddard Space Flight Center, Building 34, 8800 Greenbelt Road, Greenbelt, MD 20771, USA

³ Institut des Sciences Moléculaires d'Orsay, Univ. Paris-Sud, Bât. 210, 91405 Orsay, France

⁴ Institut de Physique Nucléaire d'Orsay, Univ. Paris-Sud, Bât. 102, 91405 Orsay, France

⁵ Laboratoire de Physique des Lasers, Atomes et Molécules, Univ. Lille 1, Bât. P5, 59655 Villeneuve d'Ascq Cedex, France

⁶ Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, Univ. Paris-Sud, Bât. 104, 91405 Orsay, France

One component of the carbonaceous dust is observed through the $3.4 \mu\text{m}$ absorption band, ubiquitous along sight lines that sample the diffuse interstellar medium of the Milky Way and other galaxies (*e.g.* Dartois *et al.* 2004; Pendleton *et al.* 1994). This feature is attributed to the stretching modes of the C-H bonds in methyl ($-\text{CH}_3$) and methylene ($-\text{CH}_2-$) groups. The corresponding bending modes (6.85 and $7.25 \mu\text{m}$) are also observed. These interstellar signatures of aliphatic hydrocarbon dust are well reproduced by hydrogenated amorphous carbons (a-C:H or HAC) that can be synthesized in laboratory to produce interstellar analogues. One of the puzzling observations concerning this carbonaceous dust component is the fact that the $3.4 \mu\text{m}$ feature is widely detected in the diffuse interstellar medium, but is not observed in dense clouds. Considering the rapid cycling of dust between these two interstellar phases, it implies an efficient mechanism for the destruction/modification of aliphatic hydrocarbon dust.

As energetic particles are known to dehydrogenate aliphatic materials, exposure of interstellar dust to cosmic rays (CR) could play a role in the disappearance of the $3.4 \mu\text{m}$ feature in dense clouds. A thorough investigation of the CR effects on the carbonaceous dust would help to fully understand its evolution. We have performed experiments of carbonaceous interstellar analogues irradiation by many swift ions. This work (Godard *et al.* 2011) represents a step further to achieve this goal and extends the work done by Mennella *et al.* (2003).

2 Experiments and results

For these experiments, two kinds of hydrocarbon samples, deposited on KBr substrates, have been used: a-C:H (produced with a hydrocarbon plasma and reproducing the spectral signatures of interstellar ones) (Godard & Dartois 2010) and soots (aromatic solids that nucleate in a propylene flame) (Pino *et al.* 2008). These materials have been exposed to many different ions (from hydrogen to iodine) and high energies (between 0.2 and 160 MeV) at the Tandem accelerator of the Institut de Physique Nucléaire (Orsay, France) and at the Laboratory for Experimental Astrophysics (INAF-Catania, Italy). These swift ions allow to reproduce a wide range of conditions comparable to the distribution of elements and energies of interstellar CR. During the experiments, the induced dehydrogenation and chemical bonding modifications were studied with *in situ* infrared spectroscopy.

The decreasing optical depth of the aliphatic C-H features with the increasing fluence (*cf.* Fig. 1) has been carefully analyzed with a model describing the mechanism of hydrogen loss in the material (Adel *et al.* 1989; Marée *et al.* 1996) and based on the fact that hydrogen leaves the irradiated material in molecular form (Möller & Scherzer 1987; Wild & Koidl 1987). We obtained the dehydrogenation parameters for each experiment (dehydrogenation cross section and asymptotic hydrogen content remaining in the irradiated material at high fluence) and their dependance with the electronic stopping power S_e (*i.e.* the energy deposited from the ion to electrons of the irradiated material, per length unit). The values obtained for a-C:H, soots, and other hydrocarbon materials found in the literature

are quite comparable, pointing out that these parameters do not seem very dependent of the H/C ratio for example. Since S_e can be calculated for each ion, these results allow thus to infer the dehydrogenation effect by each cosmic ray of given element and energy.




Fig. 1. Example of the decreasing optical depth of the aliphatic C-H stretching IR absorption feature around 2900 cm^{-1} ($3.4\text{ }\mu\text{m}$) during ion irradiation of a-C:H with 85 MeV Si^{7+} , between 0 (red upper line) and 10^{14} ions cm^{-2} (purple lower line). The fluence between two spectra is of the order of few 10^{12} ions cm^{-2} .

3 Effects of cosmic rays inferred from experiments

With our results, and by adopting an abundance and energy distribution of interstellar cosmic ray flux from the literature (<http://galprop.stanford.edu/>; Shen *et al.* 2004), we calculated the summed effect by all cosmic rays (lighter than copper) of different energies (below 10 GeV) and obtained the evolution of the interstellar aliphatic dust component (and the corresponding spectral signatures) as a function of cosmic ray exposure time. We found that the destruction of aliphatic C-H bonds by cosmic rays occurs in characteristic times of a few 10^8 years, a time longer than the characteristic lifetime of an interstellar cloud ($\sim 10^7$ years). Moreover, taking into account the non-zero asymptotic hydrogen content remaining at high fluence shows that, even for longer timescales, the destruction of interstellar a-C:H bonds by cosmic rays only cannot be complete. Even if there are uncertainties on the values of interstellar cosmic ray flux and ionisation rate¹, a higher CR flux than the one adopted could not destroy more than $\sim 60\%$ of this dust component. Thus, cosmic rays destroy interstellar a-C:H, but do not seem efficient enough to explain the disappearance of the $3.4\text{ }\mu\text{m}$ band in dense clouds.

This work confirms also that heavy ions play an important role in the destruction, even if they are less abundant. For example, we estimate that iron cosmic rays alone contribute between 5% and 40% to the destruction, although they represent only one cosmic ray over 10 000.

These results, concerning the exposure to cosmic rays deduced from experiments, help to better understand the evolution of interstellar a-C:H. They can be compared to the processing of hydrogenated amorphous carbons by UV photons

¹We have checked that the adopted cosmic ray flux was compatible with the observed interstellar ionisation rate of hydrogen by CR.

(Mennella *et al.* 2001) and H atoms (Mennella 2006) in the different interstellar regions. In diffuse interstellar medium, the formation by atomic hydrogen prevails over the destruction by UV photons. Relatively to other processing, destruction by cosmic rays is negligible in diffuse interstellar medium, but not in dense clouds where UV cannot penetrate and the hydrogen is mostly molecular. However, CR are not efficient enough to completely dehydrogenate the $3.4 \mu\text{m}$ carriers during the cloud lifetime. This interstellar component should thus be destroyed in interfaces between diffuse and dense regions, where photons still penetrate but hydrogen, already in molecular form, cannot compensate this destruction.

References

- Adel, M.E., Amir, O., Kalish, R., & Feldman, L.C., 1989, *J. Appl. Phys.*, 66, 3248
- Dartois, E., Marco, O., Muñoz-Caro, G.M., Brooks, K., Deboffle, D., & d'Hendecourt, L., 2004, *A&A*, 423, 549
- Godard, M., & Dartois, E., 2010, *A&A*, 519, A39
- Godard, M., Féraud, G., Chabot, M., *et al.*, 2011, *A&A*, 529, A146
- Marée, C., Vredenberg, A., & Habraken, F., 1996, *Mater. Chem. Phys.*, 46, 198
- Mennella, V., 2006, *ApJ*, 647, L49
- Mennella, V., Baratta, G.A., Esposito, A., Ferini, G., & Pendleton, Y.J., 2003, *ApJ*, 587, 727
- Mennella, V., Muñoz Caro, G.M., Ruiterkamp, R., *et al.*, 2001, *A&A*, 367, 355
- Möller, W., & Scherzer, B.M.U., 1987, *Appl. Phys. Lett.*, 50, 1870
- Pendleton, Y.J., Sandford, S.A., Allamandola, L.J., Tielens, A.G.G.M., & Sellgren, K., 1994, *ApJ*, 437, 683
- Pino, T., Dartois, E., Cao, A.-T., *et al.*, 2008, *A&A*, 490, 665
- Shen, C.J., Greenberg, J.M., Schutte, W.A., & van Dishoeck, E.F., 2004, *A&A*, 415, 203
- Wild, C., & Koidl, P., 1987, *Appl. Phys. Lett.*, 51, 1506