



HAL
open science

Could reasons for admission help to screen unhealthy alcohol use in emergency departments? A multicenter French study

Jonathan Chabert, Céline Lambert, Julien Cabé, Cheryl Cherpitel, Benjamin Rolland, Farès Moustafa, Patrick Lesage, Delphine Ragonnet, Julie Geneste, Emmanuel Poulet, et al.

► To cite this version:

Jonathan Chabert, Céline Lambert, Julien Cabé, Cheryl Cherpitel, Benjamin Rolland, et al.. Could reasons for admission help to screen unhealthy alcohol use in emergency departments? A multicenter French study. *Frontiers in Psychiatry*, 2023, 14, pp.1271076. 10.3389/fpsy.2023.1271076. hal-04435498

HAL Id: hal-04435498

<https://hal.science/hal-04435498>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY

Mauro Ceccanti,
Sapienza University of Rome, Italy

REVIEWED BY

Torgeir Gilje Lid,
Stavanger University Hospital, Norway
Giuliano Bertazzoni,
Sapienza University of Rome, Italy

*CORRESPONDENCE

Jonathan Chabert
✉ jchabert1@chu-clermontferrand.fr

RECEIVED 01 August 2023

ACCEPTED 16 November 2023

PUBLISHED 30 November 2023

CITATION

Chabert J, Lambert C, Cabé J, Cherpitel CJ, Rolland B, Moustafa F, Lesage P, Ragonnet D, Geneste J, Poulet E, Dematteis M, Naassila M, Chalmeton M, Llorca P-M, Pereira B, De Chazeron I and Brousse G (2023) Could reasons for admission help to screen unhealthy alcohol use in emergency departments? A multicenter French study.
Front. Psychiatry 14:1271076.
doi: 10.3389/fpsy.2023.1271076

COPYRIGHT

© 2023 Chabert, Lambert, Cabé, Cherpitel, Rolland, Moustafa, Lesage, Ragonnet, Geneste, Poulet, Dematteis, Naassila, Chalmeton, Llorca, Pereira, De Chazeron and Brousse. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Could reasons for admission help to screen unhealthy alcohol use in emergency departments? A multicenter French study

Jonathan Chabert^{1*}, Céline Lambert², Julien Cabé¹, Cheryl J. Cherpitel³, Benjamin Rolland⁴, Farès Moustafa⁵, Patrick Lesage⁶, Delphine Ragonnet⁷, Julie Geneste¹, Emmanuel Poulet⁸, Maurice Dematteis⁹, Mickael Naassila¹⁰, Maryline Chalmeton¹, Pierre-Michel Llorca¹, Bruno Pereira², Ingrid De Chazeron¹ and Georges Brousse¹

¹Service de Psychiatrie Adulte et d'Addictologie, CHU Clermont-Ferrand, CNRS, Université Clermont-Auvergne, Institut Pascal, Clermont-Ferrand, France, ²Unité de Biostatistiques, DRCl, CHU Clermont-Ferrand, Clermont-Ferrand, France, ³Alcohol Research Group, Emeryville, CA, United States, ⁴Service Universitaire d'Addictologie de Lyon, Centre Hospitalier Le Vinatier, Hospices Civils de Lyon et Université de Lyon, Lyon, France, ⁵Université Clermont Auvergne, INRAE, UNH, Clermont-Ferrand, France, ⁶Centre Hospitalier Métropole Savoie, Service des Urgences, Chambéry, France, ⁷Service Universitaire d'Addictologie de Lyon, Groupement Hospitalier Centre, Hospices Civils de Lyon, Lyon, France, ⁸Psychiatrie des Urgences - Groupement Hospitalier Edouard Herriot, EA 4615 « SIPAD », Université Lyon 1 - CH Le Vinatier, Lyon, France, ⁹Service Universitaire de Pharmaco-Addictologie, CHU Grenoble Alpes, Université Grenoble Alpes, Grenoble, France, ¹⁰INSERM UMRS1247-GRAP, Université Picardie Jules Verne, Amiens, France

Background: Many patients admitted to general emergency departments (EDs) have a pattern of drinking that could lead to future alcohol-related complications. However, it is often difficult to screen these patients in the context of emergency. The aim of this study is to analyze whether reasons for admission could help to screen patients who have an unhealthy alcohol use.

Method: Patients were recruited among six public hospital ED in France, between 2012 and 2014. During a one-month period in each hospital, anonymous questionnaires including sociodemographic questions, AUDIT-C and RAPS4-QF were administered to each patients visiting the ED. The reason for admission of each patient was noted at the end of their questionnaire by the ED practitioner.

Results: Ten thousand Four hundred twenty-one patients were included in the analysis. Patients who came to the ED for injuries and mental disorders were more likely to report unhealthy alcohol use than non-harmful use or no use. Among male patients under 65 years old admitted to the ED for a mental disorder, 24.2% drank more than four drinks (40 g ethanol) in typical day at least four time a week in the last 12 months. Among these patients, 79.7% reported daily or almost daily heavy episodic drinking (HED, 60 g ethanol), and all were positive on the RAPS4-QF.

Conclusion: This study highlights that unhealthy alcohol use is frequent among ED patients and particularly among those who come for injuries or mental disorders. Men under 65 years old with a mental disorder require special attention because of their increased prevalence of daily or almost daily HED.

KEYWORDS

alcohol drinking, alcohol-related complications, heavy episodic drinking, emergency department, alcohol use disorder, mental disorder

1 Introduction

Alcohol is the psychoactive substance most used in France and a major public health problem since it is the leading cause of hospitalization in France (1) and is responsible for approximately 41,000 premature deaths each year (2). While many users will not be harmed by their consumption during their life, many others will therefore have alcohol-induced or related complications that could lead to functional impairment and in some cases to death (3).

In this context, screening for patients whose alcohol use is at-risk for or has already resulted in complications, also called unhealthy alcohol use, seems to be essential. However, while each country has developed its own guidelines to define what constitutes unhealthy alcohol use, there is currently no international consensus on permissible thresholds for alcohol quantity and frequency. For example, the National Institute of Alcohol Abuse and Alcoholism (NIAAA) recommends drinking in moderation by limiting intake to 2 standard American drinks (28 g ethanol) or less in a day for men and 1 standard American drink (14 g ethanol) or less in a day for women (4), while Canada's Guidance recommends 1 or 2 standard Canadian drink (14 g ethanol) per week to avoid alcohol-related consequences (5). The Alcohol Use Disorders Identification Test – Consumption (AUDIT-C) (6) is a brief and democratized screening test which allows to screen unhealthy alcohol use according to the NIAAA guidelines. AUDIT-C has good sensitivity and specificity in primary care (7) and is also brief with only three questions: frequency of use, quantity and presence of heavy episodic drinking (HED), which is a pattern of drinking corresponding to the consumption of 60 or more grams of pure alcohol in a single occasion in the last month (8). AUDIT-C is however not very good to identify alcohol use disorder (AUD), and it is therefore recommended to use another questionnaire, such as the Rapid Alcohol Problems Screen 4–Quantity and Frequency (RAPS4-QF), to screen AUD. The RAPS4-QF is a six items quick self-administrated questionnaire, which has a sensitivity of 90.3% and a specificity of 84.1% for AUD (9). Some studies have even shown that the RAPS4-QF has a better sensitivity for detecting alcohol dependence than two other frequently used questionnaires: AUDIT and CAGE (10, 11). Nevertheless, despite the three questions of AUDIT-C and the six questions of RAPS4-QF, these tools are still considered too long to be incorporated into medical interviews in some time-sensitive settings, such as emergency departments (ED). Thus, many EDs practitioners do not evaluate alcohol consumption of their patients if alcohol is not related to their diagnostic or treatment, although a high prevalence of unhealthy alcohol use (almost 10%) has been found in EDs (12, 13). Furthermore, patients are significantly more likely to report heavy drinking, consequences of drinking or alcohol dependence in EDs than in medical offices (14), and as D'Onofrio et al. highlighted: “patients are often more receptive to education in the moment of crisis” (15). Thus, the ED could be an excellent site for screening patients for their alcohol use.

In 1996, Smith et al. proposed a quick scale, named the Paddington Alcohol Test (PAT), which is adapted to the ED context since it is modelled on the diagnostic approach of emergency physicians (16). The PAT, in its last version, lists the ten most prevalent presenting conditions of patients misusing alcohol (17). These presenting conditions are not specific diseases, but are combinations of symptoms, syndromes, or diseases, allowing the clinician to quickly obtain an overview of the patient's probable alcohol consumption

before having an exact diagnosis. The PAT was the alcohol screening tool most used in the United Kingdom since 41% of EDs using it in 2013 (18). This approach is quite rare in the literature, however. While many authors have focused on the relationship between alcohol consumption and some conditions frequently seen in the EDs, as falls (19–21) and injuries (22, 23), very few have evaluated whether reasons for admission could be an indicator of interest for alcohol screening.

Thus, the aim of our study is to determine whether certain reasons for admission are more frequently associated with unhealthy alcohol use. However, as unhealthy alcohol use is a broad and heterogeneous category of patients, comprising drinking patterns with different risks of developing alcohol-related complications or the type of complications (3), we then analyzed the association between certain drinking patterns and reason for ED visit.

2 Method

2.1 Study design and participants

Subjects were recruited in the ED of six public hospitals in Auvergne-Rhône-Alpes region (France), between 2012 and 2014. Anonymous questionnaires were given during one month period at each hospital, to those admitted to the ED and meeting inclusion criteria. All the participants gave their informed consent. We used a cluster random sampling design and area sampling method. Clusters were defined along geographic areas, and the month of recruiting was randomly assigned. All patients admitted in each ED unit over the course of one month were defined as a single cluster.

Inclusion criteria were aged 16 years and older, agreeing to participate, and speaking French. Patients unable to participate due to somatic or cognitive major impairments were excluded.

The protocol was approved by the Ethic Committee of Clinical Investigation Center of Auvergne-Rhône-Alpes inter-region in France (CE-CIC/GREN/12/09).

Complete study protocol has been published elsewhere (24).

2.2 Data collection

The questionnaire included sociodemographic data (age, gender...) and two self-administered alcohol screening tests, the AUDIT-C and the RAPS4-QF.

After completing the questionnaires, EDs staff completed the survey with the reason for admission of the patient in a predefined list inspired of the PAT list. When the reason for admission of the patient did not correspond to the presenting condition of the predefined list, the ED staff wrote the exact diagnosis in a box called “other.”

The eight presenting conditions listed were: “fall,” “assault,” “head injury,” “collapse,” “digestive complaint,” “chest tightness and/or palpitations,” “mental disorder” and “unwell” (suffering of unknown or imprecise origin).

2.3 Psychometric scales

The AUDIT-C is a scale evaluating alcohol drinking with three questions: “How often did you have a drink containing alcohol in the

past year?," "How many drinks did you have on a typical day when you were drinking in the past year?" and "How often did you have six or more drinks on one occasion in the past year?" For each question, a standard drink corresponds to 10g of ethanol. Each answer is rated between 0 to 4, so the maximum overall score is 12. A score ≥ 4 for men or ≥ 3 for women suggests an unhealthy alcohol use according to NIAAA.

The RAPS4-QF consists of the four questions of RAPS4: "During the last year, have you had a feeling of guilt or remorse after drinking? (Remorse)," "During the last year, has a friend or family member ever told you about things you said or did while you were drinking that you could not remember? (Amnesia, also called Blackouts)," "During the last year, have you failed to do what was normally expected from you because of drinking? (Perform)," "Do you sometimes take a drink in the morning, when you first get up? (Starter, also called eye opener)," and two additional questions on quantity and frequency: "During the last year, have you had five or more drinks on at least one occasion? (Quantity)," "During the last year, do you drink as often as once a month? (Frequency)." A positive response on any one of the RAPS4 items and/or both of the QF items is considered positive on the RAPS4-QF and suggests AUD.

2.4 Categories of conditions for analysis

The reason for admission of patients was classified into 23 categories. The first eight categories were the eight presenting conditions listed (see above). The other 15 categories were partially based on the International Classification of Disease 11th revision for Mortality and Morbidity Statistics (ICD-11 MMS). Patients who came to the EDs for obstetrical or dental conditions were excluded because there are specific EDs in France which treat these conditions, and therefore the few patients in these study with those conditions were likely not representative.

For the first stage of the analysis, based on the AUDIT-C, patients were divided into three categories: "no use," which correspond to ED's patients who did not drink alcohol the last year, "low risk alcohol use," which correspond to ED patients whose alcohol use met NIAAA guidelines, and "unhealthy alcohol use," which correspond to ED patients whose alcohol use is above NIAAA guidelines.


For the second stage of the analysis, as the unhealthy alcohol use is a heterogeneous category that presents a broad spectrum of risk of complications for patients, we decided to explore more specific pattern of alcohol use to gain precision. The first pattern is based on amount and frequency of alcohol use, and use the two first questions of the AUDIT-C to divide the patients into five categories: "non-drinker," "infrequent drinker" (those who consume less than four times a week and four or less standard drinks in a typical day), "heavy infrequent drinker" (those who consume less than four times a week and more than four standard drinks in a typical day), "frequent drinker" (those who consume four or more times a week and four or less standard drinks in a typical day) and "heavy frequent drinker" (those who consume more than four times a week and more than four standard drinks in a typical day). The quantity cut-off of four standard drinks (40g ethanol) was chosen because it synthesized both the four-drinker-per-occasion limit proposed by the WHO Collaborative group (25), and the NIAAA guidelines that consider drinking more than three standard American drinks (42g ethanol, about four standard French

drinks) for women and four standard American drinks for men (56g ethanol, about five standard French drinks) is heavy drinking (26). This threshold has also been used elsewhere as cut-off for heavy drinking (3, 27). The drinking frequency cut-off was used to differentiate non-daily drinkers from almost daily-drinkers, since daily alcohol drinking without an alcohol free day increase risk of several complications (28). This is also why many guidelines recommend one or more alcohol-free days per week (4, 8, 29). From these categories, we proposed two conditional inference trees, one based on the amount and frequency of alcohol consumption to assess heavy drinking (Figure 1), and the other based on the presence of HED (Figure 2). Conditional inference trees have the advantage of providing outcomes that can be easily used by practitioners when interviewing patients.

For the tree based on the presence of HED, the third question of the AUDIT-C was used to divide patients into three categories: "No HED," "HED" and "Chronic HED." The first cut-off is based on the WHO's definition of HED ("at least once a month" (8)), and the second cut-off is used to differentiate irregular HED from daily or almost daily heavy drinking (3).

2.5 Statistical analysis

Statistical analysis was performed using Stata software (version 15; StataCorp, College Station, Texas, United States) and R (version 4.1.3; R Foundation for Statistical Computing, Vienna, Austria). All tests were two-sided, with an alpha level set at 5%. No correction for multiple testing was applied in the analysis of outcomes or subgroup analysis (30). The patients' age was presented as mean \pm standard deviation, and the categorical variables as the number of patients, associated percentages, and 95% confidence interval (CI). Comparisons between included and non-included patients were made by the Student's t test for the patients' age, and by the Chi-squared test for categorical variables (data not shown). Comparisons between independent groups (e.g., no use vs. low risk alcohol use vs. unhealthy alcohol use) were made by ANOVA for the patients' age and by the Chi-squared test for categorical variables. When appropriate (omnibus value of p less than 0.05), a Tukey-Kramer test was performed after ANOVA, and a Marascuilo *post-hoc* procedure after Chi-squared test. Effect sizes were measured by Cramér's V and interpreted as: 0.1 = small effect, 0.3 = medium effect and 0.5 = large effect. Then, conditional inference trees were built with the "party" R package (31). The tree begins with a node called the root, representing the entire training dataset. At each node of the tree, the input variable with the strongest association to the response is recursively selected to split the data binary into two subsets. The splitting process is repeated iteratively, choosing at each step the variable that optimizes the splitting criterion. Thus, each variable (e.g., age, gender, ...) can be split at different stages of the analysis. Tree construction continues until a stopping criterion is met, such as reaching a maximum tree depth or a minimum number of samples per subsets. Two regressions tree were built with the same input variables (age in four categories, gender, and reasons for admission) but with distinct responses: the first based on the amount and frequency of alcohol consumption, and the second based on the presence of HED. The four categories for age were selected in advance. As these analyses were only descriptive, the entire sample was used (no validation set).



3 Results

Among the 23,834 patients admitted to one of the six EDs during the study period, 11,632 patients filled out the questionnaires, and 10,421 patients were included in the analysis. The excluded patients were those with obstetrical or dental conditions and patients with significant missing data on age or gender. See [Supplementary Table 1](#) for details on the included population.

3.1 No use vs. low risk alcohol use vs. unhealthy alcohol use

Similar proportion of patients were in each of the three categories: 34.3% for no use, 29.9% for low-risk alcohol use and 35.8% for unhealthy alcohol use. See [Table 1](#) for the comparison of sociodemographic data between the three categories.

[Table 2](#) describes the proportion of alcohol drinking for each reason for admission. Patients who presented to ED for (ranking from the highest to the lowest percentage): "head injury," "mental disorder," "assault," "injury, poisoning or certain other consequences of external causes," "fall," "public road accident," "pathologies of the circulatory system," "chest tightness and/or palpitations," "medical examination/post-acute care" or "genitourinary pathologies," were more likely to engage in unhealthy drinking than low-risk drinking, or not to drink at all.

3.2 Regression trees

3.2.1 First tree – amount and frequency of alcohol use

The first tree describes the amount and frequency of alcohol use in ED patients according to reasons for admission, gender, and age ([Figure 1](#)). A first distinction is made between patients aged 65 years old and over and patients aged less than 65 years old.

In patients aged 65 years old and over, regardless of gender, less than 2.5% drank more than four drinks a typical day.

In patients aged less than 65 years old, there is an important distinction in alcohol drinking depending on the reason for admission and gender. There was a significant difference in the proportion of heavy frequent drinkers (more than four drinks in a typical day at least four times a week during the last 12 months) between men under 65 years old who came to the ED for a reason related to a mental disorder (24.2, 95%CI: 19.5 to 29.4%), women under 65 years old who came for a reason related to a mental disorder (8.3, 95%CI: 5.6 to 11.8%) and all others patients (2.3, 95%CI: 2.0 to 2.6%) ($p < 0.001$).

3.2.2 Second tree – heavy episodic drinking

The second tree describes HED in ED's patients according to reasons for admission, gender, and age ([Figure 2](#)). A first distinction is made according to the reason for admission, with patients who came for reasons related to a mental disorder on the one hand and patients who came for other reasons on the other.




FIGURE 2 Proportion of heavy episodic drinking among emergency department's patients according to their reason of admission, age and gender. A = "Fall," "Assault," "Head injury," "Collapse," "Digestive complaint," "Chest tightness and/or palpitations," "Unwell," "Public road accident," "Medical examination/Post-acute care," "Dermatological pathologies," "Altered general condition," "Infectious diseases," "Injury, poisoning or certain other consequences of external causes," "Endocrine, nutritional or metabolic diseases," "Pathologies of the circulatory system," "Pathologies of the respiratory system," "Nervous pathologies," "Pathologies of the visual system," "Otolaryngology pathologies," "Osteoarticular pathologies," "Genitourinary pathologies" and "Pathologies of blood"; B = "Mental disorder"; C = chronic heavy episodic drinking; D = "Pathologies of the respiratory system" and "Otolaryngology pathologies"; E = "Fall," "Assault," "Head injury" "Collapse," "Gastro-intestinal," "Chest tightness and/or palpitation," "Unwell," "Public road accident," "Medical examination/Post-acute care," "Dermatological pathologies," "Infectious diseases," "Injury, poisoning or certain other consequences of external causes," "Pathologies of the circulatory system," "Nervous pathologies," "Pathologies of the visual system," "Osteoarticular pathologies," "Genitourinary pathologies" and "Pathologies of blood"; I = irregular heavy episodic drinking; M = Men; N = no heavy episodic drinking; W = Women.

TABLE 1 Sociodemographic characteristics of emergency department's patients according to their alcohol use.

	Overall (N = 10,421)	No use (N = 3,576)	Low risk alcohol use (N = 3,111)	Unhealthy alcohol use (N = 3,734)	p-value	V
Age (years)	46.7 ± 21.7	48.7 ± 23.0	46.9 ± 20.5	44.6 ± 21.3	<0.001	–
Female gender	4,734 (45.4)	2,094 (58.6)	1,446 (46.5)	1,194 (32.0)	<0.001	0.22
Having children*	6,288 (60.6)	2,345 (65.9)	1,940 (62.6)	2,003 (53.9)	<0.001	0.11
Marital status†					<0.001	0.13
Single	3,565 (34.4)	1,089 (30.6)	942 (30.5)	1,534 (41.3)		
In relationship	1,239 (12.0)	254 (7.1)	447 (14.5)	538 (14.5)		
Married	3,793 (36.7)	1,487 (41.9)	1,192 (38.6)	1,114 (30.0)		
Divorced	935 (9.0)	322 (9.1)	274 (8.9)	339 (9.1)		
Widowed	818 (7.9)	401 (11.3)	230 (7.5)	187 (5.0)		
Education level‡					<0.001	0.18
None	2,554 (24.9)	1,351 (38.4)	572 (18.6)	631 (17.2)		
Secondary education	4,904 (47.7)	1,572 (44.7)	1,505 (48.9)	1,827 (49.7)		
Bachelor's degree	1,828 (17.8)	450 (12.9)	613 (19.9)	765 (20.8)		
Master's degree	787 (7.7)	111 (3.2)	305 (9.9)	371 (10.1)		
Doctorate's degree	197 (1.9)	31 (0.9)	81 (2.6)	85 (2.3)		

Data are presented as number of patients (percentages in columns), or as mean ± standard deviation.

V: Cramér's V in absolute value (interpreted as: 0.1 = small effect, 0.3 = medium effect and 0.5 = large effect).

*The number of available data was 10,374 (overall), 3,557 (no use), 3,099 (low risk alcohol use), and 3,718 (unhealthy alcohol use).

†The number of available data was 10,350 (overall), 3,553 (no use), 3,085 (low risk alcohol use), and 3,712 (unhealthy alcohol use).

‡The number of available data was 10,270 (overall), 3,515 (no use), 3,076 (low risk alcohol use), and 3,679 (unhealthy alcohol use).

TABLE 2 Reasons for admission of emergency department's patients according to their alcohol use.

	No use (N = 3,576) ¹	95% CI ²	Low risk alcohol use (N = 3,111) ¹	95% CI ²	Unhealthy alcohol use (N = 3,734) ¹	95% CI ²
Fall	485 (31.4)	29, 34%	454 (29.4)	27, 32%	604 (39.1)	37, 42%
Assault	89 (34.2)	29, 40%	60 (23.1)	18, 29%	111 (42.7)	37, 49%
Head injury	63 (23.2)	18, 29%	68 (25.0)	20, 31%	141 (51.8)	46, 58%
Collapse	266 (37.6)	34, 41%	231 (32.6)	29, 36%	211 (29.8)	26, 33%
Digestive complaint	485 (41.1)	38, 44%	374 (31.7)	29, 34%	320 (27.1)	25, 30%
Chest tightness and/or palpitations	191 (33.7)	30, 38%	167 (29.5)	26, 33%	208 (36.7)	33, 41%
Mental disorder	223 (30.8)	27, 34%	172 (23.8)	21, 27%	329 (45.4)	42, 49%
Unwell	255 (45.3)	41, 50%	157 (27.9)	24, 32%	151 (26.8)	23, 31%
Public road accident	84 (25.2)	21, 30%	119 (35.7)	31, 41%	130 (39.0)	34, 45%
Medical examination/ Post-acute care	42 (32.6)	25, 41%	42 (32.6)	25, 41%	45 (34.9)	27, 44%
Dermatological pathologies	102 (37.2)	32, 43%	83 (30.3)	25, 36%	89 (32.5)	27, 38%
Altered general condition	53 (50.0)	41, 59%	23 (21.7)	15, 31%	30 (28.3)	20, 38%
Infectious diseases	60 (37.5)	30, 46%	49 (30.6)	24, 38%	51 (31.9)	25, 40%
Injury, poisoning or certain other consequences of external causes	419 (26.4)	24, 29%	499 (31.5)	29, 34%	667 (42.1)	40, 45%
Endocrine, nutritional or metabolic diseases	29 (42.6)	31, 55%	24 (35.3)	24, 48%	15 (22.1)	13, 34%
Pathologies of the circulatory system	65 (31.2)	25, 38%	65 (31.2)	25, 38%	78 (37.5)	31, 44%
Pathologies of the respiratory system	147 (45.1)	40, 51%	73 (22.4)	18, 27%	106 (32.5)	28, 38%
Nervous pathologies	172 (37.2)	33, 42%	156 (33.8)	29, 38%	134 (29.0)	25, 33%
Pathologies of the visual system	35 (22.6)	16, 30%	69 (44.5)	37, 53%	51 (32.9)	26, 41%
Otolaryngology pathologies	56 (47.1)	38, 56%	25 (21.0)	14, 30%	38 (31.9)	24, 41%
Osteoarticular pathologies	144 (38.9)	34, 44%	104 (28.1)	24, 33%	122 (33.0)	28, 38%
Genitourinary pathologies	92 (33.9)	28, 40%	85 (31.4)	26, 37%	94 (34.7)	29, 41%
Pathologies of blood	19 (47.5)	32, 64%	12 (30.0)	17, 47%	9 (22.5)	11, 39%

¹Data are presented as number of patients (percentages in rows).

²CI is confidence interval.

Among patients who came for a reason related to a mental disorder, the proportion of chronic HED in the last 12 months was 21.3% (95%CI: 17.1 to 26.1%) for men and 5.2% (95%CI: 3.2 to 7.9%) for women. The proportion of irregular HED was 17.8% (95%CI: 13.8 to 22.3%) for men and 12.2% (95%CI: 9.1 to 15.9%) for women.

Among patients who did not come for a reason related to a mental disorder, there was a low proportion of HED for men aged 65 years old and over and women aged 40 years and over (less than 10%). However,

for men between 25 and 39 years old, the proportion of irregular HED was 22.7% (95%CI: 20.6 to 24.9%).

Among male patients younger than 25 years old, who were admitted for any condition other than "Mental disorder," "Altered General Condition," "Endocrine, nutritional or metabolic diseases," "Pathologies of the respiratory system" or "Otolaryngology pathologies," the proportion of irregular HED in the last 12 months was 33.8% (95%CI: 30.9 to 36.7%). Among those with respiratory or otolaryngology conditions the

proportion of irregular HED was 30.8% (95%CI: 14.3 to 51.8%) and chronic HED was 11.5% (95%CI: 2.4 to 30.2%).

3.3 Heavy frequent drinkers among men under the age of 65 who came for a reason related to a mental disorder

As described above, 24.2% of men under age 65 years who came for a reason related to a mental disorder were heavy frequent drinker. The proportion of chronic HED in these patients was 79.7% and the proportion of irregular HED 16.2%. All these patients were positive to the RAPS4-QF, and the majority (56.8%) drank at least 10 drinks in a typical day. Table 3 shows characteristics of these individuals.

4 Discussion

This study aimed to determine whether certain reasons for admission are more frequently associated with unhealthy alcohol use, according to NIAAA guidelines. We saw that there is no reason for admission that would allow practitioners to not screen for unhealthy alcohol use, since even for patients who go to the ED for endocrine, nutritional, or metabolic diseases, which are the reasons for the lowest rates of unhealthy alcohol use, more than one in five had unhealthy alcohol use. Furthermore, there were some reasons for admission for which patients were more likely to have unhealthy drinking than low risk drinking or not to drink at all. Finally, 24% of men, under 65, who came for a mental disorder in ED have a heavy alcohol use, more than four drinks in a typical day, more than four days a week in the last

TABLE 3 Heavy frequent drinkers among men under the age of 65 who came for a reason for admission related to a mental disorder (MMD).

	Overall (N = 10,421)	MMD (N = 74)	Others (N = 10,347)	p-value	V
<i>Age (years)</i>					
Having children*	6,288 (60.6)	42 (56.8)	6,246 (60.6)	0.50	0.007
<i>Marital status[†]</i>					
Single	3,565 (34.4)	42 (56.8)	3,523 (34.3)	<0.001	0.05
In relationship	1,239 (12.0)	8 (10.8)	1,231 (12.0)		
Married	3,793 (36.6)	10 (13.5)	3,783 (36.8)		
Divorced	935 (9.0)	12 (16.2)	923 (9.0)		
Widowed	818 (7.9)	2 (2.7)	816 (7.9)		
<i>Education level[‡]</i>					
None	2,554 (24.9)	15 (20.5)	2,539 (24.9)	0.21	0.02
Secondary education	4,904 (47.8)	42 (57.5)	4,862 (47.7)		
Bachelor's degree	1,828 (17.8)	8 (11.0)	1,820 (17.8)		
Master's degree	787 (7.7)	5 (6.8)	782 (7.7)		
Doctorate's degree	197 (1.9)	3 (4.1)	194 (1.9)		
<i>Heavy episodic drinking</i>					
None	8,852 (84.9)	3 (4.1)	8,849 (85.5)	<0.001	0.42
Irregular heavy episodic drinking	1,305 (12.5)	12 (16.2)	1,293 (12.5)		
Chronic heavy episodic drinking	264 (2.5)	59 (79.7)	205 (2.0)		
<i>Alcohol amounts a typical day[#]</i>					
1 to 2 drinks	4,120 (60.2)	0 (0.0)	4,120 (60.8)	<0.001	0.30
3 to 4 drinks	1,555 (22.7)	0 (0.0)	1,555 (23.0)		
5 to 6 drinks	623 (9.1)	16 (21.6)	607 (9.0)		
7 to 9 drinks	250 (3.7)	16 (21.6)	234 (3.5)		
≥10 drinks	297 (4.3)	42 (56.8)	255 (3.8)		
Positive RAPS4-QF	3,130 (30.0)	74 (100)	3,056 (29.5)		

Data are presented as number of patients (percentages in columns), or as mean ± standard deviation. MMD: heavy frequent drinkers among men under the age of 65 who came for a reason for admission related to a mental disorder; RAPS4-QF: rapid alcohol problems screen 4-quantity and frequency; V: Cramér's V in absolute value (interpreted as: 0.1 = small effect, 0.3 = medium effect and 0.5 = large effect).

*The number of available data was 10,374 (overall), 74 (MMD), and 10,300 (others).

[†]The number of available data was 10,350 (overall), 74 (MMD), and 10,276 (others).

[‡]The number of available data was 10,270 (overall), 73 (MMD), and 10,197 (others).

[#]the number of available data was 6,845 (overall), 74 (MMD), and 6,771 (others).

12 months. Of these patients, 80% had daily or almost daily HED, and 16% had HED at least once a month.

ED's practitioners must be particularly attentive to two categories of reasons for admission: injuries and mental disorder. In this study, at least 39% of patients who came for mental disorders, or an injury were classified in the unhealthy alcohol use category according to the AUDIT-C. The relationship between alcohol drinking and injuries is well known. Some authors found that about one third of patients admitted to the ED for injury reported drinking prior to the event (22, 32–34). This might suggest that blood alcohol concentration (BAC) or breath alcohol concentration (BrAC) could be enough to detect patients with an unhealthy alcohol use in the ED. Indeed, there is a correlation between alcohol use and risk of injury, with, according to Cherpitel et al., risk doubling with the first drink and continuing to increase with subsequent drinks (35). However, alcohol-related injuries are not limited to acute alcohol intoxication. Some damages caused by long term alcohol use, such as alcohol neuropathy (36), cerebellar impairment (37, 38), etc., can also increase the risk of injuries. These other risk factors could explain that 39% of our patients who fell had an unhealthy alcohol use, while two other studies with large samples found that only 2% of patients who came to the ED for a fall had drunk before injury (20, 21). Thus, to focus solely on BrAC and BAC could lead to missing many patients with an unhealthy alcohol use. This conclusion was also highlighted by Browne et al. who found that 30% of patients with an AUD in Australia were not identified by ED staff as having alcohol-related injury or an alcohol related problem (39).

Apart from injury, practitioners must be attentive to patients who come to the ED for a complaint related to a mental disorder. Our data suggest that almost 7% of ED patients came for a mental disorder, which is more frequent than the average of other countries (4%) (40), and that 45% of them had an AUDIT-C score in favor of unhealthy alcohol use. This high rate of unhealthy alcohol use is not surprising as AUD is highly associated with many mental disorders (41–44). This study was designed to provide an overview of alcohol use associated with the general reason for ED admission and does not allow us to discern which mental disorder(s) are most closely associated with unhealthy alcohol use. This information could prove valuable to better screen patients, especially since the literature on this subject is relatively thin. Among the few studies reported, 33% of schizophrenic or bipolar patients and 22% of depressive or anxious patients who came to psychiatric emergency had at least hazardous drinking (45).

Although, we cannot describe which mental disorder is more closely associated with unhealthy alcohol use, data here suggest that there is a difference in unhealthy alcohol use's severity according to gender and age. The regression tree showed that men under age 65 admitted to the ED with a mental disorder had a 24% risk of drinking more than four standard drinks in a typical day, at least four times a week during the last 12 months. Such high frequency and amount of alcohol use can lead to many diseases. Indeed, it is well known that the risk of developing certain diseases increases linearly, or even exponentially, with alcohol use (3). Furthermore, alcohol use can destabilize the course of an underlying mental illness. For example, alcohol use in bipolar patients leads to more mood swings (46), more rapid cycling (46–48), and a higher suicidal risk (47–50). In depression, alcohol use is associated with a higher level of depressive

symptoms (51) and a more altered quality of life (52). Moreover, we found that 80% of heavy frequent drinker admitted to the ED with a mental disorder had daily or almost daily HED and all of them were positive on RAPS4-QF. Thus, in addition to consequences related to alcohol frequency and amount of consumption, these patients have HED-related risks, such as an increased risk of coronary heart diseases (53), high blood pressure (53), sexually transmitted diseases (54, 55), and injuries (56). Furthermore, the RAPS4-QF is an instrument with 90% sensitivity and 84% specificity for screening for DSM-5 AUD. This suggest that these patients not only have an unhealthy alcohol use but also have an AUD. This is significantly higher than the rest of the sample which was 30% RAPS4-QF positive, a percentage relatively similar to that found elsewhere. (57).

Our study has some limitations. First, we used self-reported drinking to measure alcohol use, which is a common method with an average validity of 92% (58), but which could be less accurate for frequent heavy drinking individuals (59). However, in analysis here, patients who drank more than four drinks in a typical day were grouped into one category, and while it is possible that those who drink the most may report with less accuracy the actual number of drinks they consume, it would be surprising if they reported less than four drinks. Secondly, we used broad categories of reasons for admission, which can lead to a loss of information. For example, it is likely that if the gastro-intestinal category has been further delineated, some reasons for admission would have been more strongly related to unhealthy alcohol use (liver diseases). However, the aim of our study was not to describe which pathology is associated with alcohol use, because the association of alcohol with many diseases is already known (3). Our aim was to propose “red flags” for ED's staff to help them to quickly screen for unhealthy alcohol use based admission complaint. Third, all the patients with a reason for admission related to the management of an alcohol intoxication or an alcohol withdrawal syndrome were classified in the mental disorder category. This logically raises the question of whether the high prevalence of daily or almost daily HED in this category was not primarily due to these patients, and whether it is necessary to be so concerned about alcohol use for other mental disorders. We do not have the necessary data to answer this question, but it should be noted that 45% of the patients whose reason for admission was a mental disorder had an unhealthy alcohol use, which represents considerably more patients than those admitted to the ED for a withdrawal syndrome or alcohol intoxication. Furthermore, even if the heaviest drinkers were mainly patients who came for withdrawal syndrome or alcohol intoxication, the message from our data would also be interesting. Indeed, it would mean that, in ED, an acute intoxication and a withdrawal syndrome should never be considered as an acute phenomenon, but as the emerging part of a deeper problem since many of these patients have daily or almost daily HED. Fourth, we recruited patients during one month in each ED and, although each month was randomly chosen, the study took place mainly during summer, and it is possible that there may be seasonal variation in findings here. Finally, we included only patients able to answer questionnaires, it therefore did not represent all the ED's patients.

To conclude, this study highlights that unhealthy alcohol use is frequent among ED patients and particularly among those who are

admitted for injuries or mental disorders. Men under age 65 who present with a mental disorder require special attention because of their increased likelihood of reporting daily or almost daily HED.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary materials](#), further inquiries can be directed to the corresponding author/s.

Ethics statement

The studies involving humans were approved by Comité d’Ethique des Centres d’Investigation Clinique de l’inter-région Rhône-Alpes-Auvergne. The studies were conducted in accordance with the local legislation and institutional requirements. Written informed consent for participation in this study was provided by the participants’ legal guardians/next of kin.

Author contributions

JoC: Writing – original draft, Conceptualization, Data curation, Methodology, Software, Writing – review & editing. CL: Software, Validation, Writing – review & editing. JuC: Conceptualization, Methodology, Writing – review & editing. CC: Conceptualization, Writing – review & editing. BR: Conceptualization, Data curation, Writing – review & editing. FM: Conceptualization, Investigation, Methodology, Writing – review & editing. PL: Conceptualization, Investigation, Methodology, Writing – review & editing. DR: Conceptualization, Investigation, Methodology, Writing – review & editing. JG: Conceptualization, Funding acquisition, Investigation, Methodology, Writing – review & editing. EP: Conceptualization, Investigation, Methodology, Writing – review & editing. MD: Conceptualization, Investigation, Writing – review & editing. MN: Conceptualization, Investigation, Writing – review & editing. MC: Conceptualization, Data curation, Funding acquisition, Investigation, Methodology, Writing – review & editing. P-ML: Conceptualization, Funding acquisition, Investigation, Methodology, Writing – review & editing. BP: Software, Writing – review & editing. IC: Conceptualization, Data curation, Funding acquisition, Investigation, Methodology, Writing – review & editing. GB: Conceptualization, Data curation, Funding acquisition, Investigation, Methodology, Supervision, Writing – review & editing.

References

- Paille F, Reynaud M. L'alcool, une des toutes premières causes d'hospitalisation en France. *Bull Épidémiologique Hebd.* (2015) 24-25:440–9.
- Bonaldi C, Hill C. La mortalité attribuable à l'alcool en France en 2015. *Bull Épidémiologique Hebd.* (2019) 5-6:97–08.
- Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos CT. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addict Abingdon Engl.* (2003) 98:1209–28. doi: 10.1046/j.1360-0443.2003.00467.x
- U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Dietary guidelines for Americans, 2020–2025. 9th ed.* Des Moines: U.S. Department of Agriculture and U.S. Department of Health and Human Services (2020).
- Paradis C, Butt P, Shield K, Poole N, Wells S, Naimi T, et al. Canada's guidance on alcohol and health: final report. *Final Rep.* (2023)
- Bush K, Kivlahan DR, McDonnell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory care quality improvement project (ACQUIP). Alcohol use disorders identification test. *Arch Intern Med.* (1998) 158:1789–95. doi: 10.1001/archinte.158.16.1789
- Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol Clin Exp Res.* (2007) 31:1208–17. doi: 10.1111/j.1530-0277.2007.00403.x

Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. The study was funded by an inter-regional public grant (PHRC interregional Auvergne - Rhône-Alpes 2012). The funder had no role in the writing of the study protocol or in the conductance of the study.

Acknowledgments

The paper is based, in part, on data collected by the following collaborators and their staffs: A. Viallon, X. Jacob, F. Carpentier, A. Genty.

Conflict of interest

P-ML participated in advisory boards for Eisai, Ethypharm, Janssen, Lundbeck, MSD, Neuraxpharm, Novartis, Otsuka, Roche, and Rovi; received speaker's honoraria and consultation fees from Eisai, Ethypharm, Janssen, Lundbeck, MSD, Neuraxpharm, Novartis, Otsuka, Roche, and Rovi. BR received lecture or expert fees from Ethypharm, Lundbeck, Indivior, Bouchara-Recordati, Gilead, AstraZeneca, Bristol-Myers-Squibb, Otsuka, and Servier. The work provided by BR on this study was supported by a research grant from the Fondation Actions-Addictions.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2023.1271076/full#supplementary-material>

8. World Health Organization. International guide for monitoring alcohol consumption and related harm [internet]. World Health Organization (2018). [cité 2 mars 2023]. Report No.: WHO/MSD/MSB/00.4. Disponible sur
9. Cherpitel CJ, Ye Y. Performance of the RAPS4/RAPS4-QF for DSM-5 compared to DSM-IV alcohol use disorders in the general population: data from the 2000-2010 National Alcohol Surveys. *Drug Alcohol Depend.* (2015) 151:258–61. doi: 10.1016/j.drugalcdep.2015.02.025
10. Cherpitel CJ, Bazargan S. Screening for alcohol problems: comparison of the audit, RAPS4 and RAPS4-QF among African American and Hispanic patients in an inner city emergency department. *Drug Alcohol Depend.* (2003) 71:275–80. doi: 10.1016/S0376-8716(03)00140-6
11. Cherpitel CJ, Ye Y, Moskalewicz J, Swiatkiewicz G. Screening for alcohol problems in two emergency service samples in Poland: comparison of the RAPS4. *CAGE AUDIT Drug Alcohol Depend.* (2005) 80:201–7. doi: 10.1016/j.drugalcdep.2005.03.025
12. van Loon M, van Gaalen ACP, van der Linden MC, Hagestein-De BC. Evaluation of screening and brief intervention for hazardous alcohol use integrated into clinical practice in an inner-city emergency department. *Eur J Emerg Med Off J Eur Soc Emerg Med.* (2017) 24:224–9. doi: 10.1097/MEJ.0000000000000320
13. Egerton-Warburton D, Gosbell A, Moore K, Wadsworth A, Richardson D, Fatovich DM. Alcohol-related harm in emergency departments: a prospective, multi-Centre study. *Addict Abingdon Engl.* (2018) 113:623–32. doi: 10.1111/add.14109
14. Cherpitel CJ. Drinking patterns and problems: a comparison of primary care with the emergency room. *Subst Abuse.* (1999) 20:85–95. doi: 10.1080/08897079909511397
15. D'Onofrio G, Bernstein E, Bernstein J, Woolard RH, Brewer PA, Craig SA, et al. Patients with alcohol problems in the emergency department, part 1: improving detection. SAEM substance abuse task force. Society for Academic Emergency Medicine. *Acad Emerg med off J Soc. Acad Emerg Med.* (1998) 5:1200–9. doi: 10.1111/j.1553-2712.1998.tb02696.x
16. Smith SG, Touquet R, Wright S, Das GN. Detection of alcohol misusing patients in accident and emergency departments: the Paddington alcohol test (PAT). *J Accid Emerg Med.* (1996) 13:308–12. doi: 10.1136/emj.13.5.308
17. Touquet R, Brown A. PAT (2009)--revisions to the Paddington alcohol test for early identification of alcohol misuse and brief advice to reduce emergency department re-attendance. *Alcohol Alcohol Oxf Oxf.* (2009) 44:284–6. doi: 10.1093/alcalc/aggp016
18. Patton R, O'Hara P. Alcohol: signs of improvement. The 2nd national emergency department survey of alcohol identification and intervention activity. *Emerg Med J EMJ.* (2013) 30:492–5. doi: 10.1136/emered-2012-201527
19. Pöyry T, Luoto TM, Kataja A, Brander A, Tenovu O, Iverson GL, et al. Acute assessment of brain injuries in ground-level falls. *J Head Trauma Rehabil.* (2013) 28:89–97. doi: 10.1097/HTR.0b013e318250eadd
20. Shakya I, Bergen G, Haddad YK, Kakara R, Moreland BL. Fall-related emergency department visits involving alcohol among older adults. *J Saf Res.* (2020) 74:125–31. doi: 10.1016/j.jsr.2020.06.001
21. Woods C, Jones R, Usher K. The impact of unintentional alcohol-related falls on emergency departments. *Australas Emerg Care.* (2019) 22:22–7. doi: 10.1016/j.auec.2018.11.001
22. Ahmed N, Greenberg P. Assessing the impact of blood alcohol concentration on the rate of in-hospital mortality following traumatic motor vehicle crash injury: a matched analysis of the National Trauma Data Bank. *Injury.* (2019) 50:33–8. doi: 10.1016/j.injury.2018.09.022
23. Leeper S, Lahri S, Myers J, Patel M, Reddy P, Martin IBK, et al. Assault-injured youth in the emergency centres of Khayelitsha, South Africa: baseline characteristics & opportunities for intervention. *Injury.* (2019) 50:2220–7. doi: 10.1016/j.injury.2019.10.014
24. De Chazeron I, Guelon T, Chalmeton M, Carpentier F, Moustafa F, Viallon A, et al. Alcohol consumption for people admitted in French emergency departments: a protocol for a multi-center cluster trial. *J Addict Res Ther.* (2015) 6:250.
25. Pasche S, Broers B, Favrod-Coune T. Comment y voir clair face à toutes les recommandations relatives à la consommation d'alcool? *Rev Med Suisse.* (2012) 355:1831–5.
26. Hagman BT, Falk D, Litten R, Koob GF. Defining recovery from alcohol use disorder: development of an NIAAA research definition. *Am J Psychiatry.* (2022) 179:807–13. doi: 10.1176/appi.ajp.21090963
27. Rhodes KV, Rodgers M, Sommers M, Hanlon A, Chittams J, Doyle A, et al. Brief motivational intervention for intimate partner violence and heavy drinking in the emergency department: a randomized clinical trial. *JAMA.* (2015) 314:466–77. doi: 10.1001/jama.2015.8369
28. McElduff P, Dobson AJ. How much alcohol and how often? Population based case-control study of alcohol consumption and risk of a major coronary event. *BMJ.* (1997) 314:1159–64. doi: 10.1136/bmj.314.7088.1159
29. Santé publique France. *Institut national du cancer. Avis d'experts relatif à l'évolution du discours public en matière de consommation d'alcool en France.* Saint-Maurice: Santé publique France (2017).
30. Feise RJ. Do multiple outcome measures require p-value adjustment? *BMC Med Res Methodol.* (2002) 2:8. doi: 10.1186/1471-2288-2-8
31. Hothorn T, Hornik K, Zeileis A. Unbiased recursive partitioning: a conditional inference framework. *J Comput Graph Stat.* (2006) 15:651–74. doi: 10.1198/106186006X133933
32. Cherpitel CJ, Ye Y, Bond J, Rehm J, Poznyak V, Macdonald S, et al. Multi-level analysis of alcohol-related injury among emergency department patients: a cross-national study. *Addict Abingdon Engl.* (2005) 100:1840–50. doi: 10.1111/j.1360-0443.2005.01257.x
33. Friedman J, Hendrix V, Fustok J, Reza T, Madda P, Smith A, et al. Correlation of ride sharing service availability and decreased alcohol-related motor vehicle collision incidence and fatality. *J Trauma Acute Care Surg.* (2020) 89:441–7. doi: 10.1097/TA.0000000000002802
34. Nweze IC, DiGiacomo JC, Shin SS, Gupta C, Ramakrishnan R, Angus LDG. Demographic and socioeconomic factors influencing disparities in prevalence of alcohol-related injury among underserved trauma patients in a safety-net hospital. *Injury.* (2016) 47:2635–41. doi: 10.1016/j.injury.2016.10.020
35. Cherpitel CJ, Ye Y, Bond J, Borges G, Monteiro M, Chou P, et al. Alcohol attributable fraction for injury morbidity from the dose-response relationship of acute alcohol consumption: emergency department data from 18 countries. *Addict Abingdon Engl.* (2015) 110:1724–32. doi: 10.1111/add.13031
36. Behl T, Yadav HN, Sharma PL. Alcoholic neuropathy: involvement of multifaceted signalling mechanisms. *Curr Mol Pharmacol.* (2021) 14:2–10. doi: 10.2174/1874467213666200512114943
37. Piguot O, Cramsie J, Bennett HP, Kril JJ, Lye TC, Corbett AJ, et al. Contributions of age and alcohol consumption to cerebellar integrity, gait and cognition in non-demented very old individuals. *Eur Arch Psychiatry Clin Neurosci.* (2006) 256:504–11. doi: 10.1007/s00406-006-0671-5
38. Mukamal KJ. Alcohol consumption and abnormalities of brain structure and vasculature. *Am J Geriatr Cardiol.* (2004) 13:22–8. doi: 10.1111/j.1076-7460.2004.01334.x
39. Browne AL, Newton M, Gope M, Schug SA, Wood F, Allsop S. Screening for harmful alcohol use in Australian trauma settings. *Injury.* (2013) 44:110–7. doi: 10.1016/j.injury.2012.01.008
40. Barratt H, Rojas-García A, Clarke K, Moore A, Whittington C, Stockton S, et al. Epidemiology of mental health attendances at emergency departments: systematic review and Meta-analysis. *PLoS One.* (2016) 11:e0154449. doi: 10.1371/journal.pone.0154449
41. Kessler RC, Crum RM, Warner LA, Nelson CB, Schulenberg J, Anthony JC. Lifetime co-occurrence of DSM-III-R alcohol abuse and dependence with other psychiatric disorders in the National Comorbidity Survey. *Arch Gen Psychiatry.* (1997) 54:313–21. doi: 10.1001/archpsyc.1997.01830160031005
42. Mueser KT, Yarnold PR, Levinson DF, Singh H, Bellack AS, Kee K, et al. Prevalence of substance abuse in schizophrenia: demographic and clinical correlates. *Schizophr Bull.* (1990) 16:31–56. doi: 10.1093/schbul/16.1.31
43. Di Florio A, Craddock N, van den Bree M. Alcohol misuse in bipolar disorder. A systematic review and meta-analysis of comorbidity rates. *Eur Psychiatry J Assoc Eur Psychiatr.* (2014) 29:117–24. doi: 10.1016/j.eurpsy.2013.07.004
44. Puddephatt JA, Irizar P, Jones A, Gage SH, Goodwin L. Associations of common mental disorder with alcohol use in the adult general population: a systematic review and meta-analysis. *Addict Abingdon Engl.* (2022) 117:1543–72. doi: 10.1111/add.15735
45. Barry KL, Milner K, Blow FC, Impens A, Welsh D, Amash J. Screening psychiatric emergency department patients with major mental illnesses for at-risk drinking. *Psychiatr Serv Wash DC.* (2006) 57:1039–42. doi: 10.1176/ps.2006.57.7.1039
46. Rakofsky JJ, Dunlop BW. Do alcohol use disorders destabilize the course of bipolar disorder? *J Affect Disord.* (2013) 145:1–10. doi: 10.1016/j.jad.2012.06.012
47. Nery FG, Miranda-Scippa A, Nery-Fernandes F, Kapczynski F, Lafer B. Prevalence and clinical correlates of alcohol use disorders among bipolar disorder patients: results from the Brazilian bipolar research network. *Compr Psychiatry.* (2014) 55:1116–21. doi: 10.1016/j.comppsy.2014.02.006
48. Gordon-Smith K, Lewis KJS, Vallejo Añón FM, di Florio A, Perry A, Craddock N, et al. Patterns and clinical correlates of lifetime alcohol consumption in women and men with bipolar disorder: findings from the UK bipolar disorder research network. *Bipolar Disord.* (2020) 22:731–8. doi: 10.1111/bdi.12905
49. Ickick R, Gard S, M'Bailara K, Biseul I, Samalin L, Brousse G, et al. The course of bipolar disorder as a function of the presence and sequence of onset of comorbid alcohol use disorders in outpatients attending the Fondamental advanced Centres of expertise. *J Affect Disord.* (2021) 287:196–03. doi: 10.1016/j.jad.2021.03.041
50. Kyu SY, La Flair LN, Mojtabei R, Lee LC, Spivak S, Crum RM. The Association of Alcohol use Disorders with suicidal ideation and suicide attempts in a population-based sample with mood symptoms. *Arch Suicide Res Off J Int Acad Suicide Res.* (2016) 20:219–32. doi: 10.1080/13811118.2015.1004489
51. Johannessen EL, Andersson HW, Bjørngaard JH, Pape K. Anxiety and depression symptoms and alcohol use among adolescents - a cross sectional study of Norwegian secondary school students. *BMC Public Health.* (2017) 17:494. doi: 10.1186/s12889-017-4389-2
52. Levola J, Pitkänen T, Kampman O, Aalto M. The association of alcohol use and quality of life in depressed and non-depressed individuals: a cross-sectional general

population study. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil.* (2018) 27:1217–26. doi: 10.1007/s11136-017-1741-z

53. Murray RP, Connett JE, Tyas SL, Bond R, Ekuma O, Silversides CK, et al. Alcohol volume, drinking pattern, and cardiovascular disease morbidity and mortality: is there a U-shaped function? *Am J Epidemiol.* (2002) 155:242–8. doi: 10.1093/aje/155.3.242

54. Trillo AD, Merchant RC, Baird JR, Ladd GT, Liu T, Nirenberg TD. Interrelationship of alcohol misuse, HIV sexual risk and HIV screening uptake among emergency department patients. *BMC Emerg Med.* (2013) 13:9. doi: 10.1186/1471-227X-13-9

55. Wray TB, Celio MA, Kahler CW, Barnett NP, Mastroleo NR, Operario D, et al. Daily co-occurrence of alcohol use and high-risk sexual behavior among heterosexual, heavy drinking emergency department patients. *Drug Alcohol Depend.* (2015) 152:109–15. doi: 10.1016/j.drugalcdep.2015.04.011

56. Cherpitel CJ, Bond J, Ye Y, Borges G, MacDonald S, Stockwell T, et al. Alcohol-related injury in the ER: a cross-national meta-analysis from the emergency Room collaborative alcohol analysis project (ERCAAP). *J Stud Alcohol.* (2003) 64:641–9. doi: 10.15288/jsa.2003.64.641

57. Bazargan-Hejazi S, Bazargan M, Gaines T, Jemanez M. Alcohol misuse and report of recent depressive symptoms among ED patients. *Am J Emerg Med.* (2008) 26:537–44. doi: 10.1016/j.ajem.2007.08.019

58. Cherpitel CJ, Ye Y, Monteiro MG. Dose-response relative risk of injury from acute alcohol consumption in 22 countries: are women at higher risk than men? *Alcohol Alcohol Oxf Oxf.* (2019) 54:396–01. doi: 10.1093/alcal/agz018

59. Greenfield TK, Nayak MB, Bond J, Ye Y, Midanik LT. Maximum quantity consumed and alcohol-related problems: assessing the most alcohol drunk with two measures. *Alcohol Clin Exp Res.* (2006) 30:1576–82. doi: 10.1111/j.1530-0277.2006.00189.x